

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Long-term effect of asthma on the development of obesity among adults: an international cohort study, ECRHS

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1856044> since 2023-08-25T14:45:14Z

Published version:

DOI:10.1136/thoraxjnl-2021-217867

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Long-term effect of asthma on the development of obesity among adults: an international cohort study, ECRHS

Subhabrata Moitra^{1,2,3,4,*}, Anne-Elie Carsin^{2,3,4}, Michael J Abramson⁵, Simone Accordini⁶, Andre F S Amaral⁷, Josep Anto^{2,3,4}, Roberto Bono⁸, Lidia Casas Ruiz^{9,10}, Isa Cerveri¹¹, Leda Chatzi^{12,13,14}, Pascal Demoly^{15,16}, Sandra Dorado-Arenas¹⁷, Bertil Forsberg¹⁸, Frank Gilliland¹³ Thorarinn Gislason^{19,20}, Jose A Gullón²¹, Joachim Heinrich^{22,23}, Mathias Holm²⁴, Christer Janson²⁵, Rain Jogi²⁶, Francisco Gómez Real^{27,28}, Debbie Jarvis²⁹, Bénédicte Leynaert^{30,31}, Dennis Nowak²², Nicole Probst-Hensch^{32,33}, José Luis Sánchez-Ramos³⁴, Chantal Raheison-Semjen³⁵, Valerie Siroux³⁶, Stefano Guerra³⁷, Manolis Kogevinas^{2,3,4}, Judith Garcia-Aymerich^{2,3,4}.

¹ Division of Pulmonary Medicine, Department of Medicine, Faculty of Medicine & Dentistry, University of Alberta, Edmonton, Alberta, Canada

² Non-Communicable Diseases and Environment Programme, ISGlobal, Barcelona, Spain

³ Universitat Pompeu Fabra (UPF), Barcelona, Spain

⁴ CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain

⁵ School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria, Australia

⁶ Unit of Epidemiology and Medical Statistics, Department of Public Health and Community Medicine, University of Verona, Verona, Italy

⁷ National Heart and Lung Institute, Imperial College London, London, UK

⁸ Department of Public Health and Pediatrics, University of Turin, Torino, Italy

⁹ Epidemiology and Social Medicine, University of Antwerp, Antwerpen, Belgium

¹⁰ Centre for Environment and Health, Department of Public Health and Primary Care, KU Leuven, Leuven, Belgium

¹¹ Unit of Respiratory Diseases, IRCCS Policlinico San Matteo, University of Pavia, Pavia, Italy

¹² Department of Social Medicine, University of Crete, Rethimno, Greece

¹³ Department of Preventive Medicine, USC Keck School of Medicine, Los Angeles, California, USA

¹⁴ Department of Genetics & Cell Biology, Maastricht University, Maastricht, The Netherlands

¹⁵ Department of Pulmonology, Division of Allergy, University Hospital of Montpellier, Montpellier, France

¹⁶ Inserm, Sorbonne Université, Equipe, EPAR - IPLESP, Paris, France

¹⁷ Osakidetza Basque Health Service, Department of Respiratory Medicine, Galdakao University Hospital, Galdakao, Spain

¹⁸ Section of Sustainable Health, Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden

¹⁹ Department of Sleep, Landspítali - The National University Hospital of Iceland, Reykjavik, Iceland

²⁰ Faculty of Medicine, University of Iceland, Reykjavik, Iceland

²¹ Department of Pneumology, University Hospital San Agustín, Avilés, Spain

²² Institute and Outpatient Clinic for Occupational, Social and Environmental Medicine, Ludwig Maximilians University Munich, University Hospital Munich, Munich, Germany

²³ Institut of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Oberschleissheim, Germany

²⁴ Department of Public Health and Community Medicine, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden

²⁵ Department of Medical Sciences, Respiratory, Allergy and Sleep Research, Uppsala University, Uppsala, Sweden

²⁶ Lung Clinic, Tartu University Hospital, Tartu, Estonia

- ²⁷ Department of Gynaecology and Obstetrics, Haukeland University Hospital, Bergen, Norway
- ²⁸ Department of Clinical Science, University of Bergen, Bergen, Norway
- ²⁹ MRC Centre for Environment and Health, Imperial College London, London, UK
- ³⁰ Inserm - U1168, VIMA (Aging and Chronic Diseases. Epidemiological and Public Health Approaches), INSERM, Villejuif, France
- ³¹ UMR-S 1168, Université de Versailles Saint-Quentin-en-Yvelines - UVSQ, Saint-Quentin-en-Yvelines, France
- ³² Department Epidemiology and Public Health, Swiss Tropical and Public Health Institute, Basel, Switzerland
- ³³ Department of Clinical Research, University of Basel, Basel, Switzerland
- ³⁴ Department of Nursing, University of Huelva, Huelva, Spain
- ³⁵ INSERM U1219, Institute of Public Health and Epidemiology, CHU de Bordeaux, Bordeaux, France
- ³⁶ Team of Environmental Epidemiology, Inserm U1209, Univ Grenoble Alpes, La Tronche, France
- ³⁷ Arizona Respiratory Center, University of Arizona Medical Center - University Campus, Tucson, Arizona, USA

Correspondence to Dr. Subhabrata Moitra, Division of Pulmonary Medicine, Department of Medicine, Faculty of Medicine & Dentistry, University of Alberta, Edmonton, AB T6G 2R3, Canada; moitra@ualberta.ca

SM and A-EC contributed equally.

Abstract

Introduction

Obesity is a known risk factor for asthma. Although some evidence showed asthma causing obesity in children, the link between asthma and obesity has not been investigated in adults.

Methods

We used data from the European Community Respiratory Health Survey (ECRHS), a cohort study in 11 European countries and Australia in 3 waves between 1990 and 2014, at intervals of approximately 10 years. We considered two study periods: from ECRHS I (t) to ECRHS II ($t+1$), and from ECRHS II (t) to ECRHS III ($t+1$). We excluded obese (body mass index ≥ 30 kg/m²) individuals at visit t . The relative risk (RR) of obesity at $t+1$ associated with asthma at t was estimated by multivariable modified Poisson regression (lag) with repeated measurements. Additionally, we examined the association of atopy and asthma medication on the development of obesity.

Results

We included 7576 participants in the period ECRHS I-II (51.5% female, mean (SD) age of 34 (7) years) and 4976 in ECRHS II-III (51.3% female, 42 (8) years). 9% of participants became obese in ECRHS I-II and 15% in ECRHS II—III. The risk of developing obesity was higher among asthmatics than non-asthmatics (RR 1.22, 95% CI 1.07 to 1.38), and particularly higher among non-atopic than atopic (1.47; 1.17 to 1.86 vs 1.04; 0.86 to 1.27), those with longer disease duration (1.32; 1.10 to 1.59 in >20 years vs 1.12; 0.87 to 1.43 in ≤ 20 years) and those on oral corticosteroids (1.99; 1.26 to 3.15 vs 1.15; 1.03 to 1.28). Physical activity was not a mediator of this association.

Conclusion

This is the first study showing that adult asthmatics have a higher risk of developing obesity than non-asthmatics, particularly those non-atopic, of longer disease duration or on oral corticosteroids.

Key messages

What is already known on this topic

- Although an association between obesity and asthma has been well established, whether asthma can relate to the increased risk of future obesity has only been studied in children.

What this study adds

- Asthmatics, particularly those who were non-atopic, had longer disease duration or were on oral corticosteroids, were at a higher risk of developing obesity later in life. This association was not mediated by physical activity.

How this study might affect research, practice or policy

- Understanding the crosstalk between asthma and obesity would help set a proper strategy for asthma management, particularly with an emphasis on reducing adverse effects of corticosteroids in asthma treatment.

Introduction

Several studies demonstrating the coexistence of obesity and asthma across the world suggest the presence of common aetiological factors between these conditions.^{1–5} Asthma and obesity share some common socioeconomic, behavioural and environmental risk factors, all of which could trigger the expression of genes, leading to the development of either of these diseases. Previous studies focused mainly on the mechanisms by which obesity could lead to asthma. Several obesity-associated biological phenomena have been proposed, such as by altering the lung compliance (causing airflow limitation), increasing the synthesis of immune-modulators, affecting the sympathetic nervous system or by modulating gene function.^{1,2} The inverse relationship, that is, whether asthma is a risk factor for later obesity had not received much attention until recently. In a longitudinal cohort study of kindergarten and first-grade children, Chen *et al.* observed that the non-obese children who were diagnosed with asthma during the recruitment phase were at increased risk of developing obesity during the 10 years follow-up.⁶ This was confirmed in a pooled analysis of 18 European cohorts associating physician diagnosed asthma at 3–4 years with incident obesity at 8 years, the risk being higher among children with active asthma symptoms.⁷ Another recent study demonstrated a link between high peak flow variability in childhood and a steeper increase of body mass index (BMI) in adulthood.⁸ Despite studies on children and young adolescents, whether and to what extent asthma increased the risk for subsequent obesity among adults remained unclear. Asthma in adults may be causally and aetiologically more complex than childhood asthma (which is predominantly allergic asthma), therefore, the magnitude of the asthma-obesity risk is likely to be different in adults.

In line with the previously reported observations, we hypothesised that asthma could be a risk factor for obesity at a later stage of life. Therefore, we aimed to investigate the occurrence of obesity attributable to asthma in participants of the European Community Respiratory Health Survey (ECRHS) over two period of 10 years.

Methods

Design and participants

ECRHS is a population-based cohort study initiated in 1990–1994 (ECRHS-I) with over 18 000 participants aged between 20 and 44 years from 30 centres in 14 countries, which had two follow-up rounds at approximately 10-year intervals (ECRHS-II in 1999–2003 and ECRHS-III in 2010–2014). In the recruitment phase, a postal questionnaire was sent containing items about asthma symptoms and exacerbation history in the last 12 months, current medication and allergic conditions such as nasal symptoms and hay fever. From those who responded to the postal questionnaire, participants were either selected from a random sample or a symptomatic sample and were invited for more detailed investigation that included an interviewer-administered questionnaire and lung function testing.^{9,10} The questionnaires and clinical investigations were repeated in the next follow-ups (ECRHS-II and III).

The design and selection of participants of this study are presented in figure 1. We considered two study periods: from ECRHS I (t) to ECRHS II ($t+1$), and from ECRHS II (t) to ECRHS III ($t+1$). At each period we classified the participants as ‘no asthma’ or ‘current asthma’ (see definitions below). To study only incident obesity, we excluded individuals who were obese (BMI ≥ 30 kg/m²) at visit t . To avoid misclassification in the definition of the ‘current asthma’ and ‘no asthma’ groups, we excluded participants with inactive asthma (ie, self-reported doctor diagnosed asthma, with no respiratory symptoms within the last 12 months). We finally excluded subjects with missing asthma or obesity information at either t or $t+1$, giving a total of 8716 participants from 26 centres of 11 European countries and Australia. Ethical approval of the study was obtained from the ethical review boards or committees of the respective centres. All participants provided signed informed consent. Detailed

information about the ethical approval, informed consent and data management were described elsewhere.¹⁰

Definition of asthma-related variables

Asthma-related information was collected at each survey. We defined current asthma as giving a positive answer to the questions ‘Have you ever had asthma?’ and ‘Was this confirmed by a doctor?’, and having reported at least one of the following in the last 12 months: waking up by an attack of shortness of breath, any asthma attack, or use of asthma medications.^{11–13} Asthma duration at t was derived as the number of years between age at t and reported age at first asthma onset. Atopy was defined as a serum specific IgE antibody concentration of ≥ 0.35 kU_A/L to at least one of four common allergens (house dust mite, cat, timothy grass or *Cladosporium herbarum*), as defined previously.¹⁴ We also obtained information on asthma drug treatment from the questions ‘Have you used any pills, capsules, tablets or medicines, other than inhaled medicines, to help your breathing at any time in the last 12 months?’, and defined whether each patient took inhaled (ICS) oral corticosteroids (OCS) or both during the last 12 months. Only at ECRHS II, information was collected on whether OCS were used continuously, for a short-course, or as rescue medication. The modified Medical Research Council (mMRC) Dyspnoea Scale was used to define dyspnoea (considered if they had mMRC grade ≥ 2).¹⁵

Obesity

Height and weight of all participants were measured by physical examination during the clinical visit at each survey. BMI was calculated as weight in kilograms divided by the square of height in metres. Obesity was defined as BMI ≥ 30 kg/m².¹⁶

Other relevant information

At each study wave, we collected data on sociodemographic and other clinical factors using questionnaires. These included sex, age, education (age at which the participants finished their formal education) and smoking status (never, former and current) and intensity (pack-years). Information about physical activity of the participants was obtained at ECRHS-II and III from a self-completed questionnaire in which they were asked about the weekly frequency and duration of vigorous physical activity.^{17–19} Those who exercised at least two times a week for at least 1 hour were considered physically active.

Statistical analyses

Relative risks (RR) and 95% CIs for the association of current asthma, asthma with atopy, asthma duration, and asthma with medication (ICS, OCS or both), with new obesity onset were estimated using modified Poisson regression for repeated measures with time of follow-up as offset.²⁰ Exposure was the asthma status at the survey time t (lag1) and outcome was obesity at subsequent time $t+1$. Therefore, participants could contribute with one or two observations, that is, from ECRHS-I to ECRHS-II and/or from ECRHS-II to ECRHS-III (figure 1). Multivariate models were fitted after testing the following potential confounders: education, sex, age, smoking status and physical activity (between ECRHS-II and III only). Only those variables associated (ie, $p < 0.05$) both with asthma and obesity were retained. Hence, the final models were adjusted for age, sex and smoking status at time t . Subjects nested in centre were included as random effects to account for the repeated data design.

Models were presented overall and separately by sex. We performed meta-analysis to determine if there was any heterogeneity between the participating countries and between centres.

Two secondary analyses were performed: (1) to determine if physical activity could be a mediator in the asthma-obesity association, we tested three paths separately: physical activity at ECRHS-II mediating the relationship between asthma at ECRHS-I and obesity at ECRHS-II (path 1), physical activity at ECRHS-II mediating the relationship between asthma at ECRHS-I and obesity at ECRHS-II

(path 2), and physical activity at ECRHS-III mediating the relationship between asthma at ECRHS-II and obesity at ECRHS-III (path 3) and (2) to estimate the impact of smoking cessation on the association between asthma and new obesity onset, we stratified our models by active smokers (at time t), smokers who quit during the study (between ECRHS-I and II, and ECRHS-II and III), and never smokers.

Some sensitivity analyses were performed to check the robustness of our results. We repeated our analyses: (1) excluding those who did not take part in all three ECRHS waves; (2) excluding participants who reported avoiding physical activity because of respiratory symptoms; (3) excluding those who reported dyspnoea at time t ; (4) defining current asthma as the a self-report of doctor's diagnosis and at least one of the following within the last 12 months: waking up by an attack of shortness of breath, any asthma attack, use of asthma medications, wheeze, shortness of breath, waking up with chest tightness or attack of cough and (5) using BMI change between t and $t+1$ as the outcome variable in linear mixed regression. All analyses were performed in STATA (V.12.1, StataCorp).

Results

Distribution of the participants included in this analysis is presented in table 1. We studied 7576 participants in the period ECRHS I-II (51.5% female, mean (SD) age of 34 (7) years at ECRHS I) and 4976 participants in the period ECRHS II-III (51.3% female, 42 (8) years at ECRHS II). Current smokers decreased during follow-up (30% in ECRHS I-II vs 19% in ECRHS II-III). The relative number of participants with current asthma remained the same across the study waves. Overall, 9% of participants became obese from ECRHS-I to ECRHS-II and this proportion rose to 15% from ECRHS II to ECRHS-III. We observed that 37% of the study population was physically active at ECRHS-II which increased slightly to 42% at ECRHS-III.

The association between asthma and obesity is presented in online supplemental table E1 and figure 2. Compared with 8.4% and 14.6% of non-asthmatics who developed obesity at ECRHS-II and ECRHS-III, respectively, 10.7% and 16.9% asthmatics developed obesity at ECRHS-II and III. The absolute risk difference in asthmatics vs no asthmatics was 2.3% (95% CI 0.8% to 3.9%) and the RR was 1.21 (95% CI 1.07 to 1.37). After adjusting for potential confounders (age, sex and smoking status at time t), the association between asthma and the development of obesity remained significant (1.22; 95% CI 1.07 to 1.38). The association was stronger among asthmatics without atopy (1.47; 95% CI 1.17 to 1.86), asthmatics with longer disease duration (1.32; 95% CI 1.10 to 1.59), asthmatics who used OCS (1.99; 95% CI 1.26 to 3.15) and those who used both OCS and ICS (2.07; 95% CI 1.19 to 3.59). We conducted post hoc analyses to understand better the association between OCS intake and incident obesity. The comparison between patients on continuous, short course or use as rescue medication of OCS suggested a higher risk of obesity among those using continuously, but very small sample size precluded any kind of statistical testing (online supplemental table E2).

The association of asthma and obesity was similar in males (1.27; 95% CI 1.06 to 1.51) and females (1.20; 95% CI 1.01 to 1.42, online supplemental table E3). The association of asthma with new-onset obesity was homogeneous across the participating countries (figure 3) and centres (online supplemental figure E1). The overall estimates from the meta-analyses were similar to that reported in the main analysis and there was no statistical evidence of heterogeneity ($I^2=0$, p values for heterogeneity were 0.70 and 0.96 for country and centre, respectively).

No evidence of physical activity as a mediator in the asthma-obesity association was observed (online supplemental table E4). When stratified by smoking status, the risk of becoming obese in asthmatics was higher among current smokers (1.46, 95% CI 1.12 to 1.90) and among never smokers (1.27, 95% CI 1.08 to 1.49), while no association between asthma and obesity was observed among those who stopped smoking (0.95; 95% CI 0.65 to 1.37) (online supplemental table E5). In the sensitivity analyses, the estimates were comparable to the main analyses (online supplemental tables E6-E10).

Discussion

In this prospective cohort study, we found that individuals with active asthma were at higher risk of developing obesity compared with individuals without asthma, that this association was higher among non-atopic asthmatics, those with long disease duration or those using OCS, and that the association was not mediated by physical activity levels.

Our observation of accelerated weight gain among asthmatics is supported by a previous prospective study that demonstrated asthma to be associated with increased weight gain at a later stage in women,²¹ although unlike our results, they did not find any such association in men. Two recent longitudinal studies in children, one of them including more than 21 000 children from 16 European cohorts, showed that childhood asthma was a risk factor for obesity at a later age.⁶

⁷ Another recent report also demonstrated that children with high peak flow variability were on a steeper increase of BMI up to young adult life that was not associated with asthma.⁸ This evidence suggests that asthma and obesity share several common physiological pathways and a bidirectional association between asthma and obesity is plausible.

It has been debated whether there is any association between atopy and increased body weight.²² While several reports support this association,^{23 24} other studies have shown that the risk of being overweight is higher among the non-atopic than those with marked allergic sensitisation.²⁵

²⁶ Our observation of a greater risk of obesity among non-atopic asthmatics indicates that allergic sensitisation is unlikely to explain the association between asthma and obesity.

We observed that the risk of developing obesity was more pronounced among asthmatics who were on OCS than those who were not, in agreement with previous reports, in adults and children with asthma, relating the intake of OCS with abnormal weight gain.^{27–30} Unfortunately, we could not test the role of corticosteroids dose or duration due to small sample size of OCS users. The effects of long-term use of OCS on weight gain are well established in asthma and general population,^{27 31} and supported by several mechanisms, including but not limited to increased lipid uptake from the gut and storing in the peripheral tissues,³⁰ or increased insulin resistance.³² However, we also found an increased and statistically significant risk of obesity among asthmatics who reported not having taken OCS. One could argue that, if asthma really causes weight gain, this effect would be higher among those with severe asthma, so the higher risk observed among OCS users is actually the result of confounding by indication and not a result of the treatment itself.³³ This is supported by our findings of higher obesity risk among asthmatics of longer disease duration.

A potential explanation for asthma causing weight gain is through the reduction of physical activity levels. Patients with chronic respiratory diseases tend to be less physically active due to recurrent episodes of breathlessness or persistent wheezing.³⁴ However, our analyses do not support this hypothesis, as physical activity was not mediating the asthma–obesity association and this association remained after excluding those participants who did not perform physical activity due to breathlessness, in line with the previously mentioned study among asthmatic children who developed obesity on follow-up independent of physical activity.⁶

Another potential explanation would be via a role of smoking. Smoking cessation has been observed to augment weight gain at least in the short term.^{35 36} However, we found no association between asthma and incident obesity among participants who had quit smoking. On the contrary, we observed greater RRs among current smokers and non-smokers. Smoking has been observed to reduce the efficacy of steroid treatment in asthma,^{37–39} so it could be argued that asthmatics who smoke are at a higher risk of the effects of asthma (such as obesity) because of worse disease control. All these suggest that there may be mechanisms other than OCS use, reducing physical activity and smoking cessation, linking asthma to obesity, but these are yet unknown.

Our study outcomes are of significant clinical importance. First, our observation of higher obesity risk in asthmatics who took OCS, advocates the minimal and precise use of OCS in asthma.⁴⁰ Although current recommendations of OCS for asthma management, several reports also indicate that oral steroids are still prescribed indiscriminately in asthma, sometimes, even without

diagnosing the severity of the disease properly.⁴¹⁻⁴³ This could result in an increasing burden of adverse health effects in asthma, obesity being one of the major consequences. Second, the observation of a greater risk of obesity among non-atopic asthmatics directs our attention towards adult-onset asthma and supports increasing attention to the reduction of adult risk factors, such as occupational exposures that account for up to 25% of adult-onset asthma.⁴⁴ Third, we also observed a higher risk of obesity among asthmatics who were smokers. Although smoking cessation programmes have now become a part of the clinical management of respiratory and cardiovascular diseases, evidence shows that not all patients benefit from them, particularly in primary care.⁴⁵

⁴⁶ Advice on smoking cessation along with its plausible consequences (such as temporary weight gain) should also be discussed with the patients. Finally, although we did not find any evidence of physical activity mediating the asthma-obesity interaction, it must be noted that lifestyle factors also significantly influence weight gain. Thus, advice on healthy diet, proper physical activity and lifestyle modification should be warranted as part of asthma management.

The strengths of our study included its longitudinal design, allowing the exclusion of obese subjects at baseline, with a large cohort that was followed over a long period of time. Furthermore, we performed several sensitivity analyses to test the robustness of our results against assumptions about asthma or obesity misclassification, residual confounding or model misspecification. Lastly, the results were homogeneous across the participating countries which indicate that study findings can be extrapolated to other populations.

A limitation in this study is the potential bias in defining asthma and obesity status, given the lack of information on when events happened during the 10 years follow-up period. However, our results were robust to several sensitivity analyses. Although we were able to consider plausible known confounders, the effect of other lifestyle factors (such as diet, only available in a subsample of ECRHS at one single time point), psychosocial attributes, or genetic predisposition could not be analysed. Finally, we were limited to explore in detail the role of OCS on obesity risk because of the lack of detailed information on dosage, duration or their use for other diseases than asthma.

Conclusion

In summary, we demonstrated that participants with asthma were at a higher risk of developing obesity later in life and the risk was more pronounced among those asthmatics who were non-atopic, those with asthma of longer duration and those on OCS. These results support earlier results in children and warrant further clinicoepidemiological and experimental research, in order to determine the mechanisms through which asthma could trigger obesity risk among adults.

REFERENCES

- 1 Ford ES, Mannino DM, Redd SC, et al. Body mass index and asthma incidence among USA adults. *Eur Respir J* 2004;24:740–4.
- 2 Ali Z, Ulrik CS. Obesity and asthma: a coincidence or a causal relationship? A systematic review. *Respir Med* 2013;107:1287–300.
- 3 Schatz M, Zeiger RS, Zhang F, et al. Overweight/obesity and risk of seasonal asthma exacerbations. *J Allergy Clin Immunol Pract* 2013;1:618–22.
- 4 Newson RB, Jones M, Forsberg B, et al. The association of asthma, nasal allergies, and positive skin prick tests with obesity, leptin, and adiponectin. *Clin Exp Allergy* 2014;44:250–60.
- 5 Peters U, Dixon AE, Forno E. Obesity and asthma. *J Allergy Clin Immunol* 2018;141:1169–79.
- 6 Chen Z, Salam MT, Alderete TL, et al. Effects of childhood asthma on the development of obesity among school-aged children. *Am J Respir Crit Care Med* 2017;195:1181–8.
- 7 Contreras ZA, Chen Z, Roumeliotaki T, et al. Does early onset asthma increase childhood obesity risk? A pooled analysis of 16 European cohorts. *Eur Respir J* 2018;52:pii: 1800504:1800504.
- 8 Lombardi E, Stern DA, Sherrill D, et al. Peak flow variability in childhood and body mass index in adult life. *J Allergy Clin Immunol* 2019;143:1224–6.
- 9 Burney PG, Luczynska C, Chinn S, et al. The European community respiratory health survey. *Eur Respir J* 1994;7:954–60.
- 10 Jarvis D, Newson R, Janson C, et al. Prevalence of asthma-like symptoms with ageing. *Thorax* 2018;73:37–48.
- 11 de Marco R, Marcon A, Jarvis D, et al. Prognostic factors of asthma severity: a 9-year International prospective cohort study. *J Allergy Clin Immunol* 2006;117:1249–56.
- 12 Cazzoletti L, Marcon A, Corsico A, et al. Asthma severity according to global initiative for asthma and its determinants: an international study. *Int Arch Allergy Immunol* 2010;151:70–9.
- 13 Janson C, Accordini S, Cazzoletti L, et al. Pharmacological treatment of asthma in a cohort of adults during a 20-year period: results from the European community respiratory health survey I, II and III. *ERJ Open Res* 2019;5:00073-2018.
- 14 Le Moual N, Carsin A-E, Siroux V, et al. Occupational exposures and uncontrolled adult-onset asthma in the European community respiratory health survey II. *Eur Respir J* 2014;43:374–86.
- 15 Bestall JC, Paul EA, Garrod R, et al. Usefulness of the medical Research Council (MRC) dyspnoea scale as a measure of disability in patients with chronic obstructive pulmonary disease. *Thorax* 1999;54:581–6.
- 16 Chinn S, Jarvis D, Melotti R, et al. Smoking cessation, lung function, and weight gain: a follow-up study. *Lancet* 2005;365:1629–35.
- 17 Washburn RA, Goldfield SR, Smith KW, et al. The validity of self-reported exercise-induced sweating as a measure of physical activity. *Am J Epidemiol* 1990;132:107–13.
- 18 Rovio S, Kåreholt I, Helkala E-L, et al. Leisure-Time physical activity at midlife and the risk of dementia and Alzheimer’s disease. *Lancet Neurol* 2005;4:705–11.
- 19 Fuertes E, Markevych I, Jarvis D, et al. Residential air pollution does not modify the positive association between physical activity and lung function in current smokers in the ECRHS study. *Environ Int* 2018;120:364–72.
- 20 Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;159:702–6.
- 21 Hasler G, Gergen PJ, Ajdacic V, et al. Asthma and body weight change: a 20-year prospective community study of young adults. *Int J Obes* 2006;30:1111–8.
- 22 Boulet L-P. Obesity and atopy. *Clin Exp Allergy* 2015;45:75–86.
- 23 Chen Y, Rennie D, Cormier Y, et al. Association between obesity and atopy in adults. *Int Arch Allergy Immunol* 2010;153:372–7.
- 24 Luo X, Xiang J, Dong X, et al. Association between obesity and atopic disorders in Chinese adults: an individually matched case-control study. *BMC Public Health* 2013;13:12.

- 25 von Mutius E, Schwartz J, Neas LM, et al. Relation of body mass index to asthma and atopy in children: the National health and nutrition examination study III. *Thorax* 2001;56:835–8.
- 26 Visness CM, London SJ, Daniels JL, et al. Association of childhood obesity with atopic and nonatopic asthma: results from the National health and nutrition examination survey 1999-2006. *J Asthma* 2010;47:822–9.
- 27 Sullivan PW, Ghushchyan VH, Globe G, et al. Oral corticosteroid exposure and adverse effects in asthmatic patients. *J Allergy Clin Immunol* 2018;141:110–6.
- 28 Cooper V, Metcalf L, Versnel J, et al. Patient-Reported side effects, concerns and adherence to corticosteroid treatment for asthma, and comparison with physician estimates of side-effect prevalence: a UK-wide, cross-sectional study. *NPJ Prim Care Respir Med* 2015;25:15026.
- 29 Sweeney J, Patterson CC, Menzies-Gow A, et al. Comorbidity in severe asthma requiring systemic corticosteroid therapy: cross-sectional data from the optimum patient care research database and the British thoracic difficult asthma registry. *Thorax* 2016;71:339–46.
- 30 Umławska W. Adipose tissue content and distribution in children and adolescents with bronchial asthma. *Respir Med* 2015;109:200–7.
- 31 Berthon BS, MacDonald-Wicks LK, Wood LG. A systematic review of the effect of oral glucocorticoids on energy intake, appetite, and body weight in humans. *Nutr Res* 2014;34:179–90.
- 32 Carpaij OA, van den Berge M. The asthma-obesity relationship: underlying mechanisms and treatment implications. *Curr Opin Pulm Med* 2018;24:42–9.
- 33 Garcia-Aymerich J, Marrades RM, Monsó E, et al. Paradoxical results in the study of risk factors of chronic obstructive pulmonary disease (COPD) re-admission. *Respir Med* 2004;98:851–7.
- 34 van 't Hul AJ, Frouws S, van den Akker E, et al. Decreased physical activity in adults with bronchial asthma. *Respir Med* 2016;114:72–7.
- 35 Bush T, Lovejoy JC, Deprey M, et al. The effect of tobacco cessation on weight gain, obesity, and diabetes risk. *Obesity* 2016;24:1834–41.
- 36 Jain P, Danaei G, Robins JM, et al. Smoking cessation and long-term weight gain in the Framingham heart study: an application of the parametric g-formula for a continuous outcome. *Eur J Epidemiol* 2016;31:1223–9.
- 37 Chaudhuri R, Livingston E, McMahon AD, et al. Cigarette smoking impairs the therapeutic response to oral corticosteroids in chronic asthma. *Am J Respir Crit Care Med* 2003;168:1308–11.
- 38 Lazarus SC, Chinchilli VM, Rollings NJ, et al. Smoking affects response to inhaled corticosteroids or leukotriene receptor antagonists in asthma. *Am J Respir Crit Care Med* 2007;175:783–90.
- 39 Kane B, Kolsum U, Southworth T, et al. The effects of smoking on the lipopolysaccharide response and glucocorticoid sensitivity of alveolar macrophages of patients with asthma. *Chest* 2009;136:163–70.
- 40 Bourdin A, Adcock I, Berger P, et al. How can we minimise the use of regular oral corticosteroids in asthma? *Eur Respir Rev* 2020;29:190085.
- 41 Bhulani N, Lalani S, Ahmed A, et al. Knowledge of asthma management by general practitioners in Karachi, Pakistan: comparison with international guidelines. *Prim Care Respir J* 2011;20:448–51.
- 42 Lommatzsch M, Sauerbeck IS, Wilmer C, et al. Oral corticosteroid prescription for asthma by general practitioners: a three-year analysis in Germany. *Respir Med* 2021;176:106242.
- 43 Blakey J, Chung LP, McDonald VM, et al. Oral corticosteroids stewardship for asthma in adults and adolescents: a position paper from the thoracic Society of Australia and New Zealand. *Respirology* 2021;26:1112–30.
- 44 MacKinnon M, To T, Ramsey C, et al. Improving detection of work-related asthma: a review of gaps in awareness, reporting and knowledge translation. *Allergy Asthma Clin Immunol* 2020;16:73.
- 45 Coleman T, Murphy E, Cheater F. Factors influencing discussion of smoking between general practitioners and patients who smoke: a qualitative study. *Br J Gen Pract* 2000;50:207–10.

46 Houston TK, Scarinci IC, Person SD, et al. Patient smoking cessation advice by health care providers: the role of ethnicity, socioeconomic status, and health. *Am J Public Health* 2005;95:1056–61.

Table 1

Demographic profile of the study participants in ECRHS I–II and in ECRHS II–III

	ECRHS I–II n=7576	ECRHS II–III n=4976
Sex (female), n (%)	3905 (51.5)	2555 (51.3)
Age <i>t</i> (years), mean (SD)	34.0 (7.1)	41.5 (7.8)
Age <i>t+1</i> (years), mean (SD)	42.7 (7.2)	53.9 (7.1)
Education, n (%)		
Low	813 (10.8)	409 (8.5)
Medium	2715 (36.0)	1675 (34.7)
High	4024 (53.3)	2744 (56.8)
Country, n (%)		
Australia	387 (5.1)	212 (4.3)
Belgium	528 (7.0)	287 (5.8)
Estonia	250 (3.3)	132 (2.7)
France	862 (11.4)	737 (14.8)
Germany	534 (7.0)	545 (11.0)
Iceland	434 (5.7)	320 (6.4)
Italy	512 (6.8)	210 (4.2)
Norway	500 (6.6)	298 (6.0)
Spain	1353 (17.9)	806 (16.2)
Sweden	1231 (16.2)	749 (15.1)
Switzerland	483 (6.4)	403 (8.1)

	ECRHS I–II n=7576	ECRHS II–III n=4976
UK	502 (6.6)	277 (5.6)
Smoking status <i>t</i> , n (%)		
Never	3272 (43.3)	2201 (44.4)
Former	2043 (27.0)	1814 (36.6)
Current	2247 (29.7)	939 (19.0)
Current asthma <i>t</i> , n (%)		
No asthma	6859 (90.5)	4495 (90.2)
Current asthma, all	717 (9.5)	490 (9.8)
Current asthma, without atopy	271 (3.6)	190 (3.8)
Current asthma, with atopy	446 (5.9)	300 (6.0)
Current asthma, ≤20 years since onset	406 (5.4)	218 (4.4)
Current asthma, >20 years since onset	308 (4.1)	266 (5.3)
Current asthma, no ICS	494 (6.5)	260 (5.2)
Current asthma, with ICS	223 (2.9)	230 (4.6)
Current asthma, no OCS	664 (8.8)	454 (9.1)
Current asthma, with OCS	53 (0.7)	36 (0.7)
Current asthma, only ICS	183 (2.4)	199 (4.0)
Current asthma, ICS and OCS	40 (0.5)	31 (0.6)
BMI <i>t</i> (kg/m ²), mean (SD)	23.3 (2.8)	24.1 (2.9)
BMI <i>t+1</i> (kg/m ²), mean (SD)	25.0 (3.6)	26.0 (3.8)
Obesity (yes) <i>t+1</i> , n (%)	655 (8.6)	736 (14.8)

	ECRHS I–II n=7576	ECRHS II–III n=4976
Dyspnoea (yes) <i>t</i> , n (%)	256 (3.4)	182 (3.7)
Physical activity (yes) <i>t+1</i> , n (%)	2627 (37.0)	2029 (41.8)

Current asthma defined as a self-report of a doctor’s asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of breath, any asthma attack or use of asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from the study (see the Methods section).

BMI, body mass index; ECRHS, European Community Respiratory Health Survey; ICS, inhaled corticosteroids; OCS, oral corticosteroids.

Figure 1

Schematic representation of the study design. ECRHS, European Community Respiratory Health Survey.




Figure 2

Association between asthma, asthma with atopy, asthma duration and asthma with medication at t and incident obesity at $t+1$. Data presented as relative risk (RR) and 95% CI. RR was calculated from modified poisson regression with robust SEs and subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex and smoking status at time t . See full figures in online supplemental table E1. ICS, inhaled corticosteroids; OCS, oral corticosteroids.




Figure 3
 Meta-analysis results of the association between asthma at t and obesity at $t+1$ by country.

