

# Carbon and Sulphur speciation in HP-UHP fluids as obtained by electrolytic-fluid thermodynamic modelling of a UHP marble

ANDREA MAFFEIS<sup>1</sup>, JAMES A. D. CONNOLLY<sup>2</sup>, SIMONA FERRANDO<sup>3</sup>, MARIA LUCE FREZZOTTI<sup>4</sup> AND DANIELE CASTELLI<sup>1</sup>

<sup>1</sup>Università degli Studi di Torino

<sup>2</sup>Institute of Geochemistry and Petrology

<sup>3</sup>University of Torino

<sup>4</sup>Università di Milano Bicocca

Presenting Author: [andrea.maffeis@unito.it](mailto:andrea.maffeis@unito.it)

Element speciation in subduction fluids controls the transport efficiency of elements from the down-going slab to the overlying mantle wedge. Elemental speciation is controlled by  $P_T/O_2$  and bulk-rock composition (i.e., rock-buffered). Elements like C and S are characterised by a wide range of possible valance states, making them crucial in regulating the redox state of the arc mantle and the associated volcanism. However, dominant carbonate sediments (representing ~70% of the total subducted carbon) are only marginally considered in the modelled subduction processes, mostly because of the lack of relevant decarbonation reactions at HP-UHP conditions. Similarly, because S is considered to be primarily subducted at UHP conditions as sulphides or sulphates within the altered oceanic crust, the S input from dominant carbonate sediments has been rarely considered.

To address the role of dominant carbonate sediments on the C and S long term cycles, we conducted electrolytic-fluid thermodynamic modelling of the fluid phase in equilibrium with a UHP impure marble from the Dora-Maira Massif (Western Alps). This marble experienced multiple carbonated dissolution-precipitation events during active subduction at HP-UHP conditions. The study of this natural sample allows linking the thermodynamically modelled HP-UHP evolution, of both rock and fluid, to the HP-UHP mineral assemblages and related fluid inclusions. Using the bulk composition of the studied marble, we modelled the chemical evolution of the fluid along the prograde P-T path (from ~490°C-1.5 GPa to ~730°C-4.3 GPa) and of at different  $fO_2$  (between +2 and -2 from the FMQ buffer). At changing P-T- $fO_2$  conditions, C and S speciation and concentration in the fluid are different. At oxidized conditions, C and S are speciated as  $HCO_3^-$  and  $SO_4^{2-}$ , respectively. At reduced conditions, C is additionally speciated as carboxylic acids and hydrocarbons, while S is speciated as  $H_2S$  and  $HS^-$ . The dissolution of carbonate dominated sediments is an effective process for the mobilisation of both C and S, with C being more easily released at reduced conditions and S at oxidised conditions instead. Thus, dissolution is a more effective process than decarbonation and desulphurisation reactions in releasing C and S during subduction at sub-arc depths.