



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Impact of low-density polyethylene microplastic on ruminal degradability of feeds

This is the author's manuscript	
Original Citation:	
Availability:	
This version is available http://hdl.handle.net/2318/2028900	since 2024-10-30T10:17:12Z
Publisher:	
EAAP, 2024	
Terms of use:	
Open Access	
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.	

(Article begins on next page)

Impact of low-density polyethylene microplastic on ruminal degradability of feeds

S. Glorio Patrucco¹, K. Abid¹, S. Barbera¹, H. Kaihara¹and S. Tassone^{*1}

¹Dept. of Agricultural, Forest and Food Sciences, University of Turin, Italy *sonia.tassone@unito.it

The primary plastic utilized in farms is low-density polyethylene (LDPE), often identified as a foreign object in the rumen of slaughtered cattle and in the form of microplastic (MPs, <5mm) both in ruminant feeds and their body tissues. Despite its widespread presence in farms, there is a notable absence of studies examining its potential impact on the ruminal degradability of feeds.

This research endeavored to examine the influence of LDPE as MPs, on the *in vitro* rumen degradability of feeds.

In 3 farms, 3 most used feeds were collected: mixed hay (Hy), corn silage (Cs), and concentrate (Cn). Feeds, dried at 60°C for 24h and ground (1mm sieve), were prepared in bags containing 0.5g on dry weight of feed, and incubated in a rumen fluid derived from Piemontese bulls. Three runs were performed. Four jars were added with 4 levels of LDPE (0, 0.5, 0.9, and 1.4% on rumen-buffer solution). Each jar contained 3 bags for each feed and each farm, for a total of 27 bags. They were incubated for 48h in the Ankom Daisy^{II} with 400mL of filtered rumen fluid, and 1600mL of buffer solutions for jar.

Results showed that LDPE at all levels did not affect feed degradability. Ruminal degradabilities (%), for 0 to 1.8% of LDPE, were: 47.6±8.92, 47.7±8.96, 49.2±8.72, and 48.5±8.76% in Hy; 59.1±3.97, 59.1±5.13, 59.1±4.86, and 59.7±4.24% in Cs; 69.2±5.80, 70.0±4.38, 71.6±5.28, and 71.6±6.35% in Cn. These results provide valuable insights into the resilience of the rumen degradation process in the face of exposure to LDPE. Future studies are needed to investigate the effects of other MP polymers to provide a deep understanding of the impacts of MPs pollution on ruminant nutrition.