

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Insect meals in a circular economy and applications in monogastric diets

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1942370> since 2023-11-08T16:58:37Z

Published version:

DOI:10.1093/af/vfad016

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Insect meals in a circular economy and applications in monogastric diets

Gasco L.¹, Renna M.², Bellezza Oddon S.¹, Rezaei Far, A.³, Naser El Deen, S.³, Veldkamp T.³

¹Department of Agricultural, Forest and Food Sciences, University of Torino, Largo P. Braccini 2, 10095 Grugliasco, Italy

²Department of Veterinary Sciences, University of Torino, Largo P. Braccini 2, 10095 Grugliasco, Italy

³Wageningen University & Research, Wageningen Livestock Research, De Elst 1, 6700 AH Wageningen, the Netherlands

* Corresponding author: Prof. Manuela Renna, Department of Veterinary Sciences, University of Turin, Largo Paolo Braccini 2, Grugliasco (TO), 10095. Phone: +390116708023. Email: manuela.renna@unito.it

Keywords: alternative proteins, animal health, aquaculture, insect-derived products, livestock feeds

Implications

- In nature, insects are part of the natural diet of aquaculture species, poultry, and pigs.
- Nutritional value of insect meals is comparable or higher than conventional protein sources in animal feed.
- Large quantities and consistent quality and chemical composition of insect meals are required for use in animal feed.
- Insect meals may improve animal health and welfare.
- Insect frass is a good fertiliser with a lower environmental load than artificial fertilisers.

Introduction

Insects are great candidates to support the sustainable development of the feed industry. In the last decade, the global industrialised insect farming has increased, aiming to deliver the market

with large amounts of insect-derived products for feed purposes (van Huis, 2022). The demand for insect protein will rise from 120,000 metric tons to 500,000 metric tons by 2030, according to an estimation by De Jong and Nikolik (2021) in a RABO Bank report. According to the report, the price of a metric ton of insect protein will decrease from EUR 3,500-5,500 during the scale-up phase (2020) to EUR 1,500-2,500 during the maturity phase (2030), making insect protein more competitive with conventional protein sources. During this transition period, the main market for insect protein, or so-called insect meal, is pet food and aquaculture but pets' share of the total market will fall from 54% in 2020 to 30% in 2030, while aquafeed's share will rise from 17 to 40%, resulting in 200,000 metric tons in 2030, which is still only 1% of the total aquafeed market. Currently, research is focusing on additional values beyond the nutritional value in the protein transition concept. Insect-derived specific compounds such as chitin, antimicrobial peptides, and medium-chain fatty acids (mainly lauric acid) have the ability for antibacterial and immunomodulating effects. Research is also focusing to make insects as feed more sustainable in a circular economy model. In EU insects can only be fed with materials of vegetal origin and some materials of animal origin such as milk, eggs and their products, honey, rendered fat or blood products from non-ruminant animals (Reg.(EU) 2022/1104). Circularity and sustainability can be given a boost if slaughterhouse or rendering derived-products, catering waste, and unsold products from supermarkets or food industries containing meat or fish will be allowed to feed the insects, once proven to be safe. These waste sources cannot be fed directly to monogastric animals and this makes insects more essential in the food chain as insects convert these waste sources into highly valuable insect-derived products which can be applied in animal nutrition. This review aims to describe the interest in applying insects in the circular economy, the production process, the nutritional and health properties of insect meals (in particular of the main insect species used for feed purposes) and the main performance results obtained with aquaculture and monogastric livestock species fed insect meals. A short overview of the interest in insect frass is also discussed. Finally, some challenges and main prospects are presented.

Circular economy, production process and nutritional value

The most promising and used insect species for feed production are the black soldier fly (*Hermetia illucens*, HI) (Figure 1), the yellow mealworm (*Tenebrio molitor*, TM) and, to a lesser extent, the common housefly (*Musca domestica*, MD) (van Huis, 2022).

Insert here Figure 1

The sustainability of insects relies on their ability to bio-convert, with low environmental impact, not otherwise valorised low-value waste streams into nutrients. For these reasons, insects in the circular economy match several Sustainable Development Goals (SDGs) (Figure 2). One factor that impairs the insect sustainability is the high energy costs needed to maintain the high temperatures for their growth. However, coupling insect factories with renewable energy sources could solve this issue.

At the end of the rearing process, the insect larvae are sieved from frass, cleaned and, if not used as live larvae mainly for poultry feeding purposes, devitalised by blanching, boiling, drying, cooling, freezing, or freeze-drying (Ravi et al., 2020). The devitalisation techniques affect the derived-product quality in terms of protein solubility and bio-availability, lipid oxidation, product colour and microbial load (Ravi et al., 2020). Insects can then be further processed to obtain two main fractions: insect-derived meals (with high levels of protein) and insect oils. As exhaustively described by Ravi et al. (2020), for HI two processing methods are used. In the dry method, dry larvae are pressed under cold ($T = 25^{\circ}\text{C}$) or hot ($T > 60^{\circ}\text{C}$) conditions, to obtain a defatted meal. The level of fat residue is dependent on the temperature and pressure used.

Insert here Figure 2

In the wet mode, fresh larvae are reduced into pulp, treated with enzymes to hydrolyse proteins and the different fractions are separated using a tricanter. The nutritional value, including amino acids, of full-fat and defatted insect meals for feed purposes largely varies depending on the

species, the life stage, and the rearing substrate. Table 1 reports a small section of published data on the composition of full-fat and defatted HI and TM meals. Other values can be found in different publications. For MD, no data covering defatted meals are present in literature.

Among the insect-derived proteins, defatted meals are of major interest because they are richer in crude protein (CP), lower in ether extract (EE) (Table 1) and more stable and easier to include in feed formulations than the full-fat ones. The nutritional value of the insect meals can vary both due to the rearing substrate and the defatting process, which could also affect the quality of the final products and – in turn – the nutrient digestibility and the animal growth performance. Therefore, particular care should be taken during processing. Finally, the nitrogen-to-protein conversion factor is an important parameter in formulating diets with insect-derived meals. An overestimation of insects' CP can result in unbalanced diets and poor growth performance (Hua, 2021).

Insert here Table 1

The important factors to be taken into account in feed formulation with insect-derived products have recently been reviewed (Gasco et al., 2023). These factors are insect species and composition, processing method, availability and consistency of supply, nutrient digestibility, anti-nutritional factors, physical pellet properties, palatability, stability, safety, costs, impact on product quality, and legislation.

Insect meals in animal feed

Aquaculture

To allow the development of aquaculture production, alternative proteins to fishmeal (FM) have been extensively studied. Nowadays the aquaculture industry uses plant-based ingredients as well as by-products of animal origin (Hua, 2021). Insect-derived products can play a major role in the development of a sustainable aquaculture and trials have been conducted to assess nutrient

digestibility, growth performance, product quality and the impact on fish and shrimp health (Hua 2021; Liland et al., 2021; Prakoso et al., 2022; Tran et al., 2022).

In carnivorous fish species, usually insect meals are included in substitution of FM. Based on recent meta-analyses, it is possible to state that inclusion levels of insect meals up to 25-30% support good growth performances (Hua, 2021; Liland, et al., 2021), but the optimal inclusion level seems to be dependent on the fish and the insect species (Hua, 2021). High HI inclusion levels ($> 30\%$) reduced the growth parameters, while TM meal seems more tolerated (Hua, 2021; Prakoso et al., 2022). Specifically, the use of TM meal in aquafeed (from 2.5 to 65% of inclusion) did not affect the growth performance (Hua, 2021). In salmonids, rainbow trout, sea bass or seabream, the digestibility of CP starts to decrease at an insect meal inclusion level of 25% (Liland et al., 2021) and this was often correlated to the increase of chitin (above 2-3%) that worsens specific growth rate and feed conversion ratio (Tran et al., 2022). Chitin binds to digestive enzymes, which may impair the nutrient absorption in the proximal intestine. In addition, insect chitin is embedded in a matrix of other components including proteins, lipids, and minerals and this could impair the accessibility of digestive enzymes to these nutrients. Despite this, studies on the CP digestibility of insect meals (defatted HI and TM meals) in rainbow trout (Gasco et al., 2022) and sea bass (Basto et al., 2020) reported general high coefficients ranging from 80 to about 93%, also when the chitin content was $\geq 4\%$, supporting the high potential of these products in aquafeed. Authors indicated how differences are not only due to insect species but also to insect meal production techniques. As far as EE digestibility is concerned, values of defatted meals are always above 92%, while lower values are reported for full-fat insect meals.

Research performed using insects in shrimp nutrition is limited and recently reviewed by Sánchez-Muros et al. (2020). No differences in *Litopenaeus vannamei* performances are reported up to 30.5% of full-fat TM meal inclusion fully substituting FM as far as a correct balance of essential amino acids is ensured. In a non-balanced formula, a percentage including up to 7% of defatted HI meal negatively affected the growth of *L. vannamei*. Compared to a dietary treatment containing FM, the inclusion of

10% of a full-fat HI in partial substitution of FM resulted in an improvement of the performances of *L. vannamei* juveniles. However, no differences emerged when full-fat TM was used even if both insect meals treatments reported high digestibility coefficients (ADC) for crude protein (ADCCP: 84 and 85%, respectively) and ether extract (ADCEE: 95 and 97%, respectively) (Shin and Lee, 2021). Richardson et al. (2021) confirmed the positive effect of HI meal (defatted) on the growth performance of *L. vannamei*. The main considerations on the use of insect meals in fish and shrimps are reported in Figure 3.

Insert here Figure 3

Poultry

Refining and innovating along the poultry value chain are crucial to further improve the sustainability of this sector. The relatively high demand for protein-rich feed ingredients in this sector makes particularly interesting to exchange conventional protein sources by insect protein sources. A recent review reported that insect protein derived from HI and MD can substitute conventional protein to a certain extent without adversely affecting production performance (Dörper et al., 2021). In general, it can be concluded that, balancing the essential amino acids profile, insect meals can be included up to 10% in diets for broilers and laying hens without affecting nutrient digestibility, growth performance, and product quality (Dörper et al., 2021). Dietary inclusion levels of insect meal over 10% resulted in inconsistent results among studies (Dörper et al., 2021). The insects can be used live (fresh), dried, and as processed forms in poultry diets. Being part of the natural diet of poultry, live larvae can improve birds' welfare. In broilers, provision of live larvae (Figure 4) at the highest frequency (4 times per day) and 10% of the estimated dietary dry matter (DM) intake determined the most prominent increase in activity and better leg health without affecting broiler performance (Ipema et al., 2020).

Insert here Figure 4

Pigs

In global compound feed production, pig feed took the second place in 2021 (24.1% of share). Regular pig feed is switching to functional and premium variants to improve the immunity of the animals as well as to reduce the risk of metabolic disorders, acidosis, injuries and infections. A shift towards more sustainable feed ingredients is foreseen to improve the sustainability of the entire pig production (Figure 5). Insects are one of the novel protein sources to substitute conventional ones and have been evaluated in a recent review which concluded that the nutrient digestibility of insect proteins was comparable to conventional protein sources (Veldkamp and Vernooij, 2021). However, nutrient digestibility of insect-based diets as well as the related effects on growth performance in pigs fed insect-based diets were different among studies (Veldkamp and Vernooij, 2021).

Insert here Figure 5

The variability in nutrient digestibility of insect-based diets and related effects on growth performance were due to changes in diet ingredients and nutrient composition when insect products were included but also due to different insect species and life stages, processing techniques, palatability, and age of the pigs used in the studies. Globally, when balancing the amino acids profile of the diet, insect products can be included up to 10% to partly replace conventional protein-rich feed ingredients in pig diets without affecting growth performance, product quality and health. Also, live HI larvae can be provided to pigs. In a study, larvae were provided to piglets (25 days old) twice a day (75 g (day 1–4) or 150 g (day 5–11)) and it was concluded that post-weaning live HI larvae provisioning had beneficial effects on piglet behaviour, by facilitating exploration behaviours and reducing the need to orally manipulate objects and pen mates. Neophobic responses towards a novel object were also reduced and performance of piglets that consumed a small amount of larvae was maintained (Ipema et al., 2021). Figure 3 reports the main aspects on the use of insect meals in poultry and pig diets.

Health effects

Insect-derived products seem able to exert positive health effect due to the presence of bioactive compounds. Indeed, as a natural defence mechanism towards the challenging environment they grew, insects develop antimicrobial peptides (AMP) that can stimulate fish, shrimps and livestock immunity. Insects contain chitin which has been mentioned as an anti-nutritional factor adversely affecting the nutrient digestion and absorption at high inclusion levels. On the other hand, chitin can stimulate the innate immune system, modulate the microbiota, and exert antioxidant and anti-inflammatory effects with positive outcomes on animal health (Veldkamp et al., 2022). Next to AMP and chitin, also medium-chain fatty acids (lauric acid) may exert antibacterial effects (Gasco et al., 2021; Veldkamp et al., 2022).

Insect-derived products can therefore sustain animal health and increase their resistance to diseases and may be used to reduce the use of antibiotics. These features can create benefits to insect-derived products and support their use in animal feed as additives (low inclusion levels) (Gasco et al., 2021; Veldkamp et al., 2022) and need to be studied in more detail. As reviewed by Gasco et al. (2021), insect meals seem to modulate and/or promote microbial diversity and – for this reason – could have a positive effect on livestock, fish and shrimp health. A rich gut microbiota increases the competition, in terms of nutrient and colonization site, with pathogens and, consequently, may improve the disease resistance. In general, insect-derived products had not negative effect on the fish microbiota while, in poultry, their positive effect is observed when the inclusion level is lower than or equal to 10%. Finally, based on the scarce literature available to date, insect-derived products seem to improve also the health of pigs and crustaceans. Authors hypothesised antibacterial and probiotic effects of insect meals on piglets and finisher pigs, respectively (Gasco et al., 2021), while HI meal positively influenced antioxidant enzyme activity and non-specific immune responses in shrimp (Shin and Lee, 2021).

Insect frass

Insect frass is what is left at the end of the insect rearing and is defined as “*a mixture of excrements derived from farmed insects, the feeding substrate, parts of farmed insects, dead eggs and with a content of dead farmed insects of not more than 5% in volume and not more than 3% in weight*” (Commission Regulation (EU) 2021/1925). Depending on their initial DM and nutrient composition, the mass substrate reduction performed by HI ranges from 30 to 80%, with 200 to 693 kg (DM) frass production per ton of waste (Lopes et al., 2022). Frass contains good amounts of carbon (C), nitrogen (N), phosphorus (P) and potassium (K) and represents a valuable fertiliser able to substitute mineral sources and introduce organic material into the soil (Schmitt and de Vries, 2020). Differences in frass composition are reported, reflecting the rearing substrate nutritional value and the insect capacity in uploading nutrients. Moreover, the C and N contents of HI frass seem more stable (about 35% and 3%) than P and K values (from 0.3% to 5.2% and from 0.2% to 4.1%) (Lopes et al., 2022; Schmitt and de Vries, 2020). Recent research reported that the improvement in plant performances is not only due to frass nutrient composition, but also to the presence of bioactive compounds (among them, chitin) and microorganisms that seem to improve nutrient utilization from plants, to promote root and plant growth, to stimulate seed germination, and to increase plant drought and stress tolerance (Barragán-Fonseca et al., 2022; Lopes et al., 2022; Schmitt and de Vries, 2020). Compared to inorganic fertilisers, frass has a lower environmental impact in some categories (minus 0.265 g SO₂ equivalents and minus 0.064 g SO₂ equivalents per kg of frass used in substitution of mineral fertilisers for terrestrial acidification and aquatic acidification, respectively) (Schmitt and de Vries, 2020). For these reasons, frass closes the loop of the circular economy applied by insects.

Issues and further directions to consider for the future

- Availability of adequate quantities and composition consistency of supplied insect meals are crucial parameters to allow the feed industry to adopt insect meals in animal formulation.

Indeed, new formulations need not only research, but also marketing actions, and both require investments.

- Due to low produced quantities and high production costs, the price of insect-based proteins is still high and in Europe not competitive when compared to FM or soybean meal. However, waste bioconversion, the decrease in dependency on less sustainable ingredients (FM, soybean meal), or the health benefits associated with the use of insect-based products, in combination with proper marketing of the final product, can justify relatively high prices.
- The quality of insect-derived products highly depends on insect species and composition, development stage at harvest, and method of devitalization and processing. Therefore, the optimal transformation process, in terms of nutrient availability and, as a consequence, of growth performance, for each insect meal should be deeply investigated.
- Safety is the most important parameter for a sector at its infant stage. Insect-derived products should be safe to be used as animal feed ingredient. Growing insects on biowaste sources that cannot be fed directly to aquaculture or livestock is one of the biggest challenges for the future to make insects the missing link in the food chain and this will give sustainability a boost.
- Bioactive compounds contained in insects represent a promising natural alternative to antimicrobial agents and a possible solution to remediate antimicrobial resistance. Moreover, the possibility of stimulating the large-scale production of insect bioactive peptides represents a promising biotech business.
- The balance between antinutritional effects and health-promoting effects of insect-derived products needs to be studied in more detail as also the mode of action to set the most optimal inclusion level of these products in aquaculture and livestock nutrition.
- More research is recommended to set the optimal inclusion level of insect-derived products for growth performance, health, and welfare. Further studies on the mode of action on beneficial health and welfare effects of inclusion of insect-derived products may give an additional value for the application of these products in animal diets.

- Frass includes bioactive compounds and microorganisms which may improve nutrients utilization by plants, promote root and plant growth, stimulate seed germination, increase plant drought and stress tolerance and has a lower environmental load than artificial fertilisers.

References

- Barragán-Fonseca, K. Y., A. Nurfikari, E. M. van de Zande, M. Wantulla, J. J. A. van Loon, W. de Boer, and M. Dicke. 2022. Insect frass and exuviae to promote plant growth and health. *Trends in Plant Science* 27:646-654. doi <https://doi.org/10.1016/j.tplants.2022.01.007>
- Basto, A., E. Matos, and L. M. Valente. 2020. Nutritional value of different insect larvae meals as protein sources for European sea bass (*Dicentrarchus labrax*) juveniles. *Aquaculture* 521:735085. doi <https://doi.org/10.1016/j.aquaculture.2020.735085>
- Benzertiha, A., B. Kierończyk, P. Kołodziejski, E. Pruszyńska–Oszmałek, M. Rawski, D. Józefiak, and A. Józefiak. 2020. *Tenebrio molitor* and *Zophobas morio* full-fat meals as functional feed additives affect broiler chickens' growth performance and immune system traits. *Poultry Science* 99:196-206. doi <https://doi.org/10.3382/ps/pez450>
- De Jong, B., and G. Nikolik. 2021. No longer crawling: Insect protein to come of age in the 2020s. RaboResearch, Rabobank, Utrecht, the Netherlands.
- Dörper, A., T. Veldkamp, and M. Dicke. 2021. Use of black soldier fly and house fly in feed to promote sustainable poultry production. *Journal of Insects as Food and Feed* 7:761-780. doi <https://doi.org/10.3920/jiff2020.0064>
- Gasco, L., S. Bellezza Oddon, G. W. Vandenberg, T. Veldkamp, and I. Biasato. 2023. Factors affecting the decision-making process of using insect-based products in animal feed formulations. *Journal of Insects as Food and Feed* (in press).
- Gasco, L., C. Caimi, A. Trocino, C. Lussiana, S. B. Oddon, V. Malfatto, R. Anedda, G. Serra, I. Biasato, A. Schiavone, F. Gai, and M. Renna. 2022. Digestibility of defatted insect meals for rainbow trout aquafeeds. *Journal of Insects as Food and Feed* 8:1385-1399. doi <https://doi.org/10.3920/jiff2021.0160>
- Gasco, L., A. Józefiak, and M. Henry. 2021. Beyond the protein concept: health aspects of using edible insects on animals. *Journal of Insects as Food and Feed* 7:715-741. doi <https://doi.org/10.3920/jiff2020.0077>
- Hall, H. N., H. V. Masey O'Neill, D. Scholey, E. Burton, M. Dickinson, and E. C. Fitches. 2018. Amino acid digestibility of larval meal (*Musca domestica*) for broiler chickens. *Poultry Science* 97:1290-1297. doi <https://doi.org/10.3382/ps/pex433>
- Heide, M. E. V. d., J. V. Nørgaard, and R. M. Engberg. 2021. Performance, nutrient digestibility and selected gut health parameters of broilers fed with black soldier fly, lesser mealworm and yellow mealworm. *Journal of Insects as Food and Feed* 7:1011-1022. doi <https://doi.org/10.3920/jiff2020.0150>
- Heuel, M., M. Kreuzer, C. Sandrock, F. Leiber, A. Mathys, B. Guggenbühl, I. D. M. Gangnat, and M. Terranova. 2022. Feeding value of black soldier fly larvae compared to soybean in methionine- and lysine-deficient laying hen diets. *Journal of Insects as Food and Feed* 8:989-999. doi <https://doi.org/10.3920/jiff2021.0178>
- Hua, K. 2021. A meta-analysis of the effects of replacing fish meals with insect meals on growth performance of fish. *Aquaculture* 530:735732. doi <https://doi.org/10.1016/j.aquaculture.2020.735732>
- Ipema, A. F., E. A. M. Bokkers, W. J. J. Gerrits, B. Kemp, and J. E. Bolhuis. 2021. Providing live black soldier fly larvae (*Hermetia illucens*) improves welfare while maintaining performance of piglets post-weaning. *Scientific Reports* 11:7371. doi <https://doi.org/10.1038/s41598-021-86765-3>
- Ipema, A. F., W. J. J. Gerrits, E. A. M. Bokkers, B. Kemp, and J. E. Bolhuis. 2020. Provisioning of live black soldier fly larvae (*Hermetia illucens*) benefits broiler activity and leg health in a frequency- and dose-dependent manner. *Applied Animal Behaviour Science* 230:105082. doi <https://doi.org/10.1016/j.applanim.2020.105082>

Leeper, A., D. Benhaïm, B. Ö. Smáráson, S. Knobloch, K. L. Ömarsson, T. Bonnafoux, M. Pipan, W. Koppe, R. Björnsdóttir, and M. Øverland. 2022. Feeding black soldier fly larvae (*Hermetia illucens*) reared on organic rest streams alters gut characteristics of Atlantic salmon (*Salmo salar*). *Journal of Insects as Food and Feed* 8:1355-1372. doi <https://doi.org/10.3920/jiff2021.0105>

Liland, N. S., P. Araujo, X. X. Xu, E.-J. Lock, G. Radhakrishnan, A. J. P. Prabhu, and I. Belghit. 2021. A meta-analysis on the nutritional value of insects in aquafeeds. *Journal of Insects as Food and Feed* 7:743-759. doi <https://doi.org/10.3920/jiff2020.0147>

Lopes, I. G., J. W. H. Yong, and C. Lalander. 2022. Frass derived from black soldier fly larvae treatment of biodegradable wastes. A critical review and future perspectives. *Waste Management* 142:65-76. doi <https://doi.org/10.1016/j.wasman.2022.02.007>

Makkar, H. P. S., G. Tran, V. Heuzé, and P. Ankers. 2014. State-of-the-art on use of insects as animal feed. *Animal Feed Science and Technology* 197:1-33. doi <https://doi.org/10.1016/j.anifeedsci.2014.07.008>

Prakoso, V. A., A. Irawan, A. Iswantari, F. Maulana, R. Samsudin, and A. Jayanegara. 2022. Evaluation of dietary inclusion of black soldier fly (*Hermetia illucens*) larvae on fish production performance: a meta-analysis. *Journal of Insects as Food and Feed* 8:1373-1384. doi <https://doi.org/10.3920/jiff2021.0159>

Ravi, H. K., A. Degrou, J. Costil, C. Trespeuch, F. Chemat, and M. A. Vian. 2020. Larvae Mediated Valorization of Industrial, Agriculture and Food Wastes: Biorefinery Concept through Bioconversion, Processes, Procedures, and Products. *Processes* 8:857. doi <https://doi.org/10.3390/pr8070857>

Richardson, A., J. Dantas-Lima, M. Lefranc, and M. Walraven. 2021. Effect of a Black Soldier Fly Ingredient on the Growth Performance and Disease Resistance of Juvenile Pacific White Shrimp (*Litopenaeus vannamei*). *Animals* 11:1450. doi <https://doi.org/10.3390/ani11051450>

Sánchez-Muros, M. J., P. Renteria, A. Vizcaino, and F. G. Barroso. 2020. Innovative protein sources in shrimp (*Litopenaeus vannamei*) feeding. *Reviews in Aquaculture* 12:186-203. doi <https://doi.org/10.1111/raq.12312>

Schmitt, E., and W. de Vries. 2020. Potential benefits of using *Hermetia illucens* frass as a soil amendment on food production and for environmental impact reduction. *Current Opinion in Green and Sustainable Chemistry* 25:100335. doi <https://doi.org/10.1016/j.cogsc.2020.03.005>

Shin, J., and K.-J. Lee. 2021. Digestibility of insect meals for Pacific white shrimp (*Litopenaeus vannamei*) and their performance for growth, feed utilization and immune responses. *PloS one* 16:e0260305. doi <https://doi.org/10.1371/journal.pone.0260305>

Silva, M. S., R. Matos, P. Araujo, E. J. Lock, R. Gopika, P. A. J. Prabhu, and I. Belghit. 2022. In vitro assessment of protein digestibility and mineral solubility of black soldier fly larvae meals for monogastric animals. *Journal of Insects as Food and Feed* 8:953-966. doi <https://doi.org/10.3920/jiff2021.0197>

Tran, H. Q., T. T. Nguyen, M. Prokešová, T. Gebauer, H. V. Doan, and V. Stejskal. 2022. Systematic review and meta-analysis of production performance of aquaculture species fed dietary insect meals. *Reviews in Aquaculture* 14:1637-1655. doi <https://doi.org/10.1111/raq.12666>

van Huis, A. 2022. Edible insects: Challenges and prospects. *Entomological Research* 52:161-177. doi <https://doi.org/10.1111/1748-5967.12582>

Veldkamp, T., L. Dong, A. Paul, and C. Govers. 2022. Bioactive properties of insect products for monogastric animals – a review. *Journal of Insects as Food and Feed* 8:1027-1040. doi <https://doi.org/10.3920/jiff2021.0031>

Veldkamp, T., and A. G. Vernooij. 2021. Use of insect products in pig diets. *Journal of Insects as Food and Feed* 7:781-793. doi <https://doi.org/10.3920/jiff2020.0091>

Weththasinghe, P., J. Ø. Hansen, M. Rawski, D. Józefiak, S. Ghimire, and M. Øverland. 2021. Insects in Atlantic salmon (*Salmo salar*) diets – comparison between full-fat, defatted, and de-chitinised meals, and oil and exoskeleton fractions. *Journal of Insects as Food and Feed* 8:1235-1247. doi <https://doi.org/10.3920/jiff2021.0094>

Figures captions

Figure 1. Black soldier fly (*Hermetia illucens*) larvae (Photo: Umberto Diecinove 2022)



Figure 2. Circular Economy and Sustainable Development Goals related to insect farming



Table 1. Nutritional value of full-fat and defatted insect meals

Nutrient	<i>Hermetia illucens</i>						<i>Tenebrio molitor</i>						<i>Musca domestica</i>		
	FF ¹	min	max	DF ²	min	max	FF ³	min	max	DF ⁴	min	max	FF ⁵	min	max
Dry matter	93.0	88.1	97.9	93.5	86.8	96.8	95.6	95.6	95.6	95.0	91.8	97.9	92.0	92.0	92.0
Crude protein	43.2	42.7	43.8	50.5	38.5	71.2	47	47	47	69.9	63.0	75.4	53.9	50.4	57.9
Crude fat	31.4	29.2	33.6	15.1	6.8	29.9	29.6	29.6	29.6	8.2	5.7	10.7	20.4	18.9	22.1
Phosphorus				1.1	0.7	1.3	0.7	0.7	0.7				1.6	1.6	1.6
Calcium				1.2	1.0	1.6	0.1	0.1	0.1				0.5	0.5	0.5
Magnesium				0.2	0.1	0.5							0.3	0.3	0.3
Chlorine				0.2	0.1	0.2									
Sodium				0.3	0.2	0.4							0.5	0.5	0.5
Ash	9.5	6.6	12.4	9.5	9.0	10.3	2.6	2.6	2.6	4.9	4.8	5.0	7.9	6.5	10.1
GE (MJ/kg DM)				22.4	19.2	25.4				23.3	23.0	23.5	22.9	22.9	22.9
Amino acids															
Essential amino acids															
Methionine	0.6	0.5	0.8	1.0	0.7	1.3	0.7	0.7	0.7	1.7	0.9	2.6	1.5	1.1	1.7
Threonine	1.6	1.3	2.0	2.2	1.8	2.5	2.5	2.5	2.5	2.9	2.8	3.0	2.9	1.8	3.6
Valine	2.2	1.7	2.8	3.2	2.3	3.9	3.3	3.3	3.3	4.0	3.5	4.6	2.6	2.0	2.9
Isoleucine	1.7	1.4	2.0	2.4	1.9	3.0	2.1	2.1	2.1	2.9	2.4	3.5	2.1	1.6	2.5
Leucine	2.8	2.3	3.3	3.9	3.1	5.3	3.8	3.8	3.8	5.2	4.7	6.0	3.6	2.7	4.2
Phenylalanine	1.8	1.4	2.1	2.3	1.9	3.1	1.9	1.9	1.9	2.4	2.2	2.7	3.4	2.3	4.1
Histidine	1.3	1.1	1.5	1.5	1.3	2.0	1.4	1.4	1.4	2.2	1.9	2.4	1.7	1.2	2.0
Lysine	2.6	2.2	3.1	3.3	2.4	4.1	2.7	2.7	2.7	4.0	3.9	4.0	4.6	3.1	4.9
Arginine	2.1	1.7	2.4	2.6	2.1	3.0	2.5	2.5	2.5	3.7	3.6	3.8	2.9	2.3	3.3
Tryptophan	0.5	0.3	0.7	0.7	0.3	1.0	0.5	0.5	0.5	0.8	0.8	0.8	3.1	0.8	4.5
Non-essential amino acids															
Cysteine	0.3	0.3	0.4	0.4	0.3	0.5	0.6	0.6	0.6	0.5	0.4	0.6	1.3	0.4	1.9
Aspartic acid	3.8	2.8	4.7	5.0	3.9	6.1	3.8	3.8	3.8	5.6	5.3	6.0	5.6	3.8	6.8
Serine	1.7	1.4	2.0	2.3	1.9	3.0	2.3	2.3	2.3	3.4	3.3	3.7	1.7	1.6	1.8
Glutamic acid	5.2	4.5	5.8	6.0	4.9	8.0	6.0	6.0	6.0	8.4	8.2	8.7	7.8	5.9	9.2
Proline	2.3	2.2	2.4	3.3	2.57	5.1	3.1	3.1	3.1	5.2	4.5	6.0	2.2	1.7	2.6
Glycine	2.2	1.7	2.7	3.0	2.31	3.9	2.7	2.7	2.7	3.6	3.4	3.7	2.7	2.1	3.1
Alanine	2.7	2.6	2.9	3.8	3.13	4.9	4.2	4.2	4.2	5.5	4.9	5.9	3.4	2.9	3.8
Tyrosine	3.2	2.9	3.4	3.0	0.00	4.2	3.14	3.1	3.1	3.2	0.0	5.1	3.6	2.4	4.4
Total amino acids	38.6	32.2	45.1	49.2	42.14	55.3	47.2	47.2	47.2	64.47	62.3	68.4	56.2	39.8	67.0

FF: full-fat insect meals; DF: defatted insect meals.

¹(Leeper et al., 2022; Weththasinghe et al., 2021)

²(Gasco et al., 2022; Heide et al., 2021; Heuel et al., 2022; Silva et al., 2022; Weththasinghe et al., 2021)

³(Benzertiha et al., 2020)

⁴(Gasco et al., 2022; Heide et al., 2021)

⁵(Hall et al., 2018; Makkar et al., 2014)

Figure 3. General aspects on the use of insect meals in diets for fish, shrimps, broilers and pigs




Figure 4. Poultry feeding live *Hermetia illucens* larvae



Figure 5. Live *Hermetia illucens* larvae can also be provided to piglets

