

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Interface between polyethylene terephthalate microplastics and microbiota activity in the ruminal environment

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/2028904> since 2024-10-30T10:29:38Z

Publisher:

Polito

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Interface between polyethylene terephthalate microplastics and microbiota activity in the ruminal environment

Khalil Abid; Salvatore Barbera; Sara Glorio Patrucco; Hatsumi Kaihara; Sonia Tassone*

Department of Agricultural, Forest and Food Sciences – University of Turin, Grugliasco, Italy

*Corresponding Author's E-mail address: sonia.tassone@unito.it

Keywords: microplastics; polyethylene terephthalate; ruminal environment; degradability and fermentative activity.

Abstract

Microplastics (MPs) enter the animal digestive system through contaminated feeds [1, 2] creating a unique interface between the rumen and microbiota, resulting in MPs degradation [2-4]. This study investigated the interaction between polyethylene terephthalate (PET) and ruminal activity to understand its influence on feed degradation and fermentation processes.

The experiment was conducted using an *in vitro* gas production (GP) system with lamb rumen fluid and buffer solution to evaluate the effects of different doses of PET contamination (0, 0.6, 1.2, 1.8% dry matter of feed) in 200 mg of concentrate on: pH; GP at different times (at 0, 2, 4, 6, 12, 24, 48, 72 and 96 hours); rumen ammonia-nitrogen (NH₃-N); dry and organic matter degradability (DMD, OMD); rumen protozoa; and microbial efficiency (PF). The experiment was triplicated and repeated over three consecutive weeks.

The presence of PET in concentrate feed within the lamb ruminal environment negatively affected degradability and fermentative processes, particularly at the highest dose. Compared to the control feed without PET addition, PET decreased DMD, OMD, ruminal protozoa, and PF (Table 1), while it increased ruminal GP (Figure 1) and NH₃-N.

The study demonstrated an interaction between PET and ruminal microbiota. These findings highlight the potential adverse impacts of PET on the ruminal fermentation and feed degradability. Further research is needed to explore the effect of PET on gastrointestinal activity and to develop strategies to mitigate the adverse effects.

References

- [1] Ramachandraiah, K. et al. *Science of Total Environment* **2022**, *844*, 157234.
- [2] Glorio Patrucco, S. et al. *International Conference on Micro and Nanoplastics in the Agri-Food Chain* **2023**.
- [3] Quartinello, F. et al. *Frontieres in bioengineering and biotechnology* **2021**, *9*, 684459.
- [4] Galyon, H. et al. *Polymers* **2022**, *14*(10), 2103.

Table 1. Effects of PET contamination in feed on the *in vitro* ruminal activity.

PET %	pH	NH ₃ -N mg/100mL	DMD mg/g	OMD mg/g	Protozoa 10 ⁵ /mL	PF
0	6.54	24.90 ^b	671.9 ^a	721.2 ^a	4.38 ^a	2.272 ^a
0.6	6.56	25.51 ^{ab}	666.8 ^{ab}	712.1 ^{ab}	4.18 ^a	2.164 ^b
1.2	6.56	25.70 ^a	652.7 ^{bc}	697.7 ^{bc}	4.24 ^a	2.077 ^c
1.8	6.53	26.56 ^a	644.4 ^c	688.3 ^c	3.87 ^b	2.057 ^c
SEM	0.041	1.022	9.36	9.99	0.235	0.0392

^{a-c} Within a column, different superscripts differ significantly (P < 0.05).




Figure 1. Effect of PET on the *in vitro* GP.