

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Natural iron ligands promote a metal-based oxidation mechanism for the Fenton reaction in water environments

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1768966> since 2021-02-04T17:53:54Z

Published version:

DOI:10.1016/j.jhazmat.2020.122413

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Natural Iron Ligands Promote a Metal-Based Oxidation

Mechanism for the Fenton Reaction in Water Environments

Giulio Farinelli,[†] Marco Minella,[‡] Marco Pazzi,[‡] Stefanos Giannakis,[§] Cesar Pulgarin,^{||} Davide Vione,^{*,‡} Alberto Tiraferri^{*,†}

[†]Department of Environment, Land and Infrastructure Engineering (DIATI), Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129, Turin, Italy

[‡]Department of Chemistry, University of Turin, Via Pietro Giuria 7, 10125 Turin, Italy

[§]Universidad Politécnica de Madrid, E.T.S. Ingenieros de Caminos, Canales y Puertos, Departamento de Hidráulica, Energía y Medio Ambiente, Unidad docente Ingeniería Sanitaria, c/ Profesor Aranguren, s/n, ES-28040 Madrid, España

^{||}SB, ISIC, Group of Advanced Oxidation Processes, Ecole Polytechnique Fédérale de Lausanne (EPFL), Station 6, 1015 Lausanne, Switzerland

Corresponding Authors

Prof. Alberto Tiraferri. E-mail: alberto.tiraferri@polito.it

Prof. Davide Vione. E-mail: davide.vione@unito.it

Abstract

The Fenton reaction is an effective advanced oxidation process occurring in nature and applied in engineering processes toward the degradation of harmful substances, including contaminants of emerging concern. The traditional Fenton application can be remarkably improved by using iron complexes with organic ligands, which allow for the degradation of contaminants at near-neutral pH and for the reduction of sludge production. This work discusses the mechanisms involved both in the classic Fenton process and in the presence of ligands that coordinate iron. Cyclohexane was selected as mechanistic probe, by following the formation of the relevant products, namely, cyclohexanol (A) and cyclohexanone (K). As expected, the classic Fenton process was associated with an A/K ratio of approximately 1, evidence of a dominant free radical behavior. Significantly, the presence of widely common natural and synthetic carboxyl ligands selectively produced mostly the alcoholic species in the first oxidation step. A ferryl-based mechanism was thus preferred when iron complexes were formed. Common iron ligands are here proven to direct the reaction pathway towards a selective metal-based catalysis. Such a system may be more easily engineered than a free radical-based one to safely remove hazardous contaminants from water and minimize the production of harmful intermediates.

Keywords: advanced oxidation; Fenton mechanism; iron ligands; free radical mechanism; metal-based catalysis.

43 **Highlights**

- 44 • Systematic work investigating the Fenton mechanism in the presence of ligands
- 45 • The presence of ligands promotes metal-based catalysis
- 46 • The oxidation mechanism depends on the concentration of the iron ligand
- 47 • Different Fenton mechanisms occur by varying the pH
- 48 • A ferryl species is reasonably involved in the oxidation pathway with ligands

49

1. INTRODUCTION

The Fenton reaction has been known since over a century (Fenton 1894; Giannakis 2019) and represents an effective advanced oxidation process to remove many recalcitrant pollutants (Barbeni et al. 1987; Lipczynskakochany 1991; Maillard et al. 1992). Nevertheless, this process has some major drawbacks when applied to contaminated waters, including the need to operate under acidic conditions, the formation of large slurries of precipitated iron in the final basification step, and the competition of pollutants with other aqueous species, e.g., carbonate and dissolved organic matter, in the reaction with non-selective free radicals (Pignatello et al. 1999; Vione et al. 2014). The reliance upon non-selective radicals decreases the efficiency of the process in the presence of interfering species, and prevents control over the formation of potentially hazardous reaction intermediates. A possible solution to tackle these limitations is the use of iron ligands, in analogy with the biological oxidations catalyzed by iron complexes, e.g., Cytochrome P450 (Cyt P450) or Taurine Dioxygenase (TauD) (Hohenberger et al. 2012). The use of similar iron(II) complexes as engineered catalysts is a promising variation of the traditional Fenton process, because it permits the degradation of persistent contaminants at near-neutral pH, while reducing the sludge production by keeping iron in solution (dos Santos et al. 2011; Song et al. 2015; Giannakis et al. 2016; Clarizia et al. 2017; Messele et al. 2019; Das et al. 2020). Nevertheless, little is known regarding the reaction pathway in these systems.

The classic description of the traditional Fenton process is based on the first reaction of the Haber-Weiss mechanism proposal (eq. 1) (Haber and Weiss 1932):



More recent investigations also proposed a mechanism that includes the formation of a transient species where iron has a formal redox state of IV (Goldstein et al. 1993; Bossmann et al. 1998). The presence of other superoxidized iron species (not only $\text{Fe}^{\text{IV}}_{(\text{aq})}$, but also $\text{Fe}^{\text{V}}_{(\text{aq})}$ and $\text{Fe}^{\text{VI}}_{(\text{aq})}$) has been additionally proven by stopped-flow experiments and UV-Vis spectroscopy (Wink et al. 1994). However, the reactivity, role, and the stability of such species is only partially known. Essential contributions were provided in the works by Bossmann et al. (1998) and Pignatello et al. (1999), who highlighted the different reactivity of ferryl and hydroxyl radicals. In particular, Bossmann and co-workers compared the products of the reaction of 2,4-xylylene in genuine $\bullet\text{OH}$ generation processes (H_2O_2 photolysis) with those obtained from Fenton reactions, showing the formation of different reactive species (Bossmann et al. 1998). Pignatello et al. (1999) provided evidence of alternative oxidants to $\bullet\text{OH}$. More recently, Minero and co-workers concluded that $\bullet\text{OH}$ (60% yield) and other species (e.g., FeO^{2+}) (40% yield) are formed simultaneously (Minero et al. 2013).


The oxidation mechanism is amply debated also when the system comprises iron complexes, e.g., in the presence of iron ligands. Rush and Koppenol investigated a variety of chelated iron complexes, concluding that a metal-oxo species was generated in neutral solutions, while $\bullet\text{OH}$ species dominated the process in acidic solutions of non-chelated iron (Rush and Koppenol 1988). Sutton et al. (1987) reached a different conclusion, proposing that free iron generates a metal-oxo species as the primary oxidant while $\bullet\text{OH}$ is dominant when chelated iron is present. A reasonable rationalization of this apparent discrepancy is that metal-oxo species and $\bullet\text{OH}$ can both be generated concurrently in Fenton systems. Indeed, Yamazaki and Piette (1990) suggested that more than one type of oxidizing intermediate is present, and that the stoichiometry $\bullet\text{OH}:\text{Fe}(\text{II})$ is also a function of the nature of the prevailing iron

chelators. Different chelating agents for Fe(II) have been reported to promote the formation of oxoiron (ferryl) species in addition to, or instead of, $\bullet\text{OH}$, thus accelerating (e.g., with fulvic acid (Southworth and Voelker 2003), oxalate (Park et al. 1997), and EDTA (Rush and Koppenol 1986)) or suppressing (e.g., with phosphates) the Fenton reaction (Iwahashi et al. 1990).

Because the direct experimental observation of the key intermediates involved in the oxidation pathways is challenging, indirect probes were developed (Klopstra et al. 2004; England et al. 2008; Oloo and Que 2013; Dong et al. 2018). Cyclohexane (Cy) was used in previous studies as an advantageous tool to discriminate between the different pathways of the Fenton reaction in organic solvents, by following the selective production of two products, namely cyclohexanol and cyclohexanone in different ratios (Oloo and Que 2013). Reactions initiated by hydroxyl radicals produce long-lived alkyl radical intermediates. These intermediates may react with dissolved molecular oxygen at diffusion-controlled rates to produce alkylperoxyl radicals, whose subsequent reaction is a Russell-type termination that gives equimolar quantities of alcohol (A) and ketone (K) (**Scheme 1a**) (Russell 1957; Meslennikov et al. 1979). Therefore, $A/K \sim 1$ suggests the occurrence of hydroxyl radical-based reaction pathways. In contrast, an A/K ratio different than 1 is indicative of a non-free radical mechanism of oxidation, i.e., the presence of metal-based oxidant species (**Scheme 1b**). However, to our knowledge, Cy oxidation has never been used as a probe to clarify in a systematic way the mechanism of the Fenton reactions in water.

By using this mechanistic tool based on Cy oxidation, in this work we (*i*) support previous findings that (*a*) the classic $\text{Fe}^{2+} + \text{H}_2\text{O}_2$ process yields $\bullet\text{OH}$ in acidic conditions, but the reactant shifts from $\bullet\text{OH}$ to ferryl at neutral pH values, and (*b*) the process combining the

commercial iron tetra-amido macrocyclic ligand compound (Fe-TAML[®]) and H₂O₂ operates through a ferryl species.



Scheme 1. Proposed mechanism for cyclohexane oxidation. **A:** free radical path with Russell termination type. **B:** a metal-based path.

One of the hypotheses of this work is that many iron ligands can promote a metal-based reaction process. As such, eight ligands were tested, namely, citric acid, tartaric acid, malic acid, quinic acid, EDTA, EDDS, and NTA, as well as Fe-TAML[®] (see Figure S1 in the Supplementary Material for its molecular structure). Therefore, we (ii) provide evidence of the nature of Fenton reactive species in the presence of several common Fe(II) ligands, also as a function of ligand concentration to control the reaction pathway. The first seven ligands are well-known iron chelators, applied to perform oxidation reactions in water at near-neutral pH. Fe-TAML[®] and the classic Fenton reagents at pH 3 are studied as standard controls for a metal-based and (supposed) free radical process, respectively (Collins 2002; de Oliveira et al. 2007; Ghosh et al. 2008). Through an active species linked to the iron-ligand complex, a metal-based mechanism can modulate the path of the reaction and generate fewer and more predictable by-products. Therefore, by verifying the involvement of a metal-based mechanism during a classic or modified Fenton process in water, one can open the route toward a safer oxidation of hazardous substances, e.g., phenols, pharmaceuticals, and pesticides.

2. EXPERIMENTAL

2.1 Chemicals

Fe(III)-TAML[®] was purchased from GreenOx Catalysts Inc. (Pittsburgh, PA, U.S.A.). Sodium phosphate tribasic was obtained from Carlo Erba (Italy). All the other reagents, buffer solutions, and solvents were purchased from Sigma-Aldrich. Water was of Milli-Q quality (TOC 2 ppb, resistivity $\geq 18.2 \text{ M}\Omega \text{ cm}$).

2.2 Reaction conditions

The reaction experiments were carried out at room temperature in a 20 mL solution for 10 min under continuous stirring, and were performed within 40 mL vials equipped by caps provided with septum. The concentrations of the catalyst (computed in terms of iron concentration), reagent (hydrogen peroxide), and substrate (cyclohexane) were 10^{-7} , 10^{-6} , and 10^{-4} mol/L, respectively, resulting in a relative ratio of 1:10:1000. While a 1:10 catalyst:oxidant ratio is typical of engineered applications, an excess of substrate was used here to avoid the subsequent oxidation of one of the major by-products, namely, cyclohexanol. Phosphate buffer (10 mM) or perchloric acid were used to fix the pH. The value of the binding constant of phosphate with iron (10^{-16}) is negligible compared to that of the various ligands investigated here. Therefore, significant formation of iron phosphate can be ruled out in favor of the formation of iron-ligand complexes. All the reactions were quenched by using *tert*-butyl alcohol (t-BuOH) as scavenger of reactive species (excess concentration of 30 mM, thus 300:1 compared to cyclohexane) for subsequent analysis (Rahhal and Richter 1988). The iron-ligand complexes were prepared in equimolar ratio in a concentrated stock solution (0.01 mM) by stirring the mixture of the iron and ligand for 5 min, and were then diluted to 0.1 μM .

2.3 Analytical conditions

The headspace, solid phase microextraction technique (HS-SPME) was chosen as extraction method before carrying out GC-MS analysis. This technique does not require solvents and allows for highly sensitive analyses. Following each reaction experiment, the vials were left in a thermostatic bath at 50 °C for 10 min to promote the transfer of all the relevant compounds into the gas-phase headspace. Then, a SPME fiber (df 75 µm, fiber assembly carboxen/polydimethylsiloxane) was injected through the septum of the cap and was left in the headspace for 10 min, before withdrawing it for the subsequent GC-MS analysis. Samples were analyzed on an Agilent 6890 GC system coupled with an Agilent 5973N mass selective detector (MSD). For the chromatographic separation, a Zebron-5MS capillary column (30 m × 250 µm × 0.25 µm) was used. The injection port temperature was 280 °C, and the oven temperature program was set as follows: 40°C for 5 min, then an increase to 310°C at a rate of 15 °C min⁻¹ (total run time 28.00 min). Helium was used as carrier gas at a constant flow of 1.2 mL min⁻¹, and the injector was held in splitless mode. The interface temperature was 280°C, the ionization energy was 70 eV, and the mass spectrometer operated in SIM mode acquiring the following fragments: 84, 56, 41 (cyclohexane); 82, 67, 57 (cyclohexanol); 98, 55, 42 (cyclohexanone). Duplicate experiments for some of the tests discussed in this study indicated high reproducibility of the results; the error associated to the data and presented below is related to the intrinsic uncertainty of the SPME technique, computed as the average among the standard deviations reported in the 525.2 method provided by the EPA.

A few control experiments were carried out to study the Fenton degradation of phenol (by both Fe²⁺ + H₂O₂ and Fe^{III}-TAML[®] + H₂O₂) using t-BuOH as scavenger, to take advantage of the ability of this compound to react with •OH faster than with electron-capture oxidants such as ferryl (Buxton et al., 1988; Rahhal and Richter 1988). The time evolution of phenol was monitored by liquid chromatography (see the Supplementary Material for additional details).

3. RESULTS AND DISCUSSION

3.1 Iron ligands promote the preferential formation of cyclohexanol

Eight widely available ligands able to complex iron in a stable fashion were tested, namely: citric acid, tartaric acid, malic acid, quinic acid, EDTA (ethylenediaminetetraacetic acid), EDDS (ethylenediamine-N,N'-disuccinic acid), NTA (nitrilotriacetate), and TAML (tetraamidomacrocyclic ligand). These ligands belong to two macro-categories: natural (citric, tartaric, malic, quinic acid) and artificial ligands (EDTA, EDDS, NTA). This choice was provisionally made to gain insight into any possible correlation between the two categories, or among ligands in the same category. The Fe-TAML[®] system is well-known to induce a metal-based oxidation process via a ferryl species, thus we expected an alcohol/ketone (A/K) product ratio different from 1 upon oxidation of cyclohexane (Collins 2002; de Oliveira et al. 2007). Conversely, the Fenton process at pH 3 generates mostly hydroxyl radicals, or at least the hydroxyl radical is the most reactive (although not the only one) species in the system. Therefore, the reaction should proceed mostly via a free radical mechanism, with an A/K ratio around 1 (Minero et al. 2013).

Figure 1 shows the A/K ratio values obtained with all the investigated ligands and in the absence of ligands, i.e., classic Fenton. The results obtained from oxidation tests are in line with expectations, thus attesting to the validity of the method. The classic Fenton process at pH 3 showed an A/K ratio around 1, while an A/K ratio significantly higher than 1 was obtained with all the other investigated ligands. Therefore, it is reasonable to hypothesize that the presence of an iron ligand in water promotes a metal-based oxidation.




Figure 1. A/K ratio observed in the oxidation of cyclohexane with different iron ligands at pH 7 (phosphate buffer) and with the classic Fenton process at pH 3 (perchloric acid) after 30 min of reaction.

These conclusions are confirmed by the results of the degradation of phenol, with $\text{Fe}^{2+} + \text{H}_2\text{O}_2$ at pH 3 and with $\text{Fe-TAML}^{\text{®}} + \text{H}_2\text{O}_2$ (see Figure S2 in Supplementary Material, as well as the related text). We carried out these experiments both in the absence and in the presence of t-BuOH, which reacts with $\bullet\text{OH}$ faster than with ferryl (Buxton et al., 1988; Rahhal and Richter 1988). However, to avoid total quenching of the system by t-BuOH, the t-BuOH:phenol ratio was 40:1 and not 300:1 as per the Cy experiments. In the case of $\text{Fe}^{2+} + \text{H}_2\text{O}_2$ t-BuOH strongly inhibited phenol degradation, while in the case of $\text{Fe-TAML}^{\text{®}}$ the effect of the alcohol was practically negligible. These findings are consistent with $\bullet\text{OH}$ being involved in phenol degradation by $\text{Fe}^{2+} + \text{H}_2\text{O}_2$ at pH 3, and with ferryl playing the same role with $\text{Fe-TAML}^{\text{®}} + \text{H}_2\text{O}_2$. Indeed, the t-BuOH scavenging experiments agree with the A/K ratios derived from the Cy degradation experiments (**Figure 1**).

It is important to check for possible variations of the A/K ratio with reaction time, to ensure that unbiased conclusions are obtained. **Figure 2** shows the kinetics of Cy degradation with citric acid (**Figure 2a**), malic acid (**Figure 2b**), and quinic acid (**Figure 2c**) as iron ligands (left Y-axis: A/K ratio, right Y-axis: Cy degradation). Cy degradation with citric and quinic acids was very fast and the process reached completion after roughly 2 min of reaction. The corresponding A/K ratio remained stable and significantly larger than 1 during the entire duration of the test (30 min). On the other hand, **Figure 2b** (malic acid) shows slower kinetics of degradation, with an A/K ratio reaching a peak value after 10 min of reaction. This slower degradation allowed for an easier monitoring of the initial preferred formation of the alcohol species ($A/K > 1$) and the subsequent oxidation of the alcohol into the ketone, which caused a slight A/K reduction following the peak. The large excess of the initial Cy consumed almost all the reactive species, thereby limiting their availability for alcohol oxidation. By monitoring the A/K time evolution, one can thus be confident that there is negligible bias linked to the further evolution of the system ($A \rightarrow K$ oxidation, or further Fenton processes involving, e.g., Fe(III) after total Fe(II) consumption) after the initial reaction step (Russell 1957).

The fact that the A/K ratio did not change much after the initial step suggests that the mechanistic conditions reflect those of the initial reaction between Fe(II) and H_2O_2 . Under our experimental conditions and based on the stoichiometry of reaction (1), this process would entail total consumption of Fe(II) that would be oxidized to Fe(III), and 10% degradation of H_2O_2 . Afterwards, Fe(III) would be recycled to Fe(II) at the expense of the remaining H_2O_2 .




Figure 2. Fraction of degraded cyclohexane with respect to the total degraded amount (circles, right axis) and trend of selectivity, i.e., A/K ratio (squares, left axis), as a function of time in a system containing hydrogen peroxide as reactant and a) citric acid, b) malic acid, or c) quinic acid as iron ligands. The pH of the aqueous system was buffered at 7 (phosphate buffer). The solid lines connecting the circles are only intended as a guide for the eye.

The influence of the reaction environment was also studied by following the A/K ratio in the oxidation of Cy at different pH values (3-7) (**Table 1**). All the individual concentrations of alcohol and ketone products obtained during the experiments are presented in the Supplementary Material (Tables S1-S3). Citric, malic, and quinic acid were chosen as iron ligands because the iron binding constant of these compounds would not change significantly within the explored pH range, thereby allowing for the pH value to solely affect the Fenton process (Supplementary Material, Figure S3). Phosphate (10 mM) was used to buffer the pH at values of 5, 6, and 7, while perchloric acid (1.16 M stock solution) was employed in tests performed at pH 3. In all these cases the A/K ratios were higher than 1, independently of the pH values. This finding suggests that a metal-based mechanism in the presence of the three ligands was observed regardless of the acidity of the solution. Although acidity has no direct effect on the mechanistic path, the conditional binding constant may be function of the pH and this indirect effect of the solution acidity is discussed below.

Table 1. A/K ratio for the oxidation of cyclohexane at different pH values with citric, malic and quinic acid as iron ligands. The pH was fixed with phosphate buffer except for pH 3 (HClO₄). The reaction time was 30 min.

	pH 3	pH 5	pH 6	pH 7
Citric acid	10±1	7.4±1.0	8.5±1.2	8.6±1.2
Malic acid	18±3	15±2	16±2	15±2
Quinic acid	28±4	31±4	23±3	28±4

3.2 Switching the mechanism from free radical to metal-based catalysis

In the previous section, we hypothesized that the mechanistic degradation path depends on the presence of the ligand, and possibly on its concentration and conditional binding constant. In order to investigate the influence of the ligand concentration and of its conditional binding constant with the metal (i.e., the value of the binding constant that takes into account the protonation of the ligand at the given pH value), EDTA and EDDS were chosen as iron ligands and applied at acidic pH. Despite their structural similarity, these two ligands behave differently in terms of their conditional binding constant as a function of pH. Specifically, while the conditional binding constant at pH 7 is high for both EDTA and EDDS ($\sim 10^{11}$ and 10^6 , respectively), at pH 3 EDDS features a low conditional binding constant (10^{-3}), while that of EDTA is still relatively high ($\sim 10^4$) (see Supplementary Material, Figure S3). Therefore, when using EDDS as iron ligand at pH 3, one expects a high amount of free iron to occur in solution, which could reasonably induce a classic Fenton process (free radical mechanism). In this series of experiments, the pH value was fixed at a value of 3 by addition of perchloric acid.

Figure 3 reports the A/K ratios observed when EDDS and EDTA were used as ligands at pH 3, at various [Ligand]:[Fe(II)] ratios. All the individual concentrations of alcohol and ketone products obtained during the experiments are presented in the Supplementary Material (Tables S4-S5). Please note that **Figure 1** summarized instead the A/K ratios measured at 1:1 [Ligand]:[Fe(II)] ratio and at pH 7. Consistently with the hypotheses, at 0:1 ligand:iron ratio (i.e., with no ligand in solution) we observed the classic Fenton process and the A/K ratio was close to 1. The A/K ratio remained always close to 1 with EDDS, regardless of its concentration. Based on the conditional binding constants, when using EDDS as iron ligand at pH 3, one expects free iron to occur in solution, which could reasonably promote a process similar to a classic $\bullet\text{OH}$ -based Fenton reaction. In contrast, with EDTA the reaction clearly

switched from a free radical mechanism to a metal-based one when the ligand concentration increased. These results strongly suggest that the presence of a bonded Fe(II)-ligand complex plays a crucial role in the direction of the mechanistic path. This parameter is a strong function of the concentration of the ligand as well as of the conditional binding constant of the complex metal/ligand. Therefore, one can generalize that the mechanistic path may be mostly imputable to the concentration and the conditional binding constant of the ligand, and that the environmental conditions have importance only if they affect complex formation.




Figure 3. A/K ratio observed after 30 min of reaction in the oxidation of cyclohexane with EDTA and EDDS as iron ligands, added at different concentrations. The runs were carried out at pH 3. The condition of no added Ligand ($[Ligand]/[Fe(II)] = 0$) corresponds to the classic Fenton process. The solid lines are only intended as a guide for the eye. The dashed line depicts the expected A/K ratio for a pure free radicals-based catalysis.

3.3 Evidence of the effect of pH on the traditional Fenton process

Finally, we provide some insight into the Fenton mechanism at different pH values. **Figure 4** shows the A/K ratio obtained with the classic Fenton process in the 3-7 pH range, fixed by use of phosphate buffer (PBS). Additional experiments were also conducted by fixing the pH at 3 and 4 using perchloric acid instead of phosphate buffer. Please note that Fe(II) was dosed at low concentration (0.1 μM) to avoid its precipitation as hydroxide, which would otherwise take place at near-neutral pH ($\text{pK}_s^{\text{Fe(OH)}_2} = 15.1$ (Harris 2006)). The results obtained in the absence of ligands imply that the classic Fenton reaction proceeds through a free radical or mixed mechanism up to pH 5. Above this value, the mechanism switched to a preferential metal-based one. This result is supported by previous reports proposing that a ferryl species is involved in the Fenton reaction at near-neutral pH (Rush et al. 1990; Bossmann et al. 2004; Bataineh et al. 2012). The presence of a ferryl species at pH 6-7 means that the classic Fenton reaction will be less reactive under near-neutral conditions, since the non-coordinated ferryl species is considered less reactive than the hydroxyl radical (Bataineh et al. 2012).




Figure 4 A/K ratio observed in the classic Fenton process performed at different pH values.

4. CONCLUSIONS

The oxidation mechanism of the Fenton reaction was here investigated with and without iron ligands in solution. Eight widely common ligands that are able to complex iron(II) were studied, namely, citric acid, tartaric acid, malic acid, quinic acid, EDTA, EDDS, and NTA, plus the Fe-TAML[®] system. The ligand performance was tested toward the oxidation of cyclohexane by following the formation of the main products, namely, cyclohexanol (A) and cyclohexanone (K). Measurement of the concentration ratio between these two species (A/K ratio parameter) during the reaction was used to provide evidence of the mechanisms involved in the oxidation of the substrate.

All the tested ligands showed high A/K ratios, which is a proper index of a metal-based behavior, including the well-known Fe-TAML[®] system that was expected to behave in this fashion. Also expected was the fact that the classic Fenton process was associated with an A/K ratio of approximately 1, evidence of a free radical process, which adds further evidence in favor of cyclohexane as suitable probe when coupled with the A/K ratio.

This work proves the ability of simple iron ligands to drive the reaction pathway towards selective metal-based catalysis, as opposed to the activity of free radicals that promote undifferentiated oxidation of substrates in solution. Selective catalysis allows for better control of the degradation pathway of harmful contaminants, to avoid the formation of toxic by-products. This study also provides a further insight into the role of pH in the classic Fenton process. The ubiquitous character of the Fenton process in nature and of iron complexes formed in water (e.g., citrate), alongside the generation of H₂O₂ in illuminated NOM-containing environments, lead to (photo)Fenton reactions during the diurnal cycles. As such, the present study holds important implications also in the elucidation of the Fenton process that occurs both in nature and in engineering applications, and sets the basis for

further investigation concerning the effectiveness of ligand-mediated oxidation of natural and anthropogenic contaminants.

ASSOCIATED CONTENT

Supporting Information. The Supporting Information is available free of charge on the ACS Publications website

AUTHOR INFORMATION

Corresponding Authors*

D.V.: Tel.: +39 0116705296. E-mail: davide.vione@unito.it

A.T.: Tel.: +39 0110907628. E-mail: alberto.tiraferri@polito.it

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This study was financially supported by Politecnico di Torino and University of Torino. DV acknowledges financial support by University of Torino and Compagnia di San Paolo (project CSTO168282-ABATEPHARM).

References

- Barbeni, M., Minero, C., Pelizzetti, E., Borgarello, E., Serpone, N., 1987. Chemical Degradation of Chlorophenols with Fenton Reagent ($\text{Fe}^{2+} + \text{H}_2\text{O}_2$). *Chemosphere*. 16, 2225-2237. [https://doi.org/10.1016/0045-6535\(87\)90281-5](https://doi.org/10.1016/0045-6535(87)90281-5)
- Bataineh, H., Pestovsky, O., Bakac, A., 2012. pH-induced mechanistic changeover from hydroxyl radicals to iron(IV) in the Fenton reaction. *Chem. Sci*. 3, 1594-1599. <https://doi.org/10.1039/c2sc20099f>
- Bossmann, S. H., Oliveros, E., Gob, S., Siegwart, S., Dahlen, E. P., Payawan, L., Straub, M., Worner, M., Braun, A. M., 1998. New evidence against hydroxyl radicals as reactive intermediates in the thermal and photochemically enhanced fenton reactions. *J. Phys. Chem. A*. 102, 5542-5550. <https://doi.org/10.1021/Jp980129j>
- Bossmann, S. H., Oliveros, E., Kantor, M., Niebler, S., Bonfill, A., Shahin, N., Worner, M., Braun, A. M., 2004. New insights into the mechanisms of the thermal Fenton reactions occurring using different iron(II)-complexes. *Water Sci. Technol*. 49, 75-80. <https://doi.org/10.2166/wst.2004.0224>
- Buxton, G. V., Greenstock, C. L., Helman, W. P., Ross, A. B., 1988. Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals ($\cdot\text{OH}/\cdot\text{O}^-$) in aqueous solution. *J. Phys. Chem. Ref. Data* 17(2), 513-886. <https://doi.org/10.1063/1.555805>
- Clarizia, L., Russo, D., Di Somma, I., Marotta, R., Andreozzi, R., 2017. Homogeneous photo-Fenton processes at near neutral pH: A review. *Appl. Catal. B-Environ*. 209, 358-371. <https://doi.org/10.1016/j.apcatb.2017.03.011>
- Collins, T. J., 2002. TAML oxidant activators: A new approach to the activation of hydrogen peroxide for environmentally significant problems. *Acc. Chem. Res*. 35, 782-790. <https://doi.org/10.1021/ar010079s>
- Das, B., Baruah, J. B., Sharma, M., Sarma, B., Varunak, G. V., Satyanarayana, L., Roy, S., Bhattacharyya, P. K., Kanta Borah, K., Bania, K. K., 2020. Self pH regulated iron(II) catalyst for radical free oxidation of benzyl alcohols. *Appl. Catal. A-Gen*. 589, 117292. <https://doi.org/10.1016/j.apcata.2019.117292>
- de Oliveira, F. T., Chanda, A., Banerjee, D., Shan, X. P., Mondal, S., Que, L., Bominaar, E. L., Munck, E., Collins, T. J., 2007. Chemical and spectroscopic evidence for an Fe-V-Oxo complex. *Science*. 315, 835-838. <https://doi.org/10.1126/science.1133417>
- Dong, J. J., Fernandez-Fueyo, E., Hollmann, F., Paul, C. E., Pesic, M., Schmidt, S., Wang, Y. H., Younes, S., Zhang, W. Y., 2018. Biocatalytic Oxidation Reactions: A Chemist's Perspective. *Angew. Chem. Int. Edit*. 57, 9238-9261. <https://doi.org/10.1002/anie.201800343>
- dos Santos, J. S., Palaretti, V., de Faria, A. L., Crevelin, E. J., de Moraes, L. A. B., Assis, M. D., 2011. Biomimetic simazine oxidation catalyzed by metalloporphyrins. *Appl. Catal. A-Gen*. 408, 163-170. <https://doi.org/10.1016/j.apcata.2011.09.023>

417 England, J., Davies, C. R., Banaru, M., White, A. J. P., Britovsek, G. J. P., 2008. Catalyst
 418 stability determines the catalytic activity of non-heme iron catalysts in the oxidation of
 419 alkanes. *Adv. Synth. Catal.* 350, 883-897. <https://doi.org/10.1002/adsc.200700462>

420 Fenton, H. G. H., 1894. LXXIII.—Oxidation of tartaric acid in presence of iron *J. Chem. Soc.*
 421 *Trans.* 65, 899-910. <https://doi.org/10.1039/CT8946500899>

422 Ghosh, A., Mitchell, D. A., Chanda, A., Ryabov, A. D., Popescu, D. L., Upham, E. C.,
 423 Collins, G. J., Collins, T. J., 2008. Catalase-Peroxidase Activity of Iron(III)-TAML
 424 Activators of Hydrogen Peroxide. *J. Am. Chem. Soc.* 130, 15116-15126.
 425 <https://doi.org/10.1021/ja8043689>

426 Giannakis, S., 2019. A review of the concepts, recent advances and niche applications of the
 427 (photo) Fenton process, beyond water/wastewater treatment: Surface functionalization,
 428 biomass treatment, combatting cancer and other medical uses. *Appl. Catal. B-Environ.* 248,
 429 309-319. <https://doi.org/10.1016/j.apcatb.2019.02.025>

430 Giannakis, S., Ruales-Lonfat, C., Rtimi, S., Thabet, S., Cotton, P., Pulgarin, C., 2016. Castles
 431 fall from inside: Evidence for dominant internal photo-catalytic mechanisms during treatment
 432 of *Saccharomyces cerevisiae* by photo-Fenton at near-neutral pH. *Appl. Catal. B-Environ.*
 433 185, 150-162. <https://doi.org/10.1016/j.apcatb.2015.12.016>

434 Goldstein, S., Meyerstein, D., Czapski, G., 1993. The Fenton Reagents. *Free Radic. Biol.*
 435 *Med.* 15, 435-445. [https://doi.org/10.1016/0891-5849\(93\)90043-T](https://doi.org/10.1016/0891-5849(93)90043-T)

436 Haber, F., Weiss, J., 1932. On the catalysis of hydroperoxide. *Naturwissenschaften.* 20, 948-
 437 950. <https://doi.org/10.1007/BF01504715>

438 Harris, D. (2006). Quantitative Chemical Analysis, W. H. Freeman.

439 Hohenberger, J., Ray, K., Meyer, K., 2012. The biology and chemistry of high-valent iron-
 440 oxo and iron-nitrido complexes. *Nat. Commun.* 3, <https://doi.org/10.1038/Ncomms1718>

441 Iwahashi, H., Ishii, T., Sugata, R., Kido, R., 1990. The Effects of Caffeic Acid and Its
 442 Related Catechols on Hydroxyl Radical Formation by 3-Hydroxyanthranilic Acid, Ferric-
 443 Chloride, and Hydrogen-Peroxide. *Arch. Biochem. Biophys.* 276, 242-247.
 444 [https://doi.org/10.1016/0003-9861\(90\)90033-U](https://doi.org/10.1016/0003-9861(90)90033-U)

445 Klopstra, M., Roelfes, G., Hage, R., Kellogg, R. M., Feringa, B. L., 2004. Non-heme iron
 446 complexes for stereoselective oxidation: Tuning of the selectivity in dihydroxylation using
 447 different solvents. *Eur. J. Inorg. Chem.* 846-856. <https://doi.org/10.1002/ejic.200300667>

448 Lipczynskakochany, E., 1991. Degradation of Aqueous Nitrophenols and Nitrobenzene by
 449 Means of the Fenton Reaction. *Chemosphere.* 22, 529-536. [https://doi.org/10.1016/0045-6535\(91\)90064-K](https://doi.org/10.1016/0045-6535(91)90064-K)

451 Maillard, C., Guillard, C., Pichat, P., 1992. Comparative Effects of the TiO_2 -Uv, H_2O_2 -Uv,
 452 H_2O_2 - Fe^{2+} Systems on the Disappearance Rate of Benzamide and 4-Hydroxybenzamide in
 453 Water. *Chemosphere.* 24, 1085-1094. [https://doi.org/10.1016/0045-6535\(92\)90199-2](https://doi.org/10.1016/0045-6535(92)90199-2)

454 Meslennikov, S. I., Galimova, L. G., Komissarov, V. D., 1979. Disproportionation Rate and
 455 Products of Cyclohexylperoxy Radicals. *B. Acad. Sci. USSR Ch+*. 28, 585-588.
 456 <https://doi.org/Doi 10.1007/Bf00924841>

457 Messele, S. A., Bengoa, C., Stuber, F. E., Giralt, J., Fortuny, A., Fabregat, A., Font, J., 2019.
 458 Enhanced Degradation of Phenol by a Fenton-Like System (Fe/EDTA/H₂O₂) at
 459 Circumneutral pH. *Catalysts*. 9, 474. <https://doi.org/10.3390/Catal9050474>

460 Minero, C., Lucchiari, M., Maurino, V., Vione, D., 2013. A quantitative assessment of the
 461 production of (OH)-O-center dot and additional oxidants in the dark Fenton reaction: Fenton
 462 degradation of aromatic amines. *RSC Adv.* 3, 26443-26450.
 463 <https://doi.org/10.1039/c3ra44585b>

464 Oloo, W. N., Que, L. (2013). Hydrocarbon Oxidations Catalyzed by Bio-Inspired Nonheme
 465 Iron and Copper Catalyst. *Comprehensive Inorganic Chemistry II*, Elsevier.

466 Park, J. S. B., Wood, P. M., Davies, M. J., Gilbert, B. C., Whitwood, A. C., 1997. A kinetic
 467 and ESR investigation of Iron(II) oxalate oxidation by hydrogen peroxide and dioxygen as a
 468 source of hydroxyl radicals. *Free Radic. Res.* 27, 447-458.
 469 <https://doi.org/10.3109/10715769709065785>

470 Pignatello, J. J., Liu, D., Huston, P., 1999. Evidence for an additional oxidant in the
 471 photoassisted Fenton reaction. *Environ. Sci. Technol.* 33, 1832-1839.
 472 <https://doi.org/10.1021/Es980969b>

473 Rahhal, S., Richter, H. W., 1988. Reduction of Hydrogen-Peroxide by the Ferrous Iron
 474 Chelate of Diethylenetriamine-N,N,N',N'',N''-Pentaacetate. *J Am Chem Soc.* 110, 3126-3133.
 475 <https://doi.org/Doi 10.1021/Ja00218a022>

476 Rush, J. D., Koppenol, W. H., 1986. Oxidizing Intermediates in the Reaction of Ferrous Edta
 477 with Hydrogen Peroxide - Reactions with Organic-Molecules and Ferrocyclochrome-C. *J.*
 478 *Biol. Chem.* 261, 6730-6733.

479 Rush, J. D., Koppenol, W. H., 1988. Reactions of Feiinta and Feiiedda with Hydrogen-
 480 Peroxide. *J. Am. Chem. Soc.* 110, 4957-4963. <https://doi.org/10.1021/Ja00223a013>

481 Rush, J. D., Maskos, Z., Koppenol, W. H., 1990. Distinction between Hydroxyl Radical and
 482 Ferryl Species. *Method. Enzymol.* 186, 148-156. [https://doi.org/10.1016/0076-](https://doi.org/10.1016/0076-6879(90)86104-4)
 483 [6879\(90\)86104-4](https://doi.org/10.1016/0076-6879(90)86104-4)

484 Russell, G., A., 1957. Deuterium-isotope Effects in the Autoxidation of Aralkyl
 485 Hydrocarbons. Mechanism of the Interaction of PEROxy Radicals. *J. Am. Chem. Soc.* 79,
 486 3871-3877. <https://doi.org/10.1021/ja01571a068>

487 Song, Q., Ma, W. H., Jia, M. K., Johnson, D., Huang, Y. P., 2015. Degradation of organic
 488 pollutants in waters by a water-insoluble iron(III) Schiff base complex. *Appl. Catal. A-Gen.*
 489 505, 70-76. <https://doi.org/10.1016/j.apcata.2015.07.028>

490 Southworth, B. A., Voelker, B. M., 2003. Hydroxyl radical production via the photo-Fenton
 491 reaction in the presence of fulvic acid. *Environ. Sci. Technol.* 37, 1130-1136.
 492 <https://doi.org/10.1021/es020757l>

493 Sutton, H. C., Vile, G. F., Winterbourn, C. C., 1987. Radical Driven Fenton Reactions -
 494 Evidence from Paraquat Radical Studies for Production of Tetravalent Iron in the Presence
 495 and Absence of Ethylenediaminetetraacetic Acid. Arch. Biochem. Biophys. 256, 462-471.
 496 [https://doi.org/10.1016/0003-9861\(87\)90603-5](https://doi.org/10.1016/0003-9861(87)90603-5)

497 Vione, D., Minella, M., Maurino, V., Minero, C., 2014. Indirect Photochemistry in Sunlit
 498 Surface Waters: Photoinduced Production of Reactive Transient Species. Chem-Eur J. 20,
 499 10590-10606. <https://doi.org/10.1002/chem.201400413>

500 Wink, D. A., Nims, R. W., Saavedra, J. E., Utermahlen, W. E., Ford, P. C., 1994. The Fenton
 501 Oxidation Mechanism - Reactivities of Biologically Relevant Substrates with 2 Oxidizing
 502 Intermediates Differ from Those Predicted for the Hydroxyl Radical. Proc. Natl. Acad. Sci.
 503 91, 6604-6608. <https://doi.org/10.1073/pnas.91.14.6604>

504 Yamazaki, I., Piette, L. H., 1990. ESR Spin-Trapping Studies on the Reaction of Fe-2+ Ions
 505 with H₂O₂-Reactive Species in Oxygen-Toxicity in Biology. J. Biol. Chem. 265, 13589-
 506 13594.

507

508

509 **Graphical Abstract**

