


## Observation of a Vector Charmoniumlike State at 4.7 GeV/c<sup>2</sup> and Search for Z<sub>cs</sub> in e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>J/ψ

M. Ablikim *et al.*\*  
(BESIII Collaboration)

 (Received 31 August 2023; accepted 30 October 2023; published 22 November 2023)

Using data samples with an integrated luminosity of 5.85 fb<sup>-1</sup> collected at center-of-mass energies from 4.61 to 4.95 GeV with the BESIII detector operating at the BEPCII storage ring, we measure the cross section for the process e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>J/ψ. A new resonance with a mass of  $M = 4708_{-15}^{+17} \pm 21$  MeV/c<sup>2</sup> and a width of  $\Gamma = 126_{-23}^{+27} \pm 30$  MeV is observed in the energy-dependent line shape of the e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>J/ψ cross section with a significance over 5σ. The K<sup>+</sup>J/ψ system is also investigated to search for charged charmoniumlike states, but no significant Z<sub>cs</sub><sup>+</sup> states are observed. Upper limits on the Born cross sections for e<sup>+</sup>e<sup>-</sup> → K<sup>-</sup>Z<sub>cs</sub>(3985)<sup>+</sup>/K<sup>-</sup>Z<sub>cs</sub>(4000)<sup>+</sup> + c.c. with Z<sub>cs</sub>(3985)<sup>±</sup>/Z<sub>cs</sub>(4000)<sup>±</sup> → K<sup>±</sup>J/ψ are reported at 90% confidence levels. The ratio of branching fractions  $\{[\mathcal{B}(Z_{cs}(3985)^+ \rightarrow K^+J/\psi)]/\mathcal{B}[Z_{cs}(3985)^+ \rightarrow (\bar{D}^0D_s^{*+} + \bar{D}^{*0}D_s^+)]\}$  is measured to be less than 0.03 at 90% confidence level.

DOI: 10.1103/PhysRevLett.131.211902

The charmonium system is an ideal place to study the perturbative and nonperturbative strong interactions of quarks and gluons at the hadronic scale. Below open-charm threshold, the spectrum of c $\bar{c}$  charmonium states is well described by a potential model [1]. All observed states have been found within expectations, and excellent agreement has been achieved between theories and experiments. Above open-charm threshold, however, there are still many missing states that have not yet been discovered [2], and, surprisingly, several unexpected states, such as the X(3872) [3], Y(4260) [4], and Z<sub>c</sub>(3900) [5,6], have been observed since 2003. These particles do not match the predictions of the potential models and are widely considered to be good candidates for exotic states [7–10].

Among them, the Y states show strong coupling to hidden-charm final states, such as the experimentally well-established Y(4260) → π<sup>+</sup>π<sup>-</sup>J/ψ [4,11–15] and Y(4360)/Y(4660) → π<sup>+</sup>π<sup>-</sup>ψ(3686) [16–20]. This is in contrast to the vector charmonium states in the same energy region. In recent electron-positron annihilation experiments running at high energies, structures are observed around 4.66 GeV in both open- and hidden-charm final states [21–28]. The BESIII experiment has reported evidence for a structure around 4.7 GeV in the

study of e<sup>+</sup>e<sup>-</sup> → K<sub>S</sub><sup>0</sup>K<sub>S</sub><sup>0</sup>J/ψ [28]. This structure could potentially correspond to the well-known Y(4660) resonance or another excited Y state. On the other hand, potential models predict the 5S and 4D charmonium states in this mass region [29–36]. At present, a comprehensive understanding of the Y states above 4.6 GeV is yet to be achieved, due to limited experimental information. More measurements are urgently needed to clarify their nature.

In addition, an isospin-1/2 charmoniumlike candidate was recently observed by BESIII in the process e<sup>+</sup>e<sup>-</sup> → KZ<sub>cs</sub>(3985), where the charged Z<sub>cs</sub>(3985)<sup>+</sup>, recoiling against a K<sup>-</sup>, was found decaying to (D<sup>0</sup>D<sub>s</sub><sup>\*+</sup> + D<sup>\*0</sup>D<sub>s</sub><sup>+</sup>) [37], and the corresponding neutral Z<sub>cs</sub>(3985)<sup>0</sup>, recoiling against a K<sub>S</sub>, was found decaying to (D<sub>s</sub><sup>+</sup>D<sup>\*-</sup> + D<sub>s</sub><sup>\*+</sup>D<sup>-</sup>) [38]. The Z<sub>cs</sub>(3985) is considered to be the strange partner of the Z<sub>c</sub>(3900) [39–42]. Charge-conjugate modes are implied here and elsewhere unless otherwise specified. BESIII also reported a search for Z<sub>cs</sub><sup>+</sup> → D<sub>s</sub><sup>\*+</sup>D<sup>\*0</sup> [43], finding an excess of Z<sub>cs</sub><sup>+</sup> candidates with a significance of 2.1σ. Meanwhile, LHCb reported tetraquark candidates Z<sub>cs</sub>(4000)<sup>+</sup>/Z<sub>cs</sub>(4220)<sup>+</sup> → K<sup>+</sup>J/ψ in an amplitude analysis of B<sup>+</sup> → K<sup>+</sup>J/ψφ [44]. Although the Z<sub>cs</sub>(3985)<sup>+</sup> and Z<sub>cs</sub>(4000)<sup>+</sup> have comparable masses, their widths are different by nearly an order of magnitude. There are still ongoing debates on whether these particles are the same state or not [41,45–52]. Therefore, to enhance our comprehension of these Z<sub>cs</sub> states and explore the possibility of new Z<sub>cs</sub> states, it is imperative to conduct further measurements of e<sup>+</sup>e<sup>-</sup> → KZ<sub>cs</sub> → K $\bar{K}$ J/ψ [39,47,51–57].

In this Letter, a measurement of the Born cross sections of the process e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>J/ψ is presented at center-of-mass (c.m.) energies from 4.61 to 4.95 GeV [58],

\*Full author list given at the end of the article.

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP<sup>3</sup>.

corresponding to an integrated luminosity of  $5.85 \text{ fb}^{-1}$  [59]. Compared with a previous measurement [60], new data above 4.6 GeV is analyzed for the first time, which enables us to investigate the  $Y$  states above 4.6 GeV with improved precision [61,62]. To achieve much lower background levels and to improve the statistics, both full reconstruction and partial reconstruction methods are applied. By investigating the line shape of the  $e^+e^- \rightarrow K^+K^-J/\psi$  cross section, we report the first observation of the charmoniumlike candidate  $Y(4710)$ . In addition, a search for the  $Z_{cs}^+ \rightarrow K^+J/\psi$  is presented using the same data sample.

The BESIII detector [63] records symmetric  $e^+e^-$  collisions provided by the BEPCII storage ring [64]. Simulated data samples produced with a GEANT4-based [65] Monte Carlo (MC) toolkit, which includes the geometric description of the BESIII detector and the detector response [66], are used to determine detection efficiencies and to estimate background contributions. Signal events for the process  $e^+e^- \rightarrow K^+K^-J/\psi$  are generated at each c.m. energy using a phase space (PHSP) model, while the decay of the  $J/\psi$  into a pair of leptons ( $\mu^+\mu^-/e^+e^-$ ) is modelled with VLL model in EVTGEN [67,68].

For  $K^+K^-J/\psi$  signal candidates, low-momentum kaons have relatively poor detection efficiency. In order to improve the selection efficiency, besides to reconstruct the signals with the both  $K^+K^-$  pair detected, we allow the case with only one of two charged kaons detected, while the  $J/\psi$  candidates are tagged with pairs of charged leptons. Charged tracks detected in the multilayer drift chamber (MDC) are required to be within a polar angle ( $\theta$ ) range of  $|\cos\theta| < 0.93$ , where  $\theta$  is defined with respect to the  $z$  axis, which is the symmetry axis of the MDC. The distance of closest approach to the interaction point must be less than 10 cm along the  $z$  axis, and less than 1 cm in the transverse plane. As kaons and leptons are kinematically well separated, two oppositely charged tracks with momentum greater than  $0.95 \text{ GeV}/c$  in the laboratory frame are assigned as  $\ell^\pm$  for data samples with  $\sqrt{s} < 4.84 \text{ GeV}$ . For  $\sqrt{s} \geq 4.84 \text{ GeV}$ , this value is slightly increased to  $1.05 \text{ GeV}/c$ . The amount of deposited energy in the electromagnetic calorimeter (EMC) is further used to separate muons from electrons. For both muon candidates, the deposited energy in the EMC is required to be less than  $0.4 \text{ GeV}$ , while it is required to be greater than  $1.0 \text{ GeV}$  for electrons. For the remaining charged tracks, particle identification (PID), which combines the measurements of the energy deposited in the MDC ( $dE/dx$ ) and the flight time in the time-of-flight system to form likelihoods  $\mathcal{L}(h)$  ( $h = K, \pi$ ) for each hadron  $h$  hypothesis, is used. Tracks are identified as kaons when the kaon hypothesis has a higher likelihood than the pion hypothesis [ $\mathcal{L}(K) > \mathcal{L}(\pi)$  and  $\mathcal{L}(K) > 0$ ].

For events in which a pair of oppositely charged kaons identified, a four-constraint (4C) kinematic fit imposing

energy-momentum conservation is applied to  $e^+e^- \rightarrow K^+K^-\ell^+\ell^-$ . To remove the radiative Bhabha events, the cosine of the opening angle of the kaon pair is required to be less than 0.98 in the  $J/\psi \rightarrow e^+e^-$  mode. The  $\mu/\pi$  misidentification background in the  $J/\psi \rightarrow \mu^+\mu^-$  mode is suppressed by requiring at least one of the muon candidates has a penetration depth greater than 30 cm in the muon counter (MUC).

For events in which only one kaon is identified, a one-constraint (1C) kinematic fit is performed under the hypothesis of  $e^+e^- \rightarrow K_{\text{miss}}^\pm K^\mp \ell^+ \ell^-$ , where the mass of the missing particle ( $K_{\text{miss}}^\pm$ ) is constrained to the known  $K^+$  mass [69]. In the  $J/\psi \rightarrow e^+e^-$  mode, the dominant background comes from Bhabha events, which are vetoed by requiring  $\cos(\theta_{e^+}) < 0.8$ ,  $\cos(\theta_{e^-}) > -0.8$  and  $|\cos(\theta_{K^\pm})| < 0.8$ . To further reject the radiative Bhabha events ( $\gamma e^+e^-$ ) with  $\gamma$  conversion, where the converted electrons are misidentified as kaons, we require  $|\cos(\alpha_{K^+K^\mp})| < 0.95$ ,  $|\cos(\alpha_{K^\pm e^\pm})| < 0.95$  and  $|\cos(\alpha_{K^\pm e^\mp})| < 0.95$ , where  $\alpha$  is the opening angle between tracks. As one kaon is missing, the  $\mu/\pi$  misidentification background is higher in the  $J/\psi \rightarrow \mu^+\mu^-$  mode. Therefore a tighter requirement is imposed by requiring the penetration depth of both muon candidates in the MUC to be greater than 30 cm.

After the above requirements, a clear  $J/\psi$  peak is observed in the lepton pair invariant mass distribution,  $M(\ell^+\ell^-)$ , as shown in Fig. 1. A study of the inclusive MC sample [28] indicates the background level is low and there is no peaking background. The  $J/\psi$  signal region is defined as  $[3.084, 3.116] \text{ GeV}/c^2$ , and the sideband regions are defined as  $[3.004, 3.068] \text{ GeV}/c^2$  and  $[3.132, 3.196] \text{ GeV}/c^2$ , which are four times as wide as the  $J/\psi$  signal region. The signal yield ( $N^{\text{sig}}$ ) is obtained by sideband subtraction, and the corresponding uncertainty is estimated with the profile likelihood method [70].




FIG. 1. The distribution of  $M(\ell^+\ell^-)$ . Dots with error bars are the selected data at c.m. energies from 4.61 to 4.95 GeV, the red histogram is the signal MC sample and the blue shaded histogram is the background from the inclusive MC sample.




FIG. 2. Fit to the dressed cross section of  $e^+e^- \rightarrow K^+K^-J/\psi$  with the coherent sum of three Breit-Wigner functions (solid curve). The dashed, double dotted, and dash-dotted curves shows the contributions from the  $Y(4710)$ ,  $Y(4230)$ , and  $Y(4500)$ , respectively. The solid dots with error bars are the cross sections from this study, and the open dots with error bars are the cross sections from Ref. [60]. The error bars represent statistical uncertainties only.

The Born cross section is calculated by

$$\sigma^{\text{Born}} = \frac{N^{\text{sig}}}{\mathcal{L}(1 + \delta) \frac{1}{|1 - \Pi|^2} \epsilon \mathcal{B}_{J/\psi}}, \quad (1)$$

where  $\mathcal{L}$  is the integrated luminosity [59],  $(1 + \delta)$  is the ISR correction factor,  $(1/|1 - \Pi|^2)$  is the correction factor for vacuum polarization [71],  $\epsilon$  is the detection efficiency,  $\mathcal{B}_{J/\psi}$  is the sum of the branching fractions of  $J/\psi \rightarrow e^+e^-$  and  $J/\psi \rightarrow \mu^+\mu^-$  [69]. The ISR correction factor and detection efficiency are obtained from the signal MC simulations and are corrected by an iterative weighting method [72]. The numbers of signal events, corrected detection efficiencies, and Born cross sections are summarized in the Supplemental Material [73]. The dressed cross sections ( $\sigma^{\text{Born}}/|1 - \Pi|^2$ ) are shown in Fig. 2 (solid dots with error bars), and a clear structure is found around  $4.75 \text{ GeV}/c^2$ .

To determine the parameters of the structures, a maximum likelihood method is used to fit the dressed cross sections obtained in this Letter and the dressed cross sections with  $\sqrt{s} < 4.61 \text{ GeV}$  [60]. The likelihood is constructed taking into consideration the fluctuations of the numbers of signal and background events. Assuming these  $K^+K^-J/\psi$  signals come from three resonances, the cross section is parameterized as a coherent sum of three relativistic Breit-Wigner functions,

$$\sigma^{\text{Dress}}(\sqrt{s}) = |B_1(\sqrt{s}) + B_2(\sqrt{s})e^{i\phi_2} + B_3(\sqrt{s})e^{i\phi_3}|, \quad (2)$$

where  $B_j = (M_j/\sqrt{s})[\sqrt{12\pi(\Gamma_{ee}\mathcal{B})_j\Gamma_j}/(s - M_j^2 + iM_j\Gamma_j)] \times \sqrt{[\Phi(\sqrt{s})/\Phi(M_j)]}$  with  $j = 1, 2$  or  $3$  is the relativistic Breit-Wigner function,  $\Phi$  is the three-body PHSP factor, and  $\phi_j$  with  $j = 2$  or  $3$  is the relative phase between the

Breit-Wigner functions. In the fit, the masses  $M_j$ , the total widths  $\Gamma_j$ , the products of the electronic partial width and the branching fraction to  $K^+K^-J/\psi$  [ $(\Gamma_{ee}\mathcal{B})_j$ ], and the relative phases  $\phi_j$  are free parameters. There are four solutions with the same masses and widths from the fit as shown in the Supplemental Material [73], and one of them is shown in Fig. 2. The masses and widths of the first and second structures [ $Y(4230)$  and  $Y(4500)$ ] are determined to be  $M_1 = 4226.0^{+1.4}_{-1.4} \text{ MeV}/c^2$ ,  $\Gamma_1 = 70.0^{+3.9}_{-3.6} \text{ MeV}$ ,  $M_2 = 4499.4^{+8.1}_{-7.6} \text{ MeV}/c^2$ , and  $\Gamma_2 = 124^{+22}_{-19} \text{ MeV}$ , respectively, which are consistent with that of Ref. [60]. The mass and width of the third structure, denoted as  $Y(4710)$ , are  $M_3 = 4708^{+17}_{-15} \text{ MeV}/c^2$  and  $\Gamma_3 = 126^{+27}_{-23} \text{ MeV}$ , respectively. Compared to the total fit result, the peak position of the third structure seems to be shifted due to the strong interference shown in Ref. [73]. The aforementioned uncertainties in the masses and widths are statistical only.

Fitting the dressed cross sections with a two-resonance model yields a worse result: the change in the likelihood value from the three-resonance model to the two-resonance model is  $|\Delta(-2 \ln L)| = 43.2$ . Taking into account the change in the number of degrees of freedom ( $\Delta \text{ndf} = 4$ ), the statistical significance for the three-resonance assumption over the two-resonance assumption is  $5.7\sigma$ . In addition, we also fit the cross sections with the coherent sum of three relativistic Breit-Wigner functions and a three-body PHSP term. This assumption improves the fit quality, but the change of the likelihood value is only  $|\Delta(-2 \ln L)| = 1.2$ , which indicates the PHSP term does not make an important contribution.

The systematic uncertainties in the cross section measurement mainly come from the luminosity, tracking efficiency, kinematic fit, MUC response, MC model, radiative correction, and branching fraction. The integrated luminosity is measured using Bhabha events with an uncertainty of 0.6% [58]. The uncertainty of tracking efficiency for the high-momentum leptons is 1.0% per track. By requiring at least one kaon to be detected, the kaon detection efficiency is very high and the uncertainty is negligible. A track helix parameter correction method, as discussed in Ref. [74], is applied to MC events during the kinematic fit. The difference in efficiencies with and without the correction, 2.1%, is assigned as the systematic uncertainty from the kinematic fit. The uncertainty from the MUC response is investigated using the  $e^+e^- \rightarrow \mu^+\mu^-$  data sample, and the difference in efficiencies between the data and MC simulation due to the hit depth requirement in the MUC for the muon candidates, 2.0%, is taken as the systematic uncertainty. Instead of using pure PHSP to model the  $K^+K^-J/\psi$  events, we also consider possible  $f \rightarrow K^+K^-$  in MC simulation based on a partial wave analysis [73], e.g.,  $f_0(980)$ ,  $f_0(1500)$ . The efficiency difference between this model and a three-body PHSP one is 5.9% (8.7%) for the data with  $\sqrt{s} \leq 4.70 \text{ GeV}$  ( $\sqrt{s} > 4.70 \text{ GeV}$ ). To estimate the systematic uncertainty

from the ISR correction factor, we replace the description of the default dressed cross section line shape with the coherent sum of three relativistic Breit-Wigner functions and a PHSP function, or change the parametrization of the Breit-Wigner function. The maximum difference due to the line shape, 0.6%, is assigned as the systematic uncertainty due to the ISR correction factor.

The uncertainty from the branching fraction of  $J/\psi \rightarrow \ell^+ \ell^-$  (0.4%) is taken from the Particle Data Group [69]. Assuming all the sources are independent, the total systematic uncertainty is calculated by adding them in quadrature, resulting in 6.9% (9.4%) for the cross section measurement at  $\sqrt{s} \leq 4.70$  GeV ( $\sqrt{s} > 4.70$  GeV).

The systematic uncertainties in the resonance parameters mainly come from the absolute c.m. energy measurement, the c.m. energy spread, the parameterization of the fit function, and the systematic uncertainty on the cross section measurement. The systematic uncertainty of the c.m. energy is common for all the energies and will propagate to the mass measurement directly. The uncertainty from the c.m. energy spread is estimated by convolving the fit formula with a Gaussian function, whose width is set as the energy dependent beam spread [75]. To estimate the uncertainty from the parameterization of the Breit-Wigner function, the  $\Gamma_j$  in the denominator of the Breit-Wigner function is replaced with a mass dependent width  $\Gamma_j[\Phi(\sqrt{s})/\Phi(M_j)]$ . The uncertainties from the cross section measurement are divided into two parts. The first one is uncommon uncertainties of the measured cross sections among the different c.m. energies, which mainly come from the MC model. The corresponding uncertainty is estimated by including the uncommon uncertainties in the dressed cross section fit, and the differences on the parameters are taken as the corresponding uncertainties. The second part, including all the other uncertainties of the measured cross sections, is common for all the energies, and only affects the parameter ( $\Gamma_{ee}\mathcal{B}$ ).

Intermediate states decaying into  $K^+ J/\psi$ , denoted as  $Z_{cs}^+$ , are of great interest due to their exotic nature. The distribution of the maximum of the invariant masses of the  $K^+ J/\psi$  and  $K^- J/\psi$  combinations,  $M_{\max}(K^\pm J/\psi)$ , is useful for identifying these exotic states [5]. A simultaneous fit to the  $M_{\max}(K^\pm J/\psi)$  spectra from datasets within the energy region  $4.63 \text{ GeV} \leq \sqrt{s} \leq 4.92 \text{ GeV}$  is performed. The datasets at  $\sqrt{s} = 4.61$  and  $4.95$  GeV are not included in the fit due to their relatively low statistics. In the fit, following the model in Ref. [38], the  $Z_{cs}^+$  component is modeled by the product of an  $S$ -wave Breit-Wigner shape with a mass-dependent width:

$$\mathcal{F}(M) \propto \left| \frac{\sqrt{q \cdot p}}{M^2 - m_0^2 + im_0 \Gamma(M)} \right|^2, \quad (3)$$

where  $\Gamma(M) = \Gamma_0 \cdot (p/p^*) \cdot (m_0/M)$ ,  $M$  is the reconstructed mass,  $m_0$  is the resonance mass,  $\Gamma_0$  is the width,




FIG. 3. The  $M_{\max}(K^\pm J/\psi)$  spectrum with the fit results overlaid. The red dashed line is the signal component of  $Z_{cs}$ , the blue dash-dotted line is PHSP (or  $f$  states) and the pink dotted line is the combinatorial background. The sideband events are scaled to the fitted size of combinatorial background, which are shown in green.

$q$  is the bachelor  $K^-$  momentum in the initial  $e^+e^-$  system, and  $p$  is the  $K^+$  momentum in the rest frame of the  $K^+ J/\psi$  system. To account for detector resolution and efficiency, we use an efficiency-weighted  $\mathcal{F}$  convolved with a resolution function based on MC simulation in the fits. Inspecting the Dalitz plots shown in the Supplemental Material [73], we add shapes from PHSP MC simulation for the fits at  $\sqrt{s} \leq 4.70$  GeV as no intermediate state is evident. For the fits at  $\sqrt{s} > 4.70$  GeV, we add shapes for  $f$  states according to partial wave analysis [73]. Both shapes are derived from a kernel estimation [76] thus no free parameters are introduced in these shapes. The  $m_0$  and  $\Gamma_0$  of the  $Z_{cs}$  as well as the yields of all components are free. The fit results are shown in Fig. 3, with a small excess of  $Z_{cs}$  over other components. The fitted mass and width are  $4.044 \pm 0.006 \text{ GeV}/c^2$  and  $0.036 \pm 0.016 \text{ GeV}$ , respectively. The uncertainties are statistical only. By comparing fits with and without  $Z_{cs}$  components, which gives  $|\Delta(-2 \ln L)| = 23.8$  and  $\Delta \text{ndf} = 12$ , the statistical significance is determined to be  $2.3\sigma$ . Despite no significant  $Z_{cs}$ , upper limits on the production of the  $Z_{cs}(3985)^+$  and  $Z_{cs}(4000)^+$  are of interest to further understand their properties and search for them in future experiments. The upper limits at 90% confidence level (CL) for  $\sigma^{\text{Bom}}[e^+e^- \rightarrow K^- Z_{cs}(3985)^+] \times \mathcal{B}[Z_{cs}(3985)^+ \rightarrow K^+ J/\psi]$  are  $\mathcal{O}(1)$  pb and the upper limits for  $\sigma^{\text{Bom}}[e^+e^- \rightarrow K^- Z_{cs}(4000)^+] \times \mathcal{B}[Z_{cs}(4000)^+ \rightarrow K^+ J/\psi]$  are  $\mathcal{O}(3)$  pb. These upper limits include systematic uncertainties and are summarized in the Supplemental Material [73].

The upper limit on the ratio of branching fractions

$$R_B \equiv \frac{\mathcal{B}[Z_{cs}(3985)^+ \rightarrow K^+ J/\psi]}{\mathcal{B}[Z_{cs}(3985)^+ \rightarrow (\bar{D}^0 D_s^{*+} + \bar{D}^{*0} D_s^+)]} \quad (4)$$

is determined at  $\sqrt{s} = 4.68$  GeV since the  $Z_{cs}(3985)^+$  is the most significant at this c.m. energy. We extract the distribution of  $\sigma^{\text{Bom}} \times \mathcal{B}[Z_{cs}(3985)^+ \rightarrow K^+ J/\psi]$  from the smeared likelihood values as shown in Ref. [73], which is denoted as  $u\{\sigma^{\text{Bom}} \times \mathcal{B}[Z_{cs}(3985)^+ \rightarrow K^+ J/\psi]\}$ . We model the distribution of  $\sigma^{\text{Bom}} \times \mathcal{B}[Z_{cs}(3985)^+ \rightarrow (\bar{D}^0 D_s^{*+} + \bar{D}^{*0} D_s^+)]$  by a Gaussian function  $G\{\sigma^{\text{Bom}} \times \mathcal{B}[Z_{cs}(3985)^+ \rightarrow (\bar{D}^0 D_s^{*+} + \bar{D}^{*0} D_s^+)]\}$  of which the mean and width are set to the reported center value and uncertainty [37]. Then the upper limit  $R_B^{\text{UL}}$  of  $R_B$  at the 90% CL is derived from the convolution of these two distributions,  $g(R_B) \equiv \int u(zR_B) \cdot G(z) dz$ , and  $R_B^{\text{UL}}$  is determined to be 0.03 by  $\int_0^{R_B^{\text{UL}}} g(R_B) dR_B / \int_0^{+\infty} g(R_B) dR_B = 0.9$ .

In summary, the cross sections of  $e^+ e^- \rightarrow K^+ K^- J/\psi$  at c.m. energies between 4.61 and 4.95 GeV are measured. Fitting the cross sections from this Letter and Ref. [60] with three resonances [ $Y(4230)$ ,  $Y(4500)$ , and  $Y(4710)$ ], we obtain the mass and width of the  $Y(4710)$  to be  $M(4710) = 4708_{-15}^{+17} \pm 21$  MeV/ $c^2$  and  $\Gamma(4710) = 126_{-23}^{+27} \pm 30$  MeV, where the first uncertainties are statistical and the second systematic. A new resonance structure  $Y(4710)$  is observed with a statistical significance over  $5\sigma$ , which is one of the heaviest vector charmonium-like states. Our new results confirm that the structure previously reported as evidence in Ref. [28] is indeed the  $Y(4710)$  resonance. Interestingly, the BESIII has recently reported a structure observed in  $D_s^{*+} D_s^{*-}$  system above 4.7 GeV [77], which could potentially correspond to the  $Y(4710)$  or be closely related. This raises the possibility that these  $Y$  states likely possess a significant strange component. In Ref. [78], it was suggested that the  $Y(4710)$  contains a significant  $1^{--}$  charmonium hybrid ( $c\bar{c}g$ ) component. It is also possible that the  $Y(4710)$  is an excited charmonium state predicted by the potential models [29–36], or arises from charmonia mixing [79–81]. This observation brings new insights into the charmonium(like) states above the open-charm threshold.

We also investigate the  $Z_{cs}$  states in the  $KJ/\psi$  system, but no significant structure is observed. Thus the upper limits of the product of the Born cross sections  $\sigma^B[e^+ e^- \rightarrow K^- Z_{cs}(3985)^+ / K^- Z_{cs}(4000)^+]$  and the branching fraction of  $Z_{cs}(3985)^+ / Z_{cs}(4000)^+ \rightarrow K^+ J/\psi$  are determined at 90% CL. The ratio of branching fractions  $\{\mathcal{B}[Z_{cs}(3985)^+ \rightarrow K^+ J/\psi] / \mathcal{B}[Z_{cs}(3985)^+ \rightarrow (\bar{D}^0 D_s^{*+} + \bar{D}^{*0} D_s^+)]\}$  is determined to be less than 0.03 at 90% CL. The suppression of the decay  $Z_{cs}(3985)^+ \rightarrow K^+ J/\psi$  disfavors the QCD sum rule calculation under the molecular state assumption in Refs. [54,57]. It supports the  $Z_{cs}(3985)^+$  and  $Z_{cs}(4000)^+$  as two different states [51]. Our measurements provide important inputs for the understanding of the nature of the  $Z_{cs}(3985)$ . To further improve studies of the potential  $Z_{cs}$  state, more statistics are necessary to conduct a partial wave analysis.

The BESIII Collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key R&D Program of China under Contracts No. 2020YFA0406300, No. 2020YFA0406400; National Natural Science Foundation of China (NSFC) under Contracts No. 11635010, No. 11735014, No. 11835012, No. 11935015, No. 11935016, No. 11935018, No. 11961141012, No. 11975141, No. 12022510, No. 12025502, No. 12035009, No. 12035013, No. 12061131003, No. 12192260, No. 12192261, No. 12192262, No. 12192263, No. 12192264, No. 12192265, No. 12221005, No. 12225509, No. 12235017; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); Joint Large-Scale Scientific Facility Funds of the NSFC and CAS under Contract No. U1832207; CAS Key Research Program of Frontier Sciences under Contracts No. QYZDJ-SSW-SLH003, No. QYZDJ-SSW-SLH040; 100 Talents Program of CAS; Project No. ZR2022JQ02 supported by Shandong Provincial Natural Science Foundation; The Institute of Nuclear and Particle Physics (INPAC) and Shanghai Key Laboratory for Particle Physics and Cosmology; ERC under Contract No. 758462; European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie Grant Agreement under Contract No. 894790; German Research Foundation DFG under Contracts No. 455635585, Collaborative Research Center CRC 1044, FOR5327, GRK 2149; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Research Foundation of Korea under Contract No. NRF-2022R1A2C1092335; National Science and Technology fund of Mongolia; National Science Research and Innovation Fund (NSRF) via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation of Thailand under Contract No. B16F640076; Polish National Science Centre under Contract No. 2019/35/O/ST2/02907; The Swedish Research Council; U.S. Department of Energy under Contract No. DE-FG02-05ER41374.

- 
- [1] W. Kwong, J. L. Rosner, and C. Quigg, *Annu. Rev. Nucl. Part. Sci.* **37**, 325 (1987).
  - [2] C. Z. Yuan, *Natl. Sci. Rev.* **8**, nwab182 (2021).
  - [3] S. K. Choi *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **91**, 262001 (2003).
  - [4] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **95**, 142001 (2005).
  - [5] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **110**, 252001 (2013).
  - [6] Z. Q. Liu *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **110**, 252002 (2013); **111**, 019901(E) (2013).

- [7] E. S. Swanson, *Phys. Rep.* **429**, 243 (2006).
- [8] H. X. Chen, W. Chen, X. Liu, and S. L. Zhu, *Phys. Rep.* **639**, 1 (2016).
- [9] F. K. Guo, C. Hanhart, U. G. Meißner, Q. Wang, Q. Zhao, and B. S. Zou, *Rev. Mod. Phys.* **90**, 015004 (2018); **94**, 029901(E) (2022).
- [10] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C. P. Shen, C. E. Thomas, A. Vairo, and C. Z. Yuan, *Phys. Rep.* **873**, 1 (2020).
- [11] Q. He *et al.* (CLEO Collaboration), *Phys. Rev. D* **74**, 091104 (2006).
- [12] C. Z. Yuan *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **99**, 182004 (2007).
- [13] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **118**, 092001 (2017).
- [14] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **102**, 012009 (2020).
- [15] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **106**, 072001 (2022).
- [16] X. L. Wang *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **99**, 142002 (2007).
- [17] X. L. Wang *et al.* (Belle Collaboration), *Phys. Rev. D* **91**, 112007 (2015).
- [18] J. P. Lees *et al.* (BABAR Collaboration), *Phys. Rev. D* **89**, 111103 (2014).
- [19] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **96**, 032004 (2017); **99**, 019903(E) (2019).
- [20] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **104**, 052012 (2021).
- [21] G. Pakhlova *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **101**, 172001 (2008).
- [22] S. Jia *et al.* (Belle Collaboration), *Phys. Rev. D* **100**, 111103 (2019).
- [23] S. Jia *et al.* (Belle Collaboration), *Phys. Rev. D* **101**, 091101 (2020).
- [24] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **106**, 052012 (2022).
- [25] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **129**, 102003 (2022).
- [26] M. Ablikim *et al.* (BESIII Collaboration), *J. High Energy Phys.* **01** (2023) 132.
- [27] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **130**, 121901 (2023).
- [28] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **107**, 092005 (2023).
- [29] G. J. Ding, J. J. Zhu, and M. L. Yan, *Phys. Rev. D* **77**, 014033 (2008).
- [30] N. Akbar, *J. Korean Phys. Soc.* **77**, 17 (2020).
- [31] R. Chaturvedi and A. K. Rai, *Int. J. Theor. Phys.* **59**, 3508 (2020).
- [32] M. A. Sultan, N. Akbar, B. Masud, and F. Akram, *Phys. Rev. D* **90**, 054001 (2014).
- [33] W. J. Deng, H. Liu, L. C. Gui, and X. H. Zhong, *Phys. Rev. D* **95**, 034026 (2017).
- [34] L. C. Gui, L. S. Lu, Q. F. Lü, X. H. Zhong, and Q. Zhao, *Phys. Rev. D* **98**, 016010 (2018).
- [35] S. Kanwal, F. Akram, B. Masud, and E. S. Swanson, *Eur. Phys. J. A* **58**, 219 (2022).
- [36] A. Bhaghyesh, *Adv. High Energy Phys.* **2021**, 9991152 (2021).
- [37] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **126**, 102001 (2021).
- [38] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **129**, 112003 (2022).
- [39] Z. Yang, X. Cao, F. K. Guo, J. Nieves, and M. P. Valderrama, *Phys. Rev. D* **103**, 074029 (2021).
- [40] L. Meng, B. Wang, and S. L. Zhu, *Phys. Rev. D* **102**, 111502(R) (2020).
- [41] L. Maiani, A. D. Polosa, and V. Riquer, *Sci. Bull.* **66**, 1616 (2021).
- [42] H. X. Chen, W. Chen, X. Liu, Y. R. Liu, and S. L. Zhu, *Rep. Prog. Phys.* **86**, 026201 (2023).
- [43] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **47**, 033001 (2023).
- [44] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **127**, 082001 (2021).
- [45] P. P. Shi, F. Huang, and W. L. Wang, *Phys. Rev. D* **103**, 094038 (2021).
- [46] J. F. Giron, R. F. Lebed, and S. R. Martinez, *Phys. Rev. D* **104**, 054001 (2021).
- [47] Q. N. Wang, W. Chen, and H. X. Chen, *Chin. Phys. C* **45**, 093102 (2021).
- [48] J. B. Wang, G. Li, C. S. An, C. R. Deng, and J. J. Xie, *Eur. Phys. J. C* **82**, 721 (2022).
- [49] M. Karliner and J. L. Rosner, *Phys. Rev. D* **104**, 034033 (2021).
- [50] S. Han and L. Y. Xiao, *Phys. Rev. D* **105**, 054008 (2022).
- [51] L. Meng, B. Wang, G. J. Wang, and S. L. Zhu, *Sci. Bull.* **66**, 2065 (2021).
- [52] Z. G. Wang, *Chin. Phys. C* **46**, 103106 (2022).
- [53] D. Y. Chen, X. Liu, and T. Matsuki, *Phys. Rev. Lett.* **110**, 232001 (2013).
- [54] J. M. Dias, X. Liu, and M. Nielsen, *Phys. Rev. D* **88**, 096014 (2013).
- [55] M. B. Voloshin, *Phys. Lett. B* **798**, 135022 (2019).
- [56] X. Cao and Z. Yang, *Eur. Phys. J. C* **82**, 161 (2022).
- [57] B. D. Wan and C. F. Qiao, *Nucl. Phys. B* **968**, 115450 (2021).
- [58] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **44**, 040001 (2020).
- [59] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **46**, 113003 (2022).
- [60] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **46**, 111002 (2022).
- [61] C. Z. Yuan *et al.* (Belle Collaboration), *Phys. Rev. D* **77**, 011105 (2008).
- [62] C. P. Shen *et al.* (Belle Collaboration), *Phys. Rev. D* **89**, 072015 (2014).
- [63] M. Ablikim *et al.* (BESIII Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 345 (2010).
- [64] C. Yu *et al.*, in *Proceedings of the 7th International Particle Accelerator Conference* (2016), p. TUYA01, <http://jacow.org/ipac2016/papers/tuya01.pdf>.
- [65] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [66] K. X. Huang, Z. J. Li, Z. Qian, J. Zhu, H. Y. Li, Y. M. Zhang, S. S. Sun, and Z. Y. You, *Nucl. Sci. Tech.* **33**, 142 (2022).
- [67] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [68] R. G. Ping, *Chin. Phys. C* **32**, 599 (2008).

- [69] R. L. Workman *et al.* (Particle Data Group), *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [70] W. A. Rolke, A. M. Lopez, and J. Conrad, *Nucl. Instrum. Methods Phys. Res., Sect. A* **551**, 493 (2005).
- [71] S. Actis *et al.* (Working Group on Radiative Corrections, Monte Carlo Generators for Low Energies Collaboration), *Eur. Phys. J. C* **66**, 585 (2010).
- [72] W. Sun, T. Liu, M. Jing, L. Wang, B. Zhong, and W. Song, *Front. Phys.* **16**, 64501 (2021).
- [73] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.131.211902> for the detailed information about the Born cross section,  $Z_{cs}$  research and partial wave analysis on the  $e^+e^- \rightarrow K^+K^-J/\psi$  process.
- [74] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **87**, 012002 (2013).
- [75] E. V. Abakumova *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **659**, 21 (2011).
- [76] K. S. Cranmer, *Comput. Phys. Commun.* **136**, 198 (2001).
- [77] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **131**, 151903 (2023).
- [78] N. Brambilla, W. K. Lai, A. Mohapatra, and A. Vairo, *Phys. Rev. D* **107**, 054034 (2023).
- [79] J. Z. Wang, R. Q. Qian, X. Liu, and T. Matsuki, *Phys. Rev. D* **101**, 034001 (2020).
- [80] J. Z. Wang, D. Y. Chen, X. Liu, and T. Matsuki, *Phys. Rev. D* **99**, 114003 (2019).
- [81] J. Z. Wang and X. Liu, *Phys. Rev. D* **107**, 054016 (2023).

M. Ablikim,<sup>1</sup> M. N. Achasov,<sup>4b</sup> P. Adlarson,<sup>75</sup> O. Afedulidis,<sup>3</sup> X. C. Ai,<sup>80</sup> R. Aliberti,<sup>35</sup> A. Amoroso,<sup>74a,74c</sup> Q. An,<sup>71,58</sup> Y. Bai,<sup>57</sup> O. Bakina,<sup>36</sup> I. Balossino,<sup>29a</sup> Y. Ban,<sup>46,g</sup> H.-R. Bao,<sup>63</sup> V. Batozskaya,<sup>1,44</sup> K. Begzsuren,<sup>32</sup> N. Berger,<sup>35</sup> M. Berlowski,<sup>44</sup> M. Bertani,<sup>28a</sup> D. Bettoni,<sup>29a</sup> F. Bianchi,<sup>74a,74c</sup> E. Bianco,<sup>74a,74c</sup> A. Bortone,<sup>74a,74c</sup> I. Boyko,<sup>36</sup> R. A. Briere,<sup>5</sup> A. Brueggemann,<sup>68</sup> H. Cai,<sup>76</sup> X. Cai,<sup>1,58</sup> A. Calcaterra,<sup>28a</sup> G. F. Cao,<sup>1,63</sup> N. Cao,<sup>1,63</sup> S. A. Cetin,<sup>62a</sup> J. F. Chang,<sup>1,58</sup> W. L. Chang,<sup>1,63</sup> G. R. Che,<sup>43</sup> G. Chelkov,<sup>36,a</sup> C. Chen,<sup>43</sup> C. H. Chen,<sup>9</sup> Chao Chen,<sup>55</sup> G. Chen,<sup>1</sup> H. S. Chen,<sup>1,63</sup> M. L. Chen,<sup>1,58,63</sup> S. J. Chen,<sup>42</sup> S. L. Chen,<sup>45</sup> S. M. Chen,<sup>61</sup> T. Chen,<sup>1,63</sup> X. R. Chen,<sup>31,63</sup> X. T. Chen,<sup>1,63</sup> Y. B. Chen,<sup>1,58</sup> Y. Q. Chen,<sup>34</sup> Z. J. Chen,<sup>25,h</sup> Z. Y. Chen,<sup>1,63</sup> S. K. Choi,<sup>10</sup> X. Chu,<sup>43</sup> G. Cibinetto,<sup>29a</sup> F. Cossio,<sup>74c</sup> J. J. Cui,<sup>50</sup> H. L. Dai,<sup>1,58</sup> J. P. Dai,<sup>78</sup> A. Dbeyssi,<sup>18</sup> R. E. de Boer,<sup>3</sup> D. Dedovich,<sup>36</sup> C. Q. Deng,<sup>72</sup> Z. Y. Deng,<sup>1</sup> A. Denig,<sup>35</sup> I. Denysenko,<sup>36</sup> M. Destefanis,<sup>74a,74c</sup> F. De Mori,<sup>74a,74c</sup> B. Ding,<sup>66,1</sup> X. X. Ding,<sup>46,g</sup> Y. Ding,<sup>34</sup> Y. Ding,<sup>40</sup> J. Dong,<sup>1,58</sup> L. Y. Dong,<sup>1,63</sup> M. Y. Dong,<sup>1,58,63</sup> X. Dong,<sup>76</sup> M. C. Du,<sup>1</sup> S. X. Du,<sup>80</sup> Z. H. Duan,<sup>42</sup> P. Egorov,<sup>36,a</sup> Y. H. Fan,<sup>45</sup> J. Fang,<sup>1,58</sup> J. Fang,<sup>59</sup> S. S. Fang,<sup>1,63</sup> W. X. Fang,<sup>1</sup> Y. Fang,<sup>1</sup> Y. Q. Fang,<sup>1,58</sup> R. Farinelli,<sup>29a</sup> L. Fava,<sup>74b,74c</sup> F. Feldbauer,<sup>3</sup> G. Felici,<sup>28a</sup> C. Q. Feng,<sup>71,58</sup> J. H. Feng,<sup>59</sup> Y. T. Feng,<sup>71,58</sup> K. Fischer,<sup>69</sup> M. Fritsch,<sup>3</sup> C. D. Fu,<sup>1</sup> J. L. Fu,<sup>63</sup> Y. W. Fu,<sup>1</sup> H. Gao,<sup>63</sup> Y. N. Gao,<sup>46,g</sup> Yang Gao,<sup>71,58</sup> S. Garbolino,<sup>74c</sup> I. Garzia,<sup>29a,29b</sup> P. T. Ge,<sup>76</sup> Z. W. Ge,<sup>42</sup> C. Geng,<sup>59</sup> E. M. Gersabeck,<sup>67</sup> A. Gilman,<sup>69</sup> K. Goetzen,<sup>13</sup> L. Gong,<sup>40</sup> W. X. Gong,<sup>1,58</sup> W. Gradl,<sup>35</sup> S. Gramigna,<sup>29a,29b</sup> M. Greco,<sup>74a,74c</sup> M. H. Gu,<sup>1,58</sup> Y. T. Gu,<sup>15</sup> C. Y. Guan,<sup>1,63</sup> Z. L. Guan,<sup>22</sup> A. Q. Guo,<sup>31,63</sup> L. B. Guo,<sup>41</sup> M. J. Guo,<sup>50</sup> R. P. Guo,<sup>49</sup> Y. P. Guo,<sup>12,f</sup> A. Guskov,<sup>36,a</sup> J. Gutierrez,<sup>27</sup> K. L. Han,<sup>63</sup> T. T. Han,<sup>1</sup> X. Q. Hao,<sup>19</sup> F. A. Harris,<sup>65</sup> K. K. He,<sup>55</sup> K. L. He,<sup>1,63</sup> F. H. Heinsius,<sup>3</sup> C. H. Heinz,<sup>35</sup> Y. K. Heng,<sup>1,58,63</sup> C. Herold,<sup>60</sup> T. Holtmann,<sup>3</sup> P. C. Hong,<sup>12,f</sup> G. Y. Hou,<sup>1,63</sup> X. T. Hou,<sup>1,63</sup> Y. R. Hou,<sup>63</sup> Z. L. Hou,<sup>1</sup> B. Y. Hu,<sup>59</sup> H. M. Hu,<sup>1,63</sup> J. F. Hu,<sup>56,i</sup> T. Hu,<sup>1,58,63</sup> Y. Hu,<sup>1</sup> G. S. Huang,<sup>71,58</sup> K. X. Huang,<sup>59</sup> L. Q. Huang,<sup>31,63</sup> X. T. Huang,<sup>50</sup> Y. P. Huang,<sup>1</sup> T. Hussain,<sup>73</sup> F. Hölzken,<sup>3</sup> N. Hüsken,<sup>27,35</sup> N. in der Wiesche,<sup>68</sup> M. Irshad,<sup>71,58</sup> J. Jackson,<sup>27</sup> S. Janchiv,<sup>32</sup> J. H. Jeong,<sup>10</sup> Q. Ji,<sup>1</sup> Q. P. Ji,<sup>19</sup> W. Ji,<sup>1,63</sup> X. B. Ji,<sup>1,63</sup> X. L. Ji,<sup>1,58</sup> Y. Y. Ji,<sup>50</sup> X. Q. Jia,<sup>50</sup> Z. K. Jia,<sup>71,58</sup> D. Jiang,<sup>1,63</sup> H. B. Jiang,<sup>76</sup> P. C. Jiang,<sup>46,g</sup> S. S. Jiang,<sup>39</sup> T. J. Jiang,<sup>16</sup> X. S. Jiang,<sup>1,58,63</sup> Y. Jiang,<sup>63</sup> J. B. Jiao,<sup>50</sup> J. K. Jiao,<sup>34</sup> Z. Jiao,<sup>23</sup> S. Jin,<sup>42</sup> Y. Jin,<sup>66</sup> M. Q. Jing,<sup>1,63</sup> X. M. Jing,<sup>63</sup> T. Johansson,<sup>75</sup> S. Kabana,<sup>33</sup> N. Kalantar-Nayestanaki,<sup>64</sup> X. L. Kang,<sup>9</sup> X. S. Kang,<sup>40</sup> M. Kavatsyuk,<sup>64</sup> B. C. Ke,<sup>80</sup> V. Khachatryan,<sup>27</sup> A. Khoukaz,<sup>68</sup> R. Kiuchi,<sup>1</sup> O. B. Kolcu,<sup>62a</sup> B. Kopf,<sup>3</sup> M. Kuessner,<sup>3</sup> X. Kui,<sup>1,63</sup> A. Kupsc,<sup>44,75</sup> W. Kühn,<sup>37</sup> J. J. Lane,<sup>67</sup> P. Larin,<sup>18</sup> L. Lavezzi,<sup>74a,74c</sup> T. T. Lei,<sup>71,58</sup> Z. H. Lei,<sup>71,58</sup> H. Leithoff,<sup>35</sup> M. Lellmann,<sup>35</sup> T. Lenz,<sup>35</sup> C. Li,<sup>47</sup> C. Li,<sup>43</sup> C. H. Li,<sup>39</sup> Cheng Li,<sup>71,58</sup> D. M. Li,<sup>80</sup> F. Li,<sup>1,58</sup> G. Li,<sup>1</sup> H. Li,<sup>71,58</sup> H. B. Li,<sup>1,63</sup> H. J. Li,<sup>19</sup> H. N. Li,<sup>56,i</sup> Hui Li,<sup>43</sup> J. R. Li,<sup>61</sup> J. S. Li,<sup>59</sup> Ke Li,<sup>1</sup> L. J. Li,<sup>1,63</sup> L. K. Li,<sup>1</sup> Lei Li,<sup>48</sup> M. H. Li,<sup>43</sup> P. R. Li,<sup>38,k</sup> Q. M. Li,<sup>1,63</sup> Q. X. Li,<sup>50</sup> R. Li,<sup>17,31</sup> S. X. Li,<sup>12</sup> T. Li,<sup>50</sup> W. D. Li,<sup>1,63</sup> W. G. Li,<sup>1</sup> X. Li,<sup>1,63</sup> X. H. Li,<sup>71,58</sup> X. L. Li,<sup>50</sup> Xiaoyu Li,<sup>1,63</sup> Y. G. Li,<sup>46,g</sup> Z. J. Li,<sup>59</sup> Z. X. Li,<sup>15</sup> C. Liang,<sup>42</sup> H. Liang,<sup>71,58</sup> H. Liang,<sup>1,63</sup> Y. F. Liang,<sup>54</sup> Y. T. Liang,<sup>31,63</sup> G. R. Liao,<sup>14</sup> L. Z. Liao,<sup>50</sup> Y. P. Liao,<sup>1,63</sup> J. Libby,<sup>26</sup> A. Limphirat,<sup>60</sup> D. X. Lin,<sup>31,63</sup> T. Lin,<sup>1</sup> B. J. Liu,<sup>1</sup> B. X. Liu,<sup>76</sup> C. Liu,<sup>34</sup> C. X. Liu,<sup>1</sup> F. H. Liu,<sup>53</sup> Fang Liu,<sup>1</sup> Feng Liu,<sup>6</sup> G. M. Liu,<sup>56,i</sup> H. Liu,<sup>38,j,k</sup> H. B. Liu,<sup>15</sup> H. M. Liu,<sup>1,63</sup> Huanhuan Liu,<sup>1</sup> Huihui Liu,<sup>21</sup> J. B. Liu,<sup>71,58</sup> J. Y. Liu,<sup>1,63</sup> K. Liu,<sup>38,j,k</sup> K. Y. Liu,<sup>40</sup> Ke Liu,<sup>22</sup> L. Liu,<sup>71,58</sup> L. C. Liu,<sup>43</sup> Lu Liu,<sup>43</sup> M. H. Liu,<sup>12,f</sup> P. L. Liu,<sup>1</sup> Q. Liu,<sup>63</sup> S. B. Liu,<sup>71,58</sup> T. Liu,<sup>12,f</sup> W. K. Liu,<sup>43</sup> W. M. Liu,<sup>71,58</sup> X. Liu,<sup>38,j,k</sup> X. Liu,<sup>39</sup> Y. Liu,<sup>38,j,k</sup> Y. Liu,<sup>80</sup> Y. B. Liu,<sup>43</sup> Z. A. Liu,<sup>1,58,63</sup> Z. D. Liu,<sup>9</sup> Z. Q. Liu,<sup>50</sup> X. C. Lou,<sup>1,58,63</sup> F. X. Lu,<sup>59</sup> H. J. Lu,<sup>23</sup> J. G. Lu,<sup>1,58</sup> X. L. Lu,<sup>1</sup> Y. Lu,<sup>7</sup> Y. P. Lu,<sup>1,58</sup> Z. H. Lu,<sup>1,63</sup> C. L. Luo,<sup>41</sup> M. X. Luo,<sup>79</sup> T. Luo,<sup>12,f</sup> X. L. Luo,<sup>1,58</sup> X. R. Lyu,<sup>63</sup> Y. F. Lyu,<sup>43</sup> F. C. Ma,<sup>40</sup> H. Ma,<sup>78</sup>

H. L. Ma,<sup>1</sup> J. L. Ma,<sup>1,63</sup> L. L. Ma,<sup>50</sup> M. M. Ma,<sup>1,63</sup> Q. M. Ma,<sup>1</sup> R. Q. Ma,<sup>1,63</sup> X. T. Ma,<sup>1,63</sup> X. Y. Ma,<sup>1,58</sup> Y. Ma,<sup>46,g</sup> Y. M. Ma,<sup>31</sup>  
 F. E. Maas,<sup>18</sup> M. Maggiora,<sup>74a,74c</sup> S. Malde,<sup>69</sup> A. Mangoni,<sup>28b</sup> Y. J. Mao,<sup>46,g</sup> Z. P. Mao,<sup>1</sup> S. Marcello,<sup>74a,74c</sup> Z. X. Meng,<sup>66</sup>  
 J. G. Messchendorp,<sup>13,64</sup> G. Mezzadri,<sup>29a</sup> H. Miao,<sup>1,63</sup> T. J. Min,<sup>42</sup> R. E. Mitchell,<sup>27</sup> X. H. Mo,<sup>1,58,63</sup> B. Moses,<sup>27</sup>  
 N. Yu. Muchnoi,<sup>4,b</sup> J. Muskalla,<sup>35</sup> Y. Nefedov,<sup>36</sup> F. Nerling,<sup>18,d</sup> I. B. Nikolaev,<sup>4,b</sup> Z. Ning,<sup>1,58</sup> S. Nisar,<sup>11,1</sup> Q. L. Niu,<sup>38,j,k</sup>  
 W. D. Niu,<sup>55</sup> Y. Niu,<sup>50</sup> S. L. Olsen,<sup>63</sup> Q. Ouyang,<sup>1,58,63</sup> S. Pacetti,<sup>28b,28c</sup> X. Pan,<sup>55</sup> Y. Pan,<sup>57</sup> A. Pathak,<sup>34</sup> P. Patteri,<sup>28a</sup>  
 Y. P. Pei,<sup>71,58</sup> M. Pelizaeus,<sup>3</sup> H. P. Peng,<sup>71,58</sup> Y. Y. Peng,<sup>38,j,k</sup> K. Peters,<sup>13,d</sup> J. L. Ping,<sup>41</sup> R. G. Ping,<sup>1,63</sup> S. Plura,<sup>35</sup> V. Prasad,<sup>33</sup>  
 F. Z. Qi,<sup>1</sup> H. Qi,<sup>71,58</sup> H. R. Qi,<sup>61</sup> M. Qi,<sup>42</sup> T. Y. Qi,<sup>12,f</sup> S. Qian,<sup>1,58</sup> W. B. Qian,<sup>63</sup> C. F. Qiao,<sup>63</sup> J. J. Qin,<sup>72</sup> L. Q. Qin,<sup>14</sup>  
 X. S. Qin,<sup>50</sup> Z. H. Qin,<sup>1,58</sup> J. F. Qiu,<sup>1</sup> S. Q. Qu,<sup>61</sup> Z. H. Qu,<sup>72</sup> C. F. Redmer,<sup>35</sup> K. J. Ren,<sup>39</sup> A. Rivetti,<sup>74c</sup> M. Rolo,<sup>74c</sup>  
 G. Rong,<sup>1,63</sup> Ch. Rosner,<sup>18</sup> S. N. Ruan,<sup>43</sup> N. Salone,<sup>44</sup> A. Sarantsev,<sup>36,c</sup> Y. Schelhaas,<sup>35</sup> K. Schoenning,<sup>75</sup> M. Scodreggio,<sup>29a</sup>  
 K. Y. Shan,<sup>12,f</sup> W. Shan,<sup>24</sup> X. Y. Shan,<sup>71,58</sup> J. F. Shangguan,<sup>55</sup> L. G. Shao,<sup>1,63</sup> M. Shao,<sup>71,58</sup> C. P. Shen,<sup>12,f</sup> H. F. Shen,<sup>1,8</sup>  
 W. H. Shen,<sup>63</sup> X. Y. Shen,<sup>1,63</sup> B. A. Shi,<sup>63</sup> H. C. Shi,<sup>71,58</sup> J. L. Shi,<sup>12</sup> J. Y. Shi,<sup>1</sup> Q. Q. Shi,<sup>55</sup> R. S. Shi,<sup>1,63</sup> S. Y. Shi,<sup>72</sup> X. Shi,<sup>1,58</sup>  
 J. J. Song,<sup>19</sup> T. Z. Song,<sup>59</sup> W. M. Song,<sup>34,1</sup> Y. J. Song,<sup>12</sup> Y. X. Song,<sup>46,g,m</sup> S. Sosio,<sup>74a,74c</sup> S. Spataro,<sup>74a,74c</sup> F. Stielor,<sup>35</sup>  
 Y. J. Su,<sup>63</sup> G. B. Sun,<sup>76</sup> G. X. Sun,<sup>1</sup> H. Sun,<sup>63</sup> H. K. Sun,<sup>1</sup> J. F. Sun,<sup>19</sup> K. Sun,<sup>61</sup> L. Sun,<sup>76</sup> S. S. Sun,<sup>1,63</sup> T. Sun,<sup>51,e</sup> W. Y. Sun,<sup>34</sup>  
 Y. Sun,<sup>9</sup> Y. J. Sun,<sup>71,58</sup> Y. Z. Sun,<sup>1</sup> Z. Q. Sun,<sup>1,63</sup> Z. T. Sun,<sup>50</sup> C. J. Tang,<sup>54</sup> G. Y. Tang,<sup>1</sup> J. Tang,<sup>59</sup> Y. A. Tang,<sup>76</sup> L. Y. Tao,<sup>72</sup>  
 Q. T. Tao,<sup>25,h</sup> M. Tat,<sup>69</sup> J. X. Teng,<sup>71,58</sup> V. Thoren,<sup>75</sup> W. H. Tian,<sup>59</sup> Y. Tian,<sup>31,63</sup> Z. F. Tian,<sup>76</sup> I. Uman,<sup>62b</sup> Y. Wan,<sup>55</sup>  
 S. J. Wang,<sup>50</sup> B. Wang,<sup>1</sup> B. L. Wang,<sup>63</sup> Bo Wang,<sup>71,58</sup> D. Y. Wang,<sup>46,g</sup> F. Wang,<sup>72</sup> H. J. Wang,<sup>38,j,k</sup> J. P. Wang,<sup>50</sup> K. Wang,<sup>1,58</sup>  
 L. L. Wang,<sup>1</sup> M. Wang,<sup>50</sup> Meng Wang,<sup>1,63</sup> N. Y. Wang,<sup>63</sup> S. Wang,<sup>38,j,k</sup> S. Wang,<sup>12,f</sup> T. Wang,<sup>12,f</sup> T. J. Wang,<sup>43</sup> W. Wang,<sup>59</sup>  
 W. Wang,<sup>72</sup> W. P. Wang,<sup>71,58</sup> X. Wang,<sup>46,g</sup> X. F. Wang,<sup>38,j,k</sup> X. J. Wang,<sup>39</sup> X. L. Wang,<sup>12,f</sup> X. N. Wang,<sup>1</sup> Y. Wang,<sup>61</sup>  
 Y. D. Wang,<sup>45</sup> Y. F. Wang,<sup>1,58,63</sup> Y. L. Wang,<sup>19</sup> Y. N. Wang,<sup>45</sup> Y. Q. Wang,<sup>1</sup> Yaqian Wang,<sup>17</sup> Yi Wang,<sup>61</sup> Z. Wang,<sup>1,58</sup>  
 Z. L. Wang,<sup>72</sup> Z. Y. Wang,<sup>1,63</sup> Ziyi Wang,<sup>63</sup> D. Wei,<sup>70</sup> D. H. Wei,<sup>14</sup> F. Weidner,<sup>68</sup> S. P. Wen,<sup>1</sup> Y. R. Wen,<sup>39</sup> U. Wiedner,<sup>3</sup>  
 G. Wilkinson,<sup>69</sup> M. Wolke,<sup>75</sup> L. Wollenberg,<sup>3</sup> C. Wu,<sup>39</sup> J. F. Wu,<sup>1,8</sup> L. H. Wu,<sup>1</sup> L. J. Wu,<sup>1,63</sup> X. Wu,<sup>12,f</sup> X. H. Wu,<sup>34</sup> Y. Wu,<sup>71</sup>  
 Y. H. Wu,<sup>55</sup> Y. J. Wu,<sup>31</sup> Z. Wu,<sup>1,58</sup> L. Xia,<sup>71,58</sup> X. M. Xian,<sup>39</sup> B. H. Xiang,<sup>1,63</sup> T. Xiang,<sup>46,g</sup> D. Xiao,<sup>38,j,k</sup> G. Y. Xiao,<sup>42</sup>  
 S. Y. Xiao,<sup>1</sup> Y. L. Xiao,<sup>12,f</sup> Z. J. Xiao,<sup>41</sup> C. Xie,<sup>42</sup> X. H. Xie,<sup>46,g</sup> Y. Xie,<sup>50</sup> Y. G. Xie,<sup>1,58</sup> Y. H. Xie,<sup>6</sup> Z. P. Xie,<sup>71,58</sup>  
 T. Y. Xing,<sup>1,63</sup> C. F. Xu,<sup>1,63</sup> C. J. Xu,<sup>59</sup> G. F. Xu,<sup>1</sup> H. Y. Xu,<sup>66</sup> Q. J. Xu,<sup>16</sup> Q. N. Xu,<sup>30</sup> W. Xu,<sup>1</sup> W. L. Xu,<sup>66</sup> X. P. Xu,<sup>55</sup>  
 Y. C. Xu,<sup>77</sup> Z. P. Xu,<sup>42</sup> Z. S. Xu,<sup>63</sup> F. Yan,<sup>12,f</sup> L. Yan,<sup>12,f</sup> W. B. Yan,<sup>71,58</sup> W. C. Yan,<sup>80</sup> X. Q. Yan,<sup>1</sup> H. J. Yang,<sup>51,e</sup> H. L. Yang,<sup>34</sup>  
 H. X. Yang,<sup>1</sup> Tao Yang,<sup>1</sup> Y. Yang,<sup>12,f</sup> Y. F. Yang,<sup>43</sup> Y. X. Yang,<sup>1,63</sup> Yifan Yang,<sup>1,63</sup> Z. W. Yang,<sup>38,j,k</sup> Z. P. Yao,<sup>50</sup> M. Ye,<sup>1,58</sup>  
 M. H. Ye,<sup>8</sup> J. H. Yin,<sup>1</sup> Z. Y. You,<sup>59</sup> B. X. Yu,<sup>1,58,63</sup> C. X. Yu,<sup>43</sup> G. Yu,<sup>1,63</sup> J. S. Yu,<sup>25,h</sup> T. Yu,<sup>72</sup> X. D. Yu,<sup>46,g</sup> C. Z. Yuan,<sup>1,63</sup>  
 J. Yuan,<sup>34</sup> L. Yuan,<sup>2</sup> S. C. Yuan,<sup>1</sup> Y. Yuan,<sup>1,63</sup> Z. Y. Yuan,<sup>59</sup> C. X. Yue,<sup>39</sup> A. A. Zafar,<sup>73</sup> F. R. Zeng,<sup>50</sup> S. H. Zeng,<sup>72</sup>  
 X. Zeng,<sup>12,f</sup> Y. Zeng,<sup>25,h</sup> Y. J. Zeng,<sup>59</sup> Y. J. Zeng,<sup>1,63</sup> X. Y. Zhai,<sup>34</sup> Y. C. Zhai,<sup>50</sup> Y. H. Zhan,<sup>59</sup> A. Q. Zhang,<sup>1,63</sup> B. L. Zhang,<sup>1,63</sup>  
 B. X. Zhang,<sup>1</sup> D. H. Zhang,<sup>43</sup> G. Y. Zhang,<sup>19</sup> H. Zhang,<sup>71</sup> H. C. Zhang,<sup>1,58,63</sup> H. H. Zhang,<sup>34</sup> H. H. Zhang,<sup>59</sup>  
 H. Q. Zhang,<sup>1,58,63</sup> H. Y. Zhang,<sup>1,58</sup> J. Zhang,<sup>59</sup> J. Zhang,<sup>80</sup> J. J. Zhang,<sup>52</sup> J. L. Zhang,<sup>20</sup> J. Q. Zhang,<sup>41</sup> J. W. Zhang,<sup>1,58,63</sup>  
 J. X. Zhang,<sup>38,j,k</sup> J. Y. Zhang,<sup>1</sup> J. Z. Zhang,<sup>1,63</sup> Jianyu Zhang,<sup>63</sup> L. M. Zhang,<sup>61</sup> Lei Zhang,<sup>42</sup> P. Zhang,<sup>1,63</sup> Q. Y. Zhang,<sup>39,80</sup>  
 Shuihan Zhang,<sup>1,63</sup> Shulei Zhang,<sup>25,h</sup> X. D. Zhang,<sup>45</sup> X. M. Zhang,<sup>1</sup> X. Y. Zhang,<sup>50</sup> Y. Zhang,<sup>72</sup> Y. T. Zhang,<sup>80</sup>  
 Y. H. Zhang,<sup>1,58</sup> Y. M. Zhang,<sup>39</sup> Yan Zhang,<sup>71,58</sup> Yao Zhang,<sup>1</sup> Z. D. Zhang,<sup>1</sup> Z. H. Zhang,<sup>1</sup> Z. L. Zhang,<sup>34</sup> Z. Y. Zhang,<sup>43</sup>  
 Z. Y. Zhang,<sup>76</sup> G. Zhao,<sup>1</sup> J. Y. Zhao,<sup>1,63</sup> J. Z. Zhao,<sup>1,58</sup> Lei Zhao,<sup>71,58</sup> Ling Zhao,<sup>1</sup> M. G. Zhao,<sup>43</sup> R. P. Zhao,<sup>63</sup> S. J. Zhao,<sup>80</sup>  
 Y. B. Zhao,<sup>1,58</sup> Y. X. Zhao,<sup>31,63</sup> Z. G. Zhao,<sup>71,58</sup> A. Zhemchugov,<sup>36,a</sup> B. Zheng,<sup>72</sup> J. P. Zheng,<sup>1,58</sup> W. J. Zheng,<sup>1,63</sup>  
 Y. H. Zheng,<sup>63</sup> B. Zhong,<sup>41</sup> X. Zhong,<sup>59</sup> H. Zhou,<sup>50</sup> J. Y. Zhou,<sup>34</sup> L. P. Zhou,<sup>1,63</sup> X. Zhou,<sup>76</sup> X. K. Zhou,<sup>6</sup> X. R. Zhou,<sup>71,58</sup>  
 X. Y. Zhou,<sup>39</sup> Y. Z. Zhou,<sup>12,f</sup> J. Zhu,<sup>43</sup> K. Zhu,<sup>1</sup> K. J. Zhu,<sup>1,58,63</sup> L. Zhu,<sup>34</sup> L. X. Zhu,<sup>63</sup> S. H. Zhu,<sup>70</sup> S. Q. Zhu,<sup>42</sup> T. J. Zhu,<sup>12,f</sup>  
 W. J. Zhu,<sup>12,f</sup> Y. C. Zhu,<sup>71,58</sup> Z. A. Zhu,<sup>1,63</sup> J. H. Zou,<sup>1</sup> and J. Zu<sup>71,58</sup>

(BESIII Collaboration)

<sup>1</sup>*Institute of High Energy Physics, Beijing 100049, People's Republic of China*

<sup>2</sup>*Beihang University, Beijing 100191, People's Republic of China*

<sup>3</sup>*Bochum Ruhr-University, D-44780 Bochum, Germany*

<sup>4</sup>*Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia*

<sup>5</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

<sup>6</sup>*Central China Normal University, Wuhan 430079, People's Republic of China*

<sup>7</sup>*Central South University, Changsha 410083, People's Republic of China*

<sup>8</sup>*China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China*



- <sup>9</sup>China University of Geosciences, Wuhan 430074, People's Republic of China  
<sup>10</sup>Chung-Ang University, Seoul 06974, Republic of Korea  
<sup>11</sup>COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan  
<sup>12</sup>Fudan University, Shanghai 200433, People's Republic of China  
<sup>13</sup>GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany  
<sup>14</sup>Guangxi Normal University, Guilin 541004, People's Republic of China  
<sup>15</sup>Guangxi University, Nanning 530004, People's Republic of China  
<sup>16</sup>Hangzhou Normal University, Hangzhou 310036, People's Republic of China  
<sup>17</sup>Hebei University, Baoding 071002, People's Republic of China  
<sup>18</sup>Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany  
<sup>19</sup>Henan Normal University, Xinxiang 453007, People's Republic of China  
<sup>20</sup>Henan University, Kaifeng 475004, People's Republic of China  
<sup>21</sup>Henan University of Science and Technology, Luoyang 471003, People's Republic of China  
<sup>22</sup>Henan University of Technology, Zhengzhou 450001, People's Republic of China  
<sup>23</sup>Huangshan College, Huangshan 245000, People's Republic of China  
<sup>24</sup>Hunan Normal University, Changsha 410081, People's Republic of China  
<sup>25</sup>Hunan University, Changsha 410082, People's Republic of China  
<sup>26</sup>Indian Institute of Technology Madras, Chennai 600036, India  
<sup>27</sup>Indiana University, Bloomington, Indiana 47405, USA  
<sup>28a</sup>INFN Laboratori Nazionali di Frascati, INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy  
<sup>28b</sup>INFN Sezione di Perugia, I-06100 Perugia, Italy  
<sup>28c</sup>University of Perugia, I-06100 Perugia, Italy  
<sup>29a</sup>INFN Sezione di Ferrara, INFN Sezione di Ferrara, I-44122 Ferrara, Italy  
<sup>29b</sup>University of Ferrara, I-44122 Ferrara, Italy  
<sup>30</sup>Inner Mongolia University, Hohhot 010021, People's Republic of China  
<sup>31</sup>Institute of Modern Physics, Lanzhou 730000, People's Republic of China  
<sup>32</sup>Institute of Physics and Technology, Peace Avenue 54B, Ulaanbaatar 13330, Mongolia  
<sup>33</sup>Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica 1000000, Chile  
<sup>34</sup>Jilin University, Changchun 130012, People's Republic of China  
<sup>35</sup>Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany  
<sup>36</sup>Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia  
<sup>37</sup>Justus-Liebig-Universität Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany  
<sup>38</sup>Lanzhou University, Lanzhou 730000, People's Republic of China  
<sup>39</sup>Liaoning Normal University, Dalian 116029, People's Republic of China  
<sup>40</sup>Liaoning University, Shenyang 110036, People's Republic of China  
<sup>41</sup>Nanjing Normal University, Nanjing 210023, People's Republic of China  
<sup>42</sup>Nanjing University, Nanjing 210093, People's Republic of China  
<sup>43</sup>Nankai University, Tianjin 300071, People's Republic of China  
<sup>44</sup>National Centre for Nuclear Research, Warsaw 02-093, Poland  
<sup>45</sup>North China Electric Power University, Beijing 102206, People's Republic of China  
<sup>46</sup>Peking University, Beijing 100871, People's Republic of China  
<sup>47</sup>Qufu Normal University, Qufu 273165, People's Republic of China  
<sup>48</sup>Renmin University of China, Beijing 100872, People's Republic of China  
<sup>49</sup>Shandong Normal University, Jinan 250014, People's Republic of China  
<sup>50</sup>Shandong University, Jinan 250100, People's Republic of China  
<sup>51</sup>Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China  
<sup>52</sup>Shanxi Normal University, Linfen 041004, People's Republic of China  
<sup>53</sup>Shanxi University, Taiyuan 030006, People's Republic of China  
<sup>54</sup>Sichuan University, Chengdu 610064, People's Republic of China  
<sup>55</sup>Soochow University, Suzhou 215006, People's Republic of China  
<sup>56</sup>South China Normal University, Guangzhou 510006, People's Republic of China  
<sup>57</sup>Southeast University, Nanjing 211100, People's Republic of China  
<sup>58</sup>State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China  
<sup>59</sup>Sun Yat-Sen University, Guangzhou 510275, People's Republic of China  
<sup>60</sup>Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand  
<sup>61</sup>Tsinghua University, Beijing 100084, People's Republic of China  
<sup>62a</sup>Turkish Accelerator Center Particle Factory Group, Istinye University, 34010 Istanbul, Turkey  
<sup>62b</sup>Near East University, Nicosia, North Cyprus, 99138 Mersin 10, Turkey  
<sup>63</sup>University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China  
<sup>64</sup>University of Groningen, NL-9747 AA Groningen, The Netherlands

<sup>65</sup>*University of Hawaii, Honolulu, Hawaii 96822, USA*

<sup>66</sup>*University of Jinan, Jinan 250022, People's Republic of China*

<sup>67</sup>*University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom*

<sup>68</sup>*University of Muenster, Wilhelm-Klemm-Strasse 9, 48149 Muenster, Germany*

<sup>69</sup>*University of Oxford, Keble Road, Oxford OX13RH, United Kingdom*

<sup>70</sup>*University of Science and Technology Liaoning, Anshan 114051, People's Republic of China*

<sup>71</sup>*University of Science and Technology of China, Hefei 230026, People's Republic of China*

<sup>72</sup>*University of South China, Hengyang 421001, People's Republic of China*

<sup>73</sup>*University of the Punjab, Lahore-54590, Pakistan*

<sup>74a</sup>*University of Turin and INFN, University of Turin, I-10125 Turin, Italy*

<sup>74b</sup>*University of Eastern Piedmont, I-15121 Alessandria, Italy*

<sup>74c</sup>*INFN, I-10125 Turin, Italy*

<sup>75</sup>*Uppsala University, Box 516, SE-75120 Uppsala, Sweden*

<sup>76</sup>*Wuhan University, Wuhan 430072, People's Republic of China*

<sup>77</sup>*Yantai University, Yantai 264005, People's Republic of China*

<sup>78</sup>*Yunnan University, Kunming 650500, People's Republic of China*

<sup>79</sup>*Zhejiang University, Hangzhou 310027, People's Republic of China*

<sup>80</sup>*Zhengzhou University, Zhengzhou 450001, People's Republic of China*

<sup>a</sup>Also at the Moscow Institute of Physics and Technology, Moscow 141700, Russia.

<sup>b</sup>Also at the Novosibirsk State University, Novosibirsk, 630090, Russia.

<sup>c</sup>Also at the NRC "Kurchatov Institute", PNPI, 188300, Gatchina, Russia.

<sup>d</sup>Also at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany.

<sup>e</sup>Also at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People's Republic of China.

<sup>f</sup>Also at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People's Republic of China.

<sup>g</sup>Also at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People's Republic of China.

<sup>h</sup>Also at School of Physics and Electronics, Hunan University, Changsha 410082, China.

<sup>i</sup>Also at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China.

<sup>j</sup>Also at MOE Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, People's Republic of China.

<sup>k</sup>Also at Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, People's Republic of China.

<sup>l</sup>Also at the Department of Mathematical Sciences, IBA, Karachi 75270, Pakistan.

<sup>m</sup>Also at Ecole Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland.