



Branes wrapped on orbifolds and their gravitational blocks

Federico Faedo^{1,2} · Alessio Fontanarossa^{1,2} · Dario Martelli^{1,2}

Received: 23 December 2022 / Revised: 4 March 2023 / Accepted: 1 April 2023
© The Author(s) 2023

Abstract

We construct new supersymmetric $\text{AdS}_2 \times \mathbb{M}_4$ solutions of $D = 6$ gauged supergravity, where \mathbb{M}_4 are certain four-dimensional orbifolds. After uplifting to massive type IIA supergravity these correspond to the near-horizon limit of a system of N D4-branes and N_f D8-branes wrapped on \mathbb{M}_4 . In one class of solutions, $\mathbb{M}_4 = \Sigma_g \times \Sigma$ is a spindle fibered over a smooth Riemann surface of genus $g > 1$, while in another class $\mathbb{M}_4 = \Sigma \times \Sigma$ is a spindle fibered over another spindle. Both classes can be thought of as orbifold generalizations of Hirzebruch surfaces and, in the second case, we describe the solutions in terms of toric geometry. We show that the entropy associated with these solutions is reproduced by extremizing an entropy function obtained by gluing gravitational blocks, using a general recipe for orbifolds that we propose. We also discuss how our prescription can be used to define an off-shell central charge whose extremization reproduces the gravitational central charge of analogous $\text{AdS}_3 \times \mathbb{M}_4$ solutions of $D = 7$ gauged supergravity, arising from wrapping M5-branes on \mathbb{M}_4 .

Keywords AdS/CFT correspondence · Supersymmetric solutions · Orbifolds · Toric geometry

Contents

1	Introduction
2	Solutions in $D = 6$ gauged supergravity

✉ Federico Faedo
federicomichele.faedo@unito.it

Alessio Fontanarossa
alessio.fontanarossa@unito.it

Dario Martelli
dario.martelli@unito.it

¹ Dipartimento di Matematica “Giuseppe Peano”, Università di Torino, Via Carlo Alberto 10, 10123 Turin, Italy

² INFN, Sezione di Torino, Via Pietro Giuria 1, 10125 Turin, Italy

2.1	$\text{AdS}_2 \times \Sigma_g \times \Sigma_2$ solutions
2.2	$\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solutions
2.3	Truncation to $D = 4$ minimal gauged supergravity
3	Uplift to massive type IIA supergravity
3.1	$\text{AdS}_2 \times \Sigma_g \times \Sigma_2$ solutions
3.2	$\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solutions
4	Toric geometry
4.1	$\text{AdS}_2 \times S^2 \times \Sigma_2$ solutions
4.2	$\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solutions
4.3	Toric divisors and their intersection
5	The entropy function
5.1	Twisting data
5.2	The recipe
5.3	Application to the $\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solutions
5.4	Application to the $\text{AdS}_3 \times \Sigma_1 \times \Sigma_2$ solutions
6	Discussion
A	Some solutions to the Diophantine equation
B	Riemann surfaces from spindles
B.1	$\text{AdS}_2 \times \mathbb{M}_4$ solutions
B.2	$\text{AdS}_3 \times \mathbb{M}_4$ solutions
C	More toric data from explicit metrics
C.1	Hirzebruch \mathbb{F}_1
C.2	$\text{AdS}_3 \times S^2 \times \Sigma_2$ solutions
C.3	$\text{AdS}_3 \times \Sigma_1 \times \Sigma_2$ solutions
	References

1 Introduction

For two decades several examples of the AdS/CFT correspondence have been engineered considering setups involving branes wrapped on compact cycles, preserving supersymmetry via a topological twist on the brane world-volumes [44]. On the supergravity side, these give rise to supersymmetric solutions involving a warped product of an anti-de Sitter (AdS) spacetime, together with an internal compact manifold. The corresponding dual field theories are obtained from suitably twisted compactifications of the parent theories on smooth Riemann surfaces embedded holomorphically in Calabi–Yau manifolds, or different higher-dimensional calibrated submanifolds in special holonomy manifolds.

Solutions with an AdS_2 factor are particularly interesting because on general grounds these are expected to correspond to the near-horizon limit of supersymmetric black holes arising in string or M-theory. In this case, the AdS/CFT correspondence gives access to the microscopic entropy of these black holes, obtained via counting the microstates of the dual supersymmetric quantum mechanics, which becomes particularly useful when the latter arises from a twisted compactification of a higher-dimensional gauge theory. The paradigmatic example of this successful strategy is that of the four-dimensional static black holes of [13], for which a microscopic account of the entropy has been obtained in [9].

The construction presented in [30] changed somewhat the rules of the game, showing that one can obtain examples of holographic dual pairs with genuine novel features by considering D3-branes wrapped on the orbifold $\Sigma = \mathbb{WCP}^1_{[n_-, n_+]}$, where n_-, n_+ are co-prime integers, sometimes referred to as *spindle*. Perhaps the most striking

aspect of this construction is that supersymmetry is preserved via a novel mechanism different from the topological twist, that was later called anti-twist. Further examples of various branes wrapping the spindle have been discussed in [4, 10, 18, 22, 23, 27–29, 31–33, 38] and closely related constructions appeared in [5, 6, 19, 21, 39, 50–52]. In general, the only two possible ways to preserve supersymmetry on a spindle are by the anti-twist of [30] or the twist introduced in [32], which is a global version of the standard topological twist [31].

Once it has been established that it makes sense to wrap branes on spaces with orbifold singularities, it is natural to extend the exploration of the landscape of holographic dualities to branes wrapping orbifolds of dimension higher than two. Simple examples of branes wrapped on four-dimensional orbifolds of the type $\mathbb{M}_4 = \Sigma_g \times \Sigma$ were discussed in [10] for M5-branes and in [27, 33] for D4-branes (see also [53]). In this paper, we will focus mainly on D4-branes, extending the results of [27] in various ways. In particular, we will present two classes of new supersymmetric solutions of gauged $D = 6$ supergravity of the type $\text{AdS}_2 \times \mathbb{M}_4$, where \mathbb{M}_4 are four-dimensional orbifolds. After uplifting to massive type IIA supergravity these describe a system of N D4-branes and N_f D8-branes wrapped on \mathbb{M}_4 , and are expected to correspond to the near-horizon limit of novel supersymmetric AdS_6 black holes. Previously, supersymmetric $\text{AdS}_2 \times M_4$ solutions of the same theory, where M_4 is either a negatively curved Kähler–Einstein manifold or the product of two Riemann surfaces $\Sigma_{g_1} \times \Sigma_{g_2}$ (with $g_1, g_2 > 1$), were discussed in [35] and [49], with the latter reference also presenting numerical solutions for supersymmetric AdS_6 black holes with $\Sigma_{g_1} \times \Sigma_{g_2}$ horizons. Below, we will construct two new classes of solutions. In one class, $\mathbb{M}_4 = \Sigma_g \times \Sigma$ is a spindle fibered over a smooth Riemann surface of genus $g > 1$, while in the other class $\mathbb{M}_4 = \Sigma \times \Sigma$ is a spindle fibered over another spindle. Both classes can be thought of as generalizations of Hirzebruch surfaces and analogous $\text{AdS}_3 \times \mathbb{M}_4$ solutions of $D = 7$ gauged supergravity, arising from wrapping M5-branes on \mathbb{M}_4 , appeared in [17].

One of the motivations for this paper was to begin investigating entropy functions whose extremization reproduces the entropy associated with the corresponding supergravity solutions, comprising orbifolds of dimension higher than two. In the two-dimensional case, entropy functions associated with compactifications on the spindle were discussed in [16, 32, 38] and a conjectural extension to more general off-shell free energies was put forward in [27]. In particular, for all cases, precise formulas for the off-shell free energies and for the associated constraints obeyed by the magnetic fluxes and fugacities were given in terms of data at the north and south poles of the spindle. These are the fixed points of the Killing vector generating its $U(1)$ azimuthal symmetry and the contributions to the off-shell free energies arising at these fixed points are referred to as gravitational blocks [37]. Here we will extend the conjecture of [27] presenting an *orbifold entropy function* whose contributions (blocks) arise from fixed points of a $U(1)^2$ action on \mathbb{M}_4 , together with a recipe for determining the appropriate constraints.

The key for formulating this proposal is to provide a toric description of our solutions. Despite the fact that our orbifolds are not symplectic and therefore there are no moment maps with associated polytope, assuming a $U(1)^2$ action, we can still define a set of *toric data* from the co-dimension two loci where one $U(1) \subset U(1)^2$

degenerates. In particular, these can be determined by studying the Killing vectors of the explicit metrics on \mathbb{M}_4 . We can then associate a compact polytope defined using these data as normal vectors to the edges and identify the vertices as fixed points of the torus action, as in symplectic toric geometry. Our proposal for the entropy function reproduces the results of the explicit supergravity solutions presented here, but we formulate it in a way that is in principle applicable to more generic toric orbifolds, including toric manifolds, for which explicit solutions have not yet been found.

Finally, we discuss how our prescription can be extended to describe analogous supersymmetric $\text{AdS}_3 \times \mathbb{M}_4$ solutions of $D = 7$ gauged supergravity arising from wrapping M5-branes on \mathbb{M}_4 . In these cases, one can employ the method of integrating the M5-branes anomaly polynomial on \mathbb{M}_4 to obtain an off-shell central charge of the corresponding $d = 2$, $\mathcal{N} = (0, 2)$ SCFTs and our proposal reproduces the results available in the literature. In particular, when \mathbb{M}_4 is actually a toric manifold, our recipe reduces to the construction of [36], whereas for the specific case of $\mathbb{M}_4 = \Sigma \times \Sigma$ we recover the results of [17].

The rest of the paper is organized as follows. In Sect. 2, we present the new solutions in $D = 6$ and demonstrate their supersymmetry by solving the relevant Killing spinor equations. In Sect. 3, we uplift our solutions to massive type IIA supergravity and calculate their associated geometric entropies. In Sect. 4, we work out the toric data of the relevant solutions. In Sect. 5, we present our conjectural entropy functions and use these to recover the entropies of the above solutions, as well as the central charge associated with the solutions in [17]. Appendix A contains a discussion about the different quantization conditions on the fluxes and their resolution. In Appendix B, we present a special limit in which a Riemann surface can be retrieved from a spindle. In Appendix C, we derive toric data from different metrics, including the solutions of [17].

Note added: as this work was being finalized we became aware that there would be significant overlap with the results of [20], which appeared on the arXiv on the same day.

2 Solutions in $D = 6$ gauged supergravity

In this paper, we discuss solutions of a $D = 6$ gauged supergravity with gauge group $U(1)^2$, comprising two gauge fields A_1, A_2 , a two-form B and two real scalar fields $\vec{\varphi} = (\varphi_1, \varphi_2)$. This model can also be obtained as a sub-sector of an extension of Romans $F(4)$ gauged supergravity [47], coupled to three vector multiplets [2, 26]. The bosonic part of the action reads¹

$$S_{6D} = \frac{1}{16\pi G_{(6)}} \int d^6x \sqrt{-g} \left[R - V - \frac{1}{2} |d\vec{\varphi}|^2 - \frac{1}{2} \sum_{i=1}^2 X_i^{-2} |F_i|^2 - \frac{1}{8} (X_1 X_2)^2 |H|^2 \right. \\ \left. - \frac{m^2}{4} (X_1 X_2)^{-1} |B|^2 - \frac{1}{16} \frac{\varepsilon^{\mu\nu\rho\sigma\tau\lambda}}{\sqrt{-g}} B_{\mu\nu} \left(F_{1\rho\sigma} F_{2\tau\lambda} + \frac{m^2}{12} B_{\rho\sigma} B_{\tau\lambda} \right) \right], \quad (2.1)$$

¹ Here and in what follows we define, for any p -form ω , $|\omega|^2 = \frac{1}{p!} \omega_{\mu_1 \dots \mu_p} \omega^{\mu_1 \dots \mu_p}$.

where $F_i = dA_i$, $H = dB$ and the scalar fields $\vec{\varphi}$ are parameterized as

$$X_i = e^{-\frac{1}{2}\vec{a}_i \cdot \vec{\varphi}} \quad \text{with} \quad \vec{a}_1 = (2^{1/2}, 2^{-1/2}), \quad \vec{a}_2 = (-2^{1/2}, 2^{-1/2}). \quad (2.2)$$

The scalar potential is

$$V = m^2 X_0^2 - 4g^2 X_1 X_2 - 4mg X_0(X_1 + X_2), \quad (2.3)$$

with g the gauge coupling and m the mass parameter, and where for later convenience we defined $X_0 = (X_1 X_2)^{-3/2}$.

A solution to the equations of motion of the model is supersymmetric if and only if it satisfies also the following set of Killing spinor equations [26]:

$$\begin{aligned} \mathcal{D}_\mu \epsilon^A + \frac{1}{8}[g(X_1 + X_2) + mX_0]\Gamma_\mu \epsilon^A \\ + \frac{1}{32}[m(X_1 X_2)^{-1/2} B_{\nu\lambda} \Gamma_7 \delta_B^A \\ + i(X_1^{-1} F_1 + X_2^{-1} F_2)_{\nu\lambda} (\sigma^3)^A_B](\Gamma_\mu^{\nu\lambda} - 6\delta_\mu^{\nu\lambda}) \epsilon^B \\ - \frac{1}{96}(X_1 X_2) H_{\nu\lambda\rho} \Gamma_7 (\Gamma_\mu^{\nu\lambda\rho} - 3\delta_\mu^{\nu\lambda\rho}) \epsilon^A = 0, \end{aligned} \quad (2.4)$$

$$\begin{aligned} \frac{1}{4}(X_1^{-1} \partial_\mu X_1 + X_2^{-1} \partial_\mu X_2) \Gamma^\mu \epsilon^A - \frac{1}{8}[g(X_1 + X_2) - 3mX_0] \epsilon^A \\ + \frac{1}{32}[m(X_1 X_2)^{-1/2} B_{\mu\nu} \Gamma_7 \delta_B^A - i(X_1^{-1} F_1 + X_2^{-1} F_2)_{\mu\nu} (\sigma^3)^A_B] \Gamma^{\mu\nu} \epsilon^B \\ + \frac{1}{96}(X_1 X_2) H_{\mu\nu\lambda} \Gamma_7 \Gamma^{\mu\nu\lambda} \epsilon^A = 0, \end{aligned} \quad (2.5)$$

$$\begin{aligned} \frac{1}{2}(X_1^{-1} \partial_\mu X_1 - X_2^{-1} \partial_\mu X_2) \Gamma^\mu (\sigma^3)^A_B \epsilon^B - g(X_1 - X_2) (\sigma^3)^A_B \epsilon^B \\ - \frac{i}{4}(X_1^{-1} F_1 - X_2^{-1} F_2)_{\mu\nu} \Gamma^{\mu\nu} \epsilon^A = 0, \end{aligned} \quad (2.6)$$

where

$$\mathcal{D}_\mu \epsilon^A \equiv \partial_\mu \epsilon^A + \frac{1}{4} \omega_\mu^{ab} \Gamma_{ab} \epsilon^A - \frac{i}{2} g(A_1 + A_2)_\mu (\sigma^3)^A_B \epsilon^B. \quad (2.7)$$

These follow from setting to zero the supersymmetry variations of the fermionic fields of the theory with three vector multiplets [26], that do not vanish automatically in the sub-truncation that we are considering. Here $(\sigma^3)^A_B$ is the usual third Pauli matrix, $\{\Gamma_a, \Gamma_b\} = 2\eta_{ab}$ and $\Gamma_7 \equiv \Gamma^0 \Gamma^1 \Gamma^2 \Gamma^3 \Gamma^4 \Gamma^5$. The $SU(2)$ indices A, B are raised and lowered as $\epsilon^A = \varepsilon^{AB} \epsilon_B$ and $\epsilon_A = \epsilon^B \varepsilon_{BA}$, where $\varepsilon_{AB} = -\varepsilon_{BA}$ and its inverse matrix ε^{AB} is defined such that $\varepsilon^{AB} \varepsilon_{AC} = \delta_C^B$. The supersymmetry parameter ϵ^A is an eight-component symplectic-Majorana spinor; hence, it satisfies the condition

$$\varepsilon^{AB} \epsilon_B^* = \mathcal{B}_6 \epsilon_A, \quad (2.8)$$

where \mathcal{B}_6 is related to the six-dimensional charge conjugation matrix C_6 by $\mathcal{B}_6 = -i C_6 \Gamma^0$.

2.1 $\text{AdS}_2 \times \Sigma_g \times \mathbb{S}^2$ solutions

The first local solutions to the supergravity action (2.1) that we consider are of the type $\text{AdS}_2 \times \Sigma_g \times \mathbb{S}^2$, which consist of a two-dimensional spindle \mathbb{S}^2 non-trivially fibered over a Riemann surface Σ_g , with genus $g > 1$. When $g = 0$ a solution to the equations of motion still exists, but it is not supersymmetric. Specifically, the backgrounds are described by the following set of fields solving the equations of motion

$$\begin{aligned} ds^2 &= (y^2 h_1 h_2)^{1/4} \left[\frac{1}{4} ds_{\text{AdS}_2}^2 + \frac{1}{2} ds_{\Sigma_g}^2 + \frac{y^2}{F} dy^2 + \frac{F}{h_1 h_2} \left(dz - \frac{1}{2m} \omega_{\Sigma_g} \right)^2 \right], \\ A_i &= -\frac{y^3}{h_i} \left(dz - \frac{1}{2m} \omega_{\Sigma_g} \right), \quad X_i = (y^2 h_1 h_2)^{3/8} h_i^{-1}, \\ B &= \frac{y}{2m} \text{vol}(\text{AdS}_2), \end{aligned} \tag{2.9}$$

where $ds_{\text{AdS}_2}^2$ denotes the unit radius metric on AdS_2 , $ds_{\Sigma_g}^2$ the unit radius metric on Σ_g and ω_{Σ_g} is such that $d\omega_{\Sigma_g} = \text{vol}(\Sigma_g)$. The functions h_i and F are given by

$$h_i(y) = \frac{2g}{3m} y^3 + q_i, \quad F(y) = m^2 h_1 h_2 - y^4, \tag{2.10}$$

with q_1, q_2 two real parameters. A curvature singularity lies at $y = 0$, hence without loss of generality, in what follows we will restrict to $y > 0$.

The first step we take in the analysis of these solutions is to prove that they are supersymmetric by constructing the local form of the Killing spinors solving the Eqs. (2.4)–(2.6). We employ the following orthonormal frame

$$\begin{aligned} e^{\hat{a}} &= \frac{y^{1/4} (h_1 h_2)^{1/8}}{2} \hat{e}^{\hat{a}}, & e^{\check{\alpha}} &= \frac{y^{1/4} (h_1 h_2)^{1/8}}{\sqrt{2}} \check{e}^{\check{\alpha}}, \\ e^4 &= \frac{y^{5/4} (h_1 h_2)^{1/8}}{F^{1/2}} dy, & e^5 &= \frac{y^{1/4} F^{1/2}}{(h_1 h_2)^{3/8}} \left(dz - \frac{1}{2m} \omega_{\Sigma_g} \right), \end{aligned} \tag{2.11}$$

where $\hat{e}^{\hat{a}}, \hat{a} = 0, 1$, is the zweibein on AdS_2 , whose coordinates are denoted as $x^{\hat{\mu}}$, and $\check{e}^{\check{\alpha}}, \check{\alpha} = 2, 3$, is the zweibein on Σ_g , whose coordinates are denoted as $x^{\check{\mu}}$. Equation (2.4) then splits into the following system

$$\begin{aligned}
 \left(\partial_{\hat{\mu}} + \frac{1}{4} \omega_{\hat{\mu}}^{\hat{a}\hat{b}} \Gamma_{\hat{a}\hat{b}} \right) \epsilon^A + \hat{e}_{\hat{\mu}}^{\hat{a}} \left[-\frac{i}{4} \Gamma_{\hat{a}}^{45} (\sigma^3)^A{}_B \epsilon^B + \frac{1}{4} \Gamma_{\hat{a}}^{2345} \epsilon^A \right] &= 0, \\
 \partial_{\check{\mu}} \epsilon^A - \frac{1}{2} \left[(\omega_{\Sigma_{\check{g}}})_{\check{\mu}} + \frac{1}{\sqrt{2}} \Gamma_{\check{\mu}}^{45} \right] (\Gamma^{23} \epsilon^A + i (\sigma^3)^A{}_B \epsilon^B) &= 0, \\
 \partial_y \epsilon^A - \frac{1}{16y} \left[(2 + y\tilde{h}') \epsilon^A - i \frac{4y^2}{F^{1/2}} (4 - y\tilde{h}') \Gamma^5 (\sigma^3)^A{}_B \epsilon^B \right] &= 0, \\
 \partial_z \epsilon^A - i \frac{g}{2} \left(\alpha_1 + \alpha_2 - \frac{2m}{g} \right) (\sigma^3)^A{}_B \epsilon^B &= 0,
 \end{aligned} \tag{2.12}$$

where we defined $\tilde{h} \equiv \log(h_1 h_2)$ and we applied the gauge transformation $A_i \mapsto A_i + \alpha_i dz$. Equations (2.5) and (2.6) yield the same constraint

$$F^{1/2} \Gamma^4 \epsilon^A + m(h_1 h_2)^{1/2} \epsilon^A + i y^2 \Gamma^{45} (\sigma^3)^A{}_B \epsilon^B = 0. \tag{2.13}$$

We note that the equations along y and z in (2.12) and Eq. (2.13) are the same as in the $\text{AdS}_4 \times \Sigma$ system (cf. equations (3.3) and (3.4) of [27]), a fact that will play an important role later on. From the fourth equation in (2.12) we see immediately that setting

$$\alpha_1 + \alpha_2 = \frac{2m}{g} \tag{2.14}$$

leads to Killing spinors independent of z and we will adopt this choice in the remainder of this section.

Before tackling the Killing spinor equations, we note the following fact about the $\text{AdS}_4 \times \Sigma$ solution of [27]: choosing a suitable decomposition of the six-dimensional gamma matrices, it is possible to write the six-dimensional Killing spinors as tensor products of a single AdS_4 spinor and a single spinor on Σ . Specifically, we consider the decomposition

$$\Gamma^{\hat{a}} = \gamma^{\hat{a}} \otimes \rho_*, \quad \Gamma^{4,5} = I_4 \otimes \rho^{1,2}, \tag{2.15}$$

where $\gamma^{\hat{a}}, \hat{a} = 0, \dots, 3$, are the (Lorentzian) gamma matrices in $D = 4$, $\rho^{\hat{i}}, \hat{i} = 1, 2$, are the (Euclidean) gamma matrices in $D = 2$ and $\rho_* = -i \rho^1 \rho^2$ is the related chiral matrix. For $\rho^{\hat{i}}$ we choose the following representation

$$\rho^{\hat{i}} = \sigma^{\hat{i}}, \quad \rho_* = \sigma^3, \tag{2.16}$$

and we take $\mathcal{B}_2^\rho = -\sigma^2$. For consistency, the six-dimensional matrix \mathcal{B}_6 decomposes as

$$\mathcal{B}_6 = (\mathcal{B}_4 \gamma_5) \otimes (\mathcal{B}_2^\rho \rho_*), \tag{2.17}$$

where $\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$ is the four-dimensional chiral matrix. Adopting this decomposition, the symplectic-Majorana condition (2.8) and the Killing spinor equations (3.3)–(3.4) of [27] can be solved to give

$$\epsilon^1 = (\gamma_5\vartheta_{(4)}) \otimes \eta^1, \quad \epsilon^2 = \vartheta_{(4)} \otimes \eta^2, \tag{2.18}$$

where $\vartheta_{(4)} = \vartheta_{(4)}(x^{\tilde{\mu}})$ is a Majorana Killing spinor on AdS_4 and

$$\eta^1 = \xi_\eta y^{1/8} (h_1 h_2)^{-3/16} \begin{pmatrix} f_1^{1/2} \\ -f_2^{1/2} \end{pmatrix}, \quad \eta^2 = -i\xi_\eta^* y^{1/8} (h_1 h_2)^{-3/16} \begin{pmatrix} f_2^{1/2} \\ -f_1^{1/2} \end{pmatrix}. \tag{2.19}$$

Here, ξ_η is a complex constant and we have defined

$$f_1(y) \equiv m(h_1 h_2)^{1/2} + y^2, \quad f_2(y) \equiv m(h_1 h_2)^{1/2} - y^2, \tag{2.20}$$

which satisfy $F(y) = f_1(y) f_2(y)$. Notice that η^1 and η^2 are related by $\eta^2 = i\sigma^1(\eta^1)^*$, which descends from the symplectic-Majorana condition (2.8).

Going back to the $\text{AdS}_2 \times \Sigma_g \times \Sigma_2$ background, in order to solve its Killing spinor equations we adopt the same decomposition of the six-dimensional gamma matrices as in (2.15). Moreover, we decompose the four-dimensional gamma matrices $\gamma^{\hat{a}}$ as

$$\gamma^{\hat{a}} = \beta^{\hat{a}} \otimes \tau_*, \quad \gamma^{2,3} = I_2 \otimes \tau^{1,2}, \tag{2.21}$$

where $\beta^{\hat{a}}$ are the (Lorentzian) gamma matrices in $D = 2$, $\tau^{\hat{i}}$, $\hat{i} = 1, 2$, are the (Euclidean) gamma matrices in $D = 2$ and $\tau_* = -i\tau^1\tau^2$ is the related chiral matrix. Explicitly, we adopt the same representation as in (2.16)

$$\tau^i = \sigma^i, \quad \tau_* = \sigma^3, \tag{2.22}$$

and we take $\mathcal{B}_2^\tau = \sigma^1$. The four-dimensional matrices \mathcal{B}_4 and γ_5 decompose as

$$\mathcal{B}_4 = (\mathcal{B}_2^\beta \beta_*) \otimes \mathcal{B}_2^\tau, \quad \gamma_5 = \beta_* \otimes \tau_*, \tag{2.23}$$

where $\beta_* = -\beta^0\beta^1$ is the (Lorentzian) chiral matrix in $D = 2$. The ansatz for the six-dimensional Killing spinors is

$$\epsilon^A = \vartheta \otimes \chi^A \otimes \eta^A, \tag{2.24}$$

where $\vartheta = \vartheta(x^{\tilde{\mu}})$ is a Majorana Killing spinor on AdS_2 , hence $\hat{\nabla}_{\hat{\mu}}\vartheta = \frac{1}{2}\beta_{\hat{\mu}}\vartheta$ and $\vartheta^* = \mathcal{B}_2^\beta\vartheta$. $\chi^A = \chi^A(x^{\tilde{\mu}})$ are two two-component spinors defined on the Riemann surface Σ_g , while η^1 and η^2 are given in (2.19). Putting all the ingredients together,

the symplectic-Majorana condition (2.8) and the Killing spinor Eqs. (2.12)–(2.13) can be solved to give

$$\chi^1 = \begin{pmatrix} 0 \\ \xi_\chi \end{pmatrix}, \quad \chi^2 = \begin{pmatrix} -\xi_\chi^* \\ 0 \end{pmatrix}, \tag{2.25}$$

where ξ_η is a complex constant. Notice that we can express χ^2 as $\chi^2 = -i \sigma^2 (\chi^1)^*$.

We can now count the number of supersymmetries preserved by our $\text{AdS}_2 \times \Sigma_g \ltimes \mathbb{Z}_2$ solution. ϑ is a Majorana spinor, hence it has two real degrees of freedom, while the tensor product $\chi^A \otimes \eta^A$ is fully determined by the product $\xi_\chi \xi_\eta$, which has two real degrees of freedom. Therefore, there are four real independent Killing spinors, that is one quarter of the number of supersymmetries of the six-dimensional $\mathcal{N} = (1, 1)$ theory; hence, the solution is 1/4-BPS.

We now proceed to the global analysis of the local solution (2.9), for which we set $m = 2g/3$ without loss of generality². Fixed a generic point on the Riemann surface Σ_g , the metric on the spindle \mathbb{Z}_2 reads

$$ds_{\mathbb{Z}_2}^2 = \frac{y^2}{F} dy^2 + \frac{F}{h_1 h_2} dz^2. \tag{2.26}$$

In order to have a well-defined metric and positive scalars X_i we need to take $F > 0$, $h_1 > 0$, $h_2 > 0$ in a closed interval not containing the curvature singularity in $y = 0$, thus without loss of generality we restrict to $y > 0$. Moreover, denoting $[y_N, y_S]$ the range of the coordinate y , \mathbb{Z}_2 is a proper spindle given that [27]

$$\frac{gF'(y_N)}{3y_N^3} \Delta z = \frac{2\pi}{n_-}, \quad \frac{gF'(y_S)}{3y_S^3} \Delta z = -\frac{2\pi}{n_+}, \tag{2.27}$$

where n_\pm are two co-prime integers and Δz is the periodicity of the z coordinate. These two relations ensure that at the poles $y = y_{N,S}$ there are \mathbb{Z}_{n_\mp} orbifold singularities, respectively. An additional constraint comes from the quantization of the magnetic fluxes across \mathbb{Z}_2 , which arises from the requirement that gA_i be well-defined connection one-forms on $U(1)$ bundles over Σ_g . Given any point of Σ_g , we must have

$$t_i = \frac{g}{2\pi} \int_{\mathbb{Z}_2} F_i = \frac{p_i}{n_+ n_-}, \quad p_i \in \mathbb{Z}. \tag{2.28}$$

It can be shown that [27]

$$p_1 + p_2 = n_+ + n_-, \tag{2.29}$$

meaning that the R -symmetry gauge field $A_R \equiv g(A_1 + A_2)$ is a connection one-form on the line bundle $\mathcal{O}(n_+ + n_-)$ over \mathbb{Z}_2 , and therefore the integers p_i can be

² Taking $g > 0$ and $m > 0$, we can apply the rescaling (2.4) of [27], together with $B \mapsto (m/lg)^{-1/2} B$, in order to absorb m in the coupling constant.

conveniently parameterized as

$$p_1 = \frac{n_+ + n_-}{2}(1 + z), \quad p_2 = \frac{n_+ + n_-}{2}(1 - z), \quad (2.30)$$

where z is an appropriate rational number. The value of the sum $p_1 + p_2$ implies that A_R realizes on the bundle the type of twist that was dubbed “twist,” as for the D4 [27] and M5-branes [29] wrapped on a spindle.

All the conditions we presented were studied in [27], to which we refer for a detailed analysis, and it was proven that they can be satisfied if $n_- < n_+$, $p_1 < 0$, $p_2 > 0$, the expressions of q_i and $y_{N,S}$ are determined as in equations (3.39) and (3.31) of [27] and z has periodicity

$$\Delta z = \chi_2 \frac{3\pi(x^2 + 3)(\mu - x)}{8gx^2}. \quad (2.31)$$

Here, χ_2 , the Euler characteristic of Σ_2 , and μ are defined as

$$\chi_2 = \frac{n_+ + n_-}{n_+ n_-}, \quad \mu \equiv \frac{n_+ - n_-}{n_+ + n_-}, \quad (2.32)$$

and x is the only solution of the quartic equation

$$x^4 + (8z^2 - 3 - 9\mu^2)x^2 + 12\mu x - 9\mu^2 = 0 \quad (2.33)$$

lying inside the range $0 < x < 1$.

As we already mentioned, the internal space has the structure of a spindle Σ_2 fibered over a Riemann surface Σ_g . This fibration is well defined in the orbifold sense if the one-form η describing the fibration is globally defined, i.e.,

$$\frac{1}{2\pi} \int_{\Sigma_g} d\eta = t, \quad t \in \mathbb{Z}, \quad (2.34)$$

where

$$\eta \equiv \frac{2\pi}{\Delta z} \left(dz - \frac{3}{4g} \omega_{\Sigma_g} \right). \quad (2.35)$$

This requirement yields a quantization condition relating the spindle data (n_{\pm}, z) and the genus g , namely

$$t = -\frac{8x^2(g-1)}{\chi_2(x^2+3)(\mu-x)} \in \mathbb{Z}. \quad (2.36)$$

We refer to Appendix A for the analysis of this constraint.

The last conditions follow from the quantization of the fluxes F_i through two-cycles in $\Sigma_g \times \Sigma_2$. Their integration through Σ_2 were computed in (2.28), hence they are

already appropriately quantized. Next, we define the two-cycles $S_- \equiv \{y = y_N\}$ and $S_+ \equiv \{y = y_S\}$, corresponding to the two poles of Σ_2 , and compute, e.g.,

$$\begin{aligned} \mathfrak{s}_1^- &= \frac{g}{2\pi} \int_{S_-} F_1 = \frac{3}{2} \frac{y_N^3}{h_1(y_N)} (g - 1) \\ &= t\chi_2 \frac{x^3 - (2 + 2z + 3\mu)x^2 + (3 - 2z)x - 3\mu}{8x^2}, \end{aligned} \tag{2.37}$$

where we substituted t as in (2.36). Similarly to what happened in (2.34), the quantization condition requires (2.37) to be an integer. In Appendix A, we present some examples in which this constraint is satisfied. The quantization through S_+ can be addressed easily since an explicit computation shows that

$$\mathfrak{s}_i^+ = \frac{g}{2\pi} \int_{S_+} F_i = \mathfrak{s}_i^- + t \frac{p_i}{n_+ n_-}, \tag{2.38}$$

therefore the related condition is automatically satisfied due to the fact that n_{\pm} divide t , as explained in Appendix A. We also notice that (2.38) agrees with the homology relation $S_+ - S_- = t \Sigma_2$. The fluxes of F_2 can be obtained flipping the sign of z in the previous formulas and, as a consequence, the quantization of its fluxes automatically holds exchanging p_1 with p_2 .

Focusing on the R -symmetry gauge field, its fluxes read

$$\frac{1}{2\pi} \int_{S_-} F_R = 2(g - 1) - \frac{t}{n_-}, \quad \frac{1}{2\pi} \int_{S_+} F_R = 2(g - 1) + \frac{t}{n_+}, \tag{2.39}$$

and, as expected, they are correctly quantized because n_{\pm} divide t .

2.2 AdS₂ × Σ₁ × Σ₂ solutions

A second class of solutions to the model described by action (2.1) is the AdS₂ × Σ₁ × Σ₂ background, where a two-dimensional spindle Σ₂ is non-trivially fibered over a different two-dimensional spindle Σ₁. Remarkably, there exists a specific limit of the coordinates and the parameters in which the AdS₂ × Σ_g × Σ₂ solutions are retrieved, both their bosonic content and the Killing spinors. All the details and computations are left to Appendix B. Going back to the systems under exam, they read

$$\begin{aligned} ds^2 &= (y^2 h_1 h_2)^{1/4} \left[\frac{x^2}{4} ds_{\text{AdS}_2}^2 + \frac{x^2}{q} dx^2 + \frac{q}{4x^2} d\psi^2 + \frac{y^2}{F} dy^2 \right. \\ &\quad \left. + \frac{F}{h_1 h_2} \left(dz - \frac{1}{2m} \left(1 - \frac{a}{x} \right) d\psi \right)^2 \right], \\ A_i &= -\frac{y^3}{h_i} \left(dz - \frac{1}{2m} \left(1 - \frac{a}{x} \right) d\psi \right), \quad X_i = (y^2 h_1 h_2)^{3/8} h_i^{-1}, \\ B &= \frac{ay}{2m} \text{vol}(\text{AdS}_2), \end{aligned} \tag{2.40}$$

where $ds_{\text{AdS}_2}^2$ denotes the unit radius metric on AdS_2 and a is a real parameter. The functions h_i and F are the same as in (2.10), while q is given by

$$q(x) = x^4 - 4x^2 + 4ax - a^2. \tag{2.41}$$

Notice that formally taking $a = 0$ we retrieve the $\text{AdS}_4 \times \Sigma$ background studied in [27].

In order to construct the local form of the Killing spinors and to demonstrate that this solution is supersymmetric, we must first specialize the Killing spinor Eqs. (2.4)–(2.6) to our system employing the orthonormal frame

$$\begin{aligned} e^{\hat{a}} &= \frac{x y^{1/4} (h_1 h_2)^{1/8}}{2} \hat{e}^{\hat{a}}, & e^2 &= \frac{x y^{1/4} (h_1 h_2)^{1/8}}{q^{1/2}} dx, & e^3 &= \frac{q^{1/2} y^{1/4} (h_1 h_2)^{1/8}}{2x} d\psi, \\ e^4 &= \frac{y^{5/4} (h_1 h_2)^{1/8}}{F^{1/2}} dy, & e^5 &= \frac{y^{1/4} F^{1/2}}{(h_1 h_2)^{3/8}} \left(dz - \frac{1}{2m} \left(1 - \frac{a}{x} \right) d\psi \right), \end{aligned} \tag{2.42}$$

where $\hat{e}^{\hat{a}}$, $\hat{a} = 0, 1$, is the zweibein on AdS_2 , whose coordinates are denoted as $x^{\hat{\mu}}$. Equation (2.4) then splits into the following system

$$\begin{aligned} \left(\partial_{\hat{\mu}} + \frac{1}{4} \omega_{\hat{\mu}}^{\hat{a}\hat{b}} \Gamma_{\hat{a}\hat{b}} \right) \epsilon^A + \hat{e}^{\hat{a}}_{\hat{\mu}} \left[\frac{q^{1/2}}{4x} \Gamma_{\hat{a}}^2 \epsilon^A - i \frac{x}{4} \Gamma_{\hat{a}}^{45} (\sigma^3)^A_B \epsilon^B + \frac{a}{4x} \Gamma_{\hat{a}}^{2345} \epsilon^A \right] &= 0, \\ \partial_x \epsilon^A - \frac{a}{2xq^{1/2}} \Gamma^{345} \epsilon^A - i \frac{x}{2q^{1/2}} \Gamma^{245} (\sigma^3)^A_B \epsilon^B &= 0, \\ \partial_{\psi} \epsilon^A - \frac{i}{2} \left(1 - \frac{a}{x} \right) (\sigma^3)^A_B \epsilon^B + \frac{2q - xq'}{8x^3} \Gamma^{23} \epsilon^A + \frac{aq^{1/2}}{4x^3} \Gamma^{245} \epsilon^A \\ - i \frac{q^{1/2}}{4x} \Gamma^{345} (\sigma^3)^A_B \epsilon^B &= 0, \\ \partial_y \epsilon^A - \frac{1}{16y} \left[(2 + y\tilde{h}') \epsilon^A - i \frac{4y^2}{F^{1/2}} (4 - y\tilde{h}') \Gamma^5 (\sigma^3)^A_B \epsilon^B \right] &= 0, \\ \partial_z \epsilon^A - i \frac{g}{2} \left(\alpha_1 + \alpha_2 - \frac{2m}{g} \right) (\sigma^3)^A_B \epsilon^B &= 0, \end{aligned} \tag{2.43}$$

where $\tilde{h} \equiv \log(h_1 h_2)$ is defined as before and we employed the gauge transformation $A_i \mapsto A_i + \alpha_i dz$. Equations (2.5) and (2.6) yield, again, the same constraint

$$F^{1/2} \Gamma^4 \epsilon^A + m (h_1 h_2)^{1/2} \epsilon^A + i y^2 \Gamma^{45} (\sigma^3)^A_B \epsilon^B = 0. \tag{2.44}$$

Also in this case, the Killing spinor equations along y and z and the last constraint are identical to those obtained in the $\text{AdS}_4 \times \Sigma$ background. Once again we fix the gauge such that $\alpha_1 + \alpha_2 = 2m/g$, in order to have a z -independent Killing spinors.

The explicit construction of the Killing spinors proceeds in a way similar to the $\text{AdS}_2 \times \Sigma_g \times \Sigma_2$ case. First of all, we employ the following decomposition of the

six-dimensional gamma matrices (cf. Eqs. (2.15) and (2.21)):

$$\Gamma^{\hat{a}} = \beta^{\hat{a}} \otimes \tau_* \otimes \rho_*, \quad \Gamma^{2,3} = I_2 \otimes \tau^{1,2} \otimes \rho_*, \quad \Gamma^{4,5} = I_2 \otimes I_2 \otimes \rho^{1,2}, \tag{2.45}$$

which implies,

$$\mathcal{B}_6 = \mathcal{B}_2^\beta \otimes (\mathcal{B}_2^\tau \tau_*) \otimes (\mathcal{B}_2^\rho \rho_*), \tag{2.46}$$

where $\beta^{\hat{a}}$ are the (Lorentzian) gamma matrices in $D = 2$, $\rho^{\hat{i}}$ and $\tau^{\hat{i}}$, $\hat{i} = 1, 2$, are the (Euclidean) gamma matrices in $D = 2$ and ρ_* and τ_* are the related chiral matrices. In the Euclidean sectors, we adopt the representation (2.16)

$$\rho^{\hat{i}} = \tau^{\hat{i}} = \sigma^{\hat{i}}, \quad \rho_* = \tau_* = \sigma^3, \tag{2.47}$$

and we take $\mathcal{B}_2^\rho = -\sigma^2$ and $\mathcal{B}_2^\tau = \sigma^1$. The six-dimensional Killing spinors are assumed to consist of the tensor product of spinors living on AdS_2 , Σ_1 and Σ_2 . Namely,

$$\epsilon^A = \vartheta \otimes \chi^A \otimes \eta^A, \tag{2.48}$$

with $\vartheta = \vartheta(x^{\hat{\mu}})$ Majorana Killing spinor on AdS_2 , $\chi^A = \chi^A(x, \psi)$ two two-component spinors defined on the spindle Σ_1 and η^A given in (2.19). The symplectic-Majorana condition (2.8) and the Killing spinor equations (2.43)–(2.44) are satisfied given that

$$\chi^1 = \xi_\chi x^{-1/2} \begin{pmatrix} Q_1^{1/2} \\ -Q_2^{1/2} \end{pmatrix}, \quad \chi^2 = \xi_\chi^* x^{-1/2} \begin{pmatrix} Q_2^{1/2} \\ Q_1^{1/2} \end{pmatrix}, \tag{2.49}$$

where ξ_χ is a complex constant and

$$Q_1(x) \equiv x^2 - (2x - a), \quad Q_2(x) \equiv x^2 + (2x - a), \tag{2.50}$$

which satisfy $q(x) = Q_1(x) Q_2(x)$. Notice that the relation $\chi^2 = -i\sigma^2(\chi^1)^*$ holds also in this case. The count of the real degrees of freedom is identical to the $\text{AdS}_2 \times \Sigma_{\mathbb{g}} \times \Sigma_2$ case and shows that the solution is 1/4-BPS, since it preserves four real supercharges.

The first part of the global analysis of the $\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solution (2.40) proceeds along the path traced in the previous section. Chosen a generic point on the base spindle Σ_1 , the metric of Σ_2 is the same as in (2.26), therefore are identical also the conditions for it to describe a proper spindle with $\mathbb{Z}_{n_{\mp}}$ orbifold singularities. In particular, Δz must be as in (2.31). Moreover, at a generic point of Σ_1 (different from the north and south poles) the magnetic fluxes are quantized as in (2.28):

$$t_i = \frac{g}{2\pi} \int_{\Sigma_2} F_i = \frac{p_i}{n_+ n_-}, \quad p_i \in \mathbb{Z}, \tag{2.51}$$

with $p_1 + p_2 = n_+ + n_-$.

The base spindle Σ_1 , described by the metric

$$ds_{\Sigma_1}^2 = \frac{x^2}{q} dx^2 + \frac{q}{4x^2} d\psi^2, \tag{2.52}$$

was thoroughly studied in [28], thus we will take advantage of the results presented therein, setting $j = 0$. Σ_1 is indeed a spindle, characterized by the two co-prime integers m_{\pm} , with $m_- < m_+$ ³, if the parameter a and the periodicity of the coordinate ψ are

$$a = \frac{m_+^2 - m_-^2}{m_+^2 + m_-^2}, \quad \Delta\psi = \frac{\sqrt{m_+^2 + m_-^2}}{\sqrt{2}m_+m_-} 2\pi. \tag{2.53}$$

The coordinate x is restricted between the two middle roots of the quartic polynomial $q(x)$, namely $x \in [x_N, x_S]$ with

$$x_N = -1 + \frac{\sqrt{2}m_+}{\sqrt{m_+^2 + m_-^2}}, \quad x_S = 1 - \frac{\sqrt{2}m_-}{\sqrt{m_+^2 + m_-^2}}. \tag{2.54}$$

At the north and south poles of Σ_1 , namely $x = x_{N,S}$, are present $\mathbb{Z}_{m_{\mp}}$ orbifold singularities, respectively. Moreover, defined the vector field

$$A_{4d} = \frac{1}{2} \left(1 - \frac{a}{x}\right) d\psi, \tag{2.55}$$

its magnetic flux through the spindle Σ_1 is

$$\frac{1}{2\pi} \int_{\Sigma_1} F_{4d} = \frac{m_+ - m_-}{2m_+m_-}. \tag{2.56}$$

In the four-dimensional theory, this implies that $2A_{4d}$ is a connection one-form on the line bundle $\mathcal{O}(m_+ - m_-)$ over Σ_1 and, as a consequence, that we have anti-twist.

Similarly to the $\Sigma_g \times \Sigma_2$ system, the fibration of Σ_2 over Σ_1 is well-defined given that

$$\frac{1}{2\pi} \int_{\Sigma_1} d\eta = \frac{t}{m_+m_-}, \quad t \in \mathbb{Z}, \tag{2.57}$$

where, in this case,

$$\eta \equiv \frac{2\pi}{\Delta z} \left(dz - \frac{3}{4g} \left(1 - \frac{a}{x}\right) d\psi \right). \tag{2.58}$$

³ Notice that we exchanged m_- and m_+ with respect to [28].

We derive the following constraint on the quantum numbers n_{\pm}, m_{\pm} and z

$$t = -\frac{4x^2(m_+ - m_-)}{\chi_2(x^2 + 3)(\mu - x)} \in \mathbb{Z}, \tag{2.59}$$

which is solved in Appendix A.

Like we did in the previous section, we study the quantization of the fluxes of F_i , now through two-cycles in $\Sigma_1 \times \Sigma_2$. The magnetic fluxes across Σ_2 were computed in (2.28) and are, thus, already quantized. Again we define the two-cycles $S_- \equiv \{y = y_N\}$ and $S_+ \equiv \{y = y_S\}$, which corresponds to two copies of the base spindle Σ_1 , and compute

$$\begin{aligned} \mathfrak{s}_1^- &= \frac{g}{2\pi} \int_{S_-} F_1 = \frac{3}{4} \frac{y_N^3}{h_1(y_N)} \frac{m_+ - m_-}{m_+ m_-} \\ &= \frac{t\chi_2}{m_+ m_-} \frac{x^3 - (2 + 2z + 3\mu)x^2 + (3 - 2z)x - 3\mu}{8x^2}. \end{aligned} \tag{2.60}$$

In analogy to (2.37), the flux (2.60) must be an integer divided by $(m_+ m_-)$, and we refer to Appendix A for the analysis of this constraint. The same quantization condition, but with F_1 integrated across S_+ , is automatically satisfied due to the relation

$$\mathfrak{s}_i^+ = \frac{g}{2\pi} \int_{S_+} F_i = \mathfrak{s}_i^- + \frac{t}{m_+ m_-} \frac{p_i}{n_+ n_-}, \tag{2.61}$$

which can be both computed explicitly and obtained from the homology relation $S_+ - S_- = \frac{t}{m_+ m_-} \Sigma_2$. On the other hand, the fluxes of F_2 can be obtained switching the sign of z , i.e., exchanging p_1 with p_2 , in the previous formulas and, as a consequence, the quantization follows directly. The fluxes of the R -symmetry gauge potential read

$$\begin{aligned} \frac{1}{2\pi} \int_{S_-} F_R &= \frac{m_+ - m_-}{m_+ m_-} - \frac{t}{m_+ m_- n_-}, \\ \frac{1}{2\pi} \int_{S_+} F_R &= \frac{m_+ - m_-}{m_+ m_-} + \frac{t}{m_+ m_- n_+}, \end{aligned} \tag{2.62}$$

and are well quantized, i.e., they are integers divided by $m_+ m_-$, because n_{\pm} divide t .

2.3 Truncation to $D = 4$ minimal gauged supergravity

An accurate inspection of the $\text{AdS}_2 \times \Sigma_g \times \Sigma_2$ and $\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solutions given in (2.9) and (2.40), respectively, hints at the existence of an underlying consistent truncation of matter-coupled $D = 6$, $U(1)^2$ gauged supergravity on a spindle Σ_2 , down to $D = 4$, $\mathcal{N} = 2$ minimal gauged supergravity. Even though we do not demonstrate the consistency of this truncation⁴, we conjecture its validity in analogy

⁴ This truncation has been proved to be consistent and to preserve supersymmetry in [20].

with the results of [17] and we will provide supporting evidence. For completeness, we report here the action describing the four-dimensional theory

$$S_{4D} = \frac{1}{16\pi G_{(4)}} \int d^4x \sqrt{-g} \left({}^{(4)}R + 6 - \mathcal{F}_{\mu\nu} \mathcal{F}^{\mu\nu} \right), \quad (2.63)$$

along with the corresponding Killing spinor equation

$$\left(\nabla_\mu - i \mathcal{A}_\mu + \frac{1}{2} \gamma_\mu + \frac{i}{4} \mathcal{F}_{\rho\sigma} \gamma^{\rho\sigma} \gamma_\mu \right) \epsilon = 0, \quad (2.64)$$

where $\mathcal{F} = d\mathcal{A}$, $\{\gamma_a, \gamma_b\} = 2\eta_{ab}$ and the AdS₄ radius has been set to 1.

Given a solution of $D = 4$ minimal gauged supergravity comprising a metric $ds_{(4)}^2$ and a gauge potential \mathcal{A} , we propose the following ansatz for the six-dimensional solution:

$$\begin{aligned} ds_{(6)}^2 &= (y^2 h_1 h_2)^{1/4} \left[ds_{(4)}^2 + \frac{y^2}{F} dy^2 + \frac{F}{h_1 h_2} \left(dz - \frac{1}{m} \mathcal{A} \right)^2 \right], \\ A_i &= -\frac{y^3}{h_i} \left(dz - \frac{1}{m} \mathcal{A} \right), \quad X_i = (y^2 h_1 h_2)^{3/8} h_i^{-1}, \\ B &= \frac{2y}{m} \star_4 \mathcal{F}. \end{aligned} \quad (2.65)$$

Here, \star_4 is the Hodge dual with respect to the metric $ds_{(4)}^2$ and (cf. (2.10))

$$h_i(y) = \frac{2g}{3m} y^3 + q_i, \quad F(y) = m^2 h_1 h_2 - y^4. \quad (2.66)$$

This ansatz may be justified comparing the two six-dimensional solutions already mentioned with the following AdS₂ × Σ_g [14] and AdS₂ × Σ₁ [28] backgrounds

$$ds_{(4)}^2 = \frac{1}{4} ds_{\text{AdS}_2}^2 + \frac{1}{2} ds_{\Sigma_g}^2, \quad \mathcal{A} = \frac{1}{2} \omega_{\Sigma_g}, \quad (2.67)$$

$$ds_{(4)}^2 = \frac{x^2}{4} ds_{\text{AdS}_2}^2 + \frac{x^2}{q} dx^2 + \frac{q}{4x^2} d\psi^2, \quad \mathcal{A} = \frac{1}{2} \left(1 - \frac{a}{x} \right) d\psi, \quad (2.68)$$

with $q(x) = x^4 - 4x^2 + 4ax - a^2$. In both cases, plugging the four-dimensional solutions into the ansatz (2.65) gives the corresponding ones in $D = 6$. An additional example is given by the AdS₄ × Σ₂ solution of [27], which can be obtained starting from the AdS₄ vacuum.

In addition to the truncation of the bosonic sector, we also propose an ansatz connecting the Killing spinors of the two supergravity theories in $D = 4$ and $D = 6$. This conjecture is driven both by the construction presented in [17] and by the fact, already observed, that some of the Killing spinor equations are the same for the two background considered in this paper, as well as for the AdS₄ × Σ₂ solution.

The truncation ansatz for the Killing spinors heavily depends on the choice of the decomposition of the six-dimensional gamma matrices. In what follows we will consider (2.15)

$$\Gamma^{\tilde{a}} = \gamma^{\tilde{a}} \otimes \sigma^3, \quad \Gamma^{4,5} = I_4 \otimes \sigma^{1,2}, \tag{2.69}$$

where $\gamma^{\tilde{a}}, \tilde{a} = 0, \dots, 3$, are the (Lorentzian) gamma matrices in $D = 4$. For consistency, \mathcal{B}_6 decomposes as

$$\mathcal{B}_6 = (\mathcal{B}_4 \gamma_5) \otimes (\mathcal{B}_2 \sigma^3), \tag{2.70}$$

where $\gamma_5 = i \gamma^0 \gamma^1 \gamma^2 \gamma^3$ is the four-dimensional chiral matrix and $\mathcal{B}_2 = -\sigma^2$. We conjecture that the first Killing spinor ϵ^1 takes the expression

$$\epsilon^1 = \zeta \otimes \eta, \tag{2.71}$$

where ζ is a Killing spinor of $D = 4$ minimal gauged supergravity and η is given by⁵

$$\eta = \xi y^{1/8} (h_1 h_2)^{-3/16} \begin{pmatrix} f_1^{1/2} \\ -f_2^{1/2} \end{pmatrix}, \tag{2.72}$$

where ξ is a complex constant and

$$f_1(y) \equiv m(h_1 h_2)^{1/2} + y^2, \quad f_2(y) \equiv m(h_1 h_2)^{1/2} - y^2, \tag{2.73}$$

satisfying $F(y) = f_1(y) f_2(y)$. The second Killing spinor ϵ^2 is determined by the symplectic-Majorana condition (2.8), explicitly

$$\epsilon^2 = i (\gamma_5 \zeta^c) \otimes (\sigma^1 \eta^*), \tag{2.74}$$

where we defined the charge conjugate spinor $\zeta^c = \mathcal{B}_4^{-1} \zeta^*$.

As for the bosonic part of the truncation, we can justify this ansatz making contact with the Killing spinors of $D = 4$ minimal gauged supergravity. Assuming, additionally, the decomposition of the four-dimensional gamma matrices in (2.21), the Killing spinors of the $\text{AdS}_2 \times \Sigma_g$ [14] and $\text{AdS}_2 \times \Sigma_1$ [32] (see also [28]) backgrounds are, respectively,⁶

$$\zeta = \vartheta \otimes \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \tag{2.75}$$

$$\zeta = \vartheta \otimes x^{-1/2} \begin{pmatrix} Q_1^{1/2} \\ -Q_2^{1/2} \end{pmatrix}, \tag{2.76}$$

⁵ Here we relabeled η^1 with η and ξ_η with ξ with respect to (2.19).

⁶ Here, ζ is a superposition of the two Killing spinors $\tilde{\epsilon}_1$ and $\tilde{\epsilon}_2$ of [32].

where ϑ is a Majorana Killing spinor on AdS_2 and we defined $Q_1(x) \equiv x^2 - (2x - a)$ and $Q_2(x) \equiv x^2 + (2x - a)$. Uplifting these two spinors by means of the proposed recipe, we obtain exactly the Killing spinors of the corresponding six-dimensional solution. For what concerns the $\text{AdS}_4 \times \mathbb{S}_2$ background, we must take $\zeta = \gamma_5 \vartheta_{(4)}$, with $\vartheta_{(4)}$ a Killing spinor on AdS_4 , due to a slightly different convention on the four-dimensional Killing spinor equations.

The existence and the consistency of the truncation for the Killing spinors would prove that any supersymmetric solution of $D = 4$ minimal gauged supergravity gives rise to a supersymmetric solution of $D = 6$, $U(1)^2$ gauged supergravity after uplifting on \mathbb{S}_2 via (2.65).

3 Uplift to massive type IIA supergravity

The local solutions presented in Sect. 2 can be embedded in massive type IIA supergravity by means of an appropriate ansatz. A consistent truncation for the field content of interest is unfortunately not available. However, if we set $m = 2g/3$ we can consider an enlarged version of the ansatz presented in [27]. This proposal⁷ comes as a superposition of the formulas presented in [25] and [24], which we recover in the respective limits. In the former, only one scalar is considered, but the six-dimensional two-form B is nonzero, while the latter encompasses four scalar fields, but vanishing B field. The ansatz for the metric in the string frame and the dilaton is the same as in [27]

$$ds_{\text{s.f.}}^2 = \mu_0^{-1/3} (X_1 X_2)^{-1/4} \left\{ \Delta^{1/2} ds_6^2 + g^{-2} \Delta^{-1/2} [X_0^{-1} d\mu_0^2 + X_1^{-1} (d\mu_1^2 + \mu_1^2 \sigma_1^2) + X_2^{-1} (d\mu_2^2 + \mu_2^2 \sigma_2^2)] \right\}, \quad (3.1)$$

$$e^\Phi = \mu_0^{-5/6} \Delta^{1/4} (X_1 X_2)^{-5/8}, \quad (3.2)$$

but now we have also a nonvanishing two-form field

$$B_{(2)} = \frac{1}{2} \mu_0^{2/3} B. \quad (3.3)$$

Here, ds_6^2 stands for the line element of the six-dimensional space M_6 and the one-forms $\sigma_i \equiv d\phi_i - g A_i$ are built up from the six-dimensional gauge fields. The angular coordinates ϕ_1, ϕ_2 have canonical 2π periodicities, whereas the coordinates μ_a , with $a = 0, 1, 2$, satisfy the constraint $\sum \mu_a^2 = 1$. Lastly, the warp factor reads

$$\Delta = \sum_{a=0}^2 X_a \mu_a^2. \quad (3.4)$$

⁷ A more general truncation ansatz, which does not require $m = 2g/3$, has been proposed in [20].

A possible parameterization of the coordinates μ_a is

$$\mu_0 = \sin \xi, \quad \mu_1 = \cos \xi \sin \eta, \quad \mu_2 = \cos \xi \cos \eta, \tag{3.5}$$

with $\eta \in [0, \pi/2]$ and $\xi \in (0, \pi/2]$, where the range of ξ is fixed by the necessity of having $\mu_0 > 0$. In this way, the ten-dimensional metric (3.1) parameterizes, at each point of M_6 , a four-dimensional hemisphere, denoted as \mathbb{S}^4 from now on. The metric on \mathbb{S}^4 is in general squashed, and becomes the metric on “half the round four-sphere” when $X_1 = X_2 = 1$. The RR sector of massive type IIA supergravity comprises the ten-dimensional Romans mass and the two-form flux

$$F_{(0)} = m = \frac{2g}{3}, \quad F_{(2)} = \frac{g}{3} \mu_0^{2/3} B, \tag{3.6}$$

along with the four-form flux, conveniently written in terms of its Hodge dual as⁸

$$\begin{aligned} \star_{10} F_{(4)} &= -gU \text{vol}(M_6) + \frac{1}{g^2} \sum_i X_i^{-2} \mu_i (\star_6 F_i) \wedge d\mu_i \wedge \sigma_i \\ &\quad - \frac{1}{g} \sum_a X_a^{-1} \mu_a (\star_6 dX_a) \wedge d\mu_a \\ &\quad + \frac{\mu_1 \mu_2}{3g^3} \Delta^{-1} X_0 B \wedge d\mu_1 \wedge d\mu_2 \wedge \sigma_1 \wedge \sigma_2 \\ &\quad - \frac{\mu_1 \mu_2}{2g^3} \Delta^{-1} H \wedge (X_2 \mu_2 d\mu_1 - X_1 \mu_1 d\mu_2) \wedge \sigma_1 \wedge \sigma_2, \end{aligned} \tag{3.7}$$

where we defined

$$U = 2 \sum_{a=0}^2 X_a^2 \mu_a^2 - \left[\frac{4}{3} X_0 + 2(X_1 + X_2) \right] \Delta. \tag{3.8}$$

The Hodge star operators \star_{10} and \star_6 are computed using the string frame metric (3.1) and ds_6^2 , respectively. Even though we did not prove that this truncation is consistent, we tested it successfully against our $\text{AdS}_2 \times \Sigma_g \times \bar{\Sigma}_2$ (2.9) and $\text{AdS}_2 \times \bar{\Sigma}_1 \times \bar{\Sigma}_2$ (2.40) solutions.

Provided the equations of motion of the six-dimensional gauged supergravity are satisfied, the above ansatz should solve the equations of motion of massive type IIA

⁸ The four-flux differs in a global sign from (2.16) of [27]. Although this fact does not affect the equations of motion therein, which, in that case, are invariant under $F_{(4)} \mapsto -F_{(4)}$, the expression (3.7) correctly reduces to [25] when only one scalar field is retained.

supergravity, which can be derived from the string frame action⁹

$$S_{\text{mIIA}} = \frac{1}{16\pi G_{(10)}} \left\{ \int d^{10}x \sqrt{-g} \left[e^{-2\Phi} (R + 4|d\Phi|^2 - \frac{1}{2}|H_{(3)}|^2) - \frac{1}{2}(F_{(0)}^2 + |F_{(2)}|^2 + |F_{(4)}|^2) \right] - \frac{1}{2} \int (B_{(2)} \wedge dC_{(3)} \wedge dC_{(3)} + 2F_{(0)}B_{(2)}^3 \wedge dC_{(3)} + 6F_{(0)}^2 B_{(2)}^5) \right\}, \quad (3.9)$$

where, in these conventions, the field strengths are

$$\begin{aligned} H_{(3)} &= dB_{(2)}, & F_{(2)} &= dC_{(1)} + F_{(0)}B_{(2)}, \\ F_{(4)} &= dC_{(3)} - H_{(3)} \wedge C_{(1)} + \frac{1}{2}F_{(0)}B_{(2)} \wedge B_{(2)}. \end{aligned} \quad (3.10)$$

As noticed in [27], the ten-dimensional equations of motion are invariant under a scaling symmetry, which defines an “improved uplift” and whose action is

$$\begin{aligned} ds_{\text{s.f.}}^2 &= \lambda^2 ds_{\text{s.f.}}^2, & e^{\hat{\Phi}} &= \lambda^2 e^{\Phi}, & \hat{B}_{(2)} &= \lambda^2 B_{(2)}, \\ \hat{F}_{(0)} &= \lambda^{-3} F_{(0)}, & \hat{C}_{(1)} &= \lambda^{-1} C_{(1)}, & \hat{C}_{(3)} &= \lambda C_{(3)}, \end{aligned} \quad (3.11)$$

where λ is any strictly positive constant. This additional parameter λ enters in the regularity analysis of the uplifted ten-dimensional solutions. In particular, its presence will be necessary for the fluxes to be correctly quantized [27].

In the next subsections, we will specify our proposed truncation ansatz to the solutions presented in (2.9) and (2.40). After having applied the scaling symmetry (3.11), we will quantize the fluxes, thus ensuring the global regularity of the ten-dimensional solutions.

3.1 $\text{AdS}_2 \times \Sigma_g \times \Sigma_2$ solutions

We begin this section by presenting the key ingredients for the flux quantization and the computation of the entropy. The quantization conditions are

$$\begin{aligned} (2\pi\ell_s)F_{(0)} &= n_0 \in \mathbb{N}, & \frac{1}{(2\pi\ell_s)^3} \int_{\mathbb{S}^4} F_{(4)} &= N \in \mathbb{N}, \\ & & \frac{1}{(2\pi\ell_s)^3} \int_{\mathbb{M}_4} F_{(4)} &= K \in \mathbb{N}, \end{aligned} \quad (3.12)$$

where $\mathbb{M}_4 = \Sigma_g \times \Sigma_2$. At $\xi = 0$, where $\mu_0 = 0$ and the warp factor is singular, there are an O8-plane and $N_f = 8 - n_0$ coincident D8-branes, while N D4-branes are

⁹ Here we use the shortcut $B_{(2)}^n$ to denote the wedge product of $B_{(2)}$ with itself n times, divided by $n!$.

wrapped over $\Sigma_g \times \mathbb{Z}_2$. These integers get contributions from the following fluxes

$$F_{(0)} = \frac{2g}{3\lambda^3}, \tag{3.13}$$

$$\begin{aligned}
 F_{(4)} = & -\frac{\lambda\mu_0^{1/3}h_1h_2}{g^3\Delta_h} \left\{ \frac{U_h}{\Delta_h} \frac{\mu_1\mu_2}{\mu_0} d\mu_1 \wedge d\mu_2 \wedge \sigma_1 \wedge \sigma_2 \right. \\
 & - \sum_{i \neq j} \left[g F_i \wedge d\phi_j - \frac{y^3(h'_i - 3y^{-1}h_i)h_j}{\Delta_h h_i} \mu_i^2 dy \wedge \sigma_i \wedge \sigma_j \right] \\
 & \left. \wedge (\mu_0\mu_j d\mu_j - y^3h_j^{-1}\mu_j^2 d\mu_0) \right\} \\
 & + \frac{\lambda y}{2} \mu_0^{4/3} \text{vol}(\Sigma_g) \wedge dy \wedge dz - \frac{\lambda y^2}{3} \mu_0^{1/3} \text{vol}(\Sigma_g) \wedge dz \wedge d\mu_0, \tag{3.14}
 \end{aligned}$$

while the metric and the dilaton enter in the computation of the entropy

$$\begin{aligned}
 ds_{\text{s.f.}}^2 = & \lambda^2 \mu_0^{-1/3} y^{-1} \Delta_h^{1/2} \left\{ \frac{1}{4} ds_{\text{AdS}_2}^2 + ds_{\Sigma_g}^2 + \frac{y^2}{F} dy^2 + \frac{F}{h_1 h_2} \left(dz - \frac{1}{2m} \omega_{\Sigma_g} \right)^2 \right. \\
 & \left. + g^{-2} y \Delta_h^{-1} [y^3 d\mu_0^2 + h_1 (d\mu_1^2 + \mu_1^2 \sigma_1^2) + h_2 (d\mu_2^2 + \mu_2^2 \sigma_2^2)] \right\}, \tag{3.15}
 \end{aligned}$$

$$e^\Phi = \lambda^2 \mu_0^{-5/6} y^{-3/2} \Delta_h^{1/4}. \tag{3.16}$$

For convenience, we defined the functions

$$\begin{aligned}
 \Delta_h = & h_1 h_2 \mu_0^2 + y^3 h_2 \mu_1^2 + y^3 h_1 \mu_2^2, \\
 U_h = & 2[(y^3 - h_1)(y^3 - h_2)\mu_0^2 - y^6] - \frac{4}{3} \Delta_h. \tag{3.17}
 \end{aligned}$$

Additionally, we report for completeness also the other ten-dimensional fields, which, however, do not play an important role in our analysis

$$B_{(2)} = \frac{\lambda^2 y}{4m} \mu_0^{2/3} \text{vol}(\text{AdS}_2), \quad F_{(2)} = \frac{g y}{6\lambda m} \mu_0^{2/3} \text{vol}(\text{AdS}_2). \tag{3.18}$$

Plugging our solution in the first two conditions of (3.12) gives

$$g^8 = \frac{1}{(2\pi \ell_s)^8} \frac{18\pi^6}{N^3 n_0}, \quad \lambda^8 = \frac{8\pi^2}{9N n_0^3}. \tag{3.19}$$

It is worth noting that, being N and n_0 integers, the second equation would be inconsistent with $\lambda = 1$. Such a problem arises from the fact that, without introducing λ , the first two constraints in (3.12) would have to be imposed on an unique dimensionless parameter, namely $(g\ell_s)$. This makes the scaling symmetry (3.11) crucial to establish

the regularity of the ten-dimensional uplifted solution. The integration in the third condition of (3.12) is performed along a representative of \mathbb{M}_4 , which we take to be at the pole of the hemisphere \mathbb{S}^4 , i.e., in $\xi = \pi/2$. The result can be implicitly expressed in terms of the integer parameters (g, n_{\pm}, z, N) by means of equations (3.31) and (3.38) of [27], and reads

$$K = N\chi_2(g-1) \frac{3[3\mu(x^2+1) - x(x^2+5)]}{8x(x^2+3)}. \quad (3.20)$$

The analysis of this quantization condition is left to Appendix A.

We now move to the computation of the entropy. The first step is to write the ten-dimensional metric in the form

$$ds_{\text{s.f.}}^2 = e^{2A}(ds_{\text{AdS}_2}^2 + ds_{M_8}^2), \quad (3.21)$$

where $ds_{M_8}^2$ is the metric on the internal space M_8 , that is the total space of an \mathbb{S}^4 bundle over \mathbb{M}_4 , namely $\mathbb{S}^4 \hookrightarrow M_8 \rightarrow \mathbb{M}_4$. There is a $U(1)^2$ symmetry acting on \mathbb{S}^4 , with gA_i connections on the associated circle bundles, and the correct quantization of the corresponding magnetic fluxes t_i and s_i^{\pm} , as discussed in Sect. 2.1, ensures that the ten-dimensional solution is a well-defined orbifold. The entropy of our solution can be read from the two-dimensional effective Newton constant $G_{(2)}$ as

$$S = \frac{1}{4G_{(2)}} = \frac{8\pi^2}{(2\pi\ell_s)^8} \int e^{8A-2\Phi} \text{vol}(M_8). \quad (3.22)$$

Applying this to metric (3.15), we obtain

$$\begin{aligned} S_{\Sigma_g \times \Sigma_2} &= \frac{8\pi^2}{(2\pi\ell_s)^8} \frac{3\pi^2\lambda^4}{20g^4} 4\pi(g-1)(y_S^3 - y_N^3)\Delta z \\ &= (g-1)F_{S^3 \times \Sigma_2}, \end{aligned} \quad (3.23)$$

where $F_{S^3 \times \Sigma_2}$ is the free energy of $d=3$, $\mathcal{N}=2$ SCFTs that arise from a system of N D4-branes and N_f D8-branes wrapped on a spindle [27]

$$F_{S^3 \times \Sigma_2} = \chi_2 \frac{\sqrt{3}\pi N^{5/2}}{5\sqrt{8-N_f}} \frac{[3\mu(x^2+1) - x(x^2+5)]^{3/2}}{x(x^2+3)(\mu-x)^{1/2}}. \quad (3.24)$$

3.2 AdS₂ × Σ₁ × Σ₂ solutions

Similarly to the previous section we present the explicit expressions of the fluxes, which are then quantized. Apart from the Romans mass, which is the same as in

(3.13), the complete uplifted solution includes the four-flux

$$\begin{aligned}
 F_{(4)} = & -\frac{\lambda\mu_0^{1/3}h_1h_2}{g^3\Delta_h} \left\{ \frac{U_h}{\Delta_h} \frac{\mu_1\mu_2}{\mu_0} d\mu_1 \wedge d\mu_2 \wedge \sigma_1 \wedge \sigma_2 \right. \\
 & - \sum_{i \neq j} \left[g F_i \wedge d\phi_j - \frac{y^3(h'_i - 3y^{-1}h_i)h_j}{\Delta_h h_i} \mu_i^2 dy \wedge \sigma_i \wedge \sigma_j \right] \\
 & \left. \wedge (\mu_0\mu_j d\mu_j - y^3 h_j^{-1} \mu_j^2 d\mu_0) \right\} \\
 & + \frac{\lambda a y}{2x^2} \mu_0^{4/3} dx \wedge d\psi \wedge dy \wedge dz - \frac{\lambda a y^2}{3x^2} \mu_0^{1/3} dx \wedge d\psi \wedge dz \wedge d\mu_0, \quad (3.25)
 \end{aligned}$$

and the metric and dilaton

$$\begin{aligned}
 ds_{s.f.}^2 = & \lambda^2 \mu_0^{-1/3} y^{-1} \Delta_h^{1/2} \left\{ \frac{x^2}{4} ds_{AdS_2}^2 + \frac{x^2}{q} dx^2 + \frac{q}{4x^2} d\psi^2 + \frac{y^2}{F} dy^2 \right. \\
 & + \frac{F}{h_1 h_2} \left(dz - \frac{1}{2m} \left(1 - \frac{a}{x} \right) d\psi \right)^2 \\
 & \left. + g^{-2} y \Delta_h^{-1} [y^3 d\mu_0^2 + h_1 (d\mu_1^2 + \mu_1^2 \sigma_1^2) + h_2 (d\mu_2^2 + \mu_2^2 \sigma_2^2)] \right\}, \quad (3.26)
 \end{aligned}$$

$$e^\Phi = \lambda^2 \mu_0^{-5/6} y^{-3/2} \Delta_h^{1/4}, \quad (3.27)$$

where Δ_h and U_h were defined in (3.17). Again, even if they are not fundamental for the computations, we also have

$$B_{(2)} = \frac{\lambda^2 a y}{4m} \mu_0^{2/3} \text{vol}(AdS_2), \quad F_{(2)} = \frac{a g y}{6\lambda m} \mu_0^{2/3} \text{vol}(AdS_2). \quad (3.28)$$

The quantization conditions are the same as in (3.12), where now $\mathbb{M}_4 = \Sigma_1 \times \Sigma_2$, and the first two give again the relations (3.19). The last one, expressed in terms of (m_\pm, n_\pm, z, N) , reads

$$K = N \chi_2 \frac{m_+ - m_-}{m_+ m_-} \frac{3[3\mu(x^2 + 1) - x(x^2 + 5)]}{8x(x^2 + 3)} \in \mathbb{N}. \quad (3.29)$$

We refer to Appendix A for a detailed analysis.

The ten-dimensional geometry is similar to the solution of the previous subsection, with the internal space M_8 having a fibration structure $S^4 \hookrightarrow M_8 \rightarrow \mathbb{M}_4$. Again, there is a $U(1)^2$ symmetry acting on S^4 , with gA_i connections on the associated circle bundles. Therefore, the uplifted solution can be considered a well-defined orbifold thanks to the correct quantization of the magnetic fluxes of gA_i , as discussed in Sect. 2.2.

The entropy of this solution can be computed following the same steps of the previous case and the final result is

$$\begin{aligned} S_{\Sigma_1 \times \Sigma_2} &= \frac{8\pi^2}{(2\pi\ell_s)^8} \frac{3\pi^2\lambda^4}{20g^4} (x_S - x_N)\Delta\psi (y_S^3 - y_N^3)\Delta z \\ &= \frac{1}{2\pi} A_{\Sigma_1} F_{S^3 \times \Sigma_2}, \end{aligned} \quad (3.30)$$

where $F_{S^3 \times \Sigma_2}$ is given in (3.24) and A_{Σ_1} is the area of the horizon of the four-dimensional supersymmetric black hole solutions with $\text{AdS}_2 \times \Sigma_1$ near-horizon studied in [28], namely

$$A_{\Sigma_1} = \frac{\sqrt{2}\sqrt{m_+^2 + m_-^2} - (m_+ + m_-)}{m_+ m_-} \pi. \quad (3.31)$$

4 Toric geometry

In this section, we will provide a description of the orbifold $\Sigma_1 \times \Sigma_2$ in terms of certain combinatorial data, that will be used in the next section to construct the entropy function. Since $\Sigma_1 \times \Sigma_2$ admits a $U(1)^2$ action, corresponding to the $U(1)^2$ isometry of the metric (2.40), it is natural to seek a description in terms of toric geometry. Below, we will discuss a method for extracting a set of toric data that is sufficient for the purposes of this paper, leaving a more detailed study of the geometry for future work. As a warm-up, we will begin with an $S^2 \times \Sigma_2$ geometry, namely a spindle fibered over a smooth two-sphere. This can be obtained formally as an analytic continuation of the $\Sigma_g \times \Sigma_2$ family and although it is not an actual supersymmetric background, it solves the equations of motion. This case can be analyzed in the framework of symplectic toric geometry and we will derive a *labeled polytope* [43] from the image of an associated moment map. From the algebraic geometry point of view, the corresponding fan of labeled normals to the polytope does not lead to a toric variety, but to a generalization known as a “stack.” The reason is that in ordinary toric varieties orbifold singularities along divisors cannot appear, as all the singularities are in co-dimension higher than one. In the broader context of stacks, the key combinatorial gadget is called a *stacky fan*, which roughly speaking is a simplicial fan with a distinguished lattice point on each ray of the fan (see, e.g., [11, 34, 42, 48]). For the more general $\Sigma_1 \times \Sigma_2$ family we will not be able to find a symplectic structure and therefore we will have to use an indirect method to extract the toric data. These will again take the form of a stacky fan and will be taken as inputs for constructing an entropy function in terms of gravitational blocks in the next section. As further illustration of our approach, in Appendix C we present the derivation of the Delzant polytope of the first Hirzebruch surface \mathbb{F}_1 , starting from the explicit metric found in [45], as well as the toric data relevant for the seven-dimensional solutions of [17].

4.1 $\text{AdS}_2 \times S^2 \times \Sigma_2$ solutions

Before studying the $\Sigma_1 \times \Sigma_2$ orbifold, in which case the analysis is more involved, we begin with a geometry characterized by a smooth two-dimensional base. Since a Riemann surface with $g > 0$ is not toric, we need to consider its analytic continuation to a two-sphere. Although non-supersymmetric, the $\text{AdS}_2 \times S^2 \times \Sigma_2$ backgrounds form a family of solutions to the equations of motion very similar to (2.9), but with a different value of the radius of the Riemann surface, now a sphere. Specifically, the space we consider is the four-dimensional toric orbifold $S^2 \times \Sigma_2$ with metric

$$ds^2_{S^2 \times \Sigma_2} = \frac{1}{6}(d\theta^2 + \sin^2\theta d\psi^2) + \frac{y^2}{F} dy^2 + \frac{F}{h_1 h_2} \left(dz + \frac{1}{2m} \cos\theta d\psi \right)^2. \tag{4.1}$$

In this case, the condition on t (2.36) in order to have a well-defined fibration reads

$$t = -\frac{1}{m} \frac{2\pi}{\Delta z} \in \mathbb{Z}, \tag{4.2}$$

where t results to be negative. For fixed values of y , metric (4.1) describes an S^3/\mathbb{Z}_{-t} , where S^3 is a squashed three-sphere written as a Hopf fibration; therefore, a natural basis of an effective two-torus action is [45]¹⁰

$$e_1 = \partial_{v_2}, \quad e_2 = \partial_\psi - \frac{t}{2} \partial_{v_2}, \tag{4.3}$$

where we introduced the 2π -periodic coordinate $v_2 = \frac{2\pi}{\Delta z} z$. In total, we have four fixed points v_ℓ ($\ell = 1, \dots, 4$) under the action of the two-torus, which are all the possible combinations obtained by pairing the poles of the sphere ($\theta = 0, \pi$) and the poles of the spindle ($y = y_{N,S}$)

$$\begin{aligned} v_1 &= \{\theta = 0, y = y_N\}, & v_2 &= \{\theta = 0, y = y_S\}, \\ v_3 &= \{\theta = \pi, y = y_S\}, & v_4 &= \{\theta = \pi, y = y_N\}. \end{aligned} \tag{4.4}$$

We now consider the conformally rescaled metric $ds^2 = \Gamma(y) ds^2_{S^2 \times \Sigma_2}$, with $\Gamma(y) > 0$, with compatible symplectic two-form

$$\omega = \Gamma(y) \left[\frac{1}{6} \sin\theta d\theta \wedge d\psi + \frac{y}{(h_1 h_2)^{1/2}} dy \wedge \left(dz + \frac{1}{2m} \cos\theta d\psi \right) \right]. \tag{4.5}$$

When $\Gamma'(y) = -\frac{3y}{m(h_1 h_2)^{1/2}} \Gamma(y)$ ¹¹, the two-form (4.5) is closed and can be written as

$$\omega = d\psi \wedge d\left[\frac{1}{6} \Gamma(y) \cos\theta \right] + dv_2 \wedge d\left[-\frac{1}{3t} \Gamma(y) \right]. \tag{4.6}$$

¹⁰ We exchanged e_1 and e_2 with respect to [45] in order to have the vectors \vec{n}_ℓ ordered counterclockwise.

¹¹ A real and positive solution to this differential equation exists and is unique, up to an irrelevant overall constant.




Fig. 1 Labeled polytope corresponding to the $S^2 \times \Sigma_2$ orbifold. The different facets are labeled with the integers m_ℓ . In the figure $t = -2$

From this expression we can derive the moment maps with respect to the basis (4.3)

$$\vec{\mu} = -\frac{1}{3t}\Gamma(y) \left(1, -\frac{t(1 + \cos \theta)}{2} \right), \tag{4.7}$$

and compute the image of the fixed points (4.4)

$$\begin{aligned} \vec{\mu}(v_1) &= -\frac{1}{3t}\Gamma(y_N) (1, -t), & \vec{\mu}(v_2) &= -\frac{1}{3t}\Gamma(y_S) (1, -t), \\ \vec{\mu}(v_3) &= -\frac{1}{3t}\Gamma(y_S) (1, 0), & \vec{\mu}(v_4) &= -\frac{1}{3t}\Gamma(y_N) (1, 0), \end{aligned} \tag{4.8}$$

which are vertices of the moment polytope. Vertices are connected by facets and we define the facet D_ℓ to be the one that joins $v_{\ell-1}$ and v_ℓ , where ℓ is modulo 4¹². In order to correctly draw the polytope, we notice that, since $\Gamma'(y) < 0$, we have $\Gamma(y_N) > \Gamma(y_S)$.

Due to the orbifold nature of the spindle, we can construct an associated labeled polytope following [43], which extends the classic Delzant construction. In particular, if the preimage under moment maps of a facet has an orbifold singularity locally modeled on $\mathbb{C}/\mathbb{Z}_{m_\ell}$, we label the facet with the corresponding integer m_ℓ , otherwise there is no label (formally, we have $m_\ell = 1$). Specifically, the $S^2 \times \Sigma_2$ orbifold is characterized by

$$\begin{aligned} D_1 = \{y = y_N\} : m_1 = n_-, & & D_2 = \{\theta = 0\} : m_2 = 1, \\ D_3 = \{y = y_S\} : m_3 = n_+, & & D_4 = \{\theta = \pi\} : m_4 = 1. \end{aligned} \tag{4.9}$$

The resulting labeled polytope is depicted in Fig. 1, and if the labels are stripped off

¹² With a slight abuse of notation, we denote as v_ℓ both the fixed points and the vertices. Moreover, we denoted as D_ℓ both the facets and the corresponding divisors, i.e., their preimages under $\vec{\mu}$.

this is exactly the polytope corresponding to the Hirzebruch surfaces \mathbb{F}_{-t} . This polytope can also be described uniquely by a set of inequalities

$$\langle \vec{\mu}, m_\ell \vec{n}_\ell \rangle \leq \lambda_\ell, \tag{4.10}$$

where $\langle \cdot, \cdot \rangle$ is the standard inner product between vectors in \mathbb{R}^2 , $\vec{\mu}$ is a generic point in the (μ_1, μ_2) -plane, λ_ℓ is a real number and each \vec{n}_ℓ is a primitive element of \mathbb{Z}^2 , which represents the outward-pointing normal vector to the facet D_ℓ . In particular, we have

$$\vec{n}_1 = (1, 0), \quad \vec{n}_2 = (t, 1), \quad \vec{n}_3 = (-1, 0), \quad \vec{n}_4 = (0, -1). \tag{4.11}$$

The method we employed to construct the labeled polytope above is rigorous and the result fits in the theory of symplectic toric orbifolds [43]. However, we will now study this toric orbifold from a different point of view, in order to derive a method that can be applied also to orbifolds which do not admit a conformally rescaled symplectic form. A direct computation using the metric (4.1) shows that there are four distinct loci D_ℓ where a linear combination of the two Killing vectors ∂_ψ and ∂_z degenerates. These are precisely the toric divisors which in the symplectic context correspond to the preimages of the four facets of the polytope and intersect at the four fixed points. Explicitly, denoting with $\xi_{(\ell)}$ the Killing vector that degenerates at D_ℓ , they are

$$\xi_{(1)} = n_- \partial_{v_2}, \quad \xi_{(2)} = \partial_\psi + \frac{t}{2} \partial_{v_2}, \quad \xi_{(3)} = n_+ \partial_{v_2}, \quad \xi_{(4)} = \partial_\psi - \frac{t}{2} \partial_{v_2}, \tag{4.12}$$

normalized so to have unitary surface gravity

$$\kappa_{\text{grav}}^2 = \frac{\partial_\mu |\xi|^2 \partial^\mu |\xi|^2}{4|\xi|^2} = 1 \quad \text{where} \quad |\xi|^2 = 0. \tag{4.13}$$

Expanded in the basis (4.3), the degenerating Killing vectors read

$$\xi_{(1)} = n_- e_1, \quad \xi_{(2)} = t e_1 + e_2, \quad \xi_{(3)} = n_+ e_1, \quad \xi_{(4)} = e_2. \tag{4.14}$$

We notice that

$$\xi_{(\ell)} = m_\ell \vec{n}_\ell \cdot (e_1, e_2), \tag{4.15}$$

where we packed the basis vectors (4.3) into a two-dimensional vector (e_1, e_2) and (\vec{n}_ℓ, m_ℓ) are the toric data of the corresponding labeled polytope.

We can use relation (4.15) to infer the toric data starting from the information about the degenerate Killing vectors. In particular, given a toric divisor D_ℓ where the Killing vector $\xi_{(\ell)}$ is degenerate, studying the (possible) orbifold singularity on D_ℓ we can obtain the corresponding integer m_ℓ . Expanding $\xi_{(\ell)}$ on a suitable basis—in which the two-torus act effectively—according to (4.15) we can deduce the normal vectors \vec{n}_ℓ , up to a global sign depending on the conventions. This sign can be chosen arbitrarily




Fig. 2 Stacky fan corresponding to the vectors (4.16). The cones generated by $\vec{\tilde{n}}_\ell$ and $\vec{\tilde{n}}_{\ell+1}$ are denoted as τ_ℓ . In this example, we took $n_+ = 3, n_- = 2$ and $t = -2$

for one of the vectors and for the others it follows requiring the vectors $\vec{\tilde{n}}_\ell$ to be ordered counterclockwise, in our conventions.

For an un-labeled, convex, rational polytope (not necessarily a Delzant polytope) one can pass to the algebraic point of view, where the primitive normals to the polytope represent the fan of a toric variety. In this context, the one-dimensional cones (rays) of the fan are in a one-to-one correspondence with the toric divisors and the order of the orbifold singularities at the intersection between two divisors can be read off from the determinant of two adjacent generators, namely $\det(\vec{\tilde{n}}_\ell, \vec{\tilde{n}}_{\ell+1})$. Orbifold singularities along divisors cannot arise in toric varieties, so the information of a labeled polytope cannot be accommodated in an ordinary fan. An equivalent way of thinking about the labels m_ℓ is by dropping the primitive requirement and considering $\vec{\tilde{n}}_\ell = m_\ell \vec{n}_\ell$ instead of \vec{n}_ℓ [1]. We will refer to the \vec{n}_ℓ as “short” vectors and to the $\vec{\tilde{n}}_\ell$ as “long” vectors. In algebraic geometry, the fan obtained from such “long” (non-primitive) normals can be understood in the broader context of stacks and it is referred to as a stacky fan. This turns out to be the correct combinatorial data needed in the construction of the entropy function in the next section. The “long” vectors characterizing the $S^2 \times \Sigma_2$ orbifold then read

$$\vec{\tilde{n}}_1 = (n_-, 0), \quad \vec{\tilde{n}}_2 = (t, 1), \quad \vec{\tilde{n}}_3 = (-n_+, 0), \quad \vec{\tilde{n}}_4 = (0, -1), \quad (4.16)$$

and the corresponding stacky fan, dual to the labeled polytope of Fig. 1, is presented in Fig. 2.

Let us make some comments on the orbifold $\Sigma_g \times \Sigma_2$, with $g > 1$. This cannot be toric in any sense; therefore, none of the previous considerations can be applied. Nevertheless, for our purposes, we can think about extending the results obtained for the $S^2 \times \Sigma_2$ toric orbifold to this case, and adopt as “toric data” the same vectors $\vec{\tilde{n}}_\ell$ (4.11) and the same integers m_ℓ (4.9). The information on the genus of the Riemann surface does not enter in any way in the “toric data,” but it will appear in the twisting procedure

of Sect. 5, as discussed in the examples. Clearly, for $\Sigma_g \times \Sigma_2$ the divisors D_2 and D_4 have no physical meaning, and likewise the four fixed points v_ℓ , but they are all artifacts needed for the application of the conjectural entropy function we will present in Sect. 5.

4.2 AdS₂ × Σ₁ × Σ₂ solutions

After the warm-up with the $S^2 \times \Sigma_2$ case, we are now ready to study the $\Sigma_1 \times \Sigma_2$ orbifold. In this case, we can not construct a conformal symplectic structure; therefore, we do not have a moment map to derive the labeled polytope. Nevertheless, we are still able to obtain the toric data that we need by analyzing where the Killing vectors of $\Sigma_1 \times \Sigma_2$ degenerate. Recall that the metric on $\Sigma_1 \times \Sigma_2$ is

$$ds_{\Sigma_1 \times \Sigma_2}^2 = \frac{x^2}{q} dx^2 + \frac{q}{4x^2} d\psi^2 + \frac{y^2}{F} dy^2 + \frac{F}{h_1 h_2} \left(dz - \frac{1}{2m} \left(1 - \frac{a}{x} \right) d\psi \right)^2. \tag{4.17}$$

A simple direct calculation shows that it has four Killing vectors degenerating at the poles of the two spindles, defining the four loci

$$D_1 = \{y = y_N\}, \quad D_2 = \{x = x_N\}, \quad D_3 = \{y = y_S\}, \quad D_4 = \{x = x_S\}, \tag{4.18}$$

intersecting at the four fixed points

$$\begin{aligned} v_1 &= \{x = x_N, y = y_N\}, & v_2 &= \{x = x_N, y = y_S\}, \\ v_3 &= \{x = x_S, y = y_S\}, & v_4 &= \{x = x_S, y = y_N\}. \end{aligned} \tag{4.19}$$

normalizing the degenerating Killing vectors so that they have unitary surface gravity, these read

$$\begin{aligned} \xi_{(1)} &= n_- \partial_{v_2}, & \xi_{(2)} &= m_- \left(\partial_{v_1} - \frac{a - x_N}{2mx_N} \frac{\Delta\psi}{\Delta z} \partial_{v_2} \right), \\ \xi_{(3)} &= n_+ \partial_{v_2}, & \xi_{(4)} &= m_+ \left(\partial_{v_1} - \frac{a - x_S}{2mx_S} \frac{\Delta\psi}{\Delta z} \partial_{v_2} \right), \end{aligned} \tag{4.20}$$

where, in addition to $v_2 = \frac{2\pi}{\Delta z} z$, we introduced also the 2π -periodic coordinate $v_1 = \frac{2\pi}{\Delta\psi} \psi$.

Reading off the labels associated with the D_ℓ turns out to be a bit tricky. We will first present a naive argument leading to slightly wrong conclusions, and then we will explain how to amend this. Each of the Killing vectors in (4.20) is in a one-to-one correspondence with a specific pole of either Σ_1 or Σ_2 , whose rotation around the axis is generated by the 2π -periodic Killing vector ∂_{v_1} or ∂_{v_2} , respectively. This suggests that the integer coefficient in front of this rotational Killing vector corresponds to a

$\mathbb{C}/\mathbb{Z}_{m_\ell}$ quotient singularity, with $m_\ell = (n_-, m_-, n_+, m_+)$. While for D_1 and D_3 this is correct, for D_2 and D_4 this conclusion is not precise, as we will show below.

In order to derive the normal vectors \vec{n}_ℓ we need to find an appropriate basis in which to expand the $\xi_{(\ell)}$. Inspired by (4.3), where ∂_{v_2} is the first basis element and $\xi_{(4)}$ plays the role of e_2 , we consider the following basis

$$e_1 = \partial_{v_2}, \quad e_2 = \partial_{v_1} - \frac{a - x_S}{2mx_S} \frac{\Delta\psi}{\Delta z} \partial_{v_2}, \tag{4.21}$$

in which the degenerating Killing vectors are decomposed as

$$\begin{aligned} \xi_{(1)} &= n_- e_1, & \xi_{(2)} &= \frac{t}{m_+} e_1 + m_- e_2, \\ \xi_{(3)} &= n_+ e_1, & \xi_{(4)} &= m_+ e_2, \end{aligned} \tag{4.22}$$

where we made use of the quantization condition on t in (2.59) in the form

$$\frac{t}{m_+ m_-} = -\frac{a(x_S - x_N)}{2mx_N x_S} \frac{\Delta\psi}{\Delta z}. \tag{4.23}$$

Note that, even though $x_{N,S}$ and a are real numbers, all the coefficients in (4.22) are rational. Taking the labels as above, for the ‘‘short’’ vectors we obtain

$$\vec{n}_1 = (1, 0), \quad \vec{n}_2 = \left(\frac{t}{m_+ m_-}, 1\right), \quad \vec{n}_3 = (-1, 0), \quad \vec{n}_4 = (0, -1), \tag{4.24}$$

where we picked the sign of \vec{n}_1 as in (4.11) and the others followed accordingly. However, since these vectors are not primitive elements of \mathbb{Z}^2 , they cannot be regarded as the normal vectors to the facets of a polytope, in the sense of [43]. Even adopting the alternative description in which the labels are included in the normal vectors, the ‘‘long’’ vectors associated with (4.24) are not valued in \mathbb{Z}^2 :

$$\vec{\hat{n}}_1 = (n_-, 0), \quad \vec{\hat{n}}_2 = \left(\frac{t}{m_+}, m_-\right), \quad \vec{\hat{n}}_3 = (-n_+, 0), \quad \vec{\hat{n}}_4 = (0, -m_+). \tag{4.25}$$

Although in the mathematical literature there are examples where the normal vectors are allowed to take rational [1, 41] as well as real values [7], the fact that the Killing vectors in (4.22) are not linear combinations of the basis $\{e_1, e_2\}$ with integer coefficients indicates that the two-torus is not acting effectively. A better basis can be obtained rotating the vectors $\vec{\hat{n}}_\ell$ through the following $SL(2, \mathbb{Q})$ matrix

$$S = \begin{pmatrix} 1 & -r_-/m_+ \\ 0 & 1 \end{pmatrix}, \tag{4.26}$$

where r_- is an integer prime to m_+ . Defining $\vec{w}_\ell \equiv S \vec{n}_\ell$, we have

$$\vec{w}_1 = (n_-, 0), \quad \vec{w}_2 = (r_+, m_-), \quad \vec{w}_3 = (-n_+, 0), \quad \vec{w}_4 = (r_-, -m_+), \tag{4.27}$$

where $r_\pm \in \mathbb{Z}$ are chosen such that $t = r_+ m_+ + r_- m_-$, and exist by Bézout’s lemma¹³, for co-prime m_\pm . The vectors \vec{w}_ℓ now take values in \mathbb{Z}^2 , with primitive \vec{w}_2, \vec{w}_4 , hence $m_2 = m_4 = 1$, and non-primitive \vec{w}_1, \vec{w}_3 , corresponding to the labels $m_1 = n_-$ and $m_3 = n_+$. We can now define a new basis $\{E_1, E_2\}$, obtained as $E_i = S_{ji}^{-1} e_j$, such that

$$\xi_{(\ell)} = \vec{w}_\ell \cdot (E_1, E_2), \tag{4.28}$$

and the degenerating Killing vectors take the form

$$\begin{aligned} \xi_{(1)} &= n_- E_1, & \xi_{(2)} &= m_- E_2 + r_+ E_1, \\ \xi_{(3)} &= n_+ E_1, & \xi_{(4)} &= m_+ E_2 - r_- E_1. \end{aligned} \tag{4.29}$$

All the coefficients of this decomposition are now integers, indicating that in this basis the torus action is effective. For completeness we also write the basis vectors in terms of the Killing vectors ∂_{v_1} and ∂_{v_2} ¹⁴

$$E_1 = \partial_{v_2}, \quad E_2 = \partial_{v_1} + \frac{r_+ + r_-}{m_+ - m_-} \partial_{v_2}. \tag{4.30}$$

The combinatorial data can be presented in the form of a stacky fan, as depicted in Fig. 3a. Note that for $n_+ = n_- = 1$ this reduces to the ‘‘Hirzebruch orbifold’’ discussed in [54] in the context of toric stacks. We can then read off the order of the quotient singularities at the intersections of the four divisors, corresponding to the four cones τ_ℓ , which are given by $\mathbb{C}^2 / \mathbb{Z}_{\det(\vec{n}_\ell, \vec{n}_{\ell+1})}$ and thus read $\mathbb{C}^2 / \mathbb{Z}_{n_- m_-}, \mathbb{C}^2 / \mathbb{Z}_{n_+ m_+}, \mathbb{C}^2 / \mathbb{Z}_{n_+ m_+}, \mathbb{C}^2 / \mathbb{Z}_{n_- m_-}$, for $\ell = 1, \dots, 4$, respectively. From the toric description, we see that the total space $\Sigma_1 \times \Sigma_2$ is smooth not only away from the poles of Σ_2 , but also at the poles of Σ_1 , in agreement with [17].

Despite the fact that we do not have a moment map, it is of course possible to formally define a labeled polytope, encapsulating the same combinatorial data. This is obtained from the stacky fan in Fig. 3a, considering the vectors generating the rays of the fan and interpreting them as normal vectors to each facet of the polytope. The labels are read off from the ‘‘multiplicities’’ of the non-primitive vectors. The result is the rational convex polytope depicted in Fig. 3b, described by the labels $m_\ell = (n_-, 1, n_+, 1)$ and the primitive normal vectors $\vec{w}_\ell = \vec{w}_\ell / m_\ell$, which together are packed into what we shall refer to as toric data (\vec{w}_ℓ, m_ℓ) .

¹³ By Bézout’s lemma, if m_+ and m_- are co-prime, there exist $a_\pm \in \mathbb{Z}$ such that $a_+ m_+ + a_- m_- = 1$. Then we take $r_+ = t a_+$ and $r_- = t a_-$.

¹⁴ Notice that E_2 is finite also when $m_+ = m_- = 1$. Indeed, in this limit we have $r_+ + r_- = t$, which is proportional to $m_+ - m_-$ (cf. (2.59)), thus eliminating the vanishing denominator.




Fig. 3 Toric data of the $\Sigma_1 \times \Sigma_2$ orbifold corresponding to the vectors (4.27). In this example, we took $n_+ = 5, n_- = 4, m_+ = 3, m_- = 2, r_+ = -1, r_- = 1$ and $t = -1$

4.3 Toric divisors and their intersection

In this section, we will discuss toric divisors with the aim of computing their intersection matrix, that will be a key ingredient in the construction of the entropy function. We will begin recalling some basic facts about divisors in the context of toric varieties (thus, ignoring the labels) and then we will indicate the modifications needed to pass to the broader setting in which our toric orbifolds fit. Recall that all the toric varieties defined by fans are normal and therefore all the (orbifold) singularities are in complex co-dimension higher than one. For toric surfaces \mathbb{M}_4 (i.e., in real dimension 4), this implies that only points can be singular, while there cannot be singularities along divisors. Divisors that are torus-invariant are called *toric divisors* and are in a one-to-one correspondence with the primitive generators of the rays of the fan. These are in general Weil divisors, but for toric surfaces, since all the fans are necessarily simplicial, all Weil divisors D have an integer multiple ID that is a Cartier divisor—namely they are \mathbb{Q} -Cartier divisors¹⁵. Below, we will consider only such \mathbb{Q} -Cartier toric divisors, associated with the primitive generators of a fan, \vec{v}_ℓ , that will be denoted as D_ℓ . As in the previous sections, $\ell = 1, \dots, n$ runs over the generators of the fan in cyclic order. Furthermore, associated with any toric divisor D_ℓ there is an equivariant line bundle L_ℓ , whose first Chern class $c_1(L_\ell) \in H^2(\mathbb{M}_4, \mathbb{Q})$ is Poincaré dual to D_ℓ . While the second co-homology group of \mathbb{M}_4 , $H^2(\mathbb{M}_4, \mathbb{Q})$, is $(n - 2)$ -dimensional, its equivariant extension, generated by the $c_1(L_\ell)$, has clearly dimension n .

The intersection matrix of the toric divisors is defined as

$$D_{\ell, \ell'} \equiv D_\ell \cdot D_{\ell'} = \int_{\mathbb{M}_4} c_1(L_\ell) \wedge c_1(L_{\ell'}), \tag{4.31}$$

¹⁵ Varieties with such a property are called \mathbb{Q} -factorial.

and by a standard calculation in toric geometry, it reads

$$D_{\ell,\ell'}[\vec{v}_\ell] = D_{\ell,\ell'} = \begin{cases} \frac{1}{\det(\vec{v}_{\ell-1}, \vec{v}_\ell)} & \text{if } \ell' = \ell - 1, \\ \frac{1}{\det(\vec{v}_\ell, \vec{v}_{\ell+1})} & \text{if } \ell' = \ell + 1, \\ -\frac{\det(\vec{v}_{\ell-1}, \vec{v}_{\ell+1})}{\det(\vec{v}_{\ell-1}, \vec{v}_\ell) \det(\vec{v}_\ell, \vec{v}_{\ell+1})} & \text{if } \ell' = \ell, \\ 0 & \text{otherwise,} \end{cases} \tag{4.32}$$

where we used the notation $D_{\ell,\ell'}[\vec{v}_\ell]$ to emphasize that it is the intersection matrix of the toric divisors associated with a fan with generators \vec{v}_ℓ . It can be easily proven that this satisfies the relation

$$\sum_{\ell'} D_{\ell,\ell'} \vec{v}_{\ell'} = 0, \tag{4.33}$$

implying that $D_{\ell,\ell'}$, viewed as an $n \times n$ matrix, has two null eigenvalues, and it is therefore not invertible. This of course is consistent with the fact that $\dim(H^2(\mathbb{M}_4, \mathbb{Q})) = n - 2$ implies that only $n - 2$ toric divisors D_ℓ are independent in homology.

For the special case of toric manifolds, all $\det(\vec{v}_\ell, \vec{v}_{\ell+1}) = 1$, and the above discussion reduces to that in [36]. On the other hand, for our purposes we need to incorporate the effect of the labels. We will refrain from attempting a rigorous treatment, indicating a natural way to take into account this additional information. If we have a labeled polytope/stacky fan described by “long” vectors $\vec{\hat{v}}_\ell = m_\ell \vec{v}_\ell$, it is natural to associate a divisor \hat{D}_ℓ to $\vec{\hat{v}}_\ell$, which is related to D_ℓ simply by

$$\hat{D}_\ell = \frac{1}{m_\ell} D_\ell. \tag{4.34}$$

In particular, this implies that for any two-form Λ the integrals over the two sets of divisors are related as

$$m_\ell \int_{\hat{D}_\ell} \Lambda = \int_{D_\ell} \Lambda, \tag{4.35}$$

and the intersection matrix of the divisors \hat{D}_ℓ is given by

$$\hat{D}_{\ell,\ell'} \equiv D_{\ell,\ell'}[\vec{\hat{v}}_\ell] = \frac{1}{m_\ell m_{\ell'}} D_{\ell,\ell'}[\vec{v}_\ell]. \tag{4.36}$$

Correspondingly, the first Chern classes of the dual line bundles are related as $c_1(L_\ell) = m_\ell c_1(\hat{L}_\ell)$. Since we can use either $c_1(L_\ell)$ or $c_1(\hat{L}_\ell)$ as basis for $H^2(\mathbb{M}_4, \mathbb{Q})$, these two descriptions are essentially interchangeable.

We note that $\hat{D}_{\ell, \ell'}$ is invariant under $SL(2, \mathbb{Z})$ transformations of the basis and more generally for any transformation in $SL(2, \mathbb{R})$. This follows immediately from the fact that given any two vectors $\vec{v}_1, \vec{v}_2 \in \mathbb{R}^2$, $\det(\vec{v}_1, \vec{v}_2)$ is invariant under $SL(2, \mathbb{R})$ transformations of the two vectors. In particular, taking a generic matrix $S \in SL(2, \mathbb{R})$ and transforming the vectors as $\vec{v}'_a = S \vec{v}_a$, we have that S acts on the matrix (\vec{v}_1, \vec{v}_2) simply by matrix multiplication, i.e., $(\vec{v}'_1, \vec{v}'_2) = S(\vec{v}_1, \vec{v}_2)$, therefore

$$\det(\vec{v}'_1, \vec{v}'_2) = \det(S) \det(\vec{v}_1, \vec{v}_2) = \det(\vec{v}_1, \vec{v}_2). \tag{4.37}$$

This property implies that formally the intersection matrix of the divisors computed in the two bases \vec{w}_ℓ and \vec{n}_ℓ are identical, being related by the $SL(2, \mathbb{Q})$ transformation (4.26).

In order to make contact with the computations in supergravity, we need to describe the relation between the toric divisors defined above and the representative of the cycles, that we used to integrate fluxes in our main example of $\Sigma_1 \times \Sigma_2$. The S_\pm introduced in Sect. 2.2 were precisely two copies of the base Σ_1 at the south pole (section at infinity) and north pole (zero section) of the fiber Σ_2 , and we can identify these with the toric divisors D_1 and D_3 as

$$D_1 = S_-, \quad D_3 = S_+. \tag{4.38}$$

On the other hand, D_2 and D_4 correspond to copies of the fiber Σ_2 at the poles of the base Σ_1 . In the smooth case ($m_+ = m_- = 1$), these would be simply homologous to Σ_2 . However, the orbifold singularities of the base imply that instead

$$D_2 = X_-, \quad D_4 = X_+, \tag{4.39}$$

where $X_\pm \equiv \Sigma_2 / \mathbb{Z}_{m_\pm}$, respectively. Therefore, in order to compare the integral of a two-form Λ over X_\pm with the integral over Σ_2 at a generic point of Σ_1 we should use the relation

$$m_\pm \int_{X_\pm} \Lambda = \int_{\Sigma_2} \Lambda. \tag{4.40}$$

Since in Sect. 2.2 the fluxes were computed integrating on Σ_2 (see, e.g., (2.51)), the integrals computed using X_\pm will have to be multiplied by a factor of m_\pm to be compared with the fluxes computed in Sect. 2.2.

5 The entropy function

We conjecture that the solutions presented in Sect. 2.2 are holographically dual to one-dimensional SCQMs obtained compactifying on a $\Sigma_1 \times \Sigma_2$ the five-dimensional SCFTs dual to the solution of [12]. In this section, we provide supporting evidence for this conjecture, proposing an entropy function, whose extremization reproduces the entropy (3.30) of the (putative) black hole near-horizon $AdS_2 \times \Sigma_1 \times \Sigma_2$. Starting from

first principles, this function should be derived from the localized partition function of the $d = 5$ SCFT, placed on the background of $S^1 \times \Sigma_1 \times \Sigma_2$, and then taking the large N limit. Following the idea of “gravitational blocks” [37], we will propose a large N entropy function on $S^1 \times \Sigma_1 \times \Sigma_2$ obtained by suitably gluing the S^5 free energy of the $d = 5$ theories. With some further assumptions, this prescription can be applied also to the $\text{AdS}_2 \times \Sigma_g \times \Sigma_2$ solutions of Sect. 2.1. The prescription that we propose extends the results of [27] to a broader class of configurations, including solutions arising in $D = 7$ supergravity.

5.1 Twisting data

In this section, we consider the compactification of five- and six-dimensional SCFTs on a four-dimensional toric orbifold \mathbb{M}_4 . Specifically, we place the theory on \mathbb{M}_4 and couple it to two background gauge fields A_i , with appropriately quantized magnetic fluxes. All the information about \mathbb{M}_4 is encoded in the toric data (\vec{v}_ℓ, m_ℓ) , which can be thought of as the data defining a labeled polytope. The vectors \vec{v}_ℓ must be primitive, \mathbb{Z}^2 -valued and, in our conventions, ordered counterclockwise. The recipe we will formulate is in principle applicable to toric orbifolds (including smooth toric manifolds) with an arbitrary number $n \geq 3$ of fixed points.

Consider the two line bundles E_i on which the one-forms $-A_i$ are the connections. Since the set of $c_1(L_\ell)$ form a basis for the equivariant extension of $H^2(\mathbb{M}_4, \mathbb{Q})$, we can decompose the first Chern class of E_i as

$$c_1(E_i) = -\frac{dA_i}{2\pi} = -\sum_{\ell} \mathfrak{p}_i^{(\ell)} c_1(L_\ell), \tag{5.1}$$

where $\mathfrak{p}_i^{(\ell)} \in \mathbb{Q}$. The “physical fluxes” are defined as¹⁶

$$\mathfrak{q}_i^{(\ell)} \equiv \frac{1}{2\pi} \int_{D_\ell} F_i = -\int_{D_\ell} c_1(E_i), \tag{5.2}$$

and using (5.1) we obtain the relation

$$\mathfrak{q}_i^{(\ell)} = \sum_{\ell'} D_{\ell, \ell'} \mathfrak{p}_i^{(\ell')}, \tag{5.3}$$

where $D_{\ell, \ell'}$ is the intersection matrix defined in (4.32). We recall that $H^2(\mathbb{M}_4, \mathbb{Q})$ has dimension $n - 2$, thus the whole set of $\mathfrak{p}_i^{(\ell)}$ is a redundant parameterization. In particular, for fixed i , only $n - 2$ of the $\mathfrak{p}_i^{(\ell)}$ are linearly independent. The two additional degrees of freedom are the expression of a gauge symmetry that we will discuss in the next subsection.

¹⁶ Here and in the following, we shall rename the background gauge fields as $gA_i \mapsto A_i$, which is more natural from the field theory point of view.

A further consequence of relation (4.33) is that, combined with (5.3), it gives

$$\sum_{\ell} q_i^{(\ell)} \vec{v}_{\ell} = 0. \tag{5.4}$$

This is a vector equation and, for fixed i , we have n fluxes $q_i^{(\ell)}$ constrained by two equations, one for each component of the two-dimensional vectors \vec{v}_{ℓ} . Therefore, only $n - 2$ physical fluxes are linearly independent. In what follows, given a generic toric orbifold with arbitrary fluxes, we shall require them to satisfy the constraint (5.4). We can now go back to relation (5.3). As we explained, it represents a linear system of $\text{rank}(D_{\ell,\ell'}) = n - 2$ independent equations for $n - 2$ independent unknowns $p_i^{(\ell)}$. Eliminating the redundant equations, we can solve this system and obtain $p_i^{(\ell)}$ in terms of $q_i^{(\ell)}$, up to gauge transformations.

Equation (5.4) imposes a similar condition on the R -symmetry fluxes $q_R^{(\ell)} \equiv q_1^{(\ell)} + q_2^{(\ell)}$, specifically $\sum_{\ell} q_R^{(\ell)} \vec{v}_{\ell} = 0$. This constraint can be solved writing

$$q_R^{(\ell)} = \sum_{\ell'} D_{\ell,\ell'} \frac{\sigma^{(\ell')}}{m_{\ell'}}, \tag{5.5}$$

where m_{ℓ} are the labels of the labeled polytope associated with \mathbb{M}_4 and, a priori, $\sigma^{(\ell)}$ are n arbitrary coefficients. Denoting as E_R the R -symmetry (orbifold) line bundle, Eq. (5.5) can be rewritten as

$$c_1(E_R) = - \sum_{\ell} \sigma^{(\ell)} c_1(\hat{L}_{\ell}), \tag{5.6}$$

or, following standard practice, as $c_1(E_R) = - \sum_{\ell} \sigma^{(\ell)} \hat{D}_{\ell}$. Note that the standard topological twist corresponds to identifying E_R with the orbifold canonical line bundle $K_{\mathbb{M}_4}^{\text{orb}} = - \sum_{\ell} \hat{D}_{\ell}$ and hence $\sigma_{\text{top-twist}}^{(\ell)} = (+, \dots, +)$. See, e.g., [46] for a related discussion of orbifold line bundles.

In analogy with the case of the spindle [31], we conjecture that the only possible values of $\sigma^{(\ell)}$ are ± 1 . This hypothesis is supported by the different examples analyzed, as we shall see. In other words, we expect that it should be possible to compactify a SCFT on a toric orbifold \mathbb{M}_4 , turning on a background R -symmetry gauge field with magnetic fluxes given in (5.5), with $\sigma^{(\ell)}$ parameterizing the different supersymmetry-preserving twists. Of course, we have not proven that all these different twists preserve supersymmetry, nor that there cannot exist more general twists. It would be interesting to carry out such an analysis extending the results of [31], where it was demonstrated that, in whole generality, on a spindle the only two supersymmetry-preserving twists are the twist and the anti-twist.

5.2 The recipe

We conjecture that given a general class of SCFTs in $d = 5, 6$ compactified on a four-dimensional toric orbifold \mathbb{M}_4 , with an arbitrary twist parameterized by a set of signs $\sigma^{(\ell)}$ as in (5.5), the corresponding entropy/central charge, respectively, is determined extremizing the following *off-shell free energy*

$$F(\varphi_i, \epsilon_i; \mathfrak{q}_i^{(\ell)}, m_\ell) = k_d \sum_{\ell} \frac{\eta_d^{(\ell)} \mathcal{F}_d(\Phi_i^{(\ell)})}{d_{\ell, \ell+1} \epsilon_1^{(\ell)} \epsilon_2^{(\ell)}}. \tag{5.7}$$

The sum runs over the n fixed points v_ℓ of the toric orbifold \mathbb{M}_4 , k_d is a numerical constant, which, a posteriori, takes the values

$$k_5 = -1, \quad k_6 = \frac{64}{9}, \tag{5.8}$$

$d_{\ell, \ell+1}$ is defined as $d_{\ell, \ell'} \equiv d_{\ell, \ell'}[\vec{v}_\ell] = \det(\vec{v}_\ell, \vec{v}_{\ell'})$ and \mathcal{F}_d are the usual gravitational blocks (cf. table 2 of [27])

$$\mathcal{F}_5(\Delta_i) = -\frac{4\sqrt{2}\pi N^{5/2}}{15\sqrt{8 - N_f}} (\Delta_1 \Delta_2)^{3/2}, \quad \mathcal{F}_6(\Delta_i) = -\frac{9N^3}{256} (\Delta_1 \Delta_2)^2. \tag{5.9}$$

The variables $\Phi_i^{(\ell)}$ are defined as

$$\Phi_i^{(\ell)} = \varphi_i - \mathfrak{p}_i^{(\ell)} \epsilon_1^{(\ell)} - \mathfrak{p}_i^{(\ell+1)} \epsilon_2^{(\ell)}, \tag{5.10}$$

and the auxiliary quantities $\epsilon_1^{(\ell)}$ and $\epsilon_2^{(\ell)}$ read

$$\epsilon_1^{(\ell)} = -\frac{\det(\vec{v}_{\ell+1}, \vec{\epsilon})}{d_{\ell, \ell+1}}, \quad \epsilon_2^{(\ell)} = \frac{\det(\vec{v}_\ell, \vec{\epsilon})}{d_{\ell, \ell+1}}, \tag{5.11}$$

with $\vec{\epsilon} = (\epsilon_1, \epsilon_2)^{17}$. Here, ϵ_1 and ϵ_2 may be interpreted as the fugacities associated with the two $U(1)$ rotational symmetries and parameterize their mixing with the R -symmetry. $\eta_d^{(\ell)} = \pm$ are signs that at this stage must be tuned by hand: in particular we believe that in general¹⁸ in $d = 6$ they are all +, while for $d = 5$ they are related to the type of twist and we speculate that $\eta_5^{(\ell)} = \sigma^{(\ell)} \sigma^{(\ell+1)}$. In the next section we will see some explicit examples applying this prescription to the $\text{AdS}_2 \times \mathbb{S}_1 \times \mathbb{S}_2$ family. In this construction, φ_i and ϵ_i are variables with respect to which one has to extremize $F(\varphi_i, \epsilon_i)$, \vec{v}_ℓ and m_ℓ are data describing the toric orbifold \mathbb{M}_4 and $\mathfrak{p}_i^{(\ell)}$ are related to the physical fluxes $\mathfrak{q}_i^{(\ell)}$ through (5.3).

¹⁷ The two variables $\epsilon_{1,2}$ are different from the ones adopted in [27]. Specifically, $\epsilon_i^{\text{here}} = 2\epsilon_i^{\text{there}}$.

¹⁸ Due to different conventions they could also be all $-$, as explained at the end of Sect. 5.4.

The variables φ_i and ϵ_i are subject to the constraint

$$\varphi_1 + \varphi_2 - \det(\vec{W}, \vec{\epsilon}) = 2, \quad (5.12)$$

where \vec{W} is a two-dimensional constant vector parameterizing a “gauge invariance” of the problem, that we discuss below. This condition is inherited from the R -symmetry constraint $\Delta_1 + \Delta_2 = 2$, where Δ_i are the fugacities parameterizing the R -symmetry within the Cartan subgroup of the global symmetries of the d -dimensional parent theory. The vector \vec{W} can be determined imposing

$$\vec{p}_1^{(\ell)} + \vec{p}_2^{(\ell)} = \frac{\sigma^{(\ell)}}{m_\ell} + \det(\vec{W}, \vec{v}_\ell), \quad (5.13)$$

which is a system of n equations – one for each value of ℓ – for the two components of \vec{W} , and it is thus overdetermined. Yet, a solution always exists. By means of Eqs. (5.3) and (5.13) we can write the following chain of equalities

$$\begin{aligned} \mathfrak{q}_R^{(\ell)} &= \sum_{\ell'} D_{\ell, \ell'} (\vec{p}_1^{(\ell')} + \vec{p}_2^{(\ell')}) = \sum_{\ell'} D_{\ell, \ell'} \frac{\sigma^{(\ell')}}{m_{\ell'}} + \det(\vec{W}, \sum_{\ell'} D_{\ell, \ell'} \vec{v}_{\ell'}) \\ &= \sum_{\ell'} D_{\ell, \ell'} \frac{\sigma^{(\ell')}}{m_{\ell'}}, \end{aligned} \quad (5.14)$$

where, in the last step, we made use of (4.33). First of all, this relation proves the consistency of (5.13), since, as it should be, the physical quantities $\mathfrak{q}_R^{(\ell)}$ do not depend on the auxiliary vector \vec{W} . Moreover, it shows that $n - 2$ independent linear combinations of Eqs. (5.13), obtained contracting them with $D_{\ell, \ell'}$, are already satisfied once (5.5) is imposed. The result is a linear system of two independent equations with two unknowns, that can always be solved.

As we already mentioned, the quantity $\det(\vec{v}_1, \vec{v}_2)$ is invariant under $SL(2, \mathbb{R})$ transformations of the two vectors $\vec{v}_1, \vec{v}_2 \in \mathbb{R}^2$. This property is fundamental for the consistency of our construction. A given toric orbifold can be described by an infinite number of polytopes equivalent under $SL(2, \mathbb{Z})$ transformations of the vectors \vec{v}_ℓ . These transformations are generated by rotations of the basis vectors (e_1, e_2) that give rise to effective torus actions. On the other hand, we expect the free energy not to be affected by these transformations and, indeed, this is the case. If we perform an $SL(2, \mathbb{Z})$ rotation of the vectors \vec{v}_ℓ , the expression of the intersection matrix $D_{\ell, \ell'}$ is retained, and the explicit form of the off-shell free energy is not modified provided we apply exactly the same transformation to $\vec{\epsilon}$ and \vec{W} .

The off-shell entropy function we constructed enjoys also another symmetry, the gauge symmetry we mentioned before. Thanks to (4.33), relation (5.3) is left invariant under the “gauge transformation”

$$\tilde{\mathfrak{p}}_i^{(\ell)} = \mathfrak{p}_i^{(\ell)} + \det(\vec{\lambda}_i, \vec{v}_\ell), \quad (5.15)$$

for any two-dimensional constant vector $\vec{\lambda}_i$. This transformation affects the value of the vector \vec{W} appearing in the constraint. Specifically, since $\sigma^{(\ell)}$ and m_ℓ are fixed, we must impose

$$\tilde{p}_1^{(\ell)} + \tilde{p}_2^{(\ell)} - \det(\vec{W}, \vec{v}_\ell) = p_1^{(\ell)} + p_2^{(\ell)} - \det(\vec{W}, \vec{v}_\ell), \tag{5.16}$$

which returns $\vec{W} = \vec{W} + \vec{\lambda}_1 + \vec{\lambda}_2$. Introducing the new variables $\tilde{\varphi}_i = \varphi_i + \det(\vec{\lambda}_i, \vec{e})$ it is possible to keep the functional expression of $\Phi_i^{(\ell)}$ unmodified, i.e., $\Phi_i^{(\ell)}(\tilde{\varphi}, \tilde{p}) = \Phi_i^{(\ell)}(\varphi, p)$. Remarkably, the constraint changes accordingly

$$\tilde{\varphi}_1 + \tilde{\varphi}_2 - \det(\vec{W}, \vec{e}) = \varphi_1 + \varphi_2 - \det(\vec{W}, \vec{e}) = 2. \tag{5.17}$$

As a result, both the off-shell free energy and the constraint retain the same functional expression under the gauge transformation (5.15), thus the value of the former at its critical points will be the same.

The proposed prescription takes inspiration from [36] and extends their construction to compactifications on generic toric orbifolds. Moreover, it also applies to $d = 5$, generalizing the formulas presented in [27] for compactifications on $\Sigma_{g_1} \times \Sigma_{g_2}$ and $\Sigma \times \Sigma_g$, to the case of fibered spaces. The constrained extremization of $F(\varphi_i, \epsilon_i)$ can be performed defining the function

$$\mathcal{S}(\varphi_i, \epsilon_i, \Lambda; q_i^{(\ell)}, m_\ell) = F(\varphi_i, \epsilon_i; q_i^{(\ell)}, m_\ell) + \Lambda(\varphi_1 + \varphi_2 - \det(\vec{W}, \vec{e}) - 2) \tag{5.18}$$

and extremizing it with respect to φ_i, ϵ_i and the Lagrangian multiplier Λ . In order to solve this problem, it might be helpful to notice that (5.7) is homogeneous of degree h in φ_i and ϵ_i ($h = 1$ for $d = 5$, $h = 2$ for $d = 6$). As a consequence, by Euler’s theorem, we have $F(\varphi_i^*, \epsilon_i^*) = -\frac{2}{h} \Lambda^*$.

5.3 Application to the $\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solutions

The first example to which we can apply the prescription presented in the previous section is the $\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solution (2.40). We now briefly collect some of the results needed in the construction. The toric orbifold $\mathbb{M}_4 = \Sigma_1 \times \Sigma_2$ can be described by the toric data (see Sect. 4.2)

$$\begin{aligned} m_1 = n_-, & \quad \vec{w}_1 = (1, 0), & m_2 = 1, & \quad \vec{w}_2 = (r_+, m_-), \\ m_3 = n_+, & \quad \vec{w}_3 = (-1, 0), & m_4 = 1, & \quad \vec{w}_4 = (r_-, -m_+), \end{aligned} \tag{5.19}$$

with $t = r_+ m_+ + r_- m_-$. The vectors \vec{w}_ℓ are integers, primitive and ordered counterclockwise, as required by the recipe. Keeping in mind all the observations about

divisors made in Sect. 4.3, the physical fluxes defined in (5.2) read

$$\begin{aligned} q_i^{(1)} &= \frac{g}{2\pi} \int_{S_-} F_i = s_i^-, & q_i^{(2)} &= \frac{g}{2\pi} \int_{X_-} F_i = \frac{g}{2\pi} \frac{1}{m_-} \int_{\mathbb{Z}_2} F_i = \frac{t_i}{m_-}, \\ q_i^{(3)} &= \frac{g}{2\pi} \int_{S_+} F_i = s_i^+, & q_i^{(4)} &= \frac{g}{2\pi} \int_{X_+} F_i = \frac{g}{2\pi} \frac{1}{m_+} \int_{\mathbb{Z}_2} F_i = \frac{t_i}{m_+}, \end{aligned} \tag{5.20}$$

where s_i^+, s_i^- and t_i are given, respectively, in (2.61), (2.60) and (2.28). The intersection matrix $D_{\ell, \ell'}$, computed straightforwardly using (4.32), reads

$$D_{\ell, \ell'} = \begin{pmatrix} -\frac{t}{m_+ m_-} & \frac{1}{m_-} & 0 & \frac{1}{m_+} \\ \frac{1}{m_-} & 0 & \frac{1}{m_-} & 0 \\ 0 & \frac{1}{m_-} & \frac{t}{m_+ m_-} & \frac{1}{m_+} \\ \frac{1}{m_+} & 0 & \frac{1}{m_+} & 0 \end{pmatrix}, \tag{5.21}$$

and imposing (5.5) we obtain the vector of twists

$$\sigma^{(\ell)} = (+, +, +, -). \tag{5.22}$$

Taking advantage of the gauge symmetry, we impose $p_i^{(1)} = p_i^{(3)}$ and $p_i^{(2)}/m_- = p_i^{(4)}/m_+$. Therefore, the relation (5.3) yields the identifications

$$p_i^{(1)} = p_i^{(3)} = \frac{t_i}{2}, \quad p_i^{(2)} = \frac{m_- s_i^*}{2}, \quad p_i^{(4)} = \frac{m_+ s_i^*}{2}, \tag{5.23}$$

where we defined $s_i^* \equiv \frac{s_i^+ + s_i^-}{2}$. In particular, s_1^* explicitly reads

$$s_1^* = \frac{t\chi_2}{m_+ m_-} \frac{x^3 - 3\mu x^2 + (3 - 2z)x - 3\mu}{8x^2}, \tag{5.24}$$

and s_2^* can be obtained flipping the sign of z . With this definition, s_i^\pm can be written as

$$s_i^\pm = s_i^* \pm \frac{t}{2m_+ m_-} t_i. \tag{5.25}$$

System (5.13) can be solved to get \bar{W} , which gives the constraint (5.12)

$$\varphi_1 + \varphi_2 + \frac{n_+ - n_-}{2n_+ n_-} \epsilon_1 + \left(\frac{m_+ + m_-}{2m_+ m_-} - \frac{r_+ m_+ - r_- m_-}{m_+ m_-} \frac{n_+ - n_-}{4n_+ n_-} \right) \epsilon_2 = 2. \tag{5.26}$$

The ingredients $\epsilon_{1,2}^{(\ell)}$ and $\Phi_i^{(\ell)}$ can be constructed by means of (5.11) and (5.10), whereas the off-shell free energy (5.7) reads

$$\begin{aligned}
 F(\varphi_i, \epsilon_i; \mathfrak{s}_i^\pm, \mathfrak{t}_i, m_\ell) &= -\frac{\mathcal{F}_5(\Phi_i^{(1)})}{d_{1,2} \epsilon_1^{(1)} \epsilon_2^{(1)}} - \frac{\mathcal{F}_5(\Phi_i^{(2)})}{d_{2,3} \epsilon_1^{(2)} \epsilon_2^{(2)}} + \frac{\mathcal{F}_5(\Phi_i^{(3)})}{d_{3,4} \epsilon_1^{(3)} \epsilon_2^{(3)}} + \frac{\mathcal{F}_5(\Phi_i^{(4)})}{d_{4,1} \epsilon_1^{(4)} \epsilon_2^{(4)}},
 \end{aligned}
 \tag{5.27}$$

where the relative signs have been fixed using the prescription $\eta_5^{(\ell)} = \sigma^{(\ell)} \sigma^{(\ell+1)}$.

We can now proceed to the extremization of the off-shell free energy and, after some work, we get the critical values

$$\begin{aligned}
 \varphi_i^* &= \frac{2m_+m_-}{m_+ - m_-} \left(1 + \eta \frac{m_+ + m_-}{2m_+m_-} \frac{2\pi}{\Delta\psi} \right) \mathfrak{s}_i^*, \\
 \epsilon_1^* &= -\frac{1}{m} \frac{2\pi}{\Delta z} + \eta \frac{2(r_+ + r_-)}{m_+ - m_-} \frac{2\pi}{\Delta\psi}, \quad \epsilon_2^* = -2\eta \frac{2\pi}{\Delta\psi},
 \end{aligned}
 \tag{5.28}$$

where the sign ambiguity $\eta = \pm 1$ arises by solving the equations over the complex numbers¹⁹. Inserting these values back into (5.27), we obtain

$$F(\varphi_i^*, \epsilon_i^*; \mathfrak{s}_i^\pm, \mathfrak{t}_i, m_\ell) = \frac{-\eta\sqrt{2}\sqrt{m_+^2 + m_-^2} - (m_+ + m_-)}{2m_+m_-} F_{S^3 \times \Sigma_2}, \tag{5.29}$$

where $F_{S^3 \times \Sigma_2}$ is given in (3.24). In order to get a positive entropy, we need to pick $\eta = -1$ and in this case the result agrees with the gravitational entropy (3.30).

We propose that the procedure we formulated can be extended also to non-toric four-dimensional orbifolds, e.g., in the presence of Riemann surfaces as submanifold. In this case, we can adopt the same toric data obtained replacing Σ_g ($g > 1$) with S^2 , setting $g = 0$ whenever it occurs in (5.13) and (5.5). The genus is preserved in the explicit expression of the fluxes. Specifically, we will test our conjecture against the $AdS_2 \times \Sigma_g \times \Sigma_2$ system (2.9). The ‘‘toric data’’ are given in (4.11) and (4.9), the physical fluxes are the same as in the previous example, with $m_\pm = 1$, and now \mathfrak{s}_i^\pm can be read in (2.38) and (2.37). The fluxes $p_i^{(\ell)}$ are identified as before and \mathfrak{s}_i^* retains the same form, but, again, with $m_\pm = 1$ and t as in (2.36). The constraint on the variables φ_i and ϵ_i is

$$\varphi_1 + \varphi_2 + \frac{n_+ - n_-}{2n_+n_-} \epsilon_1 - t \frac{n_+ - n_-}{4n_+n_-} \epsilon_2 = 2, \tag{5.30}$$

¹⁹ Generically, in the presence of rotation we have to work with the complex numbers [16]; therefore, we continue to do so also in the static case.

whereas the off-shell free energy reads

$$F(\varphi_i, \epsilon_i; \mathfrak{s}_i^\pm, \mathfrak{t}_i, m_\ell) = \frac{\mathcal{F}_5(\Phi_i^{(1)})}{d_{1,2} \epsilon_1^{(1)} \epsilon_2^{(1)}} + \frac{\mathcal{F}_5(\Phi_i^{(2)})}{d_{2,3} \epsilon_1^{(2)} \epsilon_2^{(2)}} + \frac{\mathcal{F}_5(\Phi_i^{(3)})}{d_{3,4} \epsilon_1^{(3)} \epsilon_2^{(3)}} + \frac{\mathcal{F}_5(\Phi_i^{(4)})}{d_{4,1} \epsilon_1^{(4)} \epsilon_2^{(4)}}, \tag{5.31}$$

where again the relative signs were chosen to be $\eta_5^{(\ell)} = \sigma^{(\ell)} \sigma^{(\ell+1)}$. These different gluing signs are due to the different types of twist that we have. In particular, on both Σ_g and Σ_2 the twist is realized, in contrast to the previous case, in which we had anti-twist on Σ_1 . The off-shell free energy is extremized by

$$\varphi_i^* = \frac{\mathfrak{s}_i^*}{g-1}, \quad \epsilon_1^* = -\frac{1}{m} \frac{2\pi}{\Delta z}, \quad \epsilon_2^* = 0, \tag{5.32}$$

to which corresponds the critical value

$$F(\varphi_i^*, \epsilon_i^*; \mathfrak{s}_i^\pm, \mathfrak{t}_i, m_\ell) = (g-1) F_{S^3 \times \Sigma_2}. \tag{5.33}$$

Also in this case, the result returned by the extremization agrees with the entropy computed from the ten-dimensional supergravity solution (3.23).

5.4 Application to the $AdS_3 \times \Sigma_1 \times \Sigma_2$ solutions

The conjectured prescription presented at the beginning of this section can also be applied to another interesting family, the $AdS_3 \times \Sigma_1 \times \Sigma_2$ solutions constructed in [17]²⁰. The toric data of the $\Sigma_1 \times \Sigma_2$ toric orbifold are derived in Appendix C.3 and we recall them here for the reader’s convenience

$$\begin{aligned} m_1 = n_-, \quad \vec{w}_1 = (1, 0), \quad m_2 = 1, \quad \vec{w}_2 = (r_+, m_-), \\ m_3 = n_+, \quad \vec{w}_3 = (-1, 0), \quad m_4 = 1, \quad \vec{w}_4 = (r_-, -m_+), \end{aligned} \tag{5.34}$$

with $t = r_+ m_+ + r_- m_-$. Global conditions lead to

$$\begin{aligned} t &= \frac{-6(m_+ - m_-)n_+n_-(n_+ - n_-)}{[s - (p_1 + p_2)][s + 2(p_1 + p_2)]}, \\ s &= \sqrt{7(p_1^2 + p_2^2) + 2p_1p_2 - 6(n_+^2 + n_-^2)}, \end{aligned} \tag{5.35}$$

where p_1 and p_2 are two integers related to the magnetic fluxes as in (2.11) of [17]. The physical fluxes can be computed as in the previous example and, compactly, they read

$$\mathfrak{q}_i^{(\ell)} = \left(\mathfrak{s}_i^-, \frac{\mathfrak{t}_i}{m_-}, \mathfrak{s}_i^+, \frac{\mathfrak{t}_i}{m_+} \right), \tag{5.36}$$

²⁰ In order to make the analogy with the $AdS_2 \times \Sigma_1 \times \Sigma_2$ systems more manifest, we exchanged $n_+ \leftrightarrow n_-$ and $m_+ \leftrightarrow m_-$ with respect to [17].

where s_1^- corresponds to the flux computed in (3.15) of [17], s_2^- and s_i^+ can be derived from the former and $t_i = \frac{p_i}{n_+n_-}$. The gauge symmetry allows us to impose $p_i^{(1)} = p_i^{(3)}$ and $p_i^{(2)}/m_- = p_i^{(4)}/m_+$, which yields, through (5.3), the identifications

$$p_i^{(1)} = p_i^{(3)} = \frac{t_i}{2}, \quad p_i^{(2)} = \frac{m_- s_i^*}{2}, \quad p_i^{(4)} = \frac{m_+ s_i^*}{2}. \tag{5.37}$$

For later convenience we defined $s_1^* \equiv \frac{s_1^+ + s_1^-}{2}$, whose expression is

$$s_i^* = \frac{tp_i[-6p_i + 4(n_+ + n_-) - s]}{6m_+m_-n_+n_-(n_+ - n_-)}. \tag{5.38}$$

The fluxes s_i^\pm are related to s_i^* through

$$s_i^\pm = s_i^* \pm \frac{t}{2m_+m_-} t_i. \tag{5.39}$$

The constraint can be derived following the path traced in the previous case and reads

$$\varphi_1 + \varphi_2 + \frac{n_+ - n_-}{2n_+n_-} \epsilon_1 + \left(\frac{m_+ + m_-}{2m_+m_-} - \frac{r_+m_+ - r_-m_-}{m_+m_-} \frac{n_+ - n_-}{4n_+n_-} \right) \epsilon_2 = 2. \tag{5.40}$$

Notice that this expression is identical to (5.26). As we discussed in Sect. 5.2, in $d = 6$ we set $\eta_6^{(\ell)} = +$ for all ℓ . Remarkably, this implies that F is *quadratic* in the φ_i , as expected for the off-shell central charge of a two-dimensional SCFT [8]. Explicitly, this reads

$$\begin{aligned} & F(\varphi_i, \epsilon_i; s_i^\pm, t_i, m_\ell) \\ &= - \left[\frac{t_2 s_1^* + t_1 s_2^*}{8} \left(t_1 t_2 \epsilon_1^2 + \frac{(r_+^2 m_+^2 + r_-^2 m_-^2) t_1 t_2 + 2m_+^2 m_-^2 s_1^* s_2^*}{2m_+^2 m_-^2} \epsilon_2^2 \right. \right. \\ &\quad \left. \left. - \frac{r_+ m_+ - r_- m_-}{m_+ m_-} t_1 t_2 \epsilon_1 \epsilon_2 \right) + \frac{t_2 s_2^* \varphi_1^2 + t_1 s_1^* \varphi_2^2}{2} + (t_2 s_1^* + t_1 s_2^*) \varphi_1 \varphi_2 \right. \\ &\quad \left. + \frac{2m_+ m_- \epsilon_1 - (r_+ m_+ - r_- m_-) \epsilon_2}{8m_+^2 m_-^2} t_1 t_2 (t_2 \varphi_1 + t_1 \varphi_2) \right] N^3. \tag{5.41} \end{aligned}$$

The extremization procedure, realized by means of Lagrangian multipliers, gives the critical values²¹

²¹ We notice that the expression of φ_i^* , written in this seemingly cumbersome way, is identical to the corresponding quantity of the $\text{AdS}_2 \times \mathbb{S}_1 \times \mathbb{S}_2$ extremization.

$$\begin{aligned} \varphi_i^* &= \frac{2m_+m_-}{m_+ - m_-} \left(1 - \frac{m_+ + m_-}{2m_+m_-} \frac{2\pi}{\Delta\psi} \right) \mathfrak{s}_i^*, \\ \epsilon_1^* &= -\frac{4}{3} \frac{2\pi}{\Delta z} - \frac{2(r_+ + r_-)}{m_+ - m_-} \frac{2\pi}{\Delta\psi}, \quad \epsilon_2^* = 2 \frac{2\pi}{\Delta\psi}, \end{aligned} \tag{5.42}$$

which, plugged into the off-shell central charge, gives

$$F(\varphi_i^*, \epsilon_i^*; \mathfrak{s}_i^\pm, t_i, m_\ell) = \frac{4(m_+ - m_-)^3}{3m_+m_-(m_+^2 + m_+m_- + m_-^2)} a_{4d}, \tag{5.43}$$

where a_{4d} is the central charge of $d = 4, \mathcal{N} = 1$ SCFTs that arise from N M5-branes wrapped on a spindle [29]

$$a_{4d} = \frac{3p_1^2 p_2^2 (\mathfrak{s} + p_1 + p_2)}{8n_+n_-(n_+ - p_1)(p_2 - n_+)[\mathfrak{s} + 2(p_1 + p_2)]^2} N^3. \tag{5.44}$$

This matches exactly the central charge of the $\text{AdS}_3 \times \mathbb{S}_1 \times \mathbb{S}_2$ system computed in [17]. In general, we expect that the anomaly polynomial computation of [17] and the prescription presented in this paper should be equivalent and connected by a suitable gauge choice and a possible redefinition of $\epsilon_{1,2}$, mixing the two related $U(1)$ isometries.

As explained in the previous section, our recipe can be applied also to non-toric four-dimensional orbifolds. In this class, we consider $\text{AdS}_3 \times \Sigma_g \times \mathbb{S}_2$, whose ‘toric data’ are deduced in Appendix C.2 for $\text{AdS}_3 \times S^2 \times \mathbb{S}_2$ and read

$$\begin{aligned} m_1 &= n_-, & \vec{n}_1 &= (-1, 0), & m_2 &= 1, & \vec{n}_2 &= (0, -1), \\ m_3 &= n_+, & \vec{n}_3 &= (1, 0), & m_4 &= 1, & \vec{n}_4 &= (-t, 1). \end{aligned} \tag{5.45}$$

Global regularity leads to

$$t = \frac{12(g - 1)n_+n_-(n_+ - n_-)}{[\mathfrak{s} - (p_1 + p_2)][\mathfrak{s} + 2(p_1 + p_2)]}, \tag{5.46}$$

with \mathfrak{s} given in Eq. (5.35). We take the physical fluxes to be as in (5.36), where $t_i = \frac{p_i}{n_+n_-}$ is unchanged and \mathfrak{s}_i^\pm can be read from (4.7) of [17]. The gauge symmetry allows us to identify the $p_i^{(\ell)}$ as in Eq. (5.37), with $\mathfrak{s}_i^\pm = \mathfrak{s}_i^* \pm \frac{t}{2} t_i$ and

$$\mathfrak{s}_i^* = \frac{tp_i[-6p_i + 4(n_+ + n_-) - \mathfrak{s}]}{6n_+n_-(n_+ - n_-)}. \tag{5.47}$$

We can then solve Eq. (5.5) to obtain $\sigma^{(\ell)} = (+, -, +, -)$, use it to identify the auxiliary vector \vec{W} through (5.13) and write down the constraint (5.12)

$$\varphi_1 + \varphi_2 - \frac{n_+ - n_-}{2n_+n_-} \epsilon_1 - t \frac{n_+ - n_-}{4n_+n_-} \epsilon_2 = 2. \tag{5.48}$$

This is the same as in (5.30), with $t \mapsto -t$ and $\epsilon_{1,2} \mapsto -\epsilon_{1,2}$, due to the different conventions used in [17].

Taking all the $\eta_6^{(\ell)} = -$ and extremizing Eq. (5.18) with respect to φ_i , ϵ_i and Λ , we obtain the critical values

$$\varphi_i^* = -\frac{\mathfrak{s}_i^*}{g-1}, \quad \epsilon_1^* = \frac{4}{3} \frac{2\pi}{\Delta z}, \quad \epsilon_2^* = 0, \tag{5.49}$$

as well as the off-shell central charge at the extremum

$$F(\varphi_i^*, \epsilon_i^*; \mathfrak{s}_i^\pm, \mathfrak{t}_i, m_\ell) = \frac{32}{3}(g-1) a_{4d}, \tag{5.50}$$

with a_{4d} given in (5.44). This result agrees with the central charge of the $\text{AdS}_3 \times \Sigma_g \times \overline{\Sigma}_2$ system studied in [17]. The origin of the choice of the $\eta_6^{(\ell)}$ can be traced to the sign of the gauge potential of the five-dimensional solution that the authors of [17] uplifted to obtain the $\text{AdS}_3 \times \Sigma_g \times \overline{\Sigma}_2$ backgrounds. The component along AdS_3 of their Killing spinors satisfy the corresponding Killing spinor equations, but with a minus, namely $\widehat{\nabla}_{\hat{\mu}} \vartheta = -\frac{1}{2} \alpha_{\hat{\mu}} \vartheta$, which implies that in the two-dimensional dual SCFTs (2, 0) supersymmetries are preserved, instead of (0, 2) as in our case. This fact leads to a different sign in the extraction of the central charge from the anomaly polynomial.

6 Discussion

In this paper, we have presented two new families of supersymmetric AdS_2 solutions of massive type IIA supergravity, associated with D4-branes wrapped on four-dimensional orbifolds \mathbb{M}_4 , which may be viewed as two different generalizations of the Hirzebruch surfaces. In one case, $\mathbb{M}_4 = \Sigma_g \times \Sigma$ is a spindle Σ fibered over a smooth Riemann surface Σ_g of genus $g > 1$, while in the other case $\mathbb{M}_4 = \Sigma \times \Sigma$ is a spindle Σ fibered over another spindle Σ . We have argued that these are dual to $\mathcal{N} = 2$ SCQMs arising from different twisted compactifications of the $d = 5$, $\mathcal{N} = 1$ $USp(2N)$ supersymmetric gauge theories on \mathbb{M}_4 . The structure of these solutions is analogous to that of the AdS_3 solutions constructed in [17] and indeed for the $\mathbb{M}_4 = \Sigma \times \Sigma$ family we have provided a toric geometry description and formulated a conjectural extremal problem that applies to both classes.

As in [17], the solutions that we found have less parameters than the data specifying the orbifolds \mathbb{M}_4 . In particular, the twisting parameter t is not an arbitrary integer, but it is unnaturally related to the spindle data n_\pm, m_\pm . We therefore expect more general solutions to exist, although the ansatz describing them is likely to be quite different from those employed here and in [17]. Furthermore, the toric description of the $\mathbb{M}_4 = \Sigma \times \Sigma$ solutions strongly suggests that there exist solutions corresponding to toric orbifolds with an arbitrary number of fixed points, and could include the case of toric manifolds.

While in this paper we provided a basic description of the orbifolds, which was adequate for characterizing the present solutions, it would be desirable to study more

systematically their underlying structure. For example, it would be useful to perform a global analysis of the gauge fields and Killing spinors on \mathbb{M}_4 , determining the bundles of which these are sections. Relatedly, given a generic compact four-dimensional toric orbifold \mathbb{M}_4 , it would be nice to determine what are the allowed supersymmetry-preserving twists. It could also be useful to cast our solutions in terms of classifications of supersymmetric solutions of massive type IIA supergravity, or to characterize them directly in the context of $D = 6$ and $D = 7$ gauged supergravities. For example, we expect that generic \mathbb{M}_4 's will be endowed with a canonical complex structure²². Another natural extension of our results is to investigate whether supersymmetric AdS₆ black holes with \mathbb{M}_4 horizons can be constructed.

In this paper, we have proposed an entropy function constructed assembling *gravitational blocks*, extending the proposal of [27] from the spindle to a very general class of four-dimensional orbifolds. We believe that our approach will be applicable to orbifolds more general than those for which we constructed explicit solutions here. However, finding additional explicit examples may lead to a more refined formulation of our recipe. Similarly to the spindle off-shell free energies [27], it is striking that the structure we introduced applies to both $D = 6$ and $D = 7$ supergravities, with minor tweaks. In the context of $D = 7$ supergravity, our proposal reproduces the results obtained by integrating the M5-branes anomaly polynomial on compact toric four-manifolds [36] and on the $\mathbb{M}_4 = \Sigma \times \Sigma$ orbifolds [17]. On the other hand, in the context of $D = 6$ supergravity, at present we do not have access to any field theory computation and therefore our proposal gives a prediction for the large N limit of the localized partition function of the $d = 5$, $\mathcal{N} = 1$ $USp(2N)$ on $S^1 \times \mathbb{M}_4$.

Acknowledgements We thank Ugo Bruzzo, Michele Rossi and Alberto Zaffaroni for helpful discussions. We also thank the authors of [20] for informing us about their upcoming paper. The research of FF is supported by the project “Nonlinear Differential Equations” of Università degli Studi di Torino.

Funding Open access funding provided by Università degli Studi di Torino within the CRUI-CARE Agreement.

Data availability Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If

²² We note that the $\Sigma \times \Sigma$ orbifolds have been discussed in the mathematical literature in the context of toric Kähler geometry and were referred to as “Hirzebruch orbifold surfaces” [3]. In particular, the authors of [41] classified four-dimensional toric Kähler orbifolds associated with labeled polytopes with four edges (quadrilaterals) and showed that they admit an extremal Kähler metric of Calabi type if and only if the base spindle is actually a smooth two-sphere (i.e., $m_+ = m_- = 1$).

Table 1 Examples of solutions to the quartic Eq. (2.33) for $n_+ \leq 100$ and $|p_1| \leq 100$

n_+	n_-	p_1	x
11	1	-22	1/6
11	4	-6	1/5
14	3	-19	3/17
17	8	-98	1/25
17	12	-52	1/29
24	1	-81	3/25
31	3	-98	2/17
32	25	-21	1/19
36	11	-85	5/47
41	19	-19	1/6
43	3	-8	12/23
43	17	-87	1/10
53	13	-6	4/11
59	13	-39	1/4
67	50	-15	1/13
73	42	-27	3/23
80	19	-18	1/3
82	5	-42	11/29
83	7	-33	2/5
83	47	-12	2/13
90	53	-93	1/11

material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

A Some solutions to the Diophantine equation

In this appendix we address the quantization conditions we stated through the paper. Although we were not able to prove that these constraints can be satisfied appropriately tuning the parameters of the solutions, nevertheless we could find some examples in which the quantization can be realized.

A priori, there are no constraints on x apart from being solution of the quartic Eq. (2.33) and lying in the range $(0, 1)$. Nevertheless, in order to construct the integer quantities we are searching for, it is easier to work with rational values of x , if they exist. Indeed, they exist, and in Table 1 we present some of them, along with the corresponding values of the integers n_+ , n_- and p_1 (recall that $p_2 = n_+ + n_- - p_1$).

Inspired by [17] we propose an algorithm to construct a family of solution to the quantization conditions. For the combinations of parameters presented in Table 1, x is rational and positive, thus we can write it as $x = p/q$, with $p, q \in \mathbb{N}$. Then, we

define t as

$$t = -8n_+n_-\mathfrak{p}^2\mathfrak{q}u, \quad (\text{A.1})$$

with $u \in \mathbb{N}$, which makes t integer by construction and, moreover, multiple of both n_+ and n_- . We now focus on the $\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ system, the case with Riemann surface proceeds similarly. Substituting the expression of t in the constraint (2.59) we obtain

$$m_+ - m_- = 2(\mathfrak{p}^2 + 3\mathfrak{q}^2)[(n_+ - n_-)\mathfrak{q} - (n_+ + n_-)\mathfrak{p}]u, \quad (\text{A.2})$$

which can be solved to find m_{\pm} in terms of n_{\pm} , z and u . The quantization condition that follows from (2.60) is automatically satisfied given t as in (A.1), indeed it becomes

$$(m_+m_-)\mathfrak{s}_1^- = -[(n_+ + n_-)\mathfrak{p}^3 - (p_1 - 3p_2 + 6n_+)\mathfrak{p}^2\mathfrak{q} + (p_1 + 5p_2)\mathfrak{p}\mathfrak{q}^2 - 3(n_+ - n_-)\mathfrak{q}^3]u, \quad (\text{A.3})$$

which is evidently an integer. The last requirement (3.29) can be met considering that $3\mu(x^2 + 1) - x(x^2 + 5) > 0$ (cf. [27]) and that all the quantities appearing are rational, hence N can be tuned appropriately so that $K \in \mathbb{N}$.

B Riemann surfaces from spindles

In the presence of twist, it was explicitly shown in [31] that the spindle factor of the known $\text{AdS}_{D-2} \times \Sigma$ solutions, with $D = 4, 5, 7$, can be turned into a round sphere performing a particular limit of the coordinates and parameters of the system. In the same spirit, as we will see in this appendix, a similar mechanism applies to the $\text{AdS}_{D-2} \times \Sigma$ solutions with anti-twist, so far constructed only in $D = 4$ and $D = 5$. In this case, the spindle becomes the two-dimensional hyperbolic space, which can later be quotiented to give a constant curvature Riemann surface with genus $g > 1$.

B.1 $\text{AdS}_2 \times \mathbb{M}_4$ solutions

Let us begin considering the local spindle metric (2.52)

$$ds_{\Sigma_1}^2 = \frac{x^2}{q} dx^2 + \frac{q}{4x^2} d\psi^2, \quad (\text{B.1})$$

where, we recall, $q(x) = x^4 - 4x^2 + 4ax - a^2$. Performing the change of coordinates

$$x = 1 + \epsilon \xi, \quad \psi = \frac{\phi}{\epsilon}, \quad \text{with} \quad a = 1 - \epsilon^2, \quad (\text{B.2})$$

and computing the limit $\epsilon \rightarrow 0^+$, the spindle metric becomes

$$ds_{\Sigma_1}^2 = \frac{1}{2} \left(\frac{d\xi^2}{\xi^2 - 1} + (\xi^2 - 1) d\phi^2 \right) = \frac{1}{2} ds_{H^2}^2, \tag{B.3}$$

which is the local metric of the hyperbolic space H^2 , with radius squared equal to $1/2$. This space can then be quotiented to obtain a Riemann surface with $g > 1$. In order for metric (B.3) to genuinely describe H^2 , the coordinates ξ and ϕ must have the correct range of definition, i.e., $|\xi| \in (1, +\infty)$ and $\phi \in [0, 2\pi]$. To address this issue, we begin considering that the coordinate x is restricted between x_N and x_S , where [28]

$$x_N = -1 + \sqrt{1+a}, \quad x_S = 1 - \sqrt{1-a}. \tag{B.4}$$

By means of (B.2), these bounds are mapped to

$$\xi_2 = \frac{-2 + \sqrt{2 - \epsilon^2}}{\epsilon} \rightarrow -\infty, \quad \xi_3 = -\frac{\sqrt{\epsilon^2}}{\epsilon} = -1, \tag{B.5}$$

in the limit $\epsilon \rightarrow 0^+$, thus correctly giving $\xi \in (-\infty, -1)$. The periodicity of ϕ can be studied by analyzing the conditions under which Σ_1 is proper spindle [28]

$$\sqrt{1+a} \cdot \Delta\psi = \frac{2\pi}{m_-}, \quad \sqrt{1-a} \cdot \Delta\psi = \frac{2\pi}{m_+}, \tag{B.6}$$

with m_{\pm} two co-prime integers such that $m_- < m_+$. By means of the change of coordinates (B.2), these conditions yield

$$\sqrt{2 - \epsilon^2} \cdot \frac{\Delta\phi}{\epsilon} = \frac{2\pi}{m_-}, \quad \sqrt{\epsilon^2} \cdot \frac{\Delta\phi}{\epsilon} = \frac{2\pi}{m_+}, \tag{B.7}$$

hence we have $\Delta\phi = 2\pi$ if and only if, at first order,

$$m_- = \frac{\epsilon}{\sqrt{2}} \rightarrow 0, \quad m_+ = 1. \tag{B.8}$$

Consistently, plugging these values of m_{\pm} into Eq. (2.53) gives $a = 1 - \epsilon^2$, as in (B.2). The global quantities characterizing the spindle Σ_1 are not mapped directly to the corresponding ones for a Riemann surface. Indeed, if we take as an examples the Euler characteristic of Σ_1 [30]

$$\chi_1 = \frac{1}{4\pi} \int_{\Sigma_1} R_{\Sigma_1} \text{vol}(\Sigma_1) = \frac{m_+ + m_-}{m_+ m_-}, \tag{B.9}$$

we see that it diverges when $m_- \rightarrow 0$ and $m_+ = 1$. The reason for this behavior can be found in the definition of Σ_g as a quotient of the hyperbolic space, whose Ricci curvature, integrated over the whole space, diverges, since it is not compact. Therefore,

we need a way to “regularize” this limit. To this end, we notice that, adopting the substitution (B.8),

$$2(1 - g) \cdot \chi_1 = \frac{2\sqrt{2}(1 - g)}{\epsilon} + 2(1 - g), \tag{B.10}$$

that is, we obtain a simple pole at $\epsilon = 0$ plus the Euler characteristic of a Riemann surface. We propose the following prescription, that we will test against different examples: given a system comprising a Σ_g factor that can be obtained from Σ_1 through limit (B.8), the global quantities of the former can be computed multiplying by $2(1 - g)$ the corresponding quantities of the latter and taking the finite term in the Laurent series.

We can now transfer this technology to higher-dimensional spacetimes. Applying the change of coordinates (B.2)—driven by the underlying limit (B.8)—to the $\text{AdS}_2 \times \Sigma_1$ system described by (2.68) we obtain the $\text{AdS}_2 \times \Sigma_g$ background in (2.67). Likewise are related the respective Killing spinors. Indeed, in this limit Q_1 vanishes and Q_2 becomes constant, thus (2.76) turns into (2.75), apart from an irrelevant overall factor. For what concerns the total charge of the two systems, applying the recipe illustrated for the Euler characteristic χ_1 we obtain (cf. (2.56))

$$2(1 - g) \cdot \frac{1}{2\pi} \int_{\Sigma_1} F_{4d}^{\Sigma_1} = \frac{a_{-1}}{\epsilon} + \frac{1}{2\pi} \int_{\Sigma_g} F_{4d}^{\Sigma_g}, \tag{B.11}$$

where a_{-1} is the residue.

The change of coordinates (B.2) also maps the $\text{AdS}_2 \times \Sigma_1 \times \Sigma_2$ solution and Killing spinors to the $\text{AdS}_2 \times \Sigma_g \times \Sigma_2$ corresponding quantities. This relation reflects in a formal connection between the respective entropies by means of (B.8):

$$2(1 - g) \cdot S_{\Sigma_1 \times \Sigma_2} = \frac{a_{-1}}{\epsilon} + S_{\Sigma_g \times \Sigma_2} + \mathcal{O}(\epsilon), \tag{B.12}$$

where $S_{\Sigma_1 \times \Sigma_2}$ is given in (3.30) and $S_{\Sigma_g \times \Sigma_2}$ in (3.23).

B.2 $\text{AdS}_3 \times \mathbb{M}_4$ solutions

Moving to the local spindle metric presented and studied in [30]²³

$$ds_{\Sigma_1}^2 = \frac{x}{q} dx^2 + \frac{q}{36x^2} d\psi^2, \tag{B.13}$$

where $q(x) = 4x^3 - 9x^2 + 6ax - a^2$, we perform the change of coordinates

$$x = 1 + 2\epsilon \xi, \quad \psi = \frac{\phi}{\epsilon}, \quad \text{with} \quad a = 1 - 3\epsilon^2. \tag{B.14}$$

²³ To stick to our notation, we relabeled the coordinates (y, z) as (x, ψ) and the integers n_{\pm} as m_{\mp} .

In the $\epsilon \rightarrow 0^+$ limit, the spindle metric becomes

$$ds_{\Sigma_1}^2 = \frac{1}{3} \left(\frac{d\xi^2}{\xi^2 - 1} + (\xi^2 - 1) d\phi^2 \right) = \frac{1}{3} ds_{H^2}^2. \tag{B.15}$$

Once again, we can quotient the hyperbolic space to give a constant curvature Riemann surface with genus $g > 1$. It can be shown that the metric obtained in this limit describes a genuine hyperbolic space if

$$m_- = \frac{2\epsilon}{3} \rightarrow 0, \quad m_+ = 1. \tag{B.16}$$

Applying this recipe to the $\text{AdS}_3 \times \Sigma_1$ solution of [30] we have

$$ds_{5d}^2 = \frac{4}{9} ds_{\text{AdS}_3}^2 + \frac{1}{3} ds_{H^2}^2, \quad A_{5d} = \frac{1}{2} \omega_{H^2}, \tag{B.17}$$

where ω_{H^2} is such that $d\omega_{H^2} = \text{vol}(H^2)$. This is the $\text{AdS}_3 \times H^2$ near-horizon of the black string solution of $D = 5, \mathcal{N} = 2$ minimal gauged supergravity constructed in [40]²⁴. Like in the six-dimensional case, applying the change of coordinates (B.14), followed by the $\epsilon \rightarrow 0$ limit, to the $\text{AdS}_3 \times \Sigma_1 \times \Sigma_2$ solution of [17] gives the $\text{AdS}_3 \times \Sigma_g \times \Sigma_2$ background therein. Moreover, it is possible to retrieve the central charge of the latter, presented in (5.50), multiplying the central charge of the former, given in (5.43), by $2(1 - g)$, substituting m_{\pm} as in (B.16) and taking the finite part of the Laurent series. Explicitly,

$$2(1 - g) \cdot c_{\Sigma_1 \times \Sigma_2} = \frac{a_{-1}}{\epsilon} + c_{\Sigma_g \times \Sigma_2} + \mathcal{O}(\epsilon). \tag{B.18}$$

C More toric data from explicit metrics

In this appendix, we will provide further examples of how to extract toric data from explicit metrics, as discussed in the main body of the paper. The first example we consider is the Hirzebruch surface \mathbb{F}_1 , that is a well-known symplectic toric manifold. Its associated polytope is a Delzant polytope (see, e.g., [15]) and below we show how to obtain it starting from an explicit metric on \mathbb{F}_1 [45], using two methods. Namely, computing the image of the moment map associated with a symplectic structure compatible with a conformally rescaled metric and studying the Killing vectors generating the $U(1)^2$ isometry of the metric. We will then employ these methods to extract the toric data associated with the metrics presented in [17]. These are used in Sect. 5 to show that our recipe reproduces the central charge computed in [17].

²⁴ The gauge potential has a different sign with respect to [40], but the solution is equivalent and still supersymmetric.

C.1 Hirzebruch \mathbb{F}_1

We now illustrate an example of Delzant polytope and its construction studying the first Hirzebruch surface \mathbb{F}_1 , whose metric is given by (cf. equation (7.11) of [45])²⁵

$$ds_{\mathbb{F}_1}^2 = \frac{1-y}{6}(d\theta^2 + \sin^2\theta d\phi^2) + \frac{dy^2}{wq} + \frac{l^2 w q}{36F}(d\Omega + \cos\theta d\phi)^2, \tag{C.1}$$

where the expression of the functions $w(y)$, $q(y)$ and $F(y)$ can be found in [45]. The angular coordinates lie in the ranges $\theta \in [0, \pi]$, $\phi \in [0, 2\pi]$, $\Omega \in [0, 4\pi]$, while y spans the range $[y_1, y_2]$, where y_1 and y_2 are, respectively, the negative and the smallest positive roots of $q(y)$. To ensure the correct signature of the metric we require $1 - y > 0$. After having rescaled the metric as $ds^2 = \Gamma(y) ds_{\mathbb{F}_1}^2$, we consider the symplectic two-form

$$\omega = \Gamma(y) \left[\frac{1-y}{6} \sin\theta d\theta \wedge d\phi + \frac{l}{6F^{1/2}} dy \wedge (d\Omega + \cos\theta d\phi) \right]. \tag{C.2}$$

When $\Gamma(y)$ is such that $\Gamma'(y) = \frac{F^{1/2}-l}{(1-y)F^{1/2}}\Gamma(y)$, the two-form (C.2) is closed and can be written as

$$\omega = d\phi \wedge d \left[\frac{1-y}{6} \Gamma(y) \cos\theta \right] + d\Omega \wedge d \left[\frac{1-y}{6} \Gamma(y) \right]. \tag{C.3}$$

For fixed values of y , metric (C.1) describes a squashed, smooth three-sphere, corresponding to the system studied in Sect. 4 of [45] for $m = 1$. Following their arguments, we take as a basis of an effective torus action²⁶

$$e_1 = V_2, \quad e_2 = V_1 + \frac{1}{2} V_2, \tag{C.4}$$

where V_1 generates the rotations of the two-sphere about the equator with weight one, hence $V_1 = \partial_\phi$, and V_2 the rotations of the S^1 fiber, also with weight one, hence $V_2 = \partial_\nu$, where we introduced the new radial coordinate $\nu = \Omega/2$, with period 2π . Once we have chosen a basis for the torus action, we can derive the related moment maps

$$\vec{\mu} = \frac{1-y}{3} \Gamma(y) \left(1, \frac{\cos\theta + 1}{2} \right). \tag{C.5}$$

In total, we have four fixed points, corresponding to all the possible combinations pairing the poles of the base two-sphere ($\theta = 0, \pi$) and the poles of the fiber two-

²⁵ Here, we replaced ℓ with l in order to avoid confusion with the index of the fixed points.

²⁶ We exchanged e_1 and e_2 with respect to [45] in order to stick to our notation of counterclockwise ordered normal vectors.




Fig. 4 Toric data of the first Hirzebruch surface \mathbb{F}_1

sphere ($y = y_{1,2}$)

$$\begin{aligned} v_1 &= \{\theta = 0, y = y_1\}, & v_2 &= \{\theta = 0, y = y_2\}, \\ v_3 &= \{\theta = \pi, y = y_2\}, & v_4 &= \{\theta = \pi, y = y_1\}. \end{aligned} \tag{C.6}$$

The image of these points under the moment maps are

$$\begin{aligned} \vec{\mu}(v_1) &= \Xi_1(1, 1), & \vec{\mu}(v_2) &= \Xi_2(1, 1), \\ \vec{\mu}(v_3) &= \Xi_2(1, 0), & \vec{\mu}(v_4) &= \Xi_1(1, 0), \end{aligned} \tag{C.7}$$

where we defined, for convenience, the positive constants $\Xi_i = \frac{1-y_i}{3} \Gamma(y_i)$. A numerical analysis shows that $\Xi_1 > \Xi_2$, therefore the moment polytope of \mathbb{F}_1 can be drawn as in Fig. 4a. A quick investigation proves that it is a Delzant polytope in \mathbb{R}^2 .

The polytope is uniquely defined by the relations

$$\langle \vec{\mu}, \vec{n}_1 \rangle \leq \Xi_1, \quad \langle \vec{\mu}, \vec{n}_2 \rangle \leq 0, \quad \langle \vec{\mu}, \vec{n}_3 \rangle \leq -\Xi_2, \quad \langle \vec{\mu}, \vec{n}_4 \rangle \leq 0, \tag{C.8}$$

where the \mathbb{Z}^2 primitive normal vectors \vec{n}_ℓ are

$$\vec{n}_1 = (1, 0), \quad \vec{n}_2 = (-1, 1), \quad \vec{n}_3 = (-1, 0), \quad \vec{n}_4 = (0, -1). \tag{C.9}$$

The fan generated by these vectors is depicted in Fig. 4b and is dual to the Delzant polytope of Fig. 4a.

A different approach to the construction of the toric data of a given toric manifold goes through the analysis of the degenerating Killing vectors of the corresponding metric. In total, we have four Killing vectors $\xi_{(\ell)}$ degenerating at the loci D_ℓ , namely

$$\begin{aligned} D_1 = \{y = y_1\} : & \quad \xi_{(1)} = 2\partial_\Omega, & D_2 = \{\theta = 0\} : & \quad \xi_{(2)} = \partial_\phi - \partial_\Omega, \\ D_3 = \{y = y_2\} : & \quad \xi_{(3)} = 2\partial_\Omega, & D_4 = \{\theta = \pi\} : & \quad \xi_{(4)} = \partial_\phi + \partial_\Omega. \end{aligned} \tag{C.10}$$

These vectors have been normalized such that they have unitary surface gravity when approaching the degeneracy hypersurface. Expanding on the basis (C.4), the four Killing vectors read

$$\xi_{(1)} = \xi_{(3)} = e_1, \quad \xi_{(2)} = e_2 - e_1, \quad \xi_{(4)} = e_2. \quad (\text{C.11})$$

We notice that the four loci D_ℓ are in a one-to-one correspondence with the facets of the polytope in Fig. 4a and that the Killing vectors $\xi_{(\ell)}$ can be written in terms of the related normal vectors \vec{n}_ℓ as $\xi_{(\ell)} = \vec{n}_\ell \cdot (e_1, e_2)$.

C.2 $\text{AdS}_3 \times S^2 \times \Sigma_2$ solutions

We consider here the $\text{AdS}_3 \times \Sigma_g \times \Sigma_2$ geometry presented in Sect. 4 of [17] subjected, as before, to the analytic continuation $\Sigma_g \mapsto S^2$. The metric is given by²⁷

$$ds^2 = \frac{4(yP)^{1/5}}{9} \left[ds_{\text{AdS}_3}^2 + \frac{3}{4} ds_{S^2}^2 + \frac{9y}{16Q} dy^2 + \frac{9Q}{4P} \left(dz - \frac{2}{3} \omega_{S^2} \right)^2 \right], \quad (\text{C.12})$$

where $d\omega_{S^2} = -\text{vol}(S^2)$ and

$$h_i(y) = y^2 + q_i, \quad P(y) = h_1(y) h_2(y), \quad Q(y) = -y^3 + \frac{1}{4} P(y). \quad (\text{C.13})$$

We can take explicitly

$$ds_{S^2}^2 = d\theta^2 + \sin^2\theta d\psi^2, \quad \omega_{S^2} = \cos\theta d\psi. \quad (\text{C.14})$$

Since the requirement of having a globally well-defined fibration

$$\frac{1}{2\pi} \int_{S^2} d\eta = t \in \mathbb{Z}, \quad \eta \equiv \frac{2\pi}{\Delta z} \left(dz - \frac{2}{3} \omega_{S^2} \right), \quad (\text{C.15})$$

yields the relation²⁸

$$\frac{4}{3t} = \frac{\Delta z}{2\pi}, \quad (\text{C.16})$$

²⁷ In order to make the comparison with [17] easier, we relabeled ϕ and a therein as z and a and exchanged $n_+ \leftrightarrow n_-$.

²⁸ Notice that here t is a positive integer, due to the different convention of [17] in defining ω .

we can define a 2π -periodic coordinate $v_2 = \frac{2\pi}{\Delta z} z$ in terms of which the four-dimensional toric orbifold $S^2 \times \Sigma_2$ of interest is

$$ds^2_{S^2 \times \Sigma_2} = \frac{1}{3} (d\theta^2 + \sin^2\theta d\psi^2) + \frac{y}{4Q} dy^2 + \frac{Q}{P} \left(\frac{2}{t} dv_2 - \cos\theta d\psi \right)^2. \tag{C.17}$$

For fixed y this metric describes an S^3/\mathbb{Z}_t , thus we consider as a basis of an effective torus action

$$e_1 = \partial_{v_2}, \quad e_2 = \partial_\psi + \frac{t}{2} \partial_{v_2}. \tag{C.18}$$

The four fixed points are

$$\begin{aligned} v_1 &= \{\theta = 0, y = y_2\}, & v_2 &= \{\theta = 0, y = y_3\}, \\ v_3 &= \{\theta = \pi, y = y_3\}, & v_4 &= \{\theta = \pi, y = y_2\}, \end{aligned} \tag{C.19}$$

where $y_{2,3}$ are the two middle roots of $Q(y)$, with $y_2 < y_3$. These points define the four divisors

$$D_1 = \{y = y_2\}, \quad D_2 = \{\theta = 0\}, \quad D_3 = \{y = y_3\}, \quad D_4 = \{\theta = \pi\}, \tag{C.20}$$

and, with respect to the basis (C.18), the degenerate normalized Killing vectors and the orbifold labels are

$$\begin{aligned} D_1 : \quad \xi_{(1)} &= n_- e_1, & m_1 &= n_-, & D_2 : \quad \xi_{(2)} &= e_2, & m_2 &= 1, \\ D_3 : \quad \xi_{(3)} &= n_+ e_1, & m_3 &= n_+, & D_4 : \quad \xi_{(4)} &= e_2 - t e_1, & m_4 &= 1. \end{aligned} \tag{C.21}$$

We can now construct the polytope, considering the four-dimensional conformally rescaled metric $ds^2 = \Gamma(y) ds^2_{S^2 \times \Sigma_2}$ and the related symplectic two-form

$$\omega = \Gamma(y) \left[\frac{1}{3} \sin\theta d\theta \wedge d\psi + \frac{y^{1/2}}{2P^{1/2}} dy \wedge \left(dz - \frac{2}{3} \cos\theta d\psi \right) \right]. \tag{C.22}$$

When $\Gamma'(y) = \frac{y^{1/2}}{P^{1/2}} \Gamma(y)$, the two-form (C.22) is closed and can be written as

$$\omega = d\psi \wedge d \left[\frac{1}{3} \Gamma(y) \cos\theta \right] + dv_2 \wedge d \left[-\frac{2}{3t} \Gamma(y) \right]. \tag{C.23}$$

From this expression, we can derive the moment maps with respect to the basis (C.18)

$$\vec{\mu} = -\frac{2}{3t} \Gamma(y) \left(1, \frac{t(1 - \cos\theta)}{2} \right), \tag{C.24}$$

whose action on the fixed points (C.19) is

$$\begin{aligned} \bar{\mu}(v_1) &= -\frac{2}{3t} \Gamma(y_2)(1, 0), & \bar{\mu}(v_2) &= -\frac{2}{3t} \Gamma(y_3)(1, 0), \\ \bar{\mu}(v_3) &= -\frac{2}{3t} \Gamma(y_3)(1, t), & \bar{\mu}(v_4) &= -\frac{2}{3t} \Gamma(y_2)(1, t). \end{aligned} \tag{C.25}$$

Since we have chosen the same basis as in (4.3), the resultant polytope is similar to the one constructed in Sect. 4.1. Indeed, it can be read off from Fig. 1 exchanging $v_1 \leftrightarrow v_3$, $v_2 \leftrightarrow v_4$ and $n_+ \leftrightarrow n_-$. The normal vectors \vec{n}_ℓ change accordingly with respect to Eq. (4.11), i.e., $\vec{n}_1 \leftrightarrow \vec{n}_3$ and $\vec{n}_2 \leftrightarrow \vec{n}_4$, together with $t \mapsto -t$. We can quickly verify that the quantities we derived satisfy relation (4.15).

C.3 AdS₃ × Σ₁ × Σ₂ solutions

We now move to the AdS₃ × Σ₁ × Σ₂ solutions presented in Sect. 3 of [17]²⁹. The metric on the toric orbifold is given by

$$ds_{\Sigma_1 \times \Sigma_2}^2 = \frac{x}{f} dx^2 + \frac{f}{36x^2} d\psi^2 + \frac{y}{4Q} dy^2 + \frac{Q}{P} \left(dz - \frac{1}{3} \left(1 - \frac{a}{x} \right) d\psi \right)^2, \tag{C.26}$$

where functions h_i , P and Q are the same as in (C.13) and

$$f(x) = 4x^3 - 9x^2 + 6ax - a^2. \tag{C.27}$$

The coordinate x ranges between x_1 and x_2 , the two smallest roots of $f(x)$ with $x_1 < x_2$, and y lies between y_2 and y_3 as before. The fibration is globally well defined if

$$\frac{m_+ m_-}{2\pi} \int_{\Sigma_1} d\eta = t \in \mathbb{Z}, \quad \eta \equiv \frac{2\pi}{\Delta z} \left(dz - \frac{1}{3} \left(1 - \frac{a}{x} \right) d\psi \right), \tag{C.28}$$

which gives the relation

$$\frac{t}{m_+ m_-} = -\frac{a(x_2 - x_1)}{3x_1 x_2} \frac{\Delta\psi}{\Delta z}. \tag{C.29}$$

Taking inspiration from Sect. 4.2, we introduce the following basis $\{E_1, E_2\}$ for an effective torus action:

$$E_1 = \partial_{v_2}, \quad E_2 = \partial_{v_1} + \frac{r_+ + r_-}{m_+ - m_-} \partial_{v_2}, \tag{C.30}$$

²⁹ In addition to n_\pm , we also exchanged $m_+ \leftrightarrow m_-$.

where $v_1 = \frac{2\pi}{\Delta\psi}\psi$ and r_{\pm} are two integers such that $t = r_+ m_+ + r_- m_-$. In total, there are four Killing vectors degenerating at the divisors $D_1 = \{y = y_2\}$, $D_2 = \{x = x_1\}$, $D_3 = \{y = y_3\}$, $D_4 = \{x = x_2\}$, which, in the basis (C.30), read

$$\begin{aligned} \xi_{(1)} &= n_- E_1, & \xi_{(2)} &= m_- E_2 + r_+ E_1, \\ \xi_{(3)} &= n_+ E_1, & \xi_{(4)} &= m_+ E_2 - r_- E_1. \end{aligned} \tag{C.31}$$

These are formally the same Killing vectors given in (4.29). The “long” normal vectors can be obtained by means of (4.28)

$$\vec{w}_1 = (n_-, 0), \quad \vec{w}_2 = (r_+, m_-), \quad \vec{w}_3 = (-n_+, 0), \quad \vec{w}_4 = (r_-, -m_+). \tag{C.32}$$

Following the same arguments of Sect. 4.2, we derive the labels $m_\ell = (n_-, 1, n_+, 1)$ and the “short” vectors $\vec{w} = \vec{w}_\ell/m_\ell$, which together constitute the toric data (\vec{w}_ℓ, m_ℓ) describing the $\Sigma_1 \times \Sigma_2$ toric orbifold. This orbifold is characterized by the same stacky fan and labeled polytope of Fig. 3.

References

1. Abreu, M.: Kahler–Sasaki geometry of toric symplectic cones in action-angle coordinates. *Port. Math.* **67**, 121 (2010). <https://doi.org/10.4171/PM/1862>. [arXiv:0912.0492](https://arxiv.org/abs/0912.0492)
2. Andrianopoli, L., D’Auria, R., Vaula, S.: Matter coupled F(4) gauged supergravity Lagrangian. *JHEP* **05**, 065 (2001). <https://doi.org/10.1088/1126-6708/2001/05/065>
3. Apostolov, V., Calderbank, D.M.J., Gauduchon, P.: Ambitoric geometry II: Extremal toric surfaces and Einstein 4-orbifolds. *Ann. Sci. Ec. Norm. Super.* **48**, 1075 (2015). <https://doi.org/10.24033/asens.2266>. [arXiv:1302.6979](https://arxiv.org/abs/1302.6979)
4. Arav, I., Gauntlett, J.P., Roberts, M.M., Rosen, C.: Leigh–Strassler compactified on a spindle. *JHEP* **10**, 067 (2022). [https://doi.org/10.1007/JHEP10\(2022\)067](https://doi.org/10.1007/JHEP10(2022)067). [arXiv:2207.06427](https://arxiv.org/abs/2207.06427)
5. Bah, I., Bonetti, F., Minasian, R., Nardoni, E.: Holographic Duals of Argyres–Douglas Theories. *Phys. Rev. Lett.* **127**, 211601 (2021). <https://doi.org/10.1103/PhysRevLett.127.211601>. [arXiv:2105.11567](https://arxiv.org/abs/2105.11567)
6. Bah, I., Bonetti, F., Minasian, R., Nardoni, E.: M5-brane sources, holography, and Argyres–Douglas theories. *JHEP* **11**, 140 (2021). [https://doi.org/10.1007/JHEP11\(2021\)140](https://doi.org/10.1007/JHEP11(2021)140). [arXiv:2106.01322](https://arxiv.org/abs/2106.01322)
7. Battaglia, F., Prato, E., Zaffran, D.: Hirzebruch surfaces in a one-parameter family. *Boll. Unione Mat. Ital.* **12**, 293 (2019). <https://doi.org/10.1007/s40574-018-0181-1>. [arXiv:1804.08503](https://arxiv.org/abs/1804.08503)
8. Benini, F., Bobev, N.: Exact two-dimensional superconformal R-symmetry and c-extremization. *Phys. Rev. Lett.* **110**, 061601 (2013). <https://doi.org/10.1103/PhysRevLett.110.061601>. [arXiv:1211.4030](https://arxiv.org/abs/1211.4030)
9. Benini, F., Hristov, K., Zaffaroni, A.: Black hole microstates in AdS₄ from supersymmetric localization. *JHEP* **05**, 054 (2016). [https://doi.org/10.1007/JHEP05\(2016\)054](https://doi.org/10.1007/JHEP05(2016)054). [arXiv:1511.04085](https://arxiv.org/abs/1511.04085)
10. Boido, A., Ipiña, J.M.P., Sparks, J.: Twisted D3-brane and M5-brane compactifications from multi-charge spindles. *JHEP* **07**, 222 (2021). [https://doi.org/10.1007/JHEP07\(2021\)222](https://doi.org/10.1007/JHEP07(2021)222). [arXiv:2104.13287](https://arxiv.org/abs/2104.13287)
11. Borisov, L.A., Chen, L., Smith, G.G.: The orbifold Chow ring of toric Deligne–Mumford stacks. *J. Am. Math. Soc.* **18**, 193 (2005). <https://doi.org/10.1090/S0894-0347-04-00471-0>. [arXiv:math/0309229](https://arxiv.org/abs/math/0309229)
12. Brandhuber, A., Oz, Y.: The D-4 - D-8 brane system and five-dimensional fixed points. *Phys. Lett. B* **460**, 307 (1999). [https://doi.org/10.1016/S0370-2693\(99\)00763-7](https://doi.org/10.1016/S0370-2693(99)00763-7). [arXiv:hep-th/9905148](https://arxiv.org/abs/hep-th/9905148)
13. Cacciatori, S.L., Klemm, D.: Supersymmetric AdS(4) black holes and attractors. *JHEP* **01**, 085 (2010). [https://doi.org/10.1007/JHEP01\(2010\)085](https://doi.org/10.1007/JHEP01(2010)085). [arXiv:0911.4926](https://arxiv.org/abs/0911.4926)
14. Caldarelli, M.M., Klemm, D.: Supersymmetry of Anti-de Sitter black holes. *Nucl. Phys. B* **545**, 434 (1999). [https://doi.org/10.1016/S0550-3213\(98\)00846-3](https://doi.org/10.1016/S0550-3213(98)00846-3). [arXiv: hep-th/9808097](https://arxiv.org/abs/hep-th/9808097)
15. Cannas da Silva, A.: Lectures on symplectic geometry, vol. 1764 of Lecture Notes in Mathematics. Springer, Berlin (2001). <https://doi.org/10.1007/978-3-540-45330-7>

16. Cassani, D., Gauntlett, J.P., Martelli, D., Sparks, J.: Thermodynamics of accelerating and supersymmetric AdS4 black holes. *Phys. Rev. D* **104**, 086005 (2021). <https://doi.org/10.1103/PhysRevD.104.086005>. arXiv:2106.05571
17. Cheung, K.C.M., Fry, J.H.T., Gauntlett, J.P., Sparks, J.: M5-branes wrapped on four-dimensional orbifolds. *JHEP* **08**, 082 (2022). [https://doi.org/10.1007/JHEP08\(2022\)082](https://doi.org/10.1007/JHEP08(2022)082). arXiv:2204.02990
18. Couzens, C.: A tale of (M)2 twists. *JHEP* **03**, 078 (2022). [https://doi.org/10.1007/JHEP03\(2022\)078](https://doi.org/10.1007/JHEP03(2022)078). arXiv:2112.04462
19. Couzens, C., Kim, H., Kim, N., Lee, Y.: Holographic duals of M5-branes on an irregularly punctured sphere. *JHEP* **07**, 102 (2022). [https://doi.org/10.1007/JHEP07\(2022\)102](https://doi.org/10.1007/JHEP07(2022)102). arXiv:2204.13537
20. Couzens, C., Kim, H., Kim, N., Lee, Y., Suh, M.: D4-branes wrapped on four-dimensional orbifolds through consistent truncation. *JHEP* **02**, 025 (2023). [https://doi.org/10.1007/JHEP02\(2023\)025](https://doi.org/10.1007/JHEP02(2023)025). arXiv:2210.15695
21. Couzens, C., Macpherson, N.T., Passias, A.: $\mathcal{N} = (2, 2)$ AdS₃ from D3-branes wrapped on Riemann surfaces. *JHEP* **02**, 189 (2022). [https://doi.org/10.1007/JHEP02\(2022\)189](https://doi.org/10.1007/JHEP02(2022)189). [arXiv:2107.13562]
22. Couzens, C., Stemerink, K.: Universal spindles: D2's on Σ and M5's on $\Sigma \times \mathbb{H}^3$. arXiv:2207.06449
23. Couzens, C., Stemerink, K., van de Heisteeg, D.: M2-branes on discs and multi-charged spindles. *JHEP* **04**, 107 (2022). [https://doi.org/10.1007/JHEP04\(2022\)107](https://doi.org/10.1007/JHEP04(2022)107). arXiv:2110.00571
24. Cvetič, M., Gubser, S.S., Lu, H., Pope, C.N.: Symmetric potentials of gauged supergravities in diverse dimensions and Coulomb branch of gauge theories. *Phys. Rev. D* **62**, 086003 (2000). <https://doi.org/10.1103/PhysRevD.62.086003>. [arXiv:hep-th/9909121]
25. Cvetič, M., Lu, H., Pope, C.N.: Gauged six-dimensional supergravity from massive type IIA. *Phys. Rev. Lett.* **83**, 5226 (1999). <https://doi.org/10.1103/PhysRevLett.83.5226>. arXiv:hep-th/9906221
26. D'Auria, R., Ferrara, S., Vaula, S.: Matter coupled F(4) supergravity and the AdS(6)/CFT(5) correspondence. *JHEP* **10**, 013 (2000). <https://doi.org/10.1088/1126-6708/2000/10/013>. arXiv: hep-th/0006107
27. Faedo, F., Martelli, D.: D4-branes wrapped on a spindle. *JHEP* **02**, 101 (2022). [https://doi.org/10.1007/JHEP02\(2022\)101](https://doi.org/10.1007/JHEP02(2022)101). arXiv:2111.13660
28. Ferrero, P., Gauntlett, J.P., Ipiña, J.M.P., Martelli, D., Sparks, J.: Accelerating black holes and spinning spindles. *Phys. Rev. D* **104**, 046007 (2021). <https://doi.org/10.1103/PhysRevD.104.046007>. arXiv:2012.08530
29. Ferrero, P., Gauntlett, J.P., Martelli, D., Sparks, J.: M5-branes wrapped on a spindle. *JHEP* **11**, 002 (2021). [https://doi.org/10.1007/JHEP11\(2021\)002](https://doi.org/10.1007/JHEP11(2021)002). arXiv:2105.13344
30. Ferrero, P., Gauntlett, J.P., Pérez Ipiña, J.M., Martelli, D., Sparks, J.: D3-Branes Wrapped on a Spindle. *Phys. Rev. Lett.* **126**, 111601 (2021). <https://doi.org/10.1103/PhysRevLett.126.111601>. arXiv:2011.10579
31. Ferrero, P., Gauntlett, J.P., Sparks, J.: Supersymmetric spindles. *JHEP* **01**, 102 (2022). [https://doi.org/10.1007/JHEP01\(2022\)102](https://doi.org/10.1007/JHEP01(2022)102). arXiv:2112.01543
32. Ferrero, P., Inglese, M., Martelli, D., Sparks, J.: Multicharge accelerating black holes and spinning spindles. *Phys. Rev. D* **105**, 126001 (2022). <https://doi.org/10.1103/PhysRevD.105.126001>. arXiv:2109.14625
33. Giri, S.: Black holes with spindles at the horizon. *JHEP* **06**, 145 (2022). [https://doi.org/10.1007/JHEP06\(2022\)145](https://doi.org/10.1007/JHEP06(2022)145). arXiv:2112.04431
34. Hochenegger, A., Witt, F.: On complex and symplectic toric stacks. *Contributions to Algebraic Geometry* **305** (2012). <https://doi.org/10.4171/114-1/11>. arXiv:1105.6280
35. Hosseini, S.M., Hristov, K., Passias, A., Zaffaroni, A.: 6D attractors and black hole microstates. *JHEP* **12**, 001 (2018). [https://doi.org/10.1007/JHEP12\(2018\)001](https://doi.org/10.1007/JHEP12(2018)001). arXiv:1809.10685
36. Hosseini, S.M., Hristov, K., Tachikawa, Y., Zaffaroni, A.: Anomalies, Black strings and the charged Cardy formula. *JHEP* **09**, 167 (2020). [https://doi.org/10.1007/JHEP09\(2020\)167](https://doi.org/10.1007/JHEP09(2020)167). arXiv:2006.08629
37. Hosseini, S.M., Hristov, K., Zaffaroni, A.: Gluing gravitational blocks for AdS black holes. *JHEP* **12**, 168 (2019). [https://doi.org/10.1007/JHEP12\(2019\)168](https://doi.org/10.1007/JHEP12(2019)168). arXiv:1909.10550
38. Hosseini, S.M., Hristov, K., Zaffaroni, A.: Rotating multi-charge spindles and their microstates. *JHEP* **07**, 182 (2021). [https://doi.org/10.1007/JHEP07\(2021\)182](https://doi.org/10.1007/JHEP07(2021)182). arXiv:2104.11249
39. Karndumri, P., Nuchino, P.: Five-branes wrapped on topological disks from 7D N=2 gauged supergravity. *Phys. Rev. D* **105**, 066010 (2022). <https://doi.org/10.1103/PhysRevD.105.066010>. arXiv:2201.05037
40. Klemm, D., Sabra, W.A.: Supersymmetry of black strings in D = 5 gauged supergravities. *Phys. Rev. D* **62**, 024003 (2000). <https://doi.org/10.1103/PhysRevD.62.024003>. arXiv:hep-th/0001131

41. Legendre, E.: Toric geometry of convex quadrilaterals. *J. Symplectic Geom.* **9**, 343 (2011). <https://doi.org/10.4310/JSG.2011.V9.N3.A3>. arXiv:0909.4512
42. Lerman, E., Malkin, A.: Hamiltonian group actions on symplectic Deligne–Mumford stacks and toric orbifolds. *Adv. Math.* **229**, 984 (2012). <https://doi.org/10.1016/J.AiM.2011.10.013>. arXiv:0908.0903
43. Lerman, E., Tolman, S.: Hamiltonian torus actions on symplectic orbifolds and toric varieties. *Trans. Am. Math. Soc.* **349**, 4201 (1997). <https://doi.org/10.1090/S0002-9947-97-01821-7>. arXiv:dg-ga/9511008
44. Maldacena, J.M., Nunez, C.: Supergravity description of field theories on curved manifolds and a no go theorem. *Int. J. Mod. Phys. A* **16**, 822 (2001). <https://doi.org/10.1142/S0217751X01003937>. arXiv:hep-th/0007018
45. Martelli, D., Sparks, J.: Toric geometry, Sasaki-Einstein manifolds and a new infinite class of AdS/CFT duals. *Commun. Math. Phys.* **262**, 51 (2006). <https://doi.org/10.1007/s00220-005-1425-3> arXiv:hep-th/0411238
46. Martelli, D., Sparks, J.: Resolutions of non-regular Ricci-flat Kahler cones. *J. Geom. Phys.* **59**, 1175 (2009). <https://doi.org/10.1016/j.geomphys.2009.06.005>. arXiv:0707.1674
47. Romans, L.J.: The F(4) Gauged Supergravity in Six-dimensions. *Nucl. Phys. B* **269**, 691 (1986). [https://doi.org/10.1016/0550-3213\(86\)90517-1](https://doi.org/10.1016/0550-3213(86)90517-1)
48. Sakai, H.: The Symplectic Deligne-Mumford Stack Associated to a Stacky Polytope. *Results. Math.* **63**, 903 (2013). <https://doi.org/10.1007/s00025-012-0240-3>. arXiv:1009.3547
49. Suh, M.: Supersymmetric AdS_6 black holes from matter coupled $F(4)$ gauged supergravity. *JHEP* **02**, 108 (2019). [https://doi.org/10.1007/JHEP02\(2019\)108](https://doi.org/10.1007/JHEP02(2019)108). arXiv:1810.00675
50. Suh, M.: D4-D8-branes wrapped on a manifold with non-constant curvature. arXiv:2108.08326
51. Suh, M.: D3-branes and M5-branes wrapped on a topological disc. *JHEP* **03**, 043 (2022). [https://doi.org/10.1007/JHEP03\(2022\)043](https://doi.org/10.1007/JHEP03(2022)043). arXiv:2108.01105
52. Suh, M.: M2-branes wrapped on a topological disk. *JHEP* **09**, 048 (2022). [https://doi.org/10.1007/JHEP09\(2022\)048](https://doi.org/10.1007/JHEP09(2022)048). arXiv:2109.13278
53. Suh, M.: M5-branes and D4-branes wrapped on a direct product of spindle and Riemann surface. arXiv:2207.00034
54. Wang, W.: Toric Sheaves on Hirzebruch Orbifolds. *Doc. Math.* **25**, 655 (2020). <https://doi.org/10.25537/dm.2020v25.655-699>. arXiv:1803.11215

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.