

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

On the refinement matrix mask of interpolating Hermite splines

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1868918> since 2022-07-10T10:07:05Z

Published version:

DOI:10.1016/j.aml.2020.106524

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

On the refinement matrix mask of interpolating Hermite splines

Lucia Romani^a, Alberto Viscardi^{a,*}

^a*Dipartimento di Matematica, Alma Mater Studiorum Università di Bologna,
Piazza di Porta San Donato 5, 40126 Bologna, Italy*

Abstract

We propose a new computational approach for constructing the refinement matrix mask of interpolating Hermite splines of any order and with general dilation factor. Our strategy exploits the refinability properties of cardinal B-splines with simple knots and simplifies the constructive procedures proposed so far.

Keywords: Cardinal Hermite Spline Interpolation; Cardinal B-Splines; Vector Subdivision; Refinement Matrix Mask; Arbitrary Arity

1. Introduction

Hermite subdivision schemes of arity $m \in \mathbb{N} \setminus \{1\}$ form a special subclass of vector subdivision schemes which started to be investigated in the early 90's (see, e.g., [2, 3, 6, 7]) but is still subject of recent research [1, 4, 5, 9]. The distinctive feature of Hermite schemes is the fact that the components of the sequence of vector data $\{\mathbf{p}_k^{[j]} \in \mathbb{R}^{n+1}\}_{k \in \mathbb{Z}}$ generated at the j th refinement step correspond to the evaluation at $m^{-j}\mathbb{Z}$ of a certain function and its derivatives up to a certain order. In the present paper we investigate Hermite subdivision schemes capable of generating interpolating Hermite splines. Precisely, for any given initial sequence of vector data $\{\mathbf{p}_k^{[0]} \in \mathbb{R}^{n+1}\}_{k \in \mathbb{Z}}$, the vector-valued limit function $\mathbf{f} = [f_0, \dots, f_n]^T \in \mathcal{C}^0(\mathbb{R}, \mathbb{R}^{n+1})$ is formed by a first scalar-valued function $f_0 \in \mathcal{C}^n(\mathbb{R})$ which is a spline of degree $2n+1$ satisfying $f_s(k) = f_0^{(s)}(k) = p_{k,s}^{[0]}$, for all $k \in \mathbb{Z}$ and $s \in \{0, \dots, n\}$. For any arbitrary dilation factor m and order $n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$, such a Hermite scheme is described by the finite matrix sequence $\{\mathbf{A}_k \in \mathbb{R}^{(n+1) \times (n+1)}\}_{k=1-m}^{m-1}$ which is called the *refinement matrix mask* of the Hermite scheme. The refinement matrix mask of the case $m=2$ was constructed in [6] by applying suitable transformation matrices to the matrix sequence proposed in [8] to describe the refinement property of the $n+1$ B-spline basis functions of degree $2n+1$, defined over the knot vector $\{(-1)^{\langle n+1 \rangle}, (0)^{\langle n+1 \rangle}, (1)^{\langle n+1 \rangle}\}$ with all knots of multiplicity $n+1$, as indicated by $(\cdot)^{\langle n+1 \rangle}$. An alternative constructive approach, relying on the analytic expressions of the basic limit functions of the Hermite scheme, was proposed more recently in [9]. The goal of our work is to provide an easier construction of the refinement matrix mask of interpolating Hermite splines for any arbitrary order and dilation factor, by exploiting the refinability properties of degree- $(2n+1)$ cardinal B-splines with simple knots. In Sec. 2 we recall basic facts about vector subdivision schemes, in Sec. 3 we review the main properties of the subclass of interpolating Hermite spline subdivision schemes and finally, in Sec. 4, we present our novel construction.

2. Basic facts about vector subdivision schemes

A univariate (shift-invariant) *vector subdivision scheme of order* $n \in \mathbb{N}_0$ takes as input an initial sequence of vector data (*control points*)

*Corresponding author

Email addresses: lucia.romani@unibo.it (Lucia Romani), alberto.viscardi@unibo.it (Alberto Viscardi)

3. A short review of the subclass of interpolating Hermite spline subdivision schemes

The interpolating Hermite spline scheme of order $n \in \mathbb{N}_0$ is a special instance of convergent vector subdivision scheme that, for any choice of $m \in \mathbb{N} \setminus \{1\}$, generates interpolating Hermite splines of degree $2n + 1$. The following properties [4, 6] make it a special member of the class of vector subdivision schemes.

i) For all $j \in \mathbb{N}_0$, each matrix in the j th level refinement mask $\{\mathbf{A}_k^{[j]} \in \mathbb{R}^{(n+1) \times (n+1)} : k \in \mathbb{Z}\}$ satisfies

$$\mathbf{A}_k^{[j]} = \mathbf{D}^{-j-1} \mathbf{A}_k \mathbf{D}^j \quad \text{with} \quad \mathbf{D} = \begin{bmatrix} 1 & & & \\ & m^{-1} & & \\ & & \ddots & \\ & & & m^{-n} \end{bmatrix};$$

moreover, the refinement mask is compactly supported on $[1 - m, m - 1] \cap \mathbb{Z}$, i.e. \mathbf{A}_k is the zero matrix for $k > m - 1$ and $k < 1 - m$. It is easy to check that the latter implies

$$\text{supp}(\varphi_{r,s}^{[0]}) \subseteq [-1, 1], \quad \forall r, s \in \{0, \dots, n\}.$$

ii) For all $r, s \in \{0, \dots, n\}$,

$$\varphi_{r,s}^{[0]}(x) = \frac{d^r}{dx^r} \varphi_{0,s}^{[0]}(x) \quad \text{and} \quad \varphi_{r,s}^{[j]}(k) = \delta_{k,0} \delta_{r,s}, \quad \forall j \in \mathbb{N}_0, k \in \mathbb{Z}.$$

In particular, the latter requires, for every $j \in \mathbb{N}_0$, $\mathbf{A}_0^{[j]}$ to be the $(n + 1)$ -dimensional identity matrix which implies $\mathbf{A}_0 = \mathbf{D}$ due to i).

iii) For all $s \in \{0, \dots, n\}$, $\varphi_{0,s}^{[0]} \in \mathcal{C}^n(\mathbb{R})$ is a polynomial spline of degree $d = 2n + 1$.

In light of ii) and iii), for any initial data $\mathbf{p}^{[0]}$, f_0 in (6) is a \mathcal{C}^n spline of degree d and, for all $r \in \{0, \dots, n\}$,

$$f_r(x) = \frac{d^r}{dx^r} f_0(x) \quad \text{with} \quad f_r(m^{-j}k) = p_{k,r}^{[j]}, \quad j \in \mathbb{N}_0, k \in \mathbb{Z}. \quad (7)$$

4. Constructing the mask of Hermite splines from cardinal B-splines with simple knots

Let $N_d \in \mathcal{C}^{d-1}(\mathbb{R})$ be the cardinal B-spline of degree $d = 2n + 1$ supported on $[-n - 1, n + 1]$ and defined over the knot vector $\{-n - 1, -n, \dots, 0, \dots, n, n + 1\}$ with all simple knots. N_d satisfies the refinement equation (see, e.g., [10])

$$N_d(x) = \sum_{k=-(m-1)(n+1)}^{(m-1)(n+1)} b_k N_d(mx - k) \quad (8)$$

where
$$\frac{1}{m} \sum_{k=-(m-1)(n+1)}^{(m-1)(n+1)} b_k z^k = z^{-(m-1)(n+1)} \left(\frac{1 - z^m}{m(1 - z)} \right)^{d+1}, \quad z \in \mathbb{C}, |z| = 1.$$

In light of (8), for any given sequence of control points $\mathbf{c}^{[0]} = \{c_k^{[0]} \in \mathbb{R}\}_{k \in \mathbb{Z}}$, we can express the associated spline as

$$f(x) = \sum_{k \in \mathbb{Z}} c_k^{[0]} N_d(x - k) = \sum_{k \in \mathbb{Z}} c_k^{[j]} N_d(m^j x - k), \quad j \in \mathbb{N}_0,$$

$$\text{where} \quad c_h^{[j+1]} = \sum_{k \in \mathbb{Z}} b_{h-mk} c_k^{[j]}, \quad h \in \mathbb{Z}. \quad (9)$$

are finite portions of the bi-infinite matrices in (12). Moreover, by construction, the interpolating Hermite spline scheme reproduces polynomials up to degree d and cardinal B-splines of degree d .

Proof. $\mathbf{A}_0 = \mathbf{D}$ is already asked in condition ii). For the remaining matrices, we observe that the block $\overline{\mathbf{A}}$ can be extracted from the bigger matrix in (12) with row indices from $n+1$ to $n+(n+1)(m-1)$ and column indices from 0 to $2n+1=d$. Thus, on the right-hand side, we need only to consider the rows of \mathbf{N} with indices from 0 to d . Since, for every $s \in \{0, \dots, n\}$, $\text{supp}(N_d^{(s)}) = [-n-1, n+1]$, due to (13), we can also cut the columns of \mathbf{N} considering only indices from $-n$ to $n+1$, leading to

$$\mathbf{N}_R = \left[\begin{array}{c|c|c|c|c} N_d(-n) & N_d(1-n) & \cdots & N_d(n) & 0 \\ N_d^{(1)}(-n) & N_d^{(1)}(1-n) & \cdots & N_d^{(1)}(n) & 0 \\ \vdots & \vdots & & \vdots & 0 \\ N_d^{(n)}(-n) & N_d^{(n)}(1-n) & \cdots & N_d^{(n)}(n) & 0 \\ \hline 0 & N_d(-n) & \cdots & N_d(n-1) & N_d(n) \\ 0 & N_d^{(1)}(-n) & \cdots & N_d^{(1)}(n-1) & N_d^{(1)}(n) \\ 0 & \vdots & & \vdots & \vdots \\ 0 & N_d^{(n)}(-n) & \cdots & N_d^{(n)}(n-1) & N_d^{(n)}(n) \end{array} \right] \in \mathbb{R}^{(d+1) \times (d+1)}. \quad (15)$$

In a similar fashion we can cut the matrices on the left-hand side considering only the rows of \mathbf{N} with indices from $n+1$ to $n+(n+1)(m-1)$ and the columns of \mathbf{B} with indices from $-n$ to $n+1$. The only non-zero entries of \mathbf{N} contained in those rows are the ones with column index between $1-n$ and $m+n-1$, extrema included. For \mathbf{B} the only non-zero entries contained in the considered columns lie on the rows with indices from $-(m-1)(n+1)-mn$ to $(m-1)(n+1)+m(n+1)$. Thus the useful indices for the cut are the ones between α and β where

$$\begin{aligned} \alpha &= \max_{m \geq 2, n \geq 0} \{ 1-n, -(m-1)(n+1)-mn \} = 1-n, \\ \beta &= \min_{m \geq 2, n \geq 0} \{ m+n-1, (m-1)(n+1)+m(n+1) \} = m+n-1. \end{aligned}$$

From (15), it is clear that $\det(\mathbf{N}_R) \neq 0$. Thus \mathbf{N}_R^{-1} in (14) is well-defined as well as the matrices $\{\mathbf{A}_k\}_{k=1-m}^{m-1}$. Since all the matrices involved in (12) are properly slanted, (12) holds for every choice of the sequence $\mathbf{c}^{[j]}$.

Now we only have to prove that for every initial data vector $\mathbf{p}^{[0]}$, the scheme produces a \mathcal{C}^n spline of degree d and its derivatives. In view of (2) and (12), we observe that, after a subdivision step, each of the added new control vectors depends only on two consecutive control vectors from the previous step. Since the refinement rules have been chosen to reproduce degree d cardinal B-splines and thus polynomials up to degree d , the first component of the vector-valued limit function will be a polynomial of degree d on every unitary interval with integer extrema. The fact that we are interpolating the first n derivatives yields \mathcal{C}^n regularity, so concluding the proof. \square

Remark 4.2. When $m=3$ one has that $\mathbf{N}_L = \mathbf{N}_R$, for every $n \in \mathbb{N}_0$.

Since the entries of \mathbf{B} are given in (8) (see (13)) and the entries of \mathbf{N} can be easily obtained by recalling the properties of cardinal B-splines and their derivatives, Theorem 4.1 gives us an easy-to-implement method for constructing the refinement mask $\{\mathbf{A}_k \in \mathbb{R}^{(n+1) \times (n+1)}\}_{k=1-m}^{m-1}$ for any order $n \in \mathbb{N}_0$ and dilation factor $m \in \mathbb{N} \setminus \{1\}$. We conclude providing some illustrative examples.

Example 4.1 ($m=2, n=1, d=3$).

$$\{b_k\}_{k=-2}^2 = \{1, 4, 6, 4, 1\} / 8, \quad \mathbf{N}_L = \left[\begin{array}{c|c|c} 1/6 & 2/3 & 1/6 \\ \hline -1/2 & 0 & 1/2 \end{array} \right], \quad \mathbf{N}_R = \left[\begin{array}{c|c} \mathbf{N}_L & \begin{array}{c} 0 \\ 0 \end{array} \\ \hline \begin{array}{c} 0 \\ 0 \end{array} & \mathbf{N}_L \end{array} \right],$$

$$\mathbf{A}_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1/2 \end{bmatrix}, \quad \bar{\mathbf{A}} = [\mathbf{A}_1 \quad \mathbf{A}_{-1}] = \left[\begin{array}{cc|cc} 1/2 & 1/8 & 1/2 & -1/8 \\ -3/4 & -1/8 & 3/4 & -1/8 \end{array} \right].$$

Example 4.2 ($m = 2, n = 2, d = 5$).

$$\{b_k\}_{k=-3}^3 = \{1, 6, 15, 20, 15, 6, 1\} / 32,$$

$$\mathbf{N}_L = \left[\begin{array}{c|c|c|c|c} 1/120 & 13/60 & 11/20 & 13/60 & 1/120 \\ -1/24 & -5/12 & 0 & 5/12 & 1/24 \\ 1/6 & 1/3 & -1 & 1/3 & 1/6 \end{array} \right], \quad \mathbf{N}_R = \left[\begin{array}{c|c} \mathbf{N}_L & \begin{matrix} 0 \\ 0 \\ 0 \end{matrix} \\ \hline \begin{matrix} 0 \\ 0 \\ 0 \end{matrix} & \mathbf{N}_L \end{array} \right],$$

$$\mathbf{A}_0 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1/4 \end{bmatrix}, \quad \bar{\mathbf{A}} = [\mathbf{A}_1 \quad \mathbf{A}_{-1}] = \left[\begin{array}{ccc|ccc} 1/2 & 5/32 & 1/64 & 1/2 & -5/32 & 1/64 \\ -15/16 & -7/32 & -1/64 & 15/16 & -7/32 & 1/64 \\ 0 & -3/8 & -1/16 & 0 & 3/8 & -1/16 \end{array} \right].$$

Example 4.3 ($m = 3, n = 2, d = 5$).

$$\{b_k\}_{k=-6}^6 = \{1, 6, 21, 50, 90, 126, 141, 126, 90, 50, 21, 6, 1\} / 243,$$

$$\mathbf{N}_L = \mathbf{N}_R = \left[\begin{array}{c|c|c|c|c|c} 1/120 & 13/60 & 11/20 & 13/60 & 1/120 & 0 \\ -1/24 & -5/12 & 0 & 5/12 & 1/24 & 0 \\ 1/6 & 1/3 & -1 & 1/3 & 1/6 & 0 \\ \hline 0 & 1/120 & 13/60 & 11/20 & 13/60 & 1/120 \\ 0 & -1/24 & -5/12 & 0 & 5/12 & 1/24 \\ 0 & 1/6 & 1/3 & -1 & 1/3 & 1/6 \end{array} \right],$$

$$\mathbf{A}_0 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/3 & 0 \\ 0 & 0 & 1/9 \end{bmatrix}, \quad \bar{\mathbf{A}} = \begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_{-2} \\ \mathbf{A}_2 & \mathbf{A}_{-1} \end{bmatrix} = \left[\begin{array}{ccc|ccc} 64/81 & 16/81 & 4/243 & 17/81 & -2/27 & 2/243 \\ -40/81 & 0 & 2/243 & 40/81 & -13/81 & 4/243 \\ -40/81 & -32/81 & -10/243 & 40/81 & -8/81 & 1/243 \\ \hline 17/81 & 2/27 & 2/243 & 64/81 & -16/81 & 4/243 \\ -40/81 & -13/81 & -4/243 & 40/81 & 0 & -2/243 \\ 40/81 & 8/81 & 1/243 & -40/81 & 32/81 & -10/243 \end{array} \right].$$

Acknowledgements: This research has been accomplished within RITA (Research ITalian network on Approximation). The authors are members of the INdAM Research group GNCS.

References

- [1] M. Cotronei, C. Moosmüller, T. Sauer, N. Sissouno, *Level-dependent interpolatory Hermite subdivision schemes and wavelets*, Constr. Approx. 50 (2019), no. 2, 341–366.
- [2] N. Dyn, D. Levin, *Analysis of Hermite-interpolatory subdivision schemes*, Spline functions and the theory of wavelets (Montreal, PQ, 1996), 105–113, CRM Proc. Lecture Notes, 18, Amer. Math. Soc., Providence, RI, 1999.
- [3] N. Dyn, D. Levin, *Subdivision schemes in geometric modelling*, Acta Numer. 11 (2002), 73–144.
- [4] J. Fageot, S. Aziznejad, M. Unser, V. Uhlmann, *Support and approximation properties of Hermite splines*, J. Comput. Appl. Math. 368 (2020), 112503, 15 pp.
- [5] B. Jeong, J. Yoon, *Construction of Hermite subdivision schemes reproducing polynomials*, J. Math. Anal. Appl. 451 (2017), no. 1, 565–582.
- [6] B. Jüttler, U. Schwanecke, *Analysis and design of Hermite subdivision schemes*, Visual Comp 18, 326–342 (2002).
- [7] J.-L. Merrien, *A family of Hermite interpolants by bisection algorithms*, Numer. Algorithms 2 (1992), no. 2, 187–200.
- [8] G. Plonka, *Two-scale symbol and autocorrelation symbol for B-splines with multiple knots*, Adv. Comput. Math. 3 (1995), no. 1-2, 1–22.
- [9] D. Ranirina, J. de Villiers, *On Hermite vector splines and multi-wavelets*, J. Comput. Appl. Math. 349 (2019), 366–378.
- [10] M. Sabin, *Subdivision of Box-splines*. In Armin Iske, Ewald Quak and Michael S. Floater (eds.), *Tutorials on Multiresolution in Geometric Modelling*, Mathematics and Visualization, pages 3–23. Springer, 2002.