



Pulmonary vein isolation through trans-jugular approach in a patient with inferior vena cava interruption

Andrea Saglietto¹ · Gaetano Maria De Ferrari¹ · Federico Ferraris¹ · Matteo Anselmino¹

Received: 10 September 2021 / Accepted: 27 December 2021
© The Author(s) 2022

1 Case report

A 50-year-old man, with surgically corrected congenital heart disease (ostium secundum atrial septal defect associated with partial anomalous pulmonary venous return and inferior vena cava — IVC — interruption), sick sinus syndrome (for which he was implanted with a single-lead atrial pacemaker) and two ablation procedures for right incisional atrial tachycardia, was referred to our institution to undergo pulmonary vein isolation (PVI) due to drug-refractory paroxysmal atrial fibrillation (AF). The rare congenital anomaly (IVC interruption with azygos continuation) precluded conventional inferior transfemoral venous approach; thus, a superior transjugular approach was planned. The procedure was performed under conscious sedation. Transseptal puncture under transesophageal echocardiography guidance was performed via right internal jugular vein with a PREFACE^R sheath (Biosense Webster) and a manually curved Brockenbrough needle with a 120° angle to manipulate the tip downward to the fossa ovalis. Electroanatomical (EA) mapping of the left atrium (CARTO, Biosense Webster) was performed using a multipolar catheter (PENTARAY^R, Biosense Webster). CARTO VIZIGOTM Bi-Directional


Guiding Sheath was then introduced and radiofrequency delivered at the pulmonary vein ostia. Complete PVI was confirmed by disappearance of venous potentials on the multipolar mapping catheter, and validated through exit block. No periprocedural complications occurred and the patient was discharged from the hospital in sinus rhythm.

Left atrial transseptal access is typically performed by inferior transfemoral venous approach. However, in a small subgroup of patients, such as those with congenital IVC interruption, a superior approach from the right internal jugular vein or left axillary/subclavian vein is required to gain access to the left atrium [1–3]. Alternatively, thoracoscopic AF ablation may be considered. In the reported case of AF ablation through unconventional trans-jugular approach, we propose the use of a guiding sheath, visualised on the electroanatomic mapping system, to significantly facilitate ablation catheter manipulation. Radiofrequency application reaching the target site with unconventional loops, such as those shown in Fig. 1 and in the videos (Supplementary Videos 1 and 2), would have hardly been possible without real time, continuous monitoring of the spatial relationship between the steerable guiding sheath and the ablation catheter.

✉ Andrea Saglietto
andrea.saglietto@live.com

¹ Division of Cardiology, Department of Medical Sciences, University of Turin, “Città della Salute e della Scienza”, Turin, Italy

Fig. 1 3D visualization of the steerable guiding sheath enabling the ablation catheter to reach target ablation site by unconventional loops. **A** Fluoroscopic anterior views during transseptal puncture (left) and left atrium mapping (right); **B** Latero-lateral 3D EA view during ablation of the ridge between the left superior pulmonary vein and the left appendage; **C** Postero-anterior 3D EA views during right pulmonary vein ablation; **D** Antero-posterior 3D EA view during right pulmonary vein ablation



Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10840-021-01114-8>.

Acknowledgements We thank Dr. Chiara Pintor (Clinical Support Specialist, Biosense Webster—Johnson & Johnson Medical) for the technical and editorial support.

Declarations

Conflict of interest Prof. Matteo Anselmino is consultant for Biosense Webster and has received educational grants from Abbott.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not

permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Santangeli P, et al. How to perform left atrial transseptal access and catheter ablation of atrial fibrillation from a superior approach. *J Cardiovasc Electrophysiol.* 2020;31:293–9.
2. Lim HE, et al. Catheter ablation of atrial fibrillation via superior approach in patients with interruption of the inferior vena cava. *Heart Rhythm.* 2009;6:174–9.
3. Kato H, et al. Circumferential pulmonary vein ablation of atrial fibrillation via superior vena cava approach in a patient with interruption of the inferior vena cava. *Europace.* 2010;12:746–8.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.