



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

LIC-KOR Promoted Formation of Conjugate Dienes as Useful Building Blocks in Palladium-Catalyzed Syntheses

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/10324 since
Publisher:
Elsevier Science Limited: Oxford Fulfillment Center, PO Box 800, Kidlington Oxford OX5 1DX United
Published version:
DOI:10.1016/j.tet.2005.01.044
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



Available online at www.sciencedirect.com



Tetrahedron

Tetrahedron 61 (2005) 3429-3436

LIC-KOR promoted formation of conjugated dienes as useful building blocks for palladium-catalyzed syntheses

Annamaria Deagostino,^a Manuele Migliardi,^a Ernesto G. Occhiato,^{a,b} Cristina Prandi,^{a,c} Chiara Zavattaro^a and Paolo Venturello^{a,*}

^aDipartimento di Chimica Generale ed Organica Applicata dell'Università, Via Pietro Giuria, 7 I-10125 Torino, Italy

^bDipartimento di Chimica Organica 'Ugo Schiff', and ICCOM, Università di Firenze, Via della Lastruccia, 13,

I-50019 Sesto Fiorentino, Italy

^cDipartimento di Scienze Ambiente e Vita, Università del Piemonte Orientale, Spalto Marengo, 33, I-15100 Alessandria, Italy

Received 24 September 2004; revised 11 November 2004; accepted 6 January 2005

Available online 2 February 2005

Abstract—It is demonstrated that α , β -unsaturated acetals can be considered a synthetic tool for transforming carbonyl derivatives into cheap and easily accessible starting materials for the construction of various and more complex structures. The lithium–potassium mixed superbase LIC-KOR induces a conjugate elimination reaction that converts α , β -unsaturated acetals into 1*E*-1-alkoxybuta-1,3-dienes. These derivatives can be readily metalated in situ and functionalized by reaction with electrophiles. The results can be grouped in two sections: (1) the palladium-catalyzed cross-coupling reaction between alkoxydienylboronates and tetralone- or isochromanone-derived vinyl triflates; (2) the regio- and stereoselective cross coupling reaction with aryl derivatives in the presence of a palladium catalyst (Heck conditions). © 2005 Elsevier Ltd. All rights reserved.

1. Introduction

A few years ago, we reported that in the presence of Schlosser's superbase LIC-KOR¹ (LIC: butyllithium, KOR: potassium *tert*-butoxide), α , β -unsaturated acetals selectively yield 1-alkoxybuta-1,3-dienes.² The product derives from a stereoselective conjugate elimination that is initiated by the metalation reaction at the γ -allylic position of the unsaturated substrate. Moreover, butadienyl derivatives can be further functionalized at their α -position by conducting the former elimination reaction in the presence of at least two equivalents of the LIC-KOR base: the excess reagent

selectively deprotonates the elimination product at the 1-alkenyl site, giving a nucleophile that can be quenched with various electrophiles, yielding substitution or addition products.³ By resorting to this procedure, we have described different functionalized unsaturated systems.⁴ More recently, the reactivity of α -metalated alkoxydienes in the presence of trialkylboranes⁵ and trialkylborates⁶ as electrophiles has also been described (Scheme 1). The arylation of α , β -unsaturated carbonyl compounds by an Heck reaction has also been reported.⁷

In this paper we provide the details of our work on two



Scheme 1. Synthesis of trialkylborates starting from α , β -unsaturated acetals, in the presence of LIC-KOR superbase.

0040–4020/\$ - see front matter @ 2005 Elsevier Ltd. All rights reserved. doi:10.1016/j.tet.2005.01.044

Keywords: Unsaturated acetals; Mixed superbases; Pd-catalyzed reactions; Suzuki reaction; Heck reaction; Nazarov reaction. * Corresponding author. Fax: +39 0112367646; e-mail: paolo.venturello@unito.it



X = N-Cbz, N-Ts, O Y = Tf, P(O)(OPh)₂ n = 0, 1, 2

Scheme 2. Pd-catalyzed cross-coupling reaction of lactam- and lactone derived triflates or phosphates with 2-(1-ethoxybuta-1,3-dienyl)-5,5-dimethyl-[1,3,2]-dioxaborinane, and subsequent Amberlyst-15[®] catalyzed cyclization.

different uses of alkoxydienyl derivatives in synthesis: (1) the Pd-catalyzed cross-coupling reaction with tetralone- or isochromanone-derived vinyl triflates or phosphates, and (2) the Heck coupling with diazonium salts.

1.1. Pd-catalyzed cross-coupling reaction with lactamand lactone-derived vinyl triflates or phosphates

We have recently found that conjugated ethoxytrienes obtained by the Pd-catalyzed cross-coupling reaction of lactam- and lactone- derived triflates or phosphates with ethoxydienylboronates (Scheme 2) undergo cyclization in the presence of the acidic resin Amberlyst-15[®] to give byciclic fused [3.3.0], [4.3.0], and [5.3.0] systems.⁸ A Nazarov-type electrocyclization can explain the cyclization reaction.⁹ This process has been extended here to the synthesis of tricarbocyclic derivatives by exploiting commercially available α -tetralones with different patterns of substitution.

Commercially available α -tetralones **1a–c** (Scheme 3) were converted into the corresponding triflates **2a–c** by treatment with LHMDS at -78 °C and by trapping the enolates with *N*-phenyl triflimide. The triflates thus obtained proved to be stable enough to be purified by flash chromatography and successively coupled with 2-(1-ethoxybuta-1,3-dienyl)-5,5,dimethyl-[1,3,2]-dioxaborinane.⁶ The reaction was performed with (Ph₃P)₂PdCl₂ (5%) as a catalyst in THF and 2 M K₂CO₃ as a base at room temperature and was complete in 1 h. Chromatography of the crude reaction mixture afforded pure ethoxytrienes **3a–c** in good yields that were used for the subsequent intramolecular cyclization. In our previous studies we have already shown that hydrolysis conditions strongly affect the reaction outcome. As a matter of fact the mechanism could be driven towards the synthesis of α,β -unsaturated ketone (5) or towards the Nazarov adducts (4) by choosing the appropriate acidic catalyst and solvent. In 0.02 M HCl MeOH-H₂O as a solvent, the open chain ketone was recovered as the main product and could be successively transformed into the Nazarov adduct by treatment with pure TFA at room temperature, while the acidic sulfonic resin Amberlyst-15[®] promoted the electrocyclic process. Ethoxytrienes 3a-c have been hydrolyzed under inert atmosphere using Amberlyst-15[®] as a catalyst and anhydrous CH₂Cl₂ as a solvent. In all examples products 4a-c were obtained in good yields, while open chain ketones 5a-c were merely obtained in traces. In particular, 4b was recovered as a 1:1 mixture of *cis* and trans diastereomers. We were not surprised to find a lack of diastereocontrol in the electrocyclization of 3b. In our studies on the torquoselectivity of the Nazarov reaction performed on ethoxytrienes in which one double bond is embedded in a heterocyclic structure, we have found that, while substituents in 2- or 4-position of the heterocyclic moiety could affect the diastereoselection, a substituent at position 3 exerted a low remote stereocontrol.⁸ In compound **3b** the methyl group in position 3 is too far from the reacting centers to affect the sense of electrocyclic conrotation so that the two faces of the endocyclic double bond are not differentiated and both modes of conrotation are thus possible.¹⁰

The here described synthetic sequence could be applicable to the construction of new derivatives structurally connected



Scheme 3. Pd-catalyzed cross-coupling reaction of α -tetralones derived triflates with 2-(1-ethoxybuta-1,3-dienyl)-5,5,-dimethyl-[1,3,2]-dioxaborinane, and subsequent Amberlyst-15[®] catalyzed cyclization.

to some attractive natural products, in particular to biologically active substances endowed with a ring system in which a substituted aromatic nucleus is fused to a hydrindane framework,11 as in the core of the hamigerans (Fig. 1). Hamigerans are a small group natural products that were isolated from the marine sponge Hamigera tarangaensis.¹² The biological activity profile of hamigerans is quite interesting: in particular, for hamigeran B3 has been found a 100% in vitro inhibition against both herpes and polio viruses.¹³ The absolute configuration of hamigeran A1 has been determined by X-ray crystallographic analysis.¹⁴ Because of their biological properties and novel structural features, the synthesis of hamigerans represents an important synthetic target.¹⁵ For the synthesis of an analogue of hamigerans containing an heteroatom in the [4.3.0] bicyclic moiety, the same synthetic sequence was applied to the aromatic lactone 3-isochroman-2-one 1d (Scheme 4). Triflate 2d was coupled with 2-(1-ethoxybuta-1,3-dienyl)-5,5,-dimethyl-[1,3,2]-dioxaborinane under the usual conditions [(Ph₃P)₂PdCl₂ (5%), THF, 2 M K₂CO₃, 25 °C] giving triene **3d**. The hydrolysis of **3d** with Amberlyst-15[®] afforded cyclic ketone 4d along with the open chain isomer 5d that, in this case, was unexpectedly isolated in significant yield. However, it could be easily converted into 4d by treatment with pure TFA at 25 °C.





1.2. Heck coupling with diazonium salts

Substituted dienes and other unsaturated structures are very attractive as precursors of dyes, UV screens, and drugs.¹⁶ Although arylation of two-carbon vinyl fragments might sometimes result in scarcely regioselective reactions,^{17,18} highly regioselective α -arylation and α -vinylation

procedures have been reported under specific experimental conditions.¹⁹ In particular, 100% regioselective α -functionalization of vinyl ethers can be achieved by favoring the coordination of the Pd complex to the carboncarbon double bond via dissociation of the anion ligand. Such a mechanism can be promoted by using triflate as a leaving group or adding a sequestering agent of halide anions.²⁰ Otherwise, a mixture of isomers is obtained when the coordination-insertion process proceeds via dissociation of one neutral ligand. Moreover, regioselective β -arylation of vinyl ethers has been achieved when the olefinic substrate contains groups that control the palladium catalyzed reaction through chelation. Such a procedure has been reported as a useful access to arylethylamines or arylacetic acids of significant pharmaceutical value.²¹ On the other hand, the Pd-catalyzed arylation of conjugate dienes would be expected to produce arylated dienes. However, some differences and perhaps complications could be expected since π -allylic palladium species are involved in the reactions. Particularly stable π -allylic complexes in the presence of tertiary amines are formed, while the use of secondary amines leads to mixtures of polyenes and allylic amines.²² Thus, low yields and/or low selectivity are encountered when the diene does not have a carbonyl or phenyl group and the synthesis of linear conjugated polyenes remains to be improved. Recently we reported a regioselective γ -arylation of α , β -unsaturated carbonyl compounds from aryl iodides and alkoxydienes by an Heck reaction.' We observed the stereoselective formation of γ -arylated α , β -unsaturated acetals in good yields working in DMSO at 80 °C, and using K₂CO₃ as a base and Pd(OAc)₂ as a catalyst (Scheme 5(A)). A possible mechanism has been also suggested: first, the arylpalladium intermediate adds to the terminal double bond of the diene, then this intermediate arranges to the π -allylic complex and an iodide ion-acetate ligand exchange takes place. Finally, attack of the hydroxy group upon the complex displaces the palladium complex and gives the cyclic acetal.

In this paper we wish to report the results obtained studying the Pd-catalyzed cross coupling reaction of 1-alkoxy-1,3butadienes with arendiazonium salts. The Heck reaction with these substrates is mild and fast. Moreover, the use of these compounds is synthetically more convenient than the use of aryl halides, since many of them, especially iodides, are prepared from diazonium salts. Interestingly, no



Scheme 4. Pd-catalyzed cross-coupling reaction of 3-isochroman-2-one derived triflate with 2-(1-ethoxybuta-1,3-dienyl)-5,5,-dimethyl-[1,3,2]-dioxaborinane, and subsequent Amberlyst-15[®] catalyzed cyclization.





B: (a) ArN₂⁺ BF₄⁻, AcONa, 25 °C, Pd(OAc)₂, MeCN

Scheme 5. Heck reaction on 1-(3-hydropropoxy)buta-1,3-dienes.

difference in reactivity in relation to the electronic nature of the diazonium compound is observed.²³ Moreover it seems that for this type of Heck reaction, contrary to the reactions of aryl chlorides or bromides, the insertion of the palladium catalyst is not the rate determining step.²⁴ Unfortunately, the process is rather inefficient as the loading of Pd precatalyst required is usually no less than 1–2 mol%. Most probably this is associated with an inefficient preactivation of the catalyst.

The use of bases and strongly coordinating ligands in reactions with arendiazonium salts is usually detrimental as a consequence of dediazotation:²⁵ unfortunately, basic conditions are required since the starting alkoxydienes and products are acid sensitive and the formation of acids caused by β -H-elimination can be problematic.²⁶ In fact, some tests carried out without addition of base did not give the expected arylated compound in appreciable amount.

The results of the cross coupling reaction are reported in Scheme 5(B). These data have been obtained working in anhydrous MeCN (alcoholic solvents turned out inadequate) at room temperature in the presence of NaOAc as a base and of $Pd(OAc)_2$ as a catalyst.

First of all, we focused our attention on halo-substituted diazonium salts because of the interest towards differential Pd-catalyzed reaction in the synthesis of unsymmetrical substituted teraryls and oligoarylenesvinylenes considering their medicinal and NLO properties.²⁷

The reactions are regio- and stereoselective, and the aryl group adds, as expected, to the unsubstituted CH_2 site of the conjugate system. Unfortunately, the coupling yields are modest: probably they are affected by the stability of whether the diazonium salts or the dienes under the reaction conditions.

When the cross-coupling reaction was carried out on (E)-4ethoxyocta-1,3-diene **6d** the corresponding (2E)- γ -arylated α , β -unsaturated ketone **7h** was recovered (Scheme 6). The formation of the conjugate ketone can be most likely attributed to the presence of water molecules that may be present in the diazonium salt or in the base.



Conditions: (a) p-MeOC₆H₄N₂⁺ BF₄⁻, AcONa, r.t., Pd(OAc)₂, MeCN

Scheme 6. Heck reaction on (E)-4-ethoxyocta-1,3-diene.

2. Conclusions

In summary, we have developed new synthetic routes that allow the acetal group of α,β -unsaturated acetals to be considered not simply a protective function but also a useful synthetic tool. In particular, we have shown that Suzuki coupling of ethoxydienylboronates with vinyl triflates or phosphates derived from aromatic ketones or lactones is a tactically suitable strategy for the synthesis of cyclopentafused polycyclic systems that are present in many natural products. Moreover, when 1-alkoxybuta-1,3-dienes cross couple under Heck conditions with arendiazonium derivatives, the final outcome of the process formally corresponds to the direct γ -arylation of starting α,β -unsaturated acetals.

3. Experimental

3.1. General

Flasks and all equipments used for the generation and reaction of moisture-sensitive compounds are flame dried under Argon. Anhydrous CH₃CN was purchased by Fluka. All aromatic substrates, palladium(II) acetate, palladium bistriphenylphosphinobichloride, NaOAc and K₂CO₃ are commercially available and are used as received. All the acetals and ethoxydienes were prepared as previously described.⁴ The arenediazonium tetrafluoborates were prepared from commercial aromatic amines according to reported procedures.²⁸ Purification of products was carried out by preparative column chromatography on Merck Silica gel 60 with light petroleum ether (distillation range 40-60 °C)-Et₂O as eluent. ¹H NMR spectra were recorded at 200 or 400 MHz in CDCl₃, using \overline{TMS} as internal standard. Coupling constants (J) are given in Hz and coupling patterns are described by abbreviations: s (singlet), d (doublet), t (triplet), q (quartet), br s (broad singlet). ¹³C NMR spectra are recorded at 50 and 100.4 MHz in CDCl₃, and chemical shifts are determined relative to the residual solvent peak (77.36). GC-MS spectra are obtained on a mass selective detector HP 5970 B instrument operating at an ionizing voltage of 70 eV connected to a HP 5890 GC, cross linked methyl silicone capillary column ($25 \text{ m} \times 0.2 \text{ mm} \times$ 0.33 µm film thickness).

3.1.1. Trifluoromethanesulfonic acid 3,4-dihydronaphthalen-1-yl ester (2a). Typical procedure. To a solution of KHMDS (8 mL of a 0.5 M solution in toluene, 4 mmol) in THF (12 mL), cooled at -78 °C and under nitrogen atmosphere, was added a solution of 3,4-dihydro-2*H*-naphthalen-1-one (440 mg, 3 mmol) in THF (4 mL) and the resulting mixture was stirred for 1.5 h. Afterward a solution of PhNTf₂ (964 mg, 2.7 mmol) in THF (2 mL) was quickly added, leaving under stirring for 1 h at -78 °C before allowing the temperature to rise to 0 °C. Then, a 10% NaOH solution (20 mL) was added, the mixture was extracted with Et₂O (3×20 mL), washed with H₂O (2× 10 mL), and dried (K₂CO₃). After filtration and evaporation of the solvent, crude vinyl triflate **1** was obtained as a yellowish oil and purified by flash chromatography (petroleum ether: Et₂O 4:1, 610 mg, 73%): 200 MHz ¹H NMR (200 MHz, CDCl₃): δ 7.3 (m, 4H), 6.05 (t, *J*=2.9 Hz, 1H), 2.90 (t, *J*=6.5 Hz, 2H), 2.5 (m, 2H).

3.1.2. Trifluoromethanesulfonic acid 4-methyl-3,4dihydro-naphthalen-1-yl ester (2b). Flash chromatography (Et₂O-petroleum ether 1:4, 0.5% Et₃N) gave pure **2b** (531 mg, 61%, $R_{\rm f}$ 0.75) as a yellow oil. ¹H NMR (200 MHz, CDCl₃): δ 7.2 (m, 4H), 6.01 (t, J=2.9 Hz, 1H), 3.01 (hex, J=6.5 Hz, 1H), 2.75 (ddd, J=13.0, 6.5, 2.9 Hz, 1H), 2.29 (dd, J=13.0, 2.9 Hz, 1H), 1.35 (d, J=6.5 Hz, 3H).

3.1.3. Trifluoromethanesulfonic acid 5-methoxy-3,4dihydro-naphthalen-1-yl ester (2c). Flash chromatography (Et₂O-petroleum ether 1:4, 0.5% Et₃N) gave pure **2c** (678 mg, 73%, $R_{\rm f}$ 0.76) as a yellow oil. ¹H NMR (200 MHz, CDCl₃): δ 7.33 (t, J=5.9 Hz, 1H), 7.18 (d, J= 5.9 Hz, 1H), 6.97 (d, J=5.9 Hz, 1H), 6.04 (t, J=2.9 Hz, 1H), 3.95 (s, 3H), 2.95 (t, J=5.5 Hz, 2H), 2.65 (m, 1H), 2.5 (m, 1H).

3.1.4. Trifluoromethanesulfonic acid 1H-isochromen-3yl ester (2d). Flash chromatography (Et₂O–petroleum ether 3:7, 0.5% Et₃N) gave pure **2d** (414 mg, 49%, $R_{\rm f}$ 0.74) as a yellow oil. ¹H NMR (200 MHz, CDCl₃): δ 7.2 (m, 4H), 5.81 (s, 1H), 5.37 (s, 2H).

E-4-(1-Ethoxy-buta-1,3-dienyl)-1,2-dihydro-3.1.5. naphthalene (3a). Typical procedure. To a solution of crude 2a (417 mg, 1.5 mmol) in THF (4 mL) were added, under a nitrogen atmosphere, (Ph₃P)₂PdCl₂ (57 mg, 0.075 mmol), 2-(1-ethoxybuta-1,3-dienyl)-5,5,-dimethyl-[1,3,2]-dioxaborinane (315 mg, 1.5 mmol), and a 2 M aqueous K_2CO_3 solution (1.5 mL). The mixture was stirred for 2.5 h at 25 °C. H₂O (10 mL) was then added, the mixture extracted with diethyl ether $(3 \times 20 \text{ mL})$ and dried (Na_2SO_4) . Evaporation of the solvent afforded a vellow oil that was purified by flash chromatography (petroleum ether-Et₂O 4:1, 0.5% Et₃N, R_f 0.7) to give **4a** (180 mg, 53%) as a colorless oil. ¹H NMR (200 MHz, CDCl₃): δ 7.09 (s, 4H), 6.38 (dt, J = 15.0, 10.5 Hz, 1H), 6.10 (t, J = 2.9 Hz, 1H), 5.65 (d, J = 10.5 Hz, 1H), 5.06 (dd, J = 15.0, 1.8 Hz, 1H), 4.81 (dd, J=10.5, 1.8 Hz, 1H), 3.84 (q, J=7.5 Hz, 2H), 2.82 (t, J=6.5 Hz, 2H), 2.42 (m, 2H), 1.27 (t, J=7.5 Hz, 3H); ¹³C NMR (50 MHz, CDCl₃): δ 152.6, 143.6, 137.5, 136.4, 135.4, 134.0, 133.1, 130.4, 129.4, 128.0, 125.8, 119.2, 69.8, 42.6, 29.3, 19.5; MS *m*/*z* 226 (M⁺, 100), 197 (35), 181 (37), 115 (23). Anal. Calcd for C₁₆H₁₈O: C, 84.91; H, 8.02. Found: C, 84.79; H, 8.72.

3.1.6. 4-(1-Ethoxy-buta-1,3-dienyl)-1-methyl-1,2dihydro-naphthalene (3b). Flash chromatography (Et₂Opetroleum ether 1:9, 0.5% Et₃N) gave pure **3b** (165 mg, 69%, $R_{\rm f}$ =0.85) as a colorless oil. ¹H NMR (CDCl₃, 200 MHz): δ 7.3 (m, 4H), 6.30 (dt, *J*=15.0, 10.5 Hz, 1H), 6.06 (t, *J*=2.9 Hz, 1H), 5.66 (d, *J*=10.5 Hz, 1H), 5.06 (dd, *J*=15.0, 1.8 Hz, 1H), 4.81 (dd, *J*=10.5, 1.8 Hz, 1H), 3.87 (q, *J*=7.5 Hz, 2H), 2.97 (hex, *J*=6.5 Hz, 1H), 2.45 (ddd, *J*=15.0, 6.5, 2.9 Hz, 1H), 2.20 (dd, *J*=15.0, 2.9 Hz, 1H), 1.28 (m, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ 160.39, 143.20, 130.83, 130.26, 130.13, 129.65, 129.01, 128.90, 128.30, 127.55, 113.92, 108.06, 65.82, 34.08, 33.85, 22.77, 17.93; MS *m*/*z* 240 (M⁺, 100), 211 (16), 193 (13), 129 (15). Anal. Calcd for C₁₇H₂₀O: C, 84.96; H, 8.39. Found: C, 84.12; H, 8.09.

4-(1-Ethoxy-buta-1,3-dienyl)-8-methoxy-1,2-3.1.7. dihydro-naphthalene (3c). Flash chromatography (Et₂Opetroleum ether 2:8, 0.5% Et₃N) gave pure 3c (154 mg, 60%, $R_{\rm f}$ =0.80) as a colorless oil. ¹H NMR (CDCl₃, 200 MHz): δ 7.01 (t, J=5.5 Hz, 1H), 6.96 (d, J=5.5 Hz, 1H), 6.85 (d, J = 5.5 Hz, 1H), 6.72 (dt, J = 15.0, 10.5 Hz, 1H), 6.11 (t, J = 3.0 Hz, 1H), 5.66 (d, J = 10.5 Hz, 1H), 5.09 (dd, J=15.0, 1.8 Hz, 1H), 4.85 (dd, J=10.5, 1.8 Hz, 1H),3.86 (q, J=7.5 Hz, 2H), 3.84 (s, 3H), 2.83 (t, J=5.2 Hz, 2H), 2.36 (m, 2H), 1.33 (t, J=7.5 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ 160.65, 158.66, 136.84, 136.16, 134.52, 129.17, 127.52, 120.26, 113.82, 112.52, 107.78, 103.31, 67.02, 58.18, 25.14, 21.99, 18.23; MS *m/z* 256 (M⁺, 100), 225 (22), 211 (32), 199 (20), 165 (23). Anal. Calcd for C₁₇H₂₀O₂: C, 79.65; H, 7.86. Found: C, 79.21; H, 7.19.

3.1.8. 3-(1-Ethoxy-buta-1,3-dienyl)-1*H***-isochromene** (**3d**). Flash chromatography (Et₂O–petroleum ether 2:8, 0.5% Et₃N) gave pure **3d** (115 mg, 50%, $R_{\rm f}$ =0.85) as a colorless oil. ¹H NMR (CDCl₃, 200 MHz): δ 7.2 (m, 5H), 6.34 (s, 1H), 5.68 (d, *J*=10.5 Hz, 1H), 5.14 (s, 2H), 5.10 (dd, *J*=15.0, 1.8 Hz, 1H), 4.95 (dd, *J*=10.5, 1.8 Hz, 1H), 3.88 (q, *J*=7.5 Hz, 2H), 1.33 (t, *J*=7.5 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ 156.58, 153.59, 135.73, 133.70, 130.80, 129.58, 127.91, 126.63, 126.35, 117.26, 111.42, 108.59, 71.12, 66.80, 18.23; MS *m*/*z* 228 (M⁺, 89), 199 (39), 155 (34), 128 (95), 77 (100). Anal. Calcd for C₁₅H₁₆O₂: C, 78.92; H, 7.06. Found: C, 78.41; H, 7.89.

3-Methyl-2,3,4,5-tetrahydro-cyclopenta[a]-3.1.9. naphthalen-1-one (4a). Typical procedure. To a solution of 3a (180 mg, 0.80 mmol) in anhydrous DCM (7 mL) under argon atmosphere, Amberlyst- $15^{\text{\tiny (B)}}$ (2.3 mequiv/g, 18 mg) was added and the resulting mixture was stirred at 25 °C. The reaction was monitored by TLC: after 2 h the resin was filtered off through a short pad of K₂CO₃ and concentrated under vacuum. Crude products were purified by flash chromatography (petroleum ether-Et₂O 1:1, 0.5%) Et₃N, R_f 0.4) to give pure **4a** (110 mg, 69%). ¹H NMR $(CDCl_3, 400 \text{ MHz}): \delta 8.21 \text{ (d}, J = 6.2 \text{ Hz}, 1\text{H}), 7.24 \text{ (m}, 3\text{H}),$ 2.95 (br s, 3H), 2.81 (dd, J = 19.4, 5.5 Hz, 1H), 2.69 (m, 1H), 2.58 (m, 1H), 2.16 (d, J = 19.4 Hz, 1H), 1.35 (d, J = 6.5 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ 205.60, 178.91, 134.46, 134.17, 129.21, 127.90, 127.57, 126.85, 124.26, 44.85, 35.39, 27.79, 24.80, 18.74; MS *m*/*z* 198 (M⁺, 100), 183 (42), 170 (53), 155 (60). Anal. Calcd for C₁₄H₁₄O: C, 84.81; H, 7.12. Found: C, 84.21; H, 7.88.

3.1.10. 3,5-Dimethyl-2,3,4,5-tetrahydro-cyclopenta[*a*]-**naphthalen-1-one** (**4b**). Flash chromatography (Et₂O-petroleum ether 1:1, 0.5% Et₃N) gave pure **4b** as 1:1 mixture of diastereoisomers. (110 mg, 63%, $R_{\rm f}$ =0.35) as a colorless oil. ¹H NMR (CDCl₃, 400 MHz): δ 8.25 (d,

J=6.2 Hz, 2H *cis* + *trans*), 7.23 (m, 6H, *cis* + *trans*), 3.1 (m, 2H, *cis* + *trans*), 2.8 (m, 2H, *cis* + *trans*), 2.82 (dd, J=17.9, 5.1 Hz, 1H *trans*), 2.80 (dd, J=18.3, 7.5 Hz, 2H *cis* + *trans*), 2.67 (dd, J=17.9, 5.0 Hz, 1H *cis*), 2.48 (dd, J=17.9, 6.6 Hz, 1H *cis*), 2.33 (dd, J=17.9, 6.0 Hz, 1H *trans*), 2.17 (d, J=18.3 Hz, 2H, *cis* + *trans*), 1.25 (d, J=6.2 Hz, 3H), 1.24 (d, J=6.2 Hz, 3H), 1.23 (d, J=6.2 Hz, 3H), 1.22 (d, J=6.2 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz): δ 205.60, 205.33, 177.72, 177.60, 139.80, 139.62, 128.20, 126.72, 126.62, 126.32, 124.43, 44.90, 44.79, 32.74, 32.68, 32.60, 32.48, 21.68, 20.91, 19.26, 18.51; MS *m/z* 212 (M⁺, 97), 197 (47), 184 (22), 155 (100). Anal. Calcd for C₁₅H₁₆O: C, 84.87; H, 7.60. Found: C, 84.21; H, 7.88.

3.1.11. 6-Methoxy-3-methyl-2,3,4,5-tetrahydro-cyclopenta[*a*]naphthalen-1-one (4c). Flash chromatography (Et₂O-petroleum ether 3:7, 0.5% Et₃N) gave pure 4c (120 mg, 53%, $R_{\rm f}$ =0.4) as a colorless oil. ¹H NMR (CDCl₃, 200 MHz): δ 8.21 (d, J=5.5 Hz, 1H), 7.24 (t, J= 5.5 Hz, 1H), 6.85 (d, J=5.5 Hz, 1H), 3.83 (s, 3H), 2.95 (m, 3H), 2.78 (dd, J=19.4, 5.5 Hz, 1H), 2.6 (m, 2H), 2.15 (d, J=19.4 Hz, 1H), 1.35 (d, J=6.5 Hz, 3H); ¹³C NMR (CDCl₃, 50 MHz) δ 205.45, 178.22, 156.09, 127.10, 126.30, 125.51, 116.94, 110.36, 109.65, 55.44, 44.72, 35.19, 24.16, 19.82, 15.22; MS *m*/*z* 228 (M⁺, 100), 200 (40), 185 (34), 115 (28). Anal. Calcd for C₁₅H₁₆O₂: C, 78.92; H, 7.06. Found: C, 78.25; H, 7.88.

3.1.12. 3-Methyl-2,3-dihydro-5*H***-cyclopenta[***c***]isochromen-1-one (4d). Flash chromatography (Et₂Opetroleum ether 45:55, 0.5% Et₃N) gave pure 4d (118 mg, 59%, R_f=0.4) as a colorless oil. ¹H NMR (CDCl₃, 200 MHz): \delta 7.3 (m, 4H), 5.32 (d,** *J***=8.5 Hz, 2H), 3.28 (m, 1H), 2.78 (dd,** *J***=19.3, 5.5 Hz, 1H), 2.18 (d,** *J***= 19.4 Hz, 1H), 1.37 (d,** *J***=6.5 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) \delta 196.89, 144.44, 128.58, 128.28, 126.70, 126.43, 126.20, 122.70, 120.42, 67.79, 40.44, 25.70, 18.86; MS** *m***/***z* **200 (M⁺, 100), 200 (40), 185 (32), 158 (21), 115 (56), 64 (25). Anal. Calcd for C₁₃H₁₂O₂: C, 77.98; H, 6.04. Found: C, 77.55; H, 6.04.**

3.1.13. 1-(1*H***-Isochromen-3-yl)-but-2-en-1-one (5d).** (40 mg, 20%, $R_{\rm f}$ =0.75) as a colorless oil. ¹H NMR (CDCl₃, 200 MHz): δ 7.2 (m, 4H), 7.17 (dq, *J*=15.5, 6.5 Hz, 1H), 7.05 (s, 1H), 6.84 (d, *J*=15.5 Hz, 1H), 5.19 (s, 2H), 1.97 (d, *J*=6.5 Hz, 3H); ¹³C NMR (CDCl₃, 50 MHz) δ 185.14, 151.59, 145.00, 129.96, 129.65, 129.35, 128.68, 125.96, 125.84, 124.24, 111.96, 68.94, 18.77; MS *m/z* 200 (M⁺, 23), 103 (26), 77(17), 69 (100). Anal. Calcd for C₁₃H₁₂O₂: C, 77.98; H, 6.04. Found: C, 77.15; H, 6.84.

3.2. General procedure for the arylation of ethoxydienes (6a–c)

Ethoxydiene (2 mmol) was added to a mixture of the corresponding arendiazonium tetrafluoborate (1 mmol), Pd(OAc)₂ (0.05 mmol), K₂CO₃ or NaOAc (1 mmol) in anhydrous CH₃CN (6 mL) degassed with argon for 10 min. The reaction mixture was stirred under argon in a sealed tube for 15–30 min. Samples were periodically taken and tested with β -naphtol. After complete consumption of the starting tetrafluoborate, H₂O (10 mL) was added. All reactions were worked up by extraction with Et₂O (3×

20 mL) and the organic phases were washed with brine (3 \times 10 mL). The products were purified by column chromatography on SiO₂ deactivated with 1% of Et₃N. Eluent: petroleum ether/Et₂O.

3.2.1. (*E*)-2-[03-(4-Methoxyphenyl)prop-1-enyl]-1,3dioxane (7a). 70 mg, (30%) as a colorless oil (70/30). ¹H NMR (200 MHz, CDCl₃) δ 7.02 (d, *J*=8.6 Hz, 2H), 6.75 (d, *J*=8.6 Hz, 2H), 5.97 (dt, *J*=16.4, 6.9 Hz, 1H), 5.42 (ddt, *J*=16.4, 5.2, 1.5 Hz, 1H), 4.95 (d, *J*=5.2 Hz, 1H), 4.05 (m, 2H), 3.77 (m, 5H), 3.28 (d, *J*=6.9 Hz, 2H), 2.0 (m, 1H), 1.3 (m, 1H); ¹³C NMR (50 MHz, CDCl₃) δ 158.01, 134.48, 129.66, 127.94, 113.84, 100.86, 66.88, 55.23, 34.48, 25.64; *m*/*z* (EI, 70 eV rel. Int.) 234 (M⁺, 38), 147 (37), 113 (100), 87 (92), 55 (31). ν_{max} (neat)/cm⁻¹ 3050, 1650, 1635, 1251, 823, 808. (Found: C, 71.00%; H 7.93%. Calcd for C₁₄H₁₈O₃: C, 71.77%; H, 7.74%).

3.2.2. (*E*)-2-[3-(3-Methoxyphenyl)prop-1-enyl]-1,3-dioxane (7b). 86 mg, (37% yield) as a colorless oil (70/30). ¹H NMR (200 MHz, CDCl₃) δ 7.2 (m, 2H), 6.7 (m, 2H), 6.05 (dt, *J*=16.0, 6.6 Hz, 1H), 5.55 (dm, *J*=16.0 Hz, 1H), 4.95 (d, *J*=5.6 Hz, 1H), 4.10 (m, 2H), 3.80 (m, 5H), 3.38 (d, *J*=6.6 Hz, 2H), 1.5 (m, 2H); ¹³C NMR (50 MHz, CDCl₃) δ 159.70, 140.89, 133.83, 129.38, 128.33, 121.12, 114.40, 111.58, 100.78, 66.90, 55.14, 38.44, 25.67; *m/z* (EI, 70 eV rel. Int.) 234 (M⁺, 21), 158 (38), 147 (17), 113 (100), 83 (23). *v*_{max} (neat)/cm⁻¹ 3010, 2853, 1599, 1454, 1259, 1078, 993, 773. (Found: C, 71.90%; H 7.02%. Calcd for C₁₄H₁₈O₃: C, 71.77%; H, 7.74%).

3.2.3. (*E*)-2-[3-(4-Iodophenyl)prop-1-enyl]-1,3-dioxane (7c). 112 mg, (34% yield) as a colorless oil (70/30). ¹H NMR (200 MHz, CDCl₃) δ 7.65 (d, *J*=7.4 Hz, 2H), 6.95 (d, *J*=7.4 Hz, 2H), 6.05 (dt, *J*=15.4, 6.2 Hz, 1H), 5.50 (dbd, *J*=15.4, 6.2 Hz, 1H), 4.95 (d, *J*=6.2 Hz, 1H), 4.15 (m, 2H), 3.80 (m, 2H), 3.30 (d, *J*=6.5 Hz, 2H), 2.1 (m, 1H), 1.3 (m, 1H); ¹³C NMR (50 MHz, CDCl₃) δ 139.20, 137.42, 133.12, 130.79, 128.73, 100.53, 91.30, 66.85, 37.77, 25.61; *m/z* (EI, 70 eV rel. Int.) 330 (M⁺, 5), 144 (15), 113 (100), 100 (36), 55 (29). ν_{max} (neat)/cm⁻¹ 3050, 2851, 1279, 1078, 930, 800. (Found: C, 47.85%; H 4.99%. Calcd for C₁₃H₁₅IO₂: C, 47.29%; H, 4.58%).

3.2.4. (*E*)-2-[3-(4-Chlorophenyl)prop-1-enyl]-1,3-dioxane (7d). 81 mg, (34% yield) as a colorless oil (90/10). ¹H NMR (200 MHz, CDCl₃) δ 7.25 (d, *J*=7.0 Hz, 2H), 7.05 (d, *J*=7.0 Hz, 2H), 6.20 (dt, *J*=14.9, 6.0 Hz, 1H), 5.55 (dbd, *J*=14.9, 5.8 Hz, 1H), 4.95 (d, *J*=5.8 Hz, 1H), 4.15 (m, 2H), 3.85 (m, 2H), 3.35 (d, *J*=6.0 Hz, 2H), 1.6 (m, 1H), 1.3 (m, 1H); ¹³C NMR (50 MHz, CDCl₃) δ 133.33, 132.05, 130.06, 129.50, 128.67, 128.51, 100.55, 66.88, 37.65, 25.61; *m/z* (EI, 70 eV rel. Int.) 238 (M⁺, 56), 180 (46), 145 (100), 115 (50), 68 (40), 55 (50). *v*_{max} (neat)/cm⁻¹ 3080, 1599, 1259, 1078, 993, 773. (Found: C, 66.05%; H 6.65%. Calcd for C₁₃H₁₅ClO₂: C, 65.41%; H, 6.33%).

3.2.5. (*E*)-2-[3-(4-Iodophenyl)-2-methylprop-1-enyl)-5,5dimethyl-1,3-dioxane (7e). 100 mg, (27% yield) as a colorless oil (90/10). ¹H NMR (200 MHz, CDCl₃) δ 7.62 (d, *J*=8.0 Hz, 2H), 6.94 (d, *J*=8.0 Hz, 2H), 5.11 (bd, *J*= 6.3 Hz, 1H), 5.08 (d, *J*=6.3 Hz, 1H), 3.65 (m, 2H), 3.51 (m, 2H), 3.23 (s, 2H), 1.68 (s, 3H), 1.23 (s, 3H), 0.75 (s, 3H); ¹³C NMR (50 MHz, CDCl₃) δ 141.44, 139.31, 138.42, 137.98, 137.41, 131.23, 124.27, 98.75, 45.12, 30.02, 22.97, 21.95, 17.03; *m*/*z* (EI, 70 eV rel. Int.) 372 (M⁺, 3), 155 (86), 128 (14), 115 (15), 69 (100). (Found: C, 49.20%; H 5.15%. Calcd for C₁₄H₁₇IO₂: C, 48.85%; H, 4.98%).

3.2.6. (*E*)-2-[3-(4-Chlorophenyl)-2-methylprop-1-enyl)-1,3-dioxane (7f). 63 mg, (25% yield) as a colorless oil (90/10). ¹H NMR (200 MHz, CDCl₃) δ 7.35 (d, *J*=8.6 Hz, 2H), 7.05 (d, *J*=8.6 Hz, 2H), 5.25 (d, *J*=6.2 Hz, 1H), 5.15 (d, *J*=6.2 Hz, 1H), 4.15 (m, 2H), 3.80 (m, 2H), 3.25 (s, 2H), 2.3 (m, 1H), 1.65 (s, 3H), 1.2 (m, 1H); ¹³C NMR (50 MHz, CDCl₃) δ 141.25, 137.16, 130.42, 128.42, 124.53, 98.75, 66.90, 44.88, 25.58, 16.98; *m*/*z* (EI, 70 eV rel. Int.) 252 (M⁺, 2), 127 (100), 87 (13), 69 (42), 41 (16). *v*_{max} (neat)/ cm⁻¹ 3050, 1490, 1376, 1138, 1086, 994, 787. (Found: C, 67.00%; H 7.10%. Calcd for C₁₄H₁₇ClO₂: C, 66.53%; H, 6.78%).

3.2.7. (*E*)-2-[3-(4-Iodophenyl)-1-methylprop-1-enyl)-1,3dioxane (7g). 103 mg, (30% yield) as a colorless oil (90/10). ¹H NMR (200 MHz, CDCl₃) δ 7.57 (d, *J*=7.0 Hz, 2H), 6.95 (d, *J*=7.0 Hz, 2H), 5.75 (bt, *J*=6.6 Hz, 1H), 4.82 (s, 1H), 4.10 (m, 2H), 3.82 (m, 2H), 3.15 (d, *J*=6.8 Hz, 2H), 2.1 (m, 1H), 1.80 (s, 3H), 1.1 (m, 1H); ¹³C NMR (50 MHz, CDCl₃) δ 137.89, 137.42, 130.54, 126.99, 104.50, 91.00, 67.02, 33.14, 25.72, 15.00; *m/z* (EI, 70 eV rel. Int.) 344 (M⁺, 7), 127 (100), 87 (57), 69 (33), 59 (15). ν_{max} (neat)/cm⁻¹ 3050, 1483, 1377, 1151, 1107, 958, 801. (Found: C, 49.15%; H 4.75%. Calcd for C₁₄H₁₇IO₂: C, 48.85%; H, 4.98%).

3.2.8. (*E*)-1-(4-Methoxyphenyl)-ott-2-en-4-one (8). 50 mg, (22% yield) as a colorless oil (90/10). ¹H NMR (200 MHz, CDCl₃) δ 7.02 (d, *J*=7.2 Hz, 2H), 6.9 (m, 3H), 6.05 (bd, *J*=15.0 Hz, 1H), 3.80 (s, 3H), 3.45 (bd, *J*= 6.2 Hz, 2H), 2.55 (t, *J*=6.1 Hz, 2H), 1.55 (pent, *J*=6.0 Hz, 2H), 1.32 (sext, *J*=6.0 Hz, 2H), 0.9 (t, *J*=6.0 Hz, 3H); ¹³C NMR (50 MHz, CDCl₃) δ 200.66, 158.32, 145.40, 130.76, 129.66, 114.00, 55.14, 39.71, 37.74, 26.17, 22.28, 13.76; *m*/z (EI, 70 eV rel. Int.) 232 (M⁺, 66), 160 (44), 175 (34), 147 (100), 121 (29). ν_{max} (neat)/cm⁻¹ 3030, 1672, 1626, 1458, 1246, 1033, 733. (Found: C, 78.05%; H 8.20%. Calcd for C₁₅H₂₀O₂: C, 77.55%; H, 8.68%).

Acknowledgements

Authors thank the Università degli Studi di Torino and the Italian MIUR for financial support through the 'Cofinanziamento di Programmi di Ricerca di Rilevante Interesse Nazionale'.

References and notes

 Schlosser, M. J. Organomet. Chem. 1967, 8, 9–16. Schlosser, M. Mod. Synth. Methods 1992, 6, 227–271. Mordini, A. In Snieckus, V., Ed.; Advances in Carbanion Chemistry; JAI: Greenwich, CT, 1992; Vol. 1, pp 1–45. Schlosser, M.; Faigl, F.; Franzini, L.; Geneste, H.; Katsoulos, G.; Zhong, G. Pure Appl. Chem. 1994, 66, 1439–1446. Lochmann, L. Eur. J. Inorg. Chem. 2000, 1115–1126.

- 2. Venturello, P. J. Chem. Soc., Chem. Commun. 1992, 1032–1033.
- 3. Prandi, C.; Venturello, P. J. Org. Chem. **1994**, 59, 3494–3496. (and 5458–5462).
- Balma Tivola, P.; Deagostino, A.; Prandi, C.; Venturello, P. J. Chem. Soc., Perkin Trans. 1 2001, 437–441. Deagostino, A.; Prandi, C.; Venturello, P. Curr. Org. Chem. 2003, 7, 821–839.
- Balma Tivola, P.; Deagostino, A.; Prandi, C.; Venturello, P. *Chem. Commun.* 2001, 1536–1537. Deagostino, A.; Prandi, C.; Zavattaro, C.; Venturello, P. *Eur. J. Org. Chem.* 2003, 2612–2616.
- Balma Tivola, P.; Deagostino, A.; Prandi, C.; Venturello, P. Org. Lett. 2002, 4, 1275–1277.
- Deagostino, A.; Prandi, C.; Venturello, P. Org. Lett. 2003, 5, 3815–3817.
- Occhiato, E. G.; Prandi, C.; Ferrali, A.; Guarna, A.; Deagostino, A.; Venturello, P. J. Org. Chem. 2002, 67, 7144–7146. Occhiato, E. G.; Prandi, C.; Ferrali, A.; Guarna, A.; Venturello, P. J. Org. Chem. 2003, 68, 9728–9741. Prandi, C.; Ferrali, A.; Guarna, A.; Venturello, P.; Occhiato, E. G. J. Org. Chem. 2004, 69, 0000.
- Nazarov, I. N.; Torgov, I. B.; Terekhova, L. N. *Izv. Akad. Nauk* SSSR, Otd. Khim. Nauk 1942, 200. Braude, E. A.; Forbes, W. F. J. Chem. Soc. 1953, 2208–2216. Habermas, K. L.; Denmark, S. E.; Jones, T. D. Org. React. 1994, 45, 1–158. Denmark, S. E. In Trost, B. M., Fleming, I., Eds.; Comprehensive Organic Synthesis; Pergamon: Oxford, UK, 1991; Vol. 5, pp 751–784. Harrington, P. E.; Tius, M. A. J. Am. Chem. Soc. 2001, 123, 8509–8514 and references therein.
- Denmark, S. E.; Habernas, K. L.; Hite, G. A.; Jones, T. K. *Tetrahedron* 1986, 42, 2821–2829.
- 11. Nicolaou, K. C.; Gray, D. L. F. J. Am. Chem. Soc. 2004, 126, 607–612 and references therein.
- 12. Wellington, K.; Cambie, R.; Rutledge, P.; Bergquist, P. J. Nat. *Prod.* **2000**, *63*, 79–85.
- 13. Shinde, H. M. Tetrahedron Lett. 2003, 44, 7049-7053.
- Cambie, R.; Lai, A.; Kernan, M. J. Nat. Prod. **1993**, 58, 940–942. Cambie, R. C.; Richard, C. E. F.; Rutledge, P. S.; Wellington, K. D. Acta Crystallogr. **2001**, C57, 958–960.
- Clive, D. L. J.; Wang, J. *Tetrahedron Lett.* 2003, 44, 7731–7733. Nicolaou, K. C.; Gray, D. L. F. J. Am. Chem. Soc. 2004, 126, 613–627.
- Baker, R.; Bradshaw, J. W. S. In Aliphatic and Related Natural Product Chemistry; Gunstone, F. D., Ed.; Royal Society of Chemistry: London, 1983. Nicolaou, K. C.; Ramphal, J. Y.; Petasis, N. A.; Serhan, C. N. Angew. Chem., Int. Ed. Engl. 1991, 30, 1100–1116. Launay, V.; Beaude, I.; Quintard, J.-P. Bull. Soc. Chim. Fr. 1997, 134, 937–946. Dominguez, B.; Iglesia, B.; de Lera, A. R. J. Org. Chem. 1998, 63, 4135–4139. Lipshutz, B. H.; Ullman, B.; Lindsley, C.; Pecchi, S.; Buzard, D. J.; Dickson, D. J. Org. Chem. 1998, 63, 6092–6093. Peng, Z. H.; Li, Y. L.; Wu, W. L.; Liu, C. X.; Wu, Y. L. J. Chem. Soc., Perkin Trans. 1. 1996, 1057–1066. Chem. Rev. 1994, 94, 1–278. Becher, J., Schaumburg, K., Eds.; Molecular Engineering for Advanced Materials; Kluwer Academic: Dordrecht, 1995; Vol. 456, Specialist Periodical Report, Vol. 3, p 159.
- Heck, R. F., Trost, B. M., Fleming, I., Eds.; Comprehensive Organic Synthesis; Pergamon: Oxford, 1991; Vol. 4, p 833.
- Daves, G. D., Jr.; Hallberg, A. Chem. Rev. 1989, 89, 1433–1446. Andersson, C.-M.; Hallberg, A.; Daves, G. D., Jr. J. Org. Chem. 1987, 52, 3529–3536. Hegedus, L. S.; Toro,

J. L.; Miles, W. H.; Harrington, P. J. J. Org. Chem. 1987, 52, 3319–3322.

- Cabri, W.; Candiani, I.; Bedeschi, A.; Penco, S.; Santi, R. J. Org. Chem. **1992**, 57, 1481–1486. Cabri, W.; Candiani, I.; Bedeschi, A.; Santi, R. J. Org. Chem. **1990**, 55, 3654–3655. Andersson, C.-M.; Halberg, A. J. Org. Chem. **1989**, 54, 1502–1505.
- 20. Cabri, W.; Candiani, I. Acc. Chem. Res. 1995, 28, 2-7.
- Andersson, C.-M.; Larsson, J.; Hallberg, A. J. Org. Chem. 1990, 55, 5757–5761. Larhed, M.; Andersson, C.-M.; Hallberg, A. Acta Chem. Scand. 1993, 47, 212–217.
- 22. Heck, R. F. Palladium Reagents in Organic Synthesis; Academic: London, 1985.
- 23. Beller, M.; Kühlein, K. Synlett 1995, 441-442.
- 24. Heck, R. F. Org. React. 1982, 27, 345-391. Ben-David, Y.;

Portnoy, M.; Gozin, M.; Milstein, D. Organometallics 1992, 11, 1995–1996.

- Kikukawa, K.; Nagira, K.; Wada, F.; Matsuda, T. J. Chem. Soc., Perkin Trans. 1 1986, 1959–1964. Sengupta, S.; Bhattacharyya, S. J. Chem. Soc., Perkin Trans. 1 1993, 1943–1944.
- Brunner, H.; Le Cousturier de Courcy, N.; Genêt, J.-P. Tetrahedron Lett. 1999, 40, 4815–4818.
- Turksov, F.; Hughes, G.; Batsanov, A. S.; Bryce, M. R. J. Mater. Chem. 2003, 13, 1554–1557. Chapman, G. M.; Stanforth, S. P.; Tarbit, B.; Watson, M. D. J. Chem. Soc. Perkin Trans. 1 2002, 581–582. Goel, A.; Srivastava, P.; Nath, M.; Ram, V. J. Synthesis 1998, 167–170.
- 28. Roe, A. Org. React. 1949, 5, 193-227.