



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Expression of the secreted factors noggin and bone morphogenetic proteins in the subependymal layer and olfactory bulb of the adult mouse brain.

This is the author's manuscript

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/102959 since 2016-01-05T10:03:17Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on: Questa è la versione dell'autore dell'opera: [Neuroscience, 128, 2004, DOI: 10.1016/j.neuroscience.2004.06.053] ovvero [P. PERETTO, C. DATI, S. DE MARCHIS, H. H. KIM, M. UKHANOVA, A. FASOLO, F. L. MARGOLIS, 128, Elsevier Ltd, 2004, pagg.685-696] The definitive version is available at:

La versione definitiva è disponibile alla URL: [http://www.sciencedirect.com/science/article/pii/S0306452204005019]

EXPRESSION OF THE SECRETED FACTORS NOGGIN AND BONE MORPHOGENETIC PROTEINS IN THE SUBEPENDYMAL LAYER AND OLFACTORY BULB OF THE ADULT MOUSE BRAIN

P. PERETTO,^a C. DATI,^a S. DE MARCHIS,^a H. H. KIM,^b M. UKHANOVA,^b A. FASOLO^a AND F. L. MARGOLIS^{b*}

^aDepartment of Human and Animal Biology, University of Torino, 10123 Torino, Italy

^bDepartment of Anatomy and Neurobiology, University of Maryland School of Medicine, 20 Penn Street HSFII-203, Baltimore, MD 21201, USA

Abstract - The antagonism between noggin and the bone morphogenetic proteins (BMPs) plays a key role during CNS morphogenesis and differentiation. Recent studies indicate that these secreted factors are also widely expressed in the postnatal and adult mammalian brain in areas characterized by different types of neural plasticity. In particular, significant levels of noggin and BMP expression have been described in the rodent olfactory system. In the mammalian forebrain, the olfactory bulb (OB) and associated subependymal layer (SEL) are documented as sites of adult neurogenesis. Here, using multiple approaches, including the analysis of noggin-LacZ heterozygous mice, we report the expression of noggin and two members of the BMP family, BMP4 and BMP7, in these regions of the adult mammalian forebrain. We observe that along the full extent of the SEL, from the lateral ventricle to the olfactory bulb, noggin and BMP4 and 7 are mainly associated with the astrocytic glial compartment. In the OB, BMP4 and 7 proteins remain primarily associated with the SEL while strong noggin expression was also found in cells located in different OB layers (i.e. granule, external plexiform, glomerular layers). Taken together our data lead us to hypothesize that within the SEL the antagonism between noggin and BMPs, both produced by the glial tubes, act through autocrine/paracrine inductive mechanisms to maintain a neurogenetic environment all the way from the lateral ventricle to the olfactory bulb. In the OB, their expression patterns suggest multiple regulatory roles on the unusual neural plasticity exhibited by this region.

Key words: adult neurogenesis, subventricular zone, olfactory system, glial tubes, neurogenetic niche, cytokines.

The fundamental roles played during vertebrate embryonic development by the bone morphogenetic proteins (BMPs) and their modulators such as noggin have become well established in recent years (for review see Hogan, 1996; Mehler et al., 1997). In early embryogenesis, these secreted proteins are involved in neurulation and in subsequent dorso-ventral patterning within

^{*}Corresponding author. Tel: +1-410-706-8913 or +1-410-706-8914; fax: +1-410-706-2512. E-mail address: fmargoli@umaryland.edu (F. L. Margolis).

Abbreviations: BMP, bone morphogenetic protein; β-gal, β-galactosidase; BSA, bovine serum albumin; GFAP, glial fibrillary acidic protein; LV, lateral ventricle; OB, olfactory bulb; PB, phosphate-buffered saline; PFA, paraformaldehyde; PSA-NCAM, polysialylated neural cell adhesion molecule; PVDF, polyvinylidenedifluoride; RE, rostral extension; SEL, subependymal layer; SVZ, subventricular zone; TBS, Tris-buffered saline; X-gal, 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside.

the neural tube (Hogan, 1996; Dale et al., 1999; Bachiller et al., 2000; Briscoe and Ericson, 2001). Later in development, BMPs, and their regulators, receptors and antagonists, are expressed within the germinal zones of the CNS where they play instructive and inductive roles in lineage specification (Mehler et al., 1997; Li and Lo Turco, 2000; Stull et al., 2001). Recent studies have shown that different BMPs and noggin are still present in the postnatal and adult mammalian CNS (Valenzuela et al., 1995; Lim et al., 2000; Peretto et al., 2002; Fan et al., 2003). Strong noggin mRNA expression has been demonstrated in neurons of the olfactory bulb (OB) such as the tufted and mitral cells (Valenzuela et al., 1995). Notable levels of noggin mRNA have been also described in neurons of the piriform cortex, hippocampus, cerebellar Purkinje cells and ependymal cells (Valenzuela et al., 1995; Lim et al., 2000; Fan et al., 2003). Abundant expression of several BMPs and their receptors has been reported in the OB and hippocampus of adult rat and mouse (Charytoniuk et al., 2000; Fan et al., 2003). Recently, we have demonstrated that BMP4, 6 and 7 are widely distributed in the mouse rostral forebrain (Peretto et al., 2002). During embryonic development they are present in the peripheral olfactory mucosa, OB and subventricular zone (SVZ), and during postnatal and young-adult life, they are observed in the OB mitral cells and subependymal layer (SEL). Whereas the role/s of BMPs and noggin in the adult brain is/are still largely unexplored, our previous data (Peretto et al., 2002), together with recent reports (Shou et al., 1999, 2000; Lim et al., 2000; Coskun et al., 2001; Coskun and Luskin, 2002), support a potential role for these proteins in anatomical and functional plasticity of the adult brain. This view is strengthened by the observation that BMPs and their antagonists and modulators are associated with highly dynamic systems outside the nervous system, and in particular to restricted zones housing stem cells, such as the stromal cell system (Deans and Moseley, 2000), gonads, skin and gut of both vertebrates and invertebrates (reviewed by Spradling et al., 2001). Thus, BMPs, noggin and other BMP-related factors should be included within the group of molecules which participate to create a microenvironmental "niche" that controls the activity of stem cells and their progeny (Spradling et al., 2001).

It is well known that like the hippocampus, the OB of adult rodents retains high structural and functional neural plasticity, principally in terms of neurogenesis of the two main classes of OB interneurons: the granule and periglomerular cells (Luskin 1993; Lois and Alvarez-Buylla, 1994). During embryonic development these cell populations derive from the lateral ganglionic eminence (Wichterle et al., 1999; Corbin et al., 2001) and in adulthood continue to be supplied by the proliferative activity of multipotent stem cells located in the SEL. The SEL is also referred to as the adult SVZ, having originated from the embryonic SVZ (Boulder Committee, 1970). The SEL corresponds to a tissue domain which directly underlies the monolayer of ependymal cells lining the lateral ventricles (SEL-LV), and extends along the olfactory peduncles with its rostral extension (SEL-RE) reaching the center of the OB (SEL-OB). Within the SEL two principal components have been described, highly proliferating/migrating neuroblasts and glial tubes (Lois et al., 1996; Jankovski and Sotelo, 1996; Peretto et al., 1997). The neuroblasts are organized as polysialylated neural cell adhesion molecule (PSA-NCAM)positive tangentially oriented chains of cells (Bonfanti and Theodosis, 1994; Rousselot et al., 1995; Peretto et al., 1999). This organization is lost when the chains reach the OB at which point the individual neuroblasts migrate radially to the granule and glomerular cell layers (Bonfanti and Theodosis 1994; Hack et al., 2002). The astrocytic glial cell bodies and processes form a dense meshwork generating a system of channels, called glial tubes, which envelop the migrating chains during their displacement to the OB. The glial tubes progressively disappear within the OB as the chains disaggregate into single cells (Peretto et al., 1997). Recent studies suggest that the glial tubes, rather than being directly involved in cell migration (Wichterle et al., 1997), act as a physical barrier to separate the area of cell genesis and migration from the mature brain parenchyma, and are potentially involved in providing a stem cell niche and the stem cell reservoir of the SEL (Doetsch et al., 1999; Wichterle et al., 1999; Lim et al., 2000). In fact, SEL astrocytes express particular morphological and molecular features some of which are frequently found in embryonic and postnatal radial glia (Peretto et al., 1997, 1999: Alves et al., 2002) that are implicated in creating molecular barriers (i.e. tenascin-c, Jankovski and Sotelo, 1996), while others are strictly related to neurogenetic processes (expression of EphB receptors and ligands, Conover et al., 2000, expression of BMP proteins and receptors, Lim et al., 2000; Coskun et al., 2001; Coskun and Luskin, 2002; Peretto et al., 2002). Additional strong evidence supporting this role for glia in inducing adult neurogenesis comes from the demonstration that the protein noggin, expressed in ependymal cells of the LV, can create a neurogenetic niche in the SEL by antagonizing the BMPs present in the SEL astrocytes (Lim et al., 2000). In the work reported here, we have used β -galactosidase (β -gal) expression by the noggin-LacZ "knock-in" heterozy gous adult mouse (McMahon et al., 1998; Lim et al., 2000) as a reporter to monitor the distribution and cell localization of noggin within the rostral forebrain, with particular attention to the SEL and the OB. This distribution was compared with that of BMP7 and BMP4

previously demonstrated to be expressed in this region (Peretto et al., 2002). Our results demonstrate strong expression of noggin, BMP4 and BMP7 throughout the prosencephalon. Within the SEL, noggin and BMPs appear largely associated with the astrocytes forming the glial tubes. The presence of noggin protein and mRNAthe SEL region and in different layers of the OB has been confirmed, by means of immunohistochemistry, in situ hybridization and RT-PCR. Our results lead us to hypothesize that the antagonism between these factors acts through autocrine/paracrine inductive mechanisms that maintains a neurogenetic environment in the SEL all the way from the LVthe OB.

EXPERIMENTAL PROCEDURES

Tissue preparation

All the procedures involving animals were approved by the University of Maryland School of Medicine's Institutional Animal Care and Use Committee and were carried out on young/adult (1–2 months old) mice. These experiments conform to international guidelines on the ethical use of animals and were designed to minimize the numbers, and discomfort, of the mice used. For immunohistochemistry and 5-bromo-4-chloro-3-indolyl-β-Dgalactopyranoside (X-gal) staining at least 10 noggin-LacZ "knock in" heterozygous mice, and 10 wild type 129S1/SvImJ mice were analyzed. Founder noggin-lac-Z hemizygous mice were generously provided by A. McMahon (Harvard University, Cambridge, MA, USA) (McMahon et al., 1998). The mice were maintained by mating hemizygous mice to wild-type mice of

the background strain, 129S1/SvImJ (Jackson Laboratories, Bar Harbor, ME, USA). Noggin lac-Z mice were identified by the blue color generated on incubating ear punches in X-gal reagent (Cummings et al., 2000). In situ hybridization (four animals) and RT-PCR (15 animals) analyses were performed on CD-1 strain mice (Charles River, Calco, Italy). Mice were deeply anesthetized with a ketamine/xylazine solution (100 mg/kg body weight: 33 mg/kg body weight) and transcardially perfused with ice cold saline solution (0.9% NaCl) followed by a freshly prepared solution of 4% paraformaldehyde (PFA) in 0.1 M phosphate buffer (PB), pH 7.4. For X-gal histochemistry, the perfused brains were separated sagittally into two halves. One was cut coronally into ca. 2 mm slices and the other half was cut into a few sagittal slabs. For immunohistochemistry and in situ hybridization tissues were postfixed (2-4 h) in the same fixative, cryoprotected in ascending sucrose solutions (7.5%, 15%, 30%), quick frozen in cold isopentane (-80 °C), cut in serial coronal or sagittal sections (10 -25μ m), and stored in 0.01 M PB. For RT-PCR, mice were anesthetized as described above, brains promptly removed and thick coronal sections (250 µm) cut on a vibratome. Under a high magnification dissecting microscope, different levels along the extent of the SEL (SEL-LV, SEL-RE, SEL-OB), the septum and OB granule layer were carefully micro-dissected.

Immunohistochemistry

Sections were processed for immunohistochemistry either by using the biotinavidin system or immunofluorescence. All incubations were run at room temperature and washes were done in 0.01 M phosphate buffered saline (PBS) for 15 min. Tissue sections were treated overnight with the following primary antibodies diluted in PBS with 0.1%- 0.5% Triton X-100 and 1% non-immune serum of the same donor species as the secondary antiserum: rabbit-anti-β-gal (1:4000; 5Prime-3Prime, Inc., Boulder, CO, USA); mouse-anti-glial fibrillary acidic protein (GFAP; 1:200; Boehringer, Mannheim, Germany); goat-antinoggin (1 µg/ml; R&D Inc., Minneapolis, MN, USA; lot FCD09); rabbit-anti-GFAP (1:800; Sigma, St. Louis, MO, USA); goat-anti-BMP7 and goat-anti-BMP4 (1:100; Santa Cruz, CA, USA); mouse-anti-PSA-NCAM (1:1000; G. Rougon, Marseille, France). The next day sections were rinsed in PBS and, for the biotin-avidin system, incubated with the appropriate biotinylated secondary antibodies for 1 h (anti-goat 1:250 for BMP4 or 1:1000 for noggin, or antirabbit 1:250; Vector Laboratories, Peterborough, UK), rinsed, and incubated in avidin-biotin complex (1:400; Vector Laboratories, Inc.). The reaction product was visualized with 0.15 mg/ml 3,3' diaminobenzidine in PBS with 0.01% H2O2. Sections were then serially mounted onto Superfrost Plus slides (Fisher, Pittsburgh, PA, USA), air dried, dehydrated in graded alcohols, cleared in xylenes, and coverslipped using D.P.X. mounting medium (Aldrich, Milwaukee, WI, USA). For dual immunofluorescence, after incubation with the primary anti-sera, sections were washed and incubated with appropriate solutions of secondary anti-rabbit Cy3, anti-mouse Cy2, anti-mouse Cy3, antigoat Cy2 (all diluted 1:800; Jackson ImmunoResearch, West Grove, PA, USA). Sections were then coverslipped with an anti-fade mounting medium (Dako Corporation, Carpinteria, CA, USA).

Control reactions included: (i) omission of the primary antibodies, (ii) incubation of sections in BMP antisera pre-absorbed with 10 μ M of the respective immunogenic peptides, (iii) incubation with rabbit-anti- β -gal of equivalent sections from wild type 129S1/SvImJ mice, (iv) noggin antibody

specificity has been evaluated by absorbing the antibody with 1 μ g/ml of recombinant human noggin protein (Pepro Tech. Inc. Rody Hill NJ), for either immunocytochemistry or Western blot analysis.

Western blot

Protein samples were prepared from OB by homogenization in 10 volumes Tris-lysis buffer (50 mM Tris-HCl, 2 mM DTT, 1 M NaCl) and protease inhibitors (10 \[\]g/ml aprotinin, 5 \[]g/ml leupeptin, 1 mM phenylmethylsulfonyl fluoride) following by sonication. After centrifugation for 1 h at 100,000xg the supernatants were aspirated and protein concentrations determined by the Bradford method (BioRad, Hercules, CA). Extracts (40 \[\]g) were separated in 10 -20% SDS-polyacrylamide gel electrophoresis (Ready Gels; BioRad) and electro-transferred to polyvinylidenedifluoride (PVDF) membranes at 100 V for 1 h in ice. The membranes were blocked for 1 h in Tris-buffered saline (TBS) with 0.1% Tween-20 (TBS-T) containing 1% bovine serum albumin (BSA) and then were incubated overnight at 4 °C with primary noggin antibodies (0.1 µg/ml in TBS-T with 1% BSA). The membranes were then washed three times for 10 min in TBS-T and incubated for 1 h at room temperature with donkey anti-goat HRP-labeled secondary antibodies (1:40,000 dilution in TBS-T with 1% (BSA); Jackson Immunoresearch). The final wash was in TBS-T three times for 10 min and twice in TBS for 15 min. The immunoreactive protein bands were visualized using Super Signal West Pico enhanced chemiluminescence (Pierce, Rockford, IL, USA).

X-gal histochemistry

Sagittal and coronal slabs were incubated, for about 2.5 h at room temperature, in X-gal substrate solution (1 mg/ml) containing 5 mM potassium ferrocyanide, 5 mM potassium ferricyanide, 2 mM MgCl2, 0.01% sodium deoxycholate and 0.02% Nonidet P-40 (Cummings et al., 2000). When the color development was complete, the tissue was rinsed with cold PBS and the reaction was stopped by transferring the tissue to 4% buffered PFA. Sections were then photographed on a Wild dissecting microscope equipped with a 35 mm camera back. Equivalent material from wild type mice did not generate any histochemical reaction under these conditions.

RT-PCR

The expression of noggin, BMP4 and BMP7 was evaluated by a semiquantitative RT-PCR. Total RNA was extracted by the acidic phenol method (Chomczynski and Sacchi, 1987). Aliquots (1 [g) of total RNA were transcribed by the MMLV Reverse Transcriptase (APBiotech, Piscataway, NJ, USA). Genomic DNA contamination was tested by incubating the RNA in the absence of reverse transcriptase. Aliquots of the cDNA were amplified by TaqDNA polymerase (RedTaq; Sigma, St. Louis, MO, USA) polymerase chain reaction, 30 cycles. The amount of cDNA to be amplified for each sample was standardized by a previous amplification reaction for the housekeeping gene glyceraldehyde-3-phosphate dehydrogenase. Primers were selected from the mouse sequences available in GenBank (http://www.ncbi.nlm.nih.gov/entrez/) by means of Primer3 (http://www.genome.wi.mit.edu/cgi-bin/primer/primer3) and were as follows: for noggin (accession number U79163), forward CAGATGTGGCTGTGGTCAC, reverse GCAGGAACACTTACACTCGG, size 261 bp, from base 682-942; for BMP4 (X56848) forward CTCCCAAGAATCATGGACTG, reverse AAAGCAGAGCTCTCACTGGT, 468 bp, from 305 to 772; for BMP7 (NM007557), forward CCTGGGCTTACAGCTCTCTG, reverse GGTGGCGTTCATGTAGGAGT, 415 bp from 726 to 1122.

Probes

The 261 bp cDNA fragment of the noggin sequence was subcloned in pGEM-T (Promega, Italia, Milano, Italy), the antisense probe was generated using T7 RNA-polymerase following restriction digestion of the clone with PstI endonuclease and the sense (control) probe by SP6 RNA-polymerase following digestion of the clone with ApaI endonuclease. The riboprobes were labeled by transcription in the presence of DIG-UTP using the DIG RNA Labeling Kit (Roche Diagnostics, Monza, Italy).

Mouse 28S ribosomal RNA, a 1.3 kb fragment (Mariani-Costantini et al., 1998) was used as a control of RNA integrity.

In situ hybridization

Hybridization was performed essentially as previously described (Dati et al., 1996). Briefly, slides were rehydrated for 5 min in PBS, treated for 5 min at 37 °C with 5 μ g/ml Proteinase K (Sigma), acetylated and then incubated overnight at 42 °C with 1 ng/ml of DIG-labeled riboprobe in a mixture containing 60% deionized formamide, 2x SSC, 1 mg/ml yeast tRNA, 100 μ g/ml herring sperm DNA, 5x Denhardt's solution, 1 U/ml RNase inhibitor (RNAguard, APBiotech). The next day slides underwent an RNase A (Sigma) treatment, 20 μ g/ml in 10 mM Tris pH 8.0, 0.5 M NaCl, 30 min at 37 °C, and a stringency wash, 0.1x SSC, 10 min at 65 °C followed by 10 min at room temperature. The slides were then incubated overnight with an anti-DIG antibody conjugated with alkaline phosphatase (Roche) diluted 1:5000. The alkaline phosphatase activity was revealed by the DIG Nucleic Acid Detection Kit (Roche).

Preparation of figures

Light photomicrographs were digitized using a Phase One PowerPhase digital camera (Phase One, Denmark) fitted to a Leica DMRX microscope (Leica Microsystems Inc., Deerfield, IL, USA). Analysis and digital photography of the double immunofluorescence was carried out using a laser-scanning Olympus Fluoview confocal system (Olympus Italia, Milano, Italy). Digital images were brightness, color and contrast balanced, assembled into montages using Corel-Draw 11 (Corel Corporation, USA). Plates of photomicrographs were printed using a Fuji Fujix Pictrography 3000 printer (Fuji Photofilm USA, Carlstadt, NJ, USA).

RESULTS

To visualize the cell type/s expressing noggin protein in the adult mouse prosencephalon we used a previously generated transgenic model obtained by a gene replacement strategy in which most of the noggin coding region wasm replaced by the β -gal coding region of Escherichia coli (McMahon et al., 1998). This placed lac-Z under the control of the noggin promoter so that the noggin-lac-Z hemizygous mice express β -gal in the same embryonic and postnatal structures where noggin expression has been identified by in situ hybridization (Valenzuela et al., 1995; McMahon et al., 1998). We confirmed this by Western blotting and immunocytochemistry using anti-noggin antibody and by RT-PCR.

$\beta\mbox{-}Gal$ activity in the forebrain of the adult brain as a reporter of noggin expression

Abundant Lac-Z expression, as a reporter of noggin expression, was visualized throughout the whole brain in thick coronal and parasagittal brain sections (ca. 2 mm) of young/adult mice (1–2 months old) visualized by X-gal histochemistry. The staining was particularly strong in restricted areas such as the hippocampal dentate gyrus, piriform cortex, septal area, and OB (Fig. 1).

β-Gal immunostaining was used for higher resolution analysis within the anterior prosencephalon, with particular attention to the SEL and OB. Parasagittal and coronal sections (12-25 µm) cut from the LV to the OB were examined with immunofluorescence and avidin-biotin staining techniques. Strong expression of β -Gal was found throughout the full extent of the SEL and in the OB (Fig. 2A-F). At the level of the LV β -gal staining was identified in the SEL and in ependymal cells lining the ventricular cavities (Figs. 2B, 3A, C). More anteriorly, along the olfactory peduncle the staining was found primarily associated with the SEL-RE (Fig. 2A, C). Within this area, thick immunoreactive septa delineate the SEL perimeter and in part enter within the SEL itself giving rise to a web-like structure resembling the glial tubes (Figs. 2C; 3D, F, G, I, J, L). The staining appeared dense and diffuse, often preventing identification of clearly distinguishable cell bodies and processes (Figs. 2, 3). In the closely surrounding tissues, some positive cells with thin astrocytic-like processes were also observed (not shown). Some neuronal-like positive cells were also observed in the anterior olfactory nucleus (not shown). Double immunofluorescence performed with PSA-NCAM as a marker of the migrating neuroblasts, and GFAP as a marker of the glial tubes, confirmed that, from the LV to the SEL-OB, the β -gal protein is mostly localized in the SEL glial compartment (Fig. 3D–I). β -Gal expression was never observed in the PSA-NCAM-positive migrating neuroblasts, neither during their tangential chain migration (Fig. 3A-F) nor when they spread out radially as individual elements through the OB (not shown).

In the OB a dense and diffuse β -gal immunostaining is present throughout diverse layers (Fig. 2A, D-F).

Within the SEL-OB the staining is less intense, compared with that observed in more posterior SEL regions, and is still associated with the glial tubes that, at this level, are progressively disappearing. A strong immunoreaction was found in granule cells of both the main and accessory OB, in putative tufted neurons localized at different depth of the external plexiform layer and in some mitral cells. Within the glomerular layer, a diffuse reaction was observed in the neurites constituting the glomerulus and in some juxtaglomerular neurons. No staining was found in the olfactory nerve layer. In equivalent sections from control wild-type mice we never observed any β -gal staining.

Fig. 1. X-gal histochemical analysis of noggin-lacZ mouse brain slabs. (A, B) Coronal sections, (C) parasagittal section. X-gal staining is visible in different areas of the adult forebrain, some of which are characterized by neural plasticity, such as the dentate gyrus of the hippocampus (A) and the granule and the SEL of the OB (C). Abundant staining is present in the SEL-LV (arrow), septum and piriform cortex (arrowhead) (B). No staining was observed in tissue from wild-type mice. CC, corpus callosum; CP, caudate putamen; DG, dentate gyrus; Grl, granule cell layer; S, septum; TH, thalamus. Scale bars=2 mm A, B; C, 1 mm.





Fig. 2. β-gal and BMP7 immunohistochemical localization in the adult mouse rostral forebrain. (A) Parasagittal section showing β-gal protein expression in the adult SEL-RE (arrows) and OB. (B, C) Coronal sections cut respectively at the level of the LV and olfactory peduncle. Strong staining is associated with the SEL-LV area (arrows in B) and SEL-RE (C). β-Gal positive staining is also present in the ependymal monolayer and septum (B). Within the OB (D–F) β-gal is expressed in the glomerular, external plexiform, and granule cell layers. (E) Higher magnification of (D) showing positive cells in the external plexiform layer (arrows). (F) Diffuse staining is present in numerous granule cells, some of which show stronger staining (e.g. arrows). (G–I) BMP7 is associated with the whole SEL extension from the LV (arrows in G) to the OB, where itmainly restricted to the SEL area (I). aco, anterior commissure; CC, corpus callosum; CP, caudate putamen; Cx, cerebral cortex; EPl, external plexiform layer; Gl, glomerular layer; Grl, granule cell layer; S, septum. Scale bars=200 μm, A; B–I, 50 μm.

The use of a knock-in reporter-gene model to study protein expression is a powerful tool; however, ectopic expression of the reporter gene may occur confounding the interpretation. In order to address this question and to further confirm the presence of noggin protein in the SEL we performed immunocytochemical and Western blot analyses of noggin protein.



Fig. 3. Double immunostaining between β -gal, BMP7, and markers of the SEL compartment in adult mice. β -Gal (A, D, G, J), PSA-NCAM (B, E), GFAP (H), BMP7 (K) immunoreactivity in coronal sections. (A-C) LV; D-L olfactory peduncle. Double immunostaining for β -gal and anti-PSA-NCAM showed no overlap of expression (A-F). At the level of the LV an intense β -gal immunoreactivity (A, C) is present in the ependymal monolayer (e) and in glial-like cells of the SEL (arrow in C). (G-I) A high degree of colocalization is observed in double staining between β -gal and GFAP (arrows in I). (J-L) Only a partial overlap was found in dual labeling for β -gal and BMP7 (arrow in L). (C, F, I, L) Higher magnifications of a simultaneous view of the pictures in the same row obtained from 1 µm optical section. Scale bars=50 µm, A, B, D, E, G, H, J, K; C, F, I, L, 20 µm.

Noggin protein detection in the SEL confirms the specificity of the Noggin Lac-Z model

The immunohistochemical detection of noggin was performed on sections cut along the rostral prosencephalon of adult mice, where we observed a strong immunoreactivity in the same areas positive for β -gal staining in the Lac-Z model (not shown). Noggin appears to be abundantly expressed in the SEL (Fig. 4). In this region the staining appears localized to cell bodies and processes; moreover, immunoreactivity is also present in extracellular spaces (Fig. 4A, B, E, H, K), Confocal microscopic analyses on sections treated for double staining with anti-noggin and anti- β -gal (see Fig. 4A–D) showed that in the SEL area all the β -gal positive cells express noggin, thus excluding ectopic β -gal expression in this region. However, the extent of the staining obtained with anti-noggin was larger than that observed with anti- β -gal. Noggin and GFAP immunoreactivities were colocalized (Fig. 4E–G), confirming the astroglial cells as the source of noggin. To a lesser extent, we also observed



Fig. 4. Relationships between noggin, β -gal, GFAP and PSA-NCAM in the SEL-RE, and noggin protein localization in the SEL-LV. (A–D) Noggin and β -gal colocalize in cells of the SEL-RE (see arrow in B–D). (E–G) Glial GFAP-positive processes are strongly immunopositive for noggin (arrows). (H–J) A partial overlap occurs between noggin and PSA-NCAM immunoreactivities. D, G, and J have been obtained examining 1 µm optical section. (K) Noggin protein is abundant in the ependyma (e) and in the SEL-LV region. High levels of immunofluorescence are identifiable in the ependymal cells and in astroglial-like cell processes in the SEL-LV and corpus callosum (CC). Scale bars=12 µm, A; B–D, 6 µm; E–J, 8 µm; K, 20 µm.

a partial overlap between noggin and PSA-NCAM (Fig. 4H–J). This partial colocalization with PSA-NCAM immunopositive migrating cells was never observed in the case of β -gal expression, and may be the consequence of secretion and diffusion of noggin protein. PSA-NCAM antibody recognizes the polysialyl- enriched extracellular portion of NCAM, thus a partial colocalization between PSA-NCAM and Noggin may be due to overlapping of these two extracellular antigens.

We have demonstrated the specificity of the antinoggin antibody in both Western blots and in immunohistochemistry by blocking with 1 μ g/ml recombinant human noggin protein (Fig. 5).

Noggin co-localizes with BMP4 and BMP7

We have previously analyzed the expression of different BMPs within the mouse olfactory system during embryonic and postnatal life (Peretto et al., 2002). Our results demonstrated a discrete spatial and temporal expression of these cytokines within the olfactory system and in particular, we observed expression of BMP4 and 7 in the embryonic and postnatal SVZ. Here we have

characterized the distribution of these two BMPs, within the extent of the SEL and in the OB of young/adult mice, in comparison to that of their antagonist/modulator noggin.

Single immunostaining obtained with anti-BMP4 and anti-BMP7 showed very similar patterns of distribution within the SEL and OB. In the SEL, from the LV to the OB, both BMP4 (not shown) and 7 (Figs. 2G–I, 3K) give rise to a dense and diffuse meshwork-like immunostaining resembling the appearance of the glial tubes. Double immunoreactions performed with antibodies to BMPs GFAP and PSA-NCAM (not shown), demonstrated no colocalization with the chains of migrating neuroblasts, and only a partial overlap with the glial GFAP-positive SEL compartment, confirming our previous results (Peretto et al., 2002). Comparison of the staining of BMP7 with that of β -gal in sections visualized with double immunofluorescence showed only partial colocalization of these molecules (Fig. 3J–L). Although anti-BMP4 and 7 each gave a strong signal in the SEL region of the OB (Fig. 2I), they gave only a very faint signal in the olfactory nerve and glomerular layers (not shown).

Noggin mRNA expression in the SEL and OB

To further confirm the presence of noggin, BMP4, and BMP7 in the SEL and OB regions, RT-PCR analyses were performed on tissue microdissected from thick coronal sections cut at different levels of the SEL (SEL-LV, -RE,-OB) and from the OB of wild-type adult mice (see Experimental Procedures for details).

As a positive control, microdissected septum tissue, that contains a significant level of noggin mRNA (Valenzuela et al., 1995), was included in our experiments. We observed the presence



Fig. 5. Specificity of anti-noggin antibody. (A) Western blot analysis; (B and C) immunohistochemistry.(A) Lanes 1, 3, 5: 5 ng recombinant human noggin. Lane 2: 40 µg of SDS protein extract from OB spiked with 5ng recombinant human noggin. Lanes 4, 6: 40 µg of SDS protein extract from OB. After electrophoresis and transferring onto PVDF membranes lanes 1- 4 were incubated with R & D goat anti-noggin antibody (0.1 µg/ml), lanes 5 and 6 were incubated with R & D goat anti-noggin antibody preincubated with 1 µg/ml of noggin protein to block the antibody. All samples were run simultaneously. Note the presence of a small amount of noggin dimer and the elimination of the immunoreactivity by blocking the antiserum. (B, C) Consecutive parasagittal sections incubated respectively with B) R & D goat anti-noggin antibody (1 µg/ml) and C) R & D goat anti-noggin antibody preincubated with 1 µg/ml of noggin protein to block the antibody. In B a strong immunoreactivity was observed both in the SEL-RE and in the anterior olfactory nucleus. No staining is observed in the same areas in the section treated with the blocked antibody. aco, anterior commissure; AON, anterior olfactory nucleus.

of noggin and BMP mRNAs in all specimens examined (Fig. 6). In situ hybridization for noggin mRNA was performed on adult wild-type animals in coronal sections through the rostral telencephalic region. Noggin mRNA was evident in several prosencephalic areas including cells of the septum and piriform cortex (not shown).



Fig. 6. Expression of noggin, BMP4 and BMP7 mRNAs in the SEL and OB of wild-type adult mice. RT-PCR analysis of microdissected tissue fragments obtained from different levels of the SEL and from the OB showed expression of noggin, and BMP4/7 mRNAs in the SEL-LV,-RE, -OB. The septum was chosen as a positive control (Valenzuela et al., 1995).

Within the SEL a reproducible in situ signal was present from the LV to the

OB. At the level of the LV, strong signal was found also associated with the ependymal cells (not shown). These data confirmed our observation of lacZ expression in these sites as a reporter of noggin protein expression. The pattern of staining obtained in the SEL-RE (Fig. 7A) is reminiscent of the glial tubes. An intense labeling was also observed in cells of the anterior olfactory nucleus (Fig. 7A). Within the OB itself, the noggin anti-sense probe showed a pattern of expression that was similar to that seen for β -gal protein. Thus, a weak in situ signal was found associated with granule cells, while strong staining was found in putative tufted and juxtaglomerular cells and in mitral cells (Fig. 7C). Incubation with the noggin sense probe gave only background levels of hybridization, confirming the specificity of the analysis (Fig. 7B, D).

DISCUSSION

In the last decade the discovery of neural stem cells in restricted regions of the adult mammalian brain has opened exciting new opportunities for research in neuroscience (see for review Bonfanti et al., 2001). In addition to the characterization of the location and properties of these stem cells, major efforts are under way to identify the mechanisms that regulate their activity. As a result a new concept, the "niche," has been proposed (Lim et al., 2000; Spradling et al., 2001). The niche is considered to be a subset of cells and extracellular substrates that create the microenvironment capable of indefinitely sequestering one or more stem cells and that functions to control their selfrenewal and expansion in vivo (Spradling et al., 2001). Knowledge of the stem cells' microenvironment is, therefore, fundamental for understanding how a stem cell can proliferate, choose its fate and eventually integrate into mature tissue.

Noggin and BMPs are components of the stem cell niche

It is striking that strong similarities have been described between the niches present in different tissues in both ertebrates and invertebrates (reviewed in Spradling et al., 2001). From the molecular point of view, several candidate



Fig. 7. In situ hybridization for noggin mRNA on coronal sections cut at the level of the olfactory peduncle and OB. (A) Noggin mRNA expression is present in the SEL-RE (arrow) and in cells of the anterior olfactory nucleus. (C) Within the OB, mitral, external plexiform and glomerular layers show numerous labeled cells. A weak reaction was also found in the granule cell layer. In adjacent sections treated with the sense probe (B, D) no hybridization is present. Arrow in B indicates the SEL-RE area. Arrow and arrowheads in C indicate stained cells respectively in the glomerular and external plexiform layers. AON, anterior olfactory nucleus; Epl, external plexiform layer; Gl, glomerular layer; Grl, granule cell layer; Mi, mitral cell layer.

niche components have been proposed. Members of the transforming growth factor- β family such as the BMPs and their antagonist/modulator noggin have been described as critical components of this stem cell niche microenvironment (Xie and Spradling, 1998, 2000;Coskun et al., 2001; Lawson et al., 1999; Kulessa et al., 2000; Lim et al., 2000; Panchision and McKay, 2002). In addition to their morphogenetic roles during development (Hogan, 1996; McMahon et al., 1998; Dale et al., 1999; Bachiller et al., 2000), the balance between BMPs, noggin and other BMP antagonists/modulators such as chordin, seems to regulate the proliferation and specification of neural cell precursors. Indeed, their expression has been widely reported in the embryonic germinal zones where undifferentiated precursor cells exhibit temporal and site-specific responsiveness to these proteins (Gross et al., 1996; Li et al., 1998; Mabie et al., 1999; Mehler et al., 2000). Furthermore, the antagonism between noggin, expressed in the developing neocortex, and BMPs may regulate the differentiation of neocortical neurons in vivo (Bachiller et al., 2000; Li and LoTurco, 2000).

Noggin and BMPs are expressed in the adult SEL

The presence of these proteins has been also described in many regions of the adult rodent brain that are characterized by different types of neural plasticity, suggesting similar and/or additional roles in the adult CNS (Valenzuela et al., 1995; Coskun et al., 2001; Charvtoniuk et al., 2000; Lim et al., 2000; Peretto et al., 2002). Consistent with these reports we have observed the presence of noggin as well as BMP4 and 7 in the rostral forebrain of young/adult mice. The analysis of noggin localization in the SEL-OB system was mainly performed by means of a well-characterized model, the noggin-lacZ knock-in mouse (McMahon et al., 1998; Lim et al., 2000). The results obtained in the SEL were compared with the immunohistochemical detection of noggin, demonstrating the absence of ectopic β -gal expression in the transgenic model. However, differences in the patterns of staining have been found, namely β -gal expression appears to be more restricted compared with noggin. This result suggests that β -gal expressed under the control of the noggin promoter does not mimic the characteristic secretory behavior of the native noggin protein. Moreover, a significant level of β -gal expression reporting the presence of noggin, was seen in multiple areas such as the hippocampus, piriform cortex, septum, and OB, where different types of

neural plasticity have been reported (Peretto et al., 1999; Bernier et al., 2002; Garcia, 2002). A strong expression of β -gal protein was found to be specifically associated with the SEL throughout its extent and in the OB, regions marked by adult neurogenesis due to the presence of multipotent stem cells (for review see Peretto et al., 1999).

Noggin and BMPs are associated with the SEL glial compartment

Within the SEL, the staining is clearly associated with the glial compartment. Our previous results demonstrated that BMP4 and 7 immunostaining also partially overlap with markers of the glial tubes but were never associated with the migrating neuroblasts (Peretto et al., 2002). In this study, a colocalization was observed between BMP4, BMP7 and β -gal. Thus, our results demonstrate that the astrocytes forming the glial tubes can produce BMP4, 7 and their natural antagonist noggin. The diffuse staining observed for noggin immunolabeling in the SEL region supports a mechanism of production and a subsequent secretion of noggin from the astroglial component. Lim and colleagues (2000) demonstrated the presence of both BMPs, their receptor and noggin within the periventricular area, and proposed a model in which the antagonism between these factors can regulate neurogenesis in the adult SVZ, consistent with the existence of similar regulatory mechanisms during development and adulthood. In their model, noggin, which they reported to be specifically associated with ependymal cells, creates a neurogenetic environment in the adjacent SEL by blocking endogenous BMP signaling. We have observed expression of noggin in the ependymal cells too, but in addition we found a similar intensity of staining in the SEL all the way from the LV to the OB. The reason for this difference is unclear but may relate to the sensitivity of the staining techniques (X-gal staining versus anti-β-gal immunostaining). Our RT-PCR analysis performed on wild-type mice, confirmed the presence of noggin mRNA at different levels of the SEL extent. Although, we cannot exclude that our microdissected fragments also contained small portions of the tissue surrounding the SEL, the intensity of the signal obtained was comparable to that of the septum, a region where high levels of noggin mRNA has been previously described (Valenzuela et al., 1995). This was confirmed by our in situ hybridization analysis demonstrating the presence of a high level of noggin mRNA in the rostral forebrain and in particular in the SEL-LV and OB.

We can now hypothesize, as previously suggested by Lim and colleagues (2000) at the level of the LV, that the antagonism between noggin and BMPs regulates adult neurogenesis all the way from the LV to the OB. This implies that the SEL represents a quite homogeneous system along its entire extension. In support of this idea, and in addition to the anatomical evidence (see Peretto et al., 1999 for review), the existence of adult neural stem cells has recently been shown along the entire SEL area up to the core of the OB (Gritti et al., 2002), thus demonstrating that the SEL-RE is not simply the migratory pathway for the progeny of periventricular stem cells, as previously thought, but is also a source of stem cells. The expression of noggin and BMPs along the SEL from the LV to the OB may also indicate that BMP signaling can exert multiple regulatory roles on adult neurogenesis. Studies addressing neurogenetic mechanisms within the postnatal SVZ suggest that SEL neuroblasts express the cell-cycle inhibitor P19 INK4d (see Coskun and Luskin, 2002). This gene product is dynamically regulated along the migratory

pathway and enables these cells to show the unusual characteristic of retaining the capacity to divide during migration. The dynamic expression of P19 INK4d, which appears higher at the level of the SEL-OB, seems to be regulated by the BMPs, since inhibition of BMP signaling, by injecting replication-deficient retrovirus encoding a dominant negative bone morphogenetic protein receptor-Ia, results in down-regulation of P19 INK4d and initiation of a new cell cycle. This result suggests that in vivo BMP signaling must be dynamically regulated along the SEL migratory route, consistent with our demonstration of the presence of the negative BMPmodulator, noggin, from the LV to the OB.

Furthermore, it has been shown that astrocytes present in adult neurogenetic regions such as the SEL and the hippocampal formation exhibit features that allow these glial cells to regulate processes of neurogenesis. (Song et al., 2002), demonstrated that in vitro, astrocytes can direct the proliferation, as well as the fate, of hippocampal stem cells, inducing their progeny to became neurons rather than glia. These data are consistent with previous reports showing that contact with fragments, or monolayers, of astrocytes can modulate proliferation, fate and differentiation of stem cells from the cortex of developing rat and adult SVZ (Temple and Davis, 1994; Lim and Alvarez-Buylla, 1999). It seems that only younger astrocytes exhibit these characteristics (Smith et al., 1990). In this context, it is worth noting that SEL astrocytes retain some features of immature radial glia-like cells (Jankovski and Sotelo, 1996; Lois et al., 1996; Peretto et al., 1997, 1999), and subpopulations of these glial cells could act as guiescent stem cells (Doetsch et al., 1999). Thus, our results demonstrating that the SEL glial compartment can express both BMPs and noggin, which are key niche components, reinforce the emerging view that astrocytes have an active regulatory role in zones of neurogenesis in adults.

Noggin and BMPs are expressed in the OB

In the OB, noggin-LacZ expression was found in all OB layers. In particular, a strong signal was observed in mature granule cells, external plexiform layer, juxtaglomerular cells and olfactory glomeruli dendritic terminals. Nevertheless, a slight difference was observed in the expression of noggin mRNA and β-gal protein. Most mitral cellular bodies were stained in the ISH analysis but only some of them were labeled in the immunocytochemical analyses of the noggin-lacZ heterozygous mice. This may reflect differential mRNA vs. Protein turnover and/or subcellular compartmentalization. Thus, our data suggest that β -gal protein is restricted to the dendrites of the mitral cells based on the strong noggin- LacZ expression found in the OB glomeruli. A differential expression of mRNA vs. protein in the mitral cell layer has also been reported in other studies (Baker et al., 2001). Alternatively, this discrepancy could be due to a lack of expression of β -gal in some mitral cells. High levels of noggin and BMPs in the OB are justified considering the peculiar characteristic of this region. Adult OB represents a site of persistent plasticity, both at the level of the glomeruli, where a continuous renewal of the olfactory receptor neurons axons can lead to reorganization within the glomerular neuropil (Graziadei and Monti Graziadei, 1978), and in the granule cell layer, the principal site of adult OB neurogenesis (Luskin 1993; Bonfanti and Theodosis, 1994; Lois and Alvarez-Buylla, 1994). Unlike noggin, the expression of BMP4 and 7 in the OB was mainly restricted to the SEL region.

This result indicates that, in the more external layers of the OB, the balance observed between these proteins along the SEL changes and noggin protein is predominant and/or antagonizes others members of the BMPs (i.e. high amounts of BMP6 is present in the OB glomeruli, Peretto et al., 2002). In conclusion, the expression and localization of BMP4, 7 and noggin reported here, lead us to hypothesize that: i) within the SEL, the glial compartment acts through an autocrine or paracrine inductive mechanism between noggin and BMPs to maintain a neurogenetic environment, reinforcing suggestions of a role of the astrocytic glia in creating the neurogenetic niche all along the SEL extension; ii) within the OB, the persistence of high levels of noggin in different types of neurons suggests an involvement of this protein in the multiple processes of neuronal plasticity peculiar to this structure.

Acknowledgments—This work was supported by NIH grants DC- 00347 and DC-03112 (F.L.M.), Italian COFIN and Compagnia di San Paolo. We thank A. McMahon for generously supplying founder mice of the noggin-lacZ strain, G. Rougon for anti-PSA- NCAM, A. Puche for his advice and cooperation and C. Gendusa and F. Scipio for technical assistance.

REFERENCES

- Alves JA, Barone P, Engelender S, Froes MM, Menezes JR (2002) Initial stages of radial glia astrocytic transformation in the early postnatal anterior subventricular zone. J Neurobiol 52:251–265.
- Bachiller D, Klingensmith J, Kemp C, Belo JA, Anderson RM, May SR, McMahon JA, Harland RM, Rossant J, De Robertis EM (2000) The organizer factors chordin and noggin are required for mouse forebrain development. Nature 403:658 – 661.
- Baker H, Liu N, Chun HS, Saino S, Berlin R, Volpe B, Son JH (2001) Phenotypic differentiation during migration of dopaminergic progenitor cells to the olfactory bulb. J Neurosci 21:8505–8513.
- Bernier PJ, Bedard A, Vinet J, Levesque M, Parent A (2002) Newly generated neurons in the amygdala and adjoining cortex of adult primates. Proc Natl Acad Sci USA 99:11464 –11469.
- Bonfanti L, Theodosis DT (1994) Expression of polysialylated neural cell adhesion molecule by proliferating cells in the subependymal layer of the adult rat, in its rostral extension and in the olfactory bulb. Neuroscience 62:291–305.
- Bonfanti L, Gritti A, Galli R, Vescovi A (2001) Multipotent stem cells in the adult central nervous system. In: Stem cells and CNS development (Rao MS, eds), pp 49–70. Totowa: The Humana Press.
- Boulder Committee (1970) Embryonic vertebrate central nervous system: revised terminology. Anat Rec 166:257–262.
- Briscoe J, Ericson J (2001) Specification of neuronal fates in the ventral neural tube. Curr Opin Neurobiol 11:43–49.
- Charytoniuk DĀ, Traiffort E, Pinard E, Issertial O, Seylaz J, Ruat M (2000) Distribution of bone morphogenetic protein and bone morphogenetic protein receptor transcripts in the rodent nervous system and up-regulation of bone morphogenetic protein receptor type II in hippocampal dentate gyrus in a rat model of global cerebral ischemia. Neuroscience 100:33–43.
- Chomczynski P, Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. Anal Biochem 162:156 –159.

- Conover JC, Doetsch F, Garcia-Verdugo JM, Gale NW, Yancopoulos GD, Alvarez-Buylla A (2000) Disruption of Eph/ephrin signalling affects migration and proliferation in the adult subventricular zone. Nat Neurosci 3:1091–1097.
- Corbin JG, Nery S, Fishell G (2001) Telencephalic cells take a tangent: non-radial migration in the mammalian forebrain. Nat Neurosci 4:1177–1182.
- Coskun V, Luskin MB (2002) Intrinsic and extrinsic regulation of the proliferation and differentiation of cells in the rodent rostral migratory stream. J Neurosci Res 69:795–802.
- Coskun V, Venkatraman G, Yang H, Rao MS, Luskin MB (2001) Retroviral manipulation of the expression of bone morphogenetic protein receptor Ia by SVZa progenitor cells leads to changes in their p19(INK4d) expression but not in their neuronal commitment. Int J Dev Neurosci 19:219 –227.
- Cummings DM, Emge DK, Small SL, Margolis FL (2000) Pattern of olfactory bulb innervation returns after recovery from reversible peripheral deafferentation. J Comp Neurol 421:362–373.
- Dale K, Sattar N, Heemskerk J, Clark JDW, Placzek M, Dodd J (1999) Differential patterning of ventral midline cells by axial mesoderm is regulated by BMP7 and chordin. Development 126:397–408.
- Dati C, Maggiora P, Puech C, De Bortoli M, Escot C (1996) Expression of the erbB-2 proto-oncogene during differentiation of the mammary gland in the rat. Cell Tissue Res 285:403–410.
- Deans RJ, Moseley AB (2000) Mesenchymal stem cells: biology and potential clinical uses. Exp Hematol 28:875–884.
- Doetsch F, Caille I, Lim DA, Garcia-Verdugo JM, Alvarez-Buylla A (1999) Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. Cell 97:703–716.
- Fan X, Xu H, Cai W, Yang Z, Zhang J (2003) Spatial and temporal patterns of expression of Noggin and BMP4 in embryonic and postnatal rat hippocampus. Brain Res Dev Brain Res 146:51–58.
- Garcia R (2002) Stress, synaptic plasticity, and psychopathology. Rev Neurosci 13:195–208.
- Graziadei PPC, Monti Graziadei GA(1978) Continuous nerve cell renewal in the olfactory system. In: Handbook of sensory physiology (Jacobson M, eds), pp 55–83. Berlin: Springer-Verlag.
- Gritti A, Bonfanti L, Doetsch F, Caille I, Alvarez-Buylla A, Lim DA, Galli R, Verdugo JM, Herrera DG, Vescovi AL (2002) Multipotent neural stem cells reside into the rostral extension and olfactory bulb of adult rodents. J Neurosci 22:437–445.
- Gross RE, Mehler MF, Mabie PC, Zang Z, Santschi L, Kessler JA (1996) Bone morphogenetic proteins promote astroglial lineage commitment by mammalian subventricular zone progenitor cells. Neuron 17:595–606.
- Hack I, Bancila M, Loulier K, Carroll P, Cremer H (2002) Reelin is a detachment signal in tangential chain-migration during postnatal neurogenesis. Nat Neurosci 5:939–345.
- Hogan BLM (1996) Bone morphogenetic proteins: multifunctional regulators of vertebrate development. Genes Dev 10:1580 -1594.
- Jankovski A, Sotelo C (1996) Subventricular zone-olfactory bulb migratory pathway in the adult mouse: cellular composition and specificity as determined by heterochronic and heterotopic transplantation. J Comp Neurol 371:376 –396.
- Kulessa H, Turk G, Hogan BL (2000) Inhibition of BMP signalling affects

growth and differentiation in the anagen hair follicle. EMBO J 19:6664 – 6674.

- Lawson KA, Dunn NR, Roelen BA, Zeinstra LM, Davis AM, Wright CV, Korving JP, Hogan BL (1999) BMP4 is required for the generation of primordial germ cells in the mouse embryo. Genes Dev 13:424 436.
- Li W, Cogswell CA, Lo Turco JJ (1998) Neuronal differentiation of precursors in the neocortical ventricular zone is triggered by BMP. J Neurosci 18: 8853–8862.
- Li W, Lo Turco JJ (2000) Noggin is a negative regulator of neuronal differentiation in developing neocortex. Dev Neurosci 22:68 –73.
- Lim DA, Alvarez-Buylla A (1999) Interaction between astrocytes and adult subventricular zone precursors stimulates neurogenesis. Proc Natl Acad Sci USA 96: 7526 –7531.
- Lim DA, Tramontin AD, Trevejo JM, Herrera DG, Garcia-Verdugo JM, Alvarez-Buylla A (2000) Noggin antagonizes BMP signaling to create a niche for adult neurogenesis. Neuron 28:713–726.
- Lois C, Alvarez-Buylla A (1994) Long-distance neuronal migration in the adult mammalian brain. Science 264:1145–1148.
- Lois C, Garcia-Verdugo J, Alvarez-Buylla A (1996) Chain migration of neuronal precursors. Science 271:978 –981.
- Luskin MB (1993) Restricted proliferation and migration of postnatally generated neurons derived from the forebrain subventricular zone. Neuron 11:173–189.
- Mabie PC, Mehler MF, Kessler JA (1999) Multiple roles of bone morphogenetic protein signaling in the regulation of cortical cell number and phenotype. J Neurosci 19:7077–7088.
- Mariani-Costantini R, Escot C, Theillet C, Gentile A, Merlo G, Lidereau R, Callahan R (1988) In situ c-myc expression and genomic status of the c-myc locus in infiltrating ductal carcinomas of the breast. Cancer Res 48:199 – 205.
- McMahon JA, Takada S, Zimmerman LB, Fan CM, Harland RM, McMahon AP (1998) Noggin-mediated antagonism of BMP signaling is required for growth and patterning of the neural tube and somite. Genes Dev 12:1438 1452.
- Mehler MF, Mabie PC, Zhang D, Kessler JA (1997) Bone morphogenetic proteins in the nervous system. Trends Neurosci 20:309 –317.
- Mehler MF, Mabie PC, Zhu G, Gokhan S, Kessler JA (2000) Development changes in progenitor cell responsiveness to bone morphogenetic proteins differentially modulate progressive CNS lineage fate. Dev Neurosci 22:74 – 85.
- Panchision DM, McKay RD (2002) The control of neural stem cells by morphogenic signals. Curr Opin Genet Dev 12:478 487.
- Peretto P, Merighi A, Fasolo A, Bonfanti L (1997) Glial tubes in the rostral migratory stream of the adult rat. Brain Res Bull 42:9–21.
- Peretto P, Merighi A, Fasolo A, Bonfanti L (1999) The subependymal layer in rodents: a site of structural plasticity and cell migration in the adult mammalian brain. Brain Res Bull 49:221–243.
- Peretto P, Cummings D, Modena C, Behrens M, Venkatraman G, Fasolo A, Margolis FL (2002) BMP mRNA and protein expression in the developing mouse olfactory system. J Comp Neurol 451:267–278.
- Rousselot P, Lois C, Alvarez-Buylla A (1995) Embryonic (PSA) NCAM reveals chains of migrating neuroblasts between the lateral ventricle and the

olfactory bulb of adult mice. J Comp Neurol 351:51-61.

- Shou J, Rim PC, Calof AL (1999) BMPs inhibit neurogenesis by a mechanism involving degradation of a transcription factor. Nat Neurosci 2:339 –345.
- Shou J, Murray RC, Rim PC, Calof AL (2000) Opposing effects of bone morphogenetic proteins on neuron production and survival in the olfactory receptor neuron lineage. Development 127:5403–5413.
- Smith GM, Rutishauser U, Silver J, Miller RH (1990) Maturation of astrocytes in vitro alters the extent and molecular basis of neurite outgrowth. Dev Biol 138:377–390.
- Song H, Stevens CF, Gage FH (2002) Astroglia induce neurogenesis from adult neural stem cells. Nature 417:39 44.
- Spradling A, Drummond-Barbosa D, Kai T (2001) Stem cells find their niche. Nature 414:98 –104.
- Stull ND, Jung JW, Iacovitti L (2001) Induction of a dopaminergic phenotype in cultured striatal neurons by bone morphogenetic proteins. Brain Res Brain Res Dev 130:91–98.
- Temple S, Davis AA (1994) Isolated rat cortical progenitor cells are maintained in division in vitro by membrane-associated factors. Development 120:999 –1008.
- Valenzuela DM, Economides AN, Rojas E, Lamb TM, Nunez L, Jones P, Lp NY, Espinosa R. 3rd, Brannan CI, Gilbert DJ (1995) Identification of mammalian noggin and its expression in the adult nervous system. J Neurosci 15:6077– 6084.
- Wichterle H, Garcia-Verdugo JM, Alvarez-Buylla A (1997) Direct evidence for homotypic, glia-independent neuronal migration. Neuron 18:779 –791.
- Wichterle H, Garcia-Verdugo JM, Herrera DG, Alvarez-Buylla A (1999) Young neurons from medial ganglionic eminence disperse in adult and embryonic brain. Nat Neurosci 2:461–466.
- Xie T, Spradling AC (1998) Decapentaplegic is essential for the maintenance and division of germline stem cells in the Drosophila ovary. Cell 94:251– 260.
- Xie T, Spradling AC (2000) A niche maintaining germ line stem cells in the Drosophila ovary. Science 290:328 –330.