

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Microbiological characterization of artisanal Raschera PDO cheese: analysis of its indigenous lactic acid bacteria**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/41163> since

*Published version:*

DOI:10.1016/j.fm.2007.09.006

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in Food Microbiology, 25, 392-399, 2008

doi:10.1016/j.fm.2007.09.006

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), <http://www.sciencedirect.com/science/article/pii/S0740002007001037>

**Microbiological characterization of artisanal Raschera PDO cheese: analysis of its indigenous lactic acid bacteria**

Paola Dolci\*, Valentina Alessandria, Giuseppe Zeppa, Kalliopi Rantsiou, Luca Coccolin.

*Dipartimento di Valorizzazione e Protezione delle Risorse agroforestali, Settore di Microbiologia e Industrie agrarie, Università degli Studi di Torino, Via L. da Vinci 44, 10095 Grugliasco (TO), Italy.*

Running title: Microbiological characterization of Raschera cheese

\* Corresponding author. Fax: +39 11 6708549  
E-mail address: [paola.dolci@unito.it](mailto:paola.dolci@unito.it)

## **ABSTRACT**

The aim of this research was to study the bacterial populations involved in the production of artisanal Raschera PDO cheese (Italian Maritime Alps, northwest Italy) in order to collect preliminary knowledge on indigenous lactic acid bacteria (LAB). A total of 21 samples of Raschera PDO cheese, collected from six dairy farms located in the production area, were submitted to microbiological analysis. LAB were randomly isolated from M17 agar, MRS agar and KAA plates and identified by combining PCR 16S-23S rRNA gene spacer analysis, species-specific primers and 16S rRNA gene sequencing. Biodiversity of *Lactococcus lactis* subsp. *lactis* isolates was investigated by RAPD-PCR.

LAB microflora showed the highest count values among all microbial groups targeted. They reached counts of  $10^9$  colony forming unit (cfu)/g in cheese samples after 3 days of salting and 15 days of ripening. Yeast population also showed considerable count values, while enterococci and coagulase-negative cocci (CNC) did not overcome  $10^7$  cfu/g. *L. lactis* subsp. *lactis* was the species most frequently isolated from Raschera PDO samples at all different production stages while in aged cheeses *Lactobacillus paracasei* was frequently isolated. RAPD-PCR highlighted that isolates of *L. lactis* subsp. *lactis* isolated from Raschera PDO were highly homogeneous.

## **Keywords**

Lactic acid bacteria; Raschera PDO cheese; microbiological characterization; RAPD-PCR; 16S-23S rRNA gene spacer analysis.

## INTRODUCTION

Raschera PDO is a semi-hard and pressed cheese produced in Piedmont, in Monregalese Valley (Italian Maritime Alps, northwest Italy). Raschera belongs to those niche products strongly connected to Italian food tradition and recently re-discovered and recognized from the European Community with the attribution of the “Protected Denomination Origin” status (European Regulation 510/06). In the last years it has been estimated, on average, an annual production of 2500 quintals, as reported by the Farmers’ Cooperative of Raschera PDO.

In the traditional Raschera PDO cheesemaking, raw cow milk coming from two milkings is warmed to 27-30°C and left resting for 20-30 min after the addition of liquid rennet. Clotting time is established visually by the cheese maker. The curd is cut into particles, then separated from the serum and collected in a hemp cloth where are left for approximately 10 min of drainage. Finally the curd wrapped in the cloth is put into cylindrical forms and pressed to allow remaining serum to drain away for at least 12 hours. At the end, the product is ready to be dry salted and ripened in natural cells dug underground (Official Gazette, No. 92, April 2004). Here the constant temperature and humidity create the optimal conditions for the 1-6 months of ripening and consequently for the formation of the reddish rind characteristic of this cheese.

In the last years industrial dairies have been started producing Raschera PDO cheese by using mesophilic LAB commercial starter cultures. Cheese made from raw milk added of starters results in a more uniform and standardized product of better sanitary quality; however the organoleptic characteristics of the cheese bears little relation to those of its traditional counterpart (Wouters *et al.* 2002; Coppola *et al.* 2006); in fact the addition of commercial starters causes the drop of load of the indigenous microflora responsible for the development

of typical cheese flavours (Steele and Unlu 1992; Wouters *et al.* 2002). The low diversity of starter cultures used in the industrial dairy fermentation is leading to an increasing demand from cheese makers of new lactic acid bacteria (LAB) strains to be used as indigenous starters. The development of autochthonous cultures specific for Raschera PDO fermentation may ensure protection of sensorial properties and at the same time guarantees an uniform and safer product (De Reu *et al.* 2002, Fortina *et al.* 2003; La Rodriguez *et al.* 2004; Samelis *et al.* 2007).

The aim of this research was to study the indigenous bacterial population involved in the production of artisanal Raschera PDO cheese, to obtain preliminary knowledge on LAB species which carry out the traditional fermentation of this cheese. This study represents a first step towards the understanding of microbial population involved in ripening processes of Raschera PDO. Next steps will be the development of selected autochthonous cultures to be used as starters in both industrial and farm processes.

## **MATERIALS and METHODS**

### *DAIRY SAMPLES*

A total of 21 samples were obtained during the summer alpine pasture from 6 representative dairy farms not using starter cultures and located in the production area of Raschera PDO. Nine samples of cheese were taken 3 days after salting that leads to an initial microbial selection. Cheese samples at different ripening stages were also collected and precisely 6 samples at 15 days ripening, 3 samples at 30 days ripening and 3 samples at 70 days ripening. All samples were transported to the laboratory under refrigerated conditions not later than 3 h from the collection, and subjected to microbiological analysis.

## *MICROBIOLOGICAL ANALYSIS*

For microbiological analysis, 10 g of cheese samples were emulsified with sterile quarter strength Ringer solution (Oxoid, Basingstoke, UK) in a Stomacher machine. Decimal dilutions were prepared in Ringer solution and plated onto different selective media for viable counts. The following analyses were carried out: mesophilic aerobic bacteria on Plate Count Agar (Oxoid) at 30°C for 48h, mesophilic and thermophilic cocci on M17 agar (Fluka, Buchs SG, Switzerland), incubated aerobically, respectively at 30°C and 37°C for 48h, mesophilic and thermophilic lactobacilli on MRS agar (Oxoid), incubated anaerobically, respectively at 30°C and 37°C for 48h, enterococci on Kanamycin Aesculin Azide Agar (Fluka) at 37°C for 48h, coagulase-negative cocci (CNC) on Mannitol Salt Agar (Oxoid) at 30°C for 48h, coliforms on Violet Red Bile Lactose Agar (Fluka) at 37°C for 24h, yeast and moulds on Malt Agar (Oxoid) supplemented with tetracycline (1 µg/ml, Fluka) at 25°C for 96h. After counting, from M17, MRS and KAA agars, a total of 15 randomly selected colonies of LAB for each sample were isolated, in order to obtain a representative LAB population of Raschera PDO. They were then purified by two subsequent subcultures on M17 agar and MRS agar and, for long-term maintenance, finally stored at -20°C in M17 and MRS broth containing 30% glycerol, before being subjected to molecular analysis.

## *DNA EXTRACTION from PURE CULTURES*

Genomic DNA from isolates was extracted from 100 µl of an overnight culture diluted with 300 µl TE 1X buffer (10 mM Tris-HCl, 1 mM Na<sub>2</sub>EDTA, pH 8.0) as described by Mora *et al.* (2000).

## *PCR IDENTIFICATION of LAB ISOLATES*

Identification of LAB isolates was carried out by combining PCR 16S-23S rRNA gene spacer analysis (RSA), species-specific primers and 16S rRNA gene sequencing. The RSA was carried out with primers G1 and L1 (Table 1). This analysis allowed to group the isolates on

the basis of polymorphism of 16S-23S rRNA gene spacer region, and to obtain an indication of the hypothetical genus they could belong to (Coppola *et al.* 2001, Fortina *et al.* 2003). According to RSA results, species-specific PCRs and 16S rRNA gene sequencing were planned in order to determine the taxonomic positions of the isolates at species level. The primers used and their corresponding references are reported in Table 1. Isolates for which it was not possible to establish a well defined RSA profile were submitted to partial 16S rRNA gene amplification with primers p8FPL and p806R (Table 1). The amplified fragments were sent for the sequencing to MWG Biotech (Ebersberg, Germany) and the sequences obtained were aligned with those in GenBank with Blast program (Altschul *et al.* 1997) to determine the closest known relatives of isolates.

RSA reactions were performed in a final volume of 25 µl containing 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 2.5 mM MgCl<sub>2</sub>, 0.2 mM deoxynucleoside triphosphates, 0.5 U of Taq polymerase (Eppendorf, Milan, Italy), 0.5 µM concentrations of each primer. One microliter of template DNA was added to the mixture. RSA thermal cycling was as follows: an initial denaturation of 95°C for 5 min, 40 cycle of 2 min at 95°C, 1 min at 55°C, 2 min at 72°C, and 10 min of final extension. PCR products were analysed by electrophoresis in 1.5% Tris-acetate-EDTA agarose gels and a 1Kb Ladder (MBI Fermentas, St. Leon-Rot, Germany) was used as size marker.

#### *RAPD-PCR*

DNA extracted from *Lactococcus lactis* subsp. *lactis* isolates was used in RAPD-PCR analysis. Amplification reactions were carried out using primer D8635 (5'-GAGCGGCCAAAGGGAGCAGAC-3') as previously described by Lombardi *et al.* (2002). RAPD-PCR products were separated by electrophoresis on 1.5% (w/v) agarose gels in 1X TBE buffer at 120 V for 2h. Pictures of the gels were digitally captured by UVIpro Platinum 1.1 Gel Software (Eppendorf) and BioNumerics Software (Applied-Maths, Sint-Martens-

Latem, Belgium) was used for pattern analysis. Calculation of similarities of band profiles was based on the Pearson product moment correlation coefficient. Dendrograms were obtained by means of the Unweighted Pair Group Method by using Arithmetic Average (UPGMA) clustering algorithm (Vauterin and Vauterin 1992). RAPDs were performed at least twice for each strain.

## RESULTS

### MICROBIAL COUNTS.

The mean microbial counts of the 21 samples of Raschera PDO cheese at different ripening stages are summarized in Figure 1 and 2. The minimum and maximum count values are also reported for each microbial group detected.

The mesophilic aerobic bacteria count ranged from  $10^8$  cfu/g to  $10^9$  cfu/g (Figure 1) and reached the maximum value of  $4.2 \times 10^9$  cfu/g in a sample collected 3 days after salting. LAB showed values of  $10^8$  cfu/g to  $10^9$  cfu/g in samples taken after 3 days of salting and 15 days of ripening with the only exception of the lowest load  $4.0 \times 10^6$  cfu/g detected on MRS agar. LAB kept values ranging from  $10^7$  cfu/g to  $10^8$  cfu/g at 30 and at 70 days of maturation (Figure 2). Mesophilic and thermophilic LAB microflora counts as well as presumptive lactococci and lactobacilli counts revealed on M17 and MRS agar plates, did not highlight strong differences.

Enterococci were present with lower counts ( $10^6$  to  $10^7$  cfu/g) compared to lactococci and lactobacilli (Figure 2). The CNC were generally present with values of about  $10^6$  cfu/g even if they reached counts of  $10^7$  cfu/g in 15 days ripened samples (Figure 1). Yeasts also reached the highest levels ( $10^9$  cfu/g) in cheese collected after 15 days of ripening, then for decrease to  $10^7$  cfu/g in 70 days ripened samples (Figure 1). Coliforms were presents at  $10^5$  cfu/g in

samples taken after 3 days of salting and 15 days of ripening. Their loads lowered in aged cheeses however the values were higher than  $10^3$  cfu/g (Figure 1).

#### MOLECULAR IDENTIFICATION of LAB ISOLATES.

A total of 283 presumptive LAB isolates were randomly selected from agar plates and subjected to molecular identification. A first clustering identification step was reached by RSA obtaining seven different profiles (Figure 3). Cluster I was composed of 198 isolates and its profile was characterized by a unique band migrating approximately at 390 bp (Figure 3, lane 2). In cluster II, 7 isolates were characterized by a unique band migrating approximately at 450 bp (Figure 3, lane 3). Cluster III was composed of 10 isolates which the profiles were characterized by a unique band migrating approximately at 380 bp (Figure 3, lane 4).

In cluster IV, 34 isolates were grouped, showing by two bands migrating at 400 and 500 bp (Figure 3, lane 5). For the other 17 isolates two amplification bands at approximately 300 and 400 bp were obtained (cluster V) (Figure 3, lane 6). Cluster VI and VII, including 7 and 3 isolates, respectively, exhibited a three or four bands profile (Figure 3, lane 7 and 8).

Species-specific primers and 16S rRNA gene sequencing were necessary to finally identify the isolates. Within cluster I the isolates were discriminated between the two major subspecies of *L. lactis*, subsp. *lactis* and subsp. *cremoris*, on the basis of the different size of partial histidine biosynthesis operon amplified (Corroler *et al.* 1999). Cluster II and III grouped isolates belonging respectively to the species *Lactococcus garvieae* and *Streptococcus infantarius* subsp. *infantarius*. Cluster IV and V included enterococcal isolates belonging respectively to the species *Enterococcus faecium* and *Enterococcus faecalis*. The isolates within group VI were identified as *Lactobacillus paracasei*, while isolates within cluster VII were tested positive to amplification with *Lactobacillus plantarum* specific primers. The isolates belonging to cluster III showed an unknown RSA profile; thus, they were submitted to 16S rRNA gene sequencing and resulted to belong to *Streptococcus infantarius* subsp.

*infantarius* (Table 2). For 7 isolates it was not possible to establish a well defined RSA profiles, thus they were also submitted to 16S rRNA gene sequencing. Three isolates resulted to belong to the species *Enterococcus casseliflavus*, 1 to *Leuconostoc pseudomesenteroides* and 1 to *Vagococcus carniphilus*. The remaining 2 isolates were classified as species belonging to *Staphylococcus* genus (Table 2)..

Lactococci were dominant in all cheese samples at the different ripening stages; in particular *L. lactis* subsp. *lactis* was present with frequencies ranging from 61% to 86% among LAB isolates; *L. garvieae* and *L. lactis* subsp. *cremoris* were isolated with percentages not higher than 9% and 3% respectively (Figure 4). Non-starter LAB (NS-LAB) isolates were obtained from aged cheeses. *Lb. paracasei* showed a considerable presence (23%) in cheese samples collected after 30 days ripening and *Lb. plantarum* was isolated from both 30 and 70 days ripened cheeses with 3% and 6% of frequency, respectively (Figure 4). Enterococci were present with low percentages at the different ripening stages with the species *E. faecium*, *E. faecalis* and *Enterococcus casseliflavus*. *Vagococcus carniphilus* and *Leuconostoc pseudomesenteroides* were occasionally isolated while *S. infantarius* subsp. *infantarius* was found in cheeses taken 3 days after salting and at 5, 15, 30 days ripening with frequencies of 8%, 7% and 3%, respectively (Figure 4).

In Figure 5 the frequency of isolation of LAB species in correlation to each selective LAB media is reported. M17 and MRS agar showed low selectivity towards lactococcal isolates obtained from both media with comparable percentages while lactobacilli grew preferably on MRS agar. KAA medium was highly selective for enterococci. The incubation temperature used in the study did not allow to obtain a marked growth differentiation among LAB species isolated belonging to mesophilic microflora.

#### *RAPD-PCR*

RAPD-PCR was carried out to explore the genetic diversity of 183 *L. lactis* subsp. *lactis* isolates from Raschera PDO samples. Each strain showed the same RAPD profile in the RAPD analysis repetitions carried out. Elaboration of RAPD profiles obtained with primer D8635 resulted in the dendrogram shown in Figure 6. A general low biodiversity was revealed among *L. lactis* subsp. *lactis* strains. At similarity level of 80% two main clusters, I and II (Figure 6), were detected including, respectively, 108 and 63 lactococcal isolates equally distributed among cheese samples at different ripening stages, coming from the diverse dairy farms. Twelve isolates did not share any similarity with the others studied, and clustered outside the two main groups.

## DISCUSSION

Typical and traditional foods represent an important patrimony for a region and a great opportunity for the development of rural areas. This is the case of Raschera PDO cheese which is produced in a small area on Italian Maritime Alps. The thorough knowledge of a dairy product represents the first step towards the improvement of its quality and, as a consequence, the expansion of its market. In this research we better define the main microflora, which plays a significant role in the ripening process of Raschera PDO cheese. We focused on LAB which play a critical role in the development of the unique characteristics of every cheese variety.

LAB microflora showed the highest count values among all microbial groups targeted, with values comparable to those reported for other Italian ripened cheeses (Randazzo *et al.* 2002; Randazzo *et al.* 2006). LAB reached the highest levels in cheese samples after 3 days of salting and 15 days of ripening. Then they started to decrease in aged cheeses as a result of

autolysis and microbial competition events (Fox *et al.* 2004). Yeast population also showed considerable count values, while enterococci and CNC did not reach loads higher than  $10^7$  cfu/g. Coliforms decreased at the beginning of ripening process but, unexpectedly, increased in number in samples at 70 days of ripening and counts of  $10^4$ – $10^5$  cfu/g were found in the final product.

According to other authors (Ampe *et al.* 1999; Randazzo *et al.* 2002; Ercolini *et al.* 2003) it is noteworthy that some of the used media were not very selective; in fact on MRS agar plates lactococci and staphylococci isolates were also found together with lactobacilli. On the contrary KAA medium showed an high selectivity level towards enterococci.

The incubation temperature used from other authors (Fortina *et al.* 2003; Torres-Llanez *et al.* 2006) to discriminate mesophilic and thermophilic microflora in dairy environments did not allow us to obtain a marked growth differentiation among LAB species isolated from Raschera PDO.

Two-hundred seventy-four LAB isolates obtained from Raschera PDO were grouped by using RSA analysis which proved to be a useful tool to obtain preliminary information on high numbers of LAB isolates. Then the results obtained from species-specific PCR highlighted that *L. lactis* subsp. *lactis* was the species most frequently isolated in Raschera PDO samples at all different ripening stages. This is in agreement with the high occurrence, detected in many european artisanal dairy products (Cogan *et al.* 1997), of wild *L. lactis* subsp. *lactis* as predominant species in cheesemaking. According to RAPD-PCR by using D8635 primer (Andrighetto *et al.* 2004; Cocolin *et al.* 2005; Psoni *et al.* 2007), *L. lactis* subsp. *lactis* isolates found in Raschera PDO showed high similarity. They resulted to belong to a quite homogeneous population; a correlation between genotyping and provenience or ripening stage was not possible.

Other lactococcal isolates found in Raschera PDO belonged to the species *L. lactis* subsp. *cremoris* and *L. garvieae*. The latter has been always considered an important agent of septicemia in fish (Eldar *et al.* 1999) but, in the last years, researchers have found this species in various dairy sources together with *L. lactis* although in a lower proportion (Klijn *et al.* 1995; Morea *et al.* 1999; Villani *et al.*, 2001; Blaiotta *et al.*, 2002; Fortina *et al.* 2003; Fortina *et al.* in press); thus it has been hypothesized a specific role of *L. garvieae* in dairy environment (Fortina *et al.* in press).

*E. faecium*, *E. faecalis* and *E. casseliflavus* were also isolated from Raschera PDO samples. The enterococci are part of the natural flora in cheese manufactured with raw milk and are believed to contribute to the ripening process and the development of flavour and aroma (Cogan *et al.* 1997). However, their presence in dairy samples is often associated to faecal contamination during milking and storing process (Garcia Fontan *et al.* 2001). *S. infantarius* subsp. *infantarius*, isolated from a few samples of Raschera PDO, is also considered an indicator of low hygienic conditions because it is a normal inhabitant of the rumen. Occasionally *V. carniphilus*, belonging to the group of vagococci, phylogenetically and phenotypically similar to enterococci (Shewmaker *et al.* 2004), was isolated. *Lc. paramesenteroides*, a common species in dairy environment (Callon *et al.* 2004; Florez *et al.* 2006) was also found in Raschera PDO.

In aged cheese samples heterofermentative lactobacilli belonging to the so called “secondary culture” were also isolated. *Lb. paracasei* showed the highest frequencies while *Lb. plantarum* was occasionally found in few samples. Considerable amount of NS-LAB are commonly found in ripened semi-hard cheeses (Beresford *et al.* 2001) where contribute to flavour development.

The results of the present research indicate that lactococci have a critical role in the manufacturing and ripening of Raschera PDO. In particular *L. lactis* subsp. *lactis* was found

with the highest frequency values in all the samples analysed so that this species should be strongly considered in the selection of an adequate starter for the production of Raschera PDO.

## REFERENCES

Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman D.J., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25, 3389-3402.

Ampe, F., Ben Omar, N., Moizan, C., Wachter, C. and Guyot, J.P., 1999. Polyphasic study of the spatial distribution of microorganisms in Mexican pozol, a fermented maize dough, demonstrates the need for cultivation-independent methods to investigate traditional fermentations. *Applied and Environmental Microbiology* 65, 5464-5473.

Andrighetto, C., Marcazzan, G. and Lombardi, A., 2004. Use of RAPD-PCR and TGGE for the evaluation of biodiversity of whey cultures for Grana Padano cheese. *Letters in Applied Microbiology* 38, 400-405.

Beresford, T.P., Fitzsimons, N.A., Brennan, N.L. and Cogan, T.M., 2001. Recent advances in cheese microbiology. *International Dairy Journal* 11, 259-274.

Berthier, F. and Ehrlich, S.D., 1998. Rapid species identification within two groups of closely related lactobacilli using PCR primers that target the 16S/23S rRNA spacer region. *FEMS Microbiology Letters* 161, 97-106.

Blaiotta, G., Pepe, O., Mauriello, G., Villani, F., Andolfi, R. and Moschetti, G., 2002. 16S–23S rDNA intergenic spacer region polymorphism of *Lactococcus garvieae*, *Lactococcus raffinolactis* and *Lactococcus lactis* as revealed by PCR and nucleotide sequence Analysis. *Systematic and Applied Microbiology* 25, 520-527.

Callon, C., Millet, L. and Montel, L.C., 2004. Diversity of lactic acid bacteria isolated from Salers cheese. *Journal of Dairy Research* 71, 231-244.

Cheng, S., McCleskey, F.K., Gress, M.J., Petroziello, J.M., Liu, R., Namdari, H., Benigna, K., Salmen, A. and DelVecchio, V.G., 1997. A PCR assay for identification of *Enterococcus faecium*. *Journal of Clinical Microbiology* 35, 1248-1250.

Cocolin, L., Stella, S., Nappi, R., Bozzetta, E., Cantoni, C. and Comi, G., 2005. Analysis of PCR-based methods for characterization of *Listeria monocytogenes* strains isolated from different sources. *International Journal of Food Microbiology* 103, 167-178.

Cogan, T.M., Barbosa, M., Beuvier, E., Bianchi-Salvadori, B., Cocconcelli, P.S., Fernandes, I., Gomez, J., Gomez, R., Kalantzpoulos, G., Ledda, A., Medina, M., Rea, M.C. and Rodriguez, E., 1997. Characterisation of the lactic acid bacteria in artisanal dairy products. *Journal of Dairy Research* 64, 409-421.

Coppola, S., Blaiotta, G., Ercolini, D. and Moschetti, G., 2001. Molecular evaluation of microbial diversity occurring in different types of mozzarella cheese. *Journal of Applied Microbiology* 90, 414-420.

Coppola, S., Fusco, V., Andolfi, R., Aponte, M., Blaiotta, G., Ercolini, D. and Moschetti, G., 2006. Evaluation of microbial diversity during the manufacture of Fior di latte di Agerola, a traditional raw milk pasta-filata cheese of the Naples area. *Journal of Dairy research* 73, 264-272.

Corroler, D., Desmasures, N. and Guéguen, M., 1999. Correlation between polymerase chain reaction analysis of the histidine biosynthesis operon, randomly amplified polymorphic DNA analysis and phenotypic characterization of dairy *Lactococcus* isolates. *Applied and Environmental Microbiology* 67, 2011-2020.

De Reu, K., Debeuckelaere, W., Botteldoorn, N., De Block, J. and Herman, L., 2002. Hygienic parameters, toxins and pathogen occurrence in raw milk cheeses. *Journal of Food safety* 22, 183-196.

Dutka-Malen, S., Evers, S. and Courvalin, P., 1995. Detection of glycopeptide resistance genotypes and identification to the species level of clinically relevant enterococci by PCR. *Journal of Environmental Microbiology* 65, 1005-1008.

Eldar, A., Gorla, M., Ghittino, C., Zlotkin, A. and Bercovier, H., 1999. Biodiversity of *Lactococcus garvieae* strains isolated from fish in Europe, Asia and Australia. *Applied and Environmental Microbiology* 65, 1005-1008.

Ercolini, D., Hill, P.J. and Dodd, C.E.R., 2003. Bacterial community structure and location in Stilton cheese. *Applied and Environmental Microbiology* 69 (6), 3540-3548.

European Union 2006. Commission Regulation EC 510/06. March 2006. Official Journal of the European Communities. L 93/12.

Florez, A.B., Lopez-Diaz, T.M., Alvarez-Martin, P. and Mayo B., 2006. Microbial characterization of the traditional Spanish blue-veined Cabrales cheese: identification of dominant lactic acid bacteria. *European Food Research and Technology* 223, 503-508.

Fortina, M.G., Ricci, G., Acquati, A., Zeppa, G., Gandini, A. and Manachini, P.L., 2003. Genetic characterization of some lactic acid bacteria occurring in an artisanal protected denomination origin (POD) Italian cheese, the Toma piemontese. *Food Microbiology* 20, 397-404.

Fortina, M. G., Ricci, G., Foschino, G.R., Picozzi, C., Dolci, P., Zeppa, G., Cocolin, L. and Manachini, P. L., in press. Phenotypic typing, technological properties and safety aspects of *Lactococcus garvieae* strains from dairy environments. *Journal of Applied Microbiology*, doi:10.1111/j.1365-2672.2006.03265.x.

Fox, P.F., McSweeney, P.L.H., Cogan, T.M. and Guinee, T.P. (Ed.), 2004 "Cheese: Chemistry, Physics and Microbiology, Vol. 1, 3rd ed. Elsevier Ltd, London, UK.

Garcia Fontan, M.C., Franco, I., Prieto, B., Tornadijo, M.E. and Carballo, J., 2001. Microbiological changes in San Simón cheese throughout ripening and its relationship with physico-chemical parameters. *Food Microbiology* 18, 25-33.

Jensen, M.A., Webster, J.A. and Strauss, N., 1993. Rapid identification of bacteria on the basis of polymerase chain reaction-amplified ribosomal DNA spacer region polymorphisms. *Applied and Environmental Microbiology* 59, 945-952.

Klijn, N., Weerkamp, A.H. and De Vos, W.M., 1995. Detection and characterization of lactose-utilizing *Lactococcus* spp. in natural ecosystem. *Applied and Environmental Microbiology* 61, 788-792.

La Rodriguez, S.N., Veciana Nogues, M.T., Roig Sagues, A.X., Trujillo Mesa, A.J. and Vidal Carou, M.C., 2004. Evaluation of biogenic amines and microbial counts throughout the ripening of goat cheeses from pasteurized and raw milk. *Journal of Dairy Research* 71, 245-252.

Lombardi, A., Dal Maistro, L., De Dea, P., Gatti, M., Giraffa, G. and Neviani, E., 2002. A polyphasic approach to highlight genotypic and phenotypic diversities of *Lactobacillus helveticus* strains isolated from dairy starter cultures and cheese. *Journal of Dairy Research* 69, 139-149.

Mora, D., Parini, C., Fortina, M.G. and Manachini, P.L., 2000. Development of molecular RAPD marker for the identification of *Pediococcus acidilactici* strains. *Systematic and Applied Microbiology* 23, 400-408.

Morea, D., Baruzzi, F. and Cocconcelli, P.S. 1999. Molecular and physiological characterization of dominant populations in traditional Mozzarella cheese processing. *Journal of Applied Microbiology* 87, 574-582.

Official Gazette, April 2004. Ministerial Decree. Modifica del disciplinare di produzione della denominazione di origine protetta «Raschera» registrata con regolamento (CE) n. 1263/96 della Commissione del 1° luglio 1996.

Psoni, L., Kotzamanidis, C., Yiangou, M., Tzanetakis, N. and Litopoulou-Tzanetaki, 2007. Genotypic and phenotypic diversity of *Lactococcus lactis* isolates from Batzos, a Greek PDO raw goat milk cheese. *International Journal of Food Microbiology* 114, 211-220.

Randazzo, C.L., Torriani S., Akkermans, A.D.L., De Vos, W.M. and Vaughan, E.E., 2002. Diversity, dynamics, and activity of bacterial communities during production of an artisanal sicilian cheese as evaluated by 16S rRNA analysis. *Applied and Environmental Microbiology* 68, 1882-1892.

Randazzo, C.L., Vaughan, E.E. and Caggia, C., 2006. Artisanal and experimental Pecorino Siciliano cheese: microbial dynamics during manufacture assessed by culturing and PCR-DGGE analyses. *International Journal of Food Microbiology* 109, 1-8.

Samelis, J. and Kakouri, A., 2007. Microbial safety qualities of PDO Galotyri cheese manufactured at the industrial or artisan scale in Epirus, Greece. *Italian Journal of Food Science* 19, 81-90.

Shewmaker, P.L., Steigerwalt, A.G., Morey, R.E., Carvalho M.S., Elliott, J.A., Joyce, K., Barrett, T.J., Teixeira, L.M. and Facklam R.R., 2004. *Vagococcus carniphilus* sp. nov., isolated from ground beef. *International Journal of Sistematic and Evolutionary Microbiology* 54, 1505-1510.

Steele, J.L. and Unlu, G., 1992. Impact of lactic acid bacteria on cheese flavor development. *Food Technology*, 46, 128-135.

Teymoortash, A., Wollstein, A.C., Lippert, B.M., Peldszus, R. and Werner, J.A., 2002. Bacteria and Pathogenesis of Human Salivary Calculus. *Acta Oto-Laryngologica* 122, 210–214.

Torres-Llanez, M.J., Vallejo-Cordoba, B., Diaz-Cinco, M.E., Mazorra-Manzano, M.A. and Gonzalez-Cordova, A.F., 2006. Characterization of the natural microflora of artisanal Mexican Fresco cheese. *Food Control* 17, 683-690.

Vauterin, L. and Vauterin P., 1992. Computer-aided objective comparison of electrophoretic patterns for grouping and identification of microorganisms. *European Microbiology* 1, 37-41.

Zlotkin, A., Eldar, A., Ghittino, C. and Bercovier, H., 1998. Identification of *Lactococcus garvieae* by PCR. *Journal of Clinical Microbiology* 36, 983-985.

Villani, F., Aponte, M., Blaiotta, G., Mauriello, G., Pepe, O. and Moschetti, G., 2001. Detection and characterization of a bacteriocin, garviecin L1-5, produced by *Lactococcus garvieae* isolated from raw cow's milk. *Journal of Applied Microbiology* 90, 430-439.

Ward, L.J.H. and Timmins, M.J., 1999. Differentiation of *Lactobacillus casei*, *Lactobacillus paracasei* and *Lactobacillus rhamnosus* by polymerase chain reaction. *Letters in Applied Microbiology* 29, 90–92.

Wouters, J.T.M., Ayad, E.H.E., Hugenholtz, J. and Smit, G., 2002. Microbes from raw milk for fermented dairy products. *International Dairy Journal* 12, 91-109.

Table 1. Nucleotidic sequence, target gene and reference of primers used for LAB isolates identification.

<i>Primers (5' to 3')</i>	<i>Target gene</i>	<i>Reference</i>
<b>G1</b> GAAGTCGTAACAAGG <b>L1</b> CAAGGCATCCACCGT	Bacterial 16S-23S rRNA gene spacer region	Jensen <i>et al.</i> 1993
<b>p8FPL</b> AGAGTTTGATCCTGGCTCAG <b>p806R</b> CTACGGCTACCTTGTTACGA	Bacterial 16S rRNA region	Teymoortash <i>et al.</i> 2002
<b>EM1A</b> TTGAGGCAGACCAGATTGACG <b>EM1B</b> TATGACAGCGCTCCGATTCC	Nucleotide sequence conserved in <i>E. faecium</i>	Cheng <i>et al.</i> 1997
<b>E1</b> ATCAAGTACAGTTAGTCTT <b>E2</b> ACGATTCAAAGCTAACTG	<i>Ddl</i> gene in <i>E. faecalis</i>	Dutka-Malen <i>et al.</i> 1995
<b>His1</b> CTTCGTTATGATTTTACA <b>His2</b> CAATATCAACAATTCCAT	Histidine biosynthesis operon in <i>L. lactis</i>	Corroler <i>et al.</i> 1999
<b>pLG-1</b> CATAACAATGAGAATCGC <b>pLG-2</b> GCACCCTCGCGGGTTG	16S rRNA gene in <i>L. garvieae</i>	Zlotkin <i>et al.</i> 1998
<b>para</b> CATAACAATGAGAATCGC <b>Y2</b> GCACCCTCGCGGGTTG	16S rRNA gene in <i>L. paracasei</i>	Ward and Timmins 1999
<b>16</b> CATAACAATGAGAATCGC <b>Lpl</b> GCACCCTCGCGGGTTG	16S-23S rRNA gene spacer region in <i>L. plantarum</i>	Berthier <i>et al.</i> 1998

Table 2. Sequence information for isolates with unclear or unknown RSA profile.

Strain	Closest sequence relative	% Identity	GenBank accession no.
I	<i>Streptococcus infantarius</i> subsp. <i>infantarius</i>	99%	DQ232530.1
II	<i>Streptococcus infantarius</i> subsp. <i>infantarius</i>	99%	DQ232529.1
III	<i>Streptococcus infantarius</i> subsp. <i>infantarius</i>	99%	AF177729.1
IV	<i>Enterococcus casseliflavus</i>	100%	EU075043.1
V	<i>Enterococcus casseliflavus</i>	100%	DQ223887.1
VI	<i>Enterococcus casseliflavus</i>	100%	DQ839454.1
VII	<i>Vagococcus carniphilus</i>	99%	AY179329.1
VIII	<i>Leuconostoc pseudomesenteroides</i>	100%	AY929289.1
IX	<i>Staphylococcus</i> sp.	99%	AY578092.1
X	<i>Staphylococcus</i> sp.	99%	DQ356500

Figure 1. Log microbial counts (mean  $\pm$  maximum and minimum values) of non lactic microflora during ripening of Raschera PDO cheese.

Ch3S: 3 days after salting cheeses; Ch15: 15 days ripened cheeses; Ch30: 30 days ripened cheeses; Ch70: 70 days ripened cheeses.

Figure 2. Log microbial counts (mean  $\pm$  maximum and minimum values) of LAB during ripening of Raschera PDO cheese.

Ch3S: 3 days after salting cheeses; Ch15: 15 days ripened cheeses; Ch30: 30 days ripened cheeses; Ch70: 70 days ripened cheeses.

Figure 3. RSA profiles obtained from the LAB isolated from Raschera PDO cheese samples.

Lanes 1: 1 Kb DNA Ladder (MBI); Lanes 2 to 8: amplification profiles obtained by RSA analysis.

Figure 4. Frequency of isolation of LAB species from cheese samples of Raschera PDO at different ripening stages.

Ch3S: 3 days after salting cheeses; Ch15: 15 days ripening cheeses; Ch30: 30 days ripening cheeses; Ch70: 70 days ripening cheeses.

Figure 5. Prevalence of LAB species among isolates randomly selected on KAA, and M17 agar and MRS agar plates incubated at both 30 and 37°C.

Figure 6. RAPD-PCR cluster analysis of profiles obtained with primer D8635 from *Lactococcus lactis* subsp. *lactis* strains isolated from Raschera POD samples.

Figure 1




Figure 2




Figure 3




Figure 4





Figure 5






Figure 6