



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Tumor growth instability and its implications for chemotherapy

This is the author's manuscript		
Original Citation:		
Availability:		
This version is available http://hdl.handle.net/2318/102043 since		
Published version:		
DOI:10.1158/0008-5472.CAN-09-0653		
Terms of use:		
Open Access		
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.		

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on: Questa è la versione dell'autore dell'opera:

Cancer Research 69 (21) 2009 : 8507-8515

DOI: <u>10.1158/0008-5472.CAN-09-0653</u>

The definitive version is available at: La versione definitiva è disponibile alla URL:

http://cancerres.aacrjournals.org/content/69/21/8507

Tumor growth instability and its implications for chemotherapy

P. Castorina^{a,b,c}, D. Carcò^d, C. Guiot^{e,f}, and T.S. Deisboeck^g

^aDipartimento di Fisica, Universita' di Catania, Via Santa Sofia 64, Catania, Italy

^bIstituto Nazionale Fisica Nucleare -Catania, Italy

°Centro Siciliano Fisica Nucleare e Struttura della Materia-Catania, Italy

^dIstituto Oncologico del Mediterraneo - Viagrande (Ct) - Italy

^eDipartimento Neuroscienze, Torino University, Italy

^fCNISM Torino University e Politecnico, Italy

^gComplex Biosystems Modeling Laboratory, Harvard-MIT (HST) Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown, MA 02129, USA

Abstract

The chemotherapy late intensity schedule is revised to account for tumor growth instability, where a small tumor cell fraction emerges that exhibits a higher proliferation rate than the parent strain. Modeling this instability as simplified two-population dynamics we find that: 1) if this instability precedes the onset of treatment, the slope of the linear increase of the drug concentration for the standard "Norton-Simon late intensity schedule" changes and the initial value of the dose strongly depends on the balance of the two tumor cell populations and on their distinct growth rates; and, 2) if the instability trails the initial treatment, the effective chemotherapeutic drug concentration changes as well. Both cases point towards necessary improvements of the "Norton-Simon late intensity" schedule.

Introduction

Cancer therapy is aimed at targeting specific tumor (re)growth patterns to inhibit, or at least control, local tumor proliferation and metastatic dissemination. Mathematical modeling provides a promising tool by which to shed light onto these growth dynamics from a theoretical perspective, using both experimental and clinical data as input. Such works range from investigating various forms of exponential growth kinetics, e.g.,(1,2) to that of describing a Universal growth law (3), with intriguing implications of the latter to radiotherapy (4). For cytotoxic chemotherapy, a well documented case in which theoretical reasoning had an impact on the therapeutic approach is that of breast cancer where, following the "Norton-Simon" hypothesis (5), standard treatment has been adjusted to a "dose dense" regimen; that is, the agent is given at a greater dose rate (i.e., multiple pulses of chemotherapy) to optimize efficacy, and to keep toxic side effects comparatively low. The underlying hypothesis is that, in comparison to the host organ, cancerous tissue exhibits a denser structure with higher fractal mass dimension, and is composed of smaller, relatively faster growing sub-seeds, all of which follow the Gompertz law. Differing from the traditional "log-kill" concept (6), Norton and Simon then argued, based on a body of experimental and clinical studies on solid cancers, that "therapy results in a rate of regression proportional to the growth rate of an unperturbed neoplasm of that size" (7,8).

This hypothesis has been tested in a phase III clinical study led by the Cancer and Leukemia Group B (CALGB 97-41) [http://www.calgb.org/](9) This large prospective randomized trial

demonstrated that shortening the time interval between each chemotherapy cycle while maintaining the same dose size resulted in significant improvements in disease-free and overall survival in patients with node-positive breast carcinoma. More recently, also a phase I study in patients with metastatic breast cancer confirmed the therapeutic results predicted by the Norton-Simon model (10). However, the issue of drug resistance is only considered at the qualitative level within the framework of the Norton-Simon model, contrary to some leading work addressing this issue performed by Goldie and Coldman (11, 12). In a more recent theoretical work, (13) Coldman and Murray investigated a stochastic model of non-cell cycle phase specific cancer chemotherapy for exponentially growing tumours including the development of drug resistance and the effect of chemotherapy on normal cells. The authors concluded that early intensification, a feature shared with the Norton Simon model, is a common aspect of successful regimes where drug resistance is likely. We note that the role of intra-tumor heterogeneity and its implications for the evolution of drug resistance and cell kinetics has already been considered by Gardner (14), who proposed a kinetically tailored treatment (KITT model). Finally, Monro & Gaffney (15) recently proposed a model aiming at a unifying description based on well founded approximations, i.e. Gompertzian tumour growth, log cell kill by chemotherapy and Luria-Delbruck mutation to resistance (this assumption was based on the Nobel Prize winning work of Luria and Delbruck in 1943 (16), who showed that bacterial cultures developed resistance to bacteriophages at random). They were able to show that palliative continuous chemotherapy achieves optimal results for intermediate (rather than high) dosage levels with later (rather than earlier) intervention, due to the effects of competition between resistant and sensitive cells. However, their approach focused only on the different tumor response to therapy and not on the different growth rate (both before and after therapy) that represents intrinsic tumor heterogeneity.

Here, we present evidence that cancer's inherent clonal diversity results in non-homogenous growth patterns which in turn - even for its simplest form of two-population dynamics - will require a modification of the "late intensification schedule" that has been put forward by Norton-Simon (17). The next section introduces the underlying growth instability model and its relevance for the current standard schedule for both situations, i.e. if the instability arises prior or after the regimen. We also propose an approach more closely related to clinical therapy and compare the results with data on survival probability.

Methods

Let us recall that for a homogeneous system the general growth law can be written as

$$\frac{1}{N}\frac{dN}{dt} = \alpha(t)$$
[1]

where N is the number of individual cells at time t and α is their specific growth rate. The growth laws can be classified according to the order of power expansion in of the generating function (18)

$$\frac{d\alpha}{dt} = \Phi\left(\alpha\right)$$
[2]

For instance, exponential growth, the Gompertz and logistic laws as well as various power laws proposed to describe tumor growth can be obtained by using

$$\Phi(\alpha) = 0, \Phi(\alpha) = c_1 \alpha, \quad \Phi(\alpha) = c_1 \alpha + c_2 \alpha^2$$

If N is a population composed of n sub-populations, each proliferating with a different specific growth rate α_i , i = 1, ..., n,

$$\frac{1}{N_i}\frac{dN_i}{dt} = \alpha_i(t)$$
[3]

then eqn. [1] can be restored, provided α is assumed to be the average of α_i , i.e.

$$\alpha(t) = \frac{1}{N} \sum_{i=1}^{n} \alpha_i(t) N_i(t)$$
[4]

For simplicity, let us consider a cancer system where, at time δ following a progression event, a small cancer cell subpopulation emerges which grows at a distinctively different replication rate. At any time one can write

$$N(t) = N_1(t) + N_2(t)$$
[5]

Where $N_1(t)$ and $N_2(t)$ are the number of cells belonging to the two subpopulations, evolving according to eqns.:

$$\frac{1}{N}\frac{dN_1}{dt} = \alpha_1(t)$$
[6]

and

$$\frac{1}{N_2} \frac{dN_2}{dt} = \alpha_2(t) \theta(t - t_\delta)$$
^[7]

where θ ($t - t_{\delta}$) is 1 for $t \ge t_{\delta}$ and 0 for $t < t_{\delta}$.

The initial value is given by $N(0) = N_1(0)$, at $t = t_{\delta}$ the cell population $N_2(t)$ starts to grow and, since the instability emerges when a small portion of tumor cells is more rapidly growing, let us consider $N_1(t_{\delta}) >> N_2(t_{\delta})$.

To clarify the implications of the previous equations, let us assume that the two populations both follow the Gompertz law (GL), i.e.

$$\frac{1}{N_1} \frac{dN_1}{dt} = K_1^{g} \ln\left(\frac{N_1^{\infty}}{N_1}\right)$$
[8]

$$\frac{1}{N_2} \frac{dN_2}{dt} = \theta \left(t - t_{\delta} \right) K_2^g \ln\left(\frac{N_2^{\infty}}{N_2}\right)$$
[9]

This translates into the idea that the same type of cancer, in its sub-clones $N_1(t)$ and in $N_2(t)$, follows the **same** functional form of the growth law bu**tlifferent** values of the specific growth rate. In the simplest model of this instability, with $K = K_1^g = K_2^g$, the difference in the specific growth is related to the saturation value of the cell number $N(t \to \infty) = N^\infty$ given by

$$N^{\infty} = N_1^{\infty} + N_2^{\infty}$$
[10]

According to previous eqns. [8,9], different specific growth rates are obtained for $N_1^{\infty} \neq N_2^{\infty}$. In order to evaluate the effects of the choices of the parameter values, various configurations will be considered.

1. Let us first show the effects of the delay, i.e. the onset of growth of N_2 at $t_{\delta} \neq 0$, on

tumor growth assuming $N_1^{\infty} = N_2^{\infty} = \frac{N^{\infty}}{2}$, that is for equal specific growth rates. Figs.1a and 1b show the results for breast cancer with an asymptotic mass $M^{\infty} = 640g$ (the mass is assumed to be proportional to the cell number N and an initial mass of M(0) = 6.4 g. The delay t_{δ} is 100 days in fig. 1a and 500 days in fig.1b. The corresponding GL for a homogeneous cell population with the same initial and final mass is also plotted.

- 2. Let us now consider another case, in which the second cell population develops with the same delay of 500 days, but its specific replication rate (leading to a different fraction of the final tumor mass) is modified. In both cases, N_2 grows more rapidly than N_1 , which is obtained by imposing $N_1^{\infty} < N_2^{\infty}$. In fig.2a the result for $M_1^{\infty} = 0.4$ M^{∞} and $M_2^{\infty} = 0.6 M^{\infty}$ is depicted, while fig.2b shows the plot for $M_1^{\infty} = 0.2 M^{\infty}$ and $M_2^{\infty} = 0.8 M^{\infty}$ respectively.
- 3. Let us now clarify the effects of the size of the sub-system N_2 on tumor growth, at time delay t_{δ} , with respect to the sub-system N_1 . If the sub-clone N_2 contributes substantially to N^{∞} its specific proliferation rate must strongly increase to reach a predefined fraction of the total final tumor mass, and such an increase is much larger depending on how, at t_{δ} its size compares to N_1 (t_{δ}).

In fig. 3a we show, for the same set of parameters used in fig. 2a, the dramatic impact of the unstable system when $N_2(t_{\delta})$ is 10^{-6} smaller than $N_1(t_{\delta})$, for $t_{\delta} = 500$ days. The effect of increasing the delay time is shown in Fig. 3b by considering the same set of values of the parameter as in Fig. 3a and $t_{\delta} = 1000$ days. Let us note that a similar analysis can be performed easily with other growth laws, for example the Universal law proposed by West et al (19,20). The results are expected to be qualitatively the same as those obtained using the Gompertz law, which is appropriately used here to relate the effects of tumor instability to chemotherapy regimen.

To investigate the potential effects of such tumor instability on treatment, let us consider the 'Norton-Simon' hypothesis with regards to its 'late intensity' schedule for chemotherapy. Considering a monoclonal cell population that grows according to eqn.[1], and assuming that the dose-response per cell to a non-specific chemotherapeutic cycle of treatment can be described by a function C(t), the time evolution of the system is given by the following equation:

$$\frac{1}{N}\frac{dN}{dt} = \alpha(t) - C(t)$$
[11]

The tumor stops growing, and eventually regresses, when the therapeutic dose, delivered at time t^* , is larger than a critical value given by

$$C > C_{cr} = \alpha \left(t^* \right)$$
[12]

Moreover, assuming that, due to the rapeutic impact, the cancer cell number decreases exponentially with a rate ω , then

$$N(t) = N(t^*) \exp\left(-\omega(t - t^*)\right)$$
^[13]

which implies that, substituting $\frac{1}{N}\frac{dN}{dT} = -\omega$ in eqn. [11],

$$C(t) = \omega + \alpha(t)|_{N=N(t)}$$
[14]

i.e. the time-dependent dose, according to the 'Norton-Simon' hypothesis, depends on the tumor growth rate $\alpha(t)$ of a single Gompertzian growth. By substituting for N the expression given in eqn. [13] and collecting the constant and the time dependent terms, we obtain that C(t), in the case of Gompertzian growth rate, is a linearly increasing function of time:

$$C^{g}(t) = A + B(t - t^{*})$$
[15]

where $A = \omega + K^g \ln \left(\frac{N^{\infty}}{N(t^*)}\right)$ and $B = \omega K^g$ and

$$C^{g}_{cr} = K^{g} \ln\left(\frac{N^{\infty}}{N(t^{*})}\right)$$
[16]

Therefore a linear increase of the chemotherapeutic drug dose with time can reduce the number of tumor cells exponentially. However, chemotherapeutic treatments based on the previous considerations assume *single* population dynamics.

Indeed if one considers the simplest case of heterogeneity i.e. a system of two subpopulations, with in general $K_1^g \neq K_2^g$ (see eqs.[8,9]), since $N(t) = N_1(t) + N_2(t)$ the previous eqn.[13] gives

$$N(t) = N_1(t^*) \exp(-\omega(t - t^*)) + N_2(t^*) \exp(-\omega(t - t^*))$$
[17]

The cancer populations N_1 and N_2 follow Gompertzian growth kinetics with the corresponding specific rates and drug concentrations, that is

$$\frac{1}{N_1} \frac{dN_1}{dt} = \alpha_1(t) - C_1(t)$$
[18]

$$\frac{1}{N_2}\frac{dN_2}{dt} = \alpha_2(t) - C_2(t)$$
[19]

and therefore for the time evolution of the total number of tumor cells, N(t), after onset of treatment, i.e $t > t^*$ one arrives at the following equation:

$$\frac{1}{N}\frac{dN}{dt} = \alpha (t) - C(t)$$
[20]

Where

$$\bar{\alpha(t)} = \frac{N_1(t)}{N(t)} \alpha_1(t) + \frac{N_2(t)}{N(t)} \alpha_2(t)$$
[21]

and

$$\bar{C(t)} = \frac{N_1(t)}{N(t)} C_1(t) + \frac{N_2(t)}{N(t)} C_2(t)$$
[22]

We will now discuss which modification of the `late intensity schedule' can be relevant for an unstable, i.e. heterogeneous tumor exemplified by two-population dynamics, according to the previous equations.

i) Onset of tumor instability after the beginning of chemotherapy

Let us first assume that tumor instability starts after the beginning of the therapy treatment, i.e. $t_{\delta} > t^*$. In this case, for $t^* < t < t_{\delta}$, the time evolution is due to population $N_1(t)$ only and the corresponding drug concentration for a late intensity schedule is

$$C_{I}(t) = \omega + K_{1}^{g} \ln\left(\frac{N^{\infty}}{N(t^{*})}\right) + \omega K_{1}^{g}(t-t^{*}) \qquad for \quad t^{*} < t < t_{\delta}$$
[23]

Note that, in this time interval, the growth rate of population $N_1(t)$ is that of the `one-population Gompertzian law' with saturation value N^{∞} , because $N_1(t) = N(t)$.

By recalling that

$$N_{1}(t_{\delta}) = N_{1}(t^{*}) \exp(-\omega(t_{\delta} - t^{*}))$$
[24]

and

$$N(t_{\delta}) = N_1(t_{\delta}) + N_2(t_{\delta})$$
^[25]

therefore, by assuming that for $t > t_{\delta}$ the exponential decrease of the total tumor cell number elicited by the therapy is still effective, i.e.

$$N(t) = N(t_{\delta}) \exp\left(-\omega(t - t_{\delta})\right)$$
[26]

it turns out

Where

$$A_d = \omega + \frac{N_1(t_{\delta})}{N(t_{\delta})} K_1^g \ln\left(\frac{N_1^{\infty}}{N_1(t_{\delta})}\right) + \frac{N_2(t_{\delta})}{N(t_{\delta})} K_2^g \ln\left(\frac{N_2^{\infty}}{N_2(t_{\delta})}\right)$$
[28]

and

$$B_d = \frac{N_1(t_{\delta})}{N(t_{\delta})} K_1^g + \frac{N_2(t_{\delta})}{N(t_{\delta})} K_2^g$$
[29]

On the other hand, for a `one-population Gompertzian growth', with saturation value N^{∞} , one has

$$C_g = \omega + K^g \ln\left(\frac{N^{\infty}}{N(t_{\delta})}\right) + \omega K^g (t - t_{\delta})$$
[30]

ii) Onset of tumor instability prior to the beginning of chemotherapy

Let us now assume that $t^* > t_{\delta}$, i.e. the treatment (delivered at time t^* follows the onset of instability (at time t_{δ}). Accordingly, the tumor grows following the `one-population Gompertz law' up until time t_{δ} ($N(t) = N_1(t)$ for $t < t_{\delta}$) when the population $N_2(t)$ emerges with a different specific proliferation rate.

Through the aforementioned eq.(22), the drug concentration for $t < t^*$ turns out to be

$$C(t) = A_{2p} + \omega B_{2p}(t - t_{\delta})$$
^[31]

Where

$$A_{2p} = \omega + \frac{N_1(t^*)}{N(t^*)} K_1^g \ln\left(\frac{N_1^{\infty}}{N_1(t^*)}\right) + \frac{N_2(t^*)}{N(t^*)} K_2^g \ln\left(\frac{N_2^{\infty}}{N_2(t^*)}\right)$$
[32]

and

$$B_{2p} = \frac{N_1(t^*)}{N(t^*)} K_1^g + \frac{N_2(t^*)}{N(t^*)} K_2^g$$
[33]

The role of tumor instability is crucial for the time behavior of the drug concentration because it determines different populations, $N_1(t^*)$ and $N_2(t^*)$ at time t^* , and different slopes in the linear time dependence of $\overline{C(t)}$. In fact, during the interval $t_{\delta} < t < t^*$ the two populations, N_1 and N_2 , follow a Gompertzian growth with different specific rates. As we shall see in the next section, the presence of instabilities, both before and after the time at which the therapy has been administrated, postulates further improvements of the "Norton-Simon late intensity" schedule based on an uniform rate of regression, ω , of the number of cells according to eqn. [13]. The previous eqns.[27,30,31] show that, for the two population dynamics following a Gompertzian growth, the average drug concentration required to obtain such a result is different from the drug concentration for a single tumor Gompertzian population. This general result is obtained by mimicking the differential response of the two populations to the treatment by different drug concentrations C_1 and C_2 and the same ω (see eqns.[17,18,19]). However, one can consider another approach to the problem, that could, to some extent, be closer to the real setting, where a "cocktail" of drugs is given which are expected to be effective against a particular tumor on the basis of previous empirical data. Indeed, in the therapy modeled according to the previous treatment, one expects that the drug concentration is uniformly delivered among cells and that $C_1 = C_2 = C$ but the "differential" effectiveness against a cancerous clones, i.e. the reduction (for unit time) of cancer cells in the two populations, is translated in a different rate of regression for populations N_1 and N_2 , that is $\omega_1 \neq \omega_2$.

To be more precise, let us consider the growth equations

$$\frac{1}{N_1} \frac{dN_1}{dt} = \alpha_1 (t) - C(t)$$
[34]

$$\frac{1}{N_2} \frac{dN_2}{dt} = \alpha_2 (t) - C (t)$$
[35]

and let us assume that, after the time t^* at which the treatment started, the therapeutic result is

$$N_1(t) = N_1(t^*) \exp(-\omega_1(t - t^*))$$
[36]

and

$$N_{2}(t) = N_{2}(t^{*}) \exp\left(-\omega_{2}(t-t^{*})\right)$$
[37]

In this discussion the delay time t_{δ} , is neglected but it can be easily included. From the previous equations for population N_1 it turns out that (one assumes for simplicity that $K_1^g = K_2^g = K$)

$$C(t) = \omega_1 \left(1 + K(t - t^*) \right) + K \ln \left(\frac{N_1^{\infty}}{N_1(t^*)} \right)$$
[38]

which quantifies the linearly increasing drug concentration required for population N_1 to exhibit a decrease in time with **constant** rate ω_1 . Analogously, by assuming that ω_2 is time independent, for population N_2 the required drug concentration turns out to be

$$C(t) = \omega_2 \left(1 + K \left(t - t^* \right) \right) + K \ln \left(\frac{N_2^{\infty}}{N_2 \left(t^* \right)} \right)$$
[39]

In other terms, by requiring that the same dose C(t) can reduce the two populations with different constant rates, one obtains that, at any time, the following condition should be fulfilled:

$$\omega_1 \left(1 + K \left(t - t^* \right) \right) + K \ln \left(\frac{N_1^{\infty}}{N_1 \left(t^* \right)} \right) = \omega_2 \left(1 + K \left(t - t^* \right) \right) + K \ln \left(\frac{N_2^{\infty}}{N_2 \left(t^* \right)} \right)$$
[40]

Apart from the trivial solution $\omega_1 = \omega_2$ and

$$\ln \left(\frac{N_1^{\infty}}{N_1(t^*)}\right) = \ln \left(\frac{N_2^{\infty}}{N_2(t^*)}\right)$$
[41]

eqn. [40] cannot be satisfied except when relaxing the requirement that some reduction rate is time independent. This implies that eqn.[37] for the effect of the drug concentration on population N_2 , for example, must be modified with a time dependent ω_2 , i.e.

$$N_{2}(t) = N_{2}(t^{*}) \exp(-\omega_{2}(t)(t - t^{*}))$$
[42]

obtaining for ω_2 the linear differential equation:

$$\frac{1}{N_2} \frac{dN_2}{dt} = -\frac{d\omega_2}{dt} (t - t^*) - \omega_2 = K \ln \frac{(N_2^{\circ\circ})}{N_2(t^*)} + \omega_2 K (t - t^*) - \omega_1 (1 + K (t - t^*)) - K \ln \frac{(N_1^{\circ\circ})}{N_1(t^*)}$$
[43]

where the last term has been obtained by imposing that there is a decrease of population N_1 with constant rate ω_1 . By defining the parameter

$$\beta = K \ln \left(\frac{N_1^{\infty} N_2(t^*)}{N_2^{\infty} N_1(t^*)} \right)$$
[44]

the solution of the previous equation can be written as

$$\omega_{2}(t) = \omega_{1} - \beta \left(\frac{1 - \exp\left(-K\left(t - t^{*}\right)\right)}{K\left(t - t^{*}\right)} \right)$$
[45]

with the boundary condition $\omega_2(t^*) = \omega_1 - \beta$.

This result allows us to compare the predicted evolution of the two different populations with experimental data.

Results

Using the model described above, we now evaluate its effects on chemotherapeutic treatment and its predictive power by using medical data on breast cancer survival probability.

The ratio $D_{II} = \frac{C_{II}}{C_g}$ in eqns.[27,30] describes the effects in the drug therapy with respect to the single Gompertzian growth when tumor instability starts after the beginning of the therapy, i.e. $t_{\delta} > t^*$. For $t^* < t < t_{\delta}$ this ratio is 1 and therefore one can set $t^* = 0$. In Fig.4 D_{II} is plotted for

 $t_{\delta} = 60$ days and $\omega = 0.1$ per month, and different values of $\frac{K_2^g}{K_1^g}$ are considered. For a onepopulation Gompertzian growth, of course, $D_{II} = 1$. With this set of parameters D_{II} changes

less than 10%. Notice that, when $\frac{K_2^g}{K_1^g} = 1.5$, the value of D_{II} can be less than one for some time intervals. This is due to the fact that at the onset of instability the total tumor mass splits in two subpopulations with different specific proliferation rates but also with different cell numbers at t_{δ} . Therefore a depletion of the large population $N_1(t_{\delta})$ occurs and it requires some time for

the growing instability, N_2 , to compensate this initial negative step. Then ratio $D = \frac{C(t)}{C_g(t)}$ in eqns.[30,31] (which is, by definition, equal to 1 for $t < t_{\delta}$) describes the effects for onset of tumor instability prior to the beginning of chemotherapy. D strongly depends on the specific proliferation rates of the two populations and on their relative fraction of the total cancer cell population at the onset of instability and of treatment. In Fig.5a the ratio D is plotted as a

function of time for $t^* = 365$ days and $t_{\delta} = 100$ days for different values of $\frac{K_2^g}{K_1^g}$.

In Fig.5b, *D* is plotted for the same t_{δ} and t^* values, and for the same ratios $\frac{K_2^g}{K_1^g}$, but changing the relative population $\frac{N_2}{N_1}$ at the onset of instability and at saturation (i.e. when $t \to \infty$). Depending on the value of $\frac{K_2^g}{K_1^g}$, the ratio *D* increases from 20% to 70% (see figure captions for details).

We note that the values of the ratios D and D_{II} depend on many parameters, which should be properly taken into account in a clinical context. On the other hand the presence of instabilities, both before and after the time at which the therapy has been administrated, postulates further improvements of the "Norton-Simon late intensity" schedule. For this reason, in the previous section, a more general approach with different rates of regression for populations N_1 and N_2 ($\omega_1 \neq \omega_2$) has been also considered. It can be applied to more realistic clinical cases. For example, it is well known that primary breast cancer cells can express a variety of surface receptors which correlates with clinical outcome. In particular, during their proliferative activity, breast cancer cell populations may start expressing receptors for hormones, mainly for estrogens and progesterone (21,22). Let us denote, in a breast cancer, the original cell population N_1 , that fails to express estrogen receptor, as ER-, and the population N_2 , that starts to express such receptors, as ER+. To be able to compare our model with experimental data, one has to relate the survival probability of a patient at timeP(t), after the end of chemotherapy which occurred at time t_e , assuming a survival tumor cell fraction at time computed according to eqns.[36,37], in which $t^* = 0$: therefore

$$\alpha_1 = K ln \left(\frac{N_1^{\infty}}{N_1(0)} \right) + \omega_1 K t_e$$
[46]

and

$$\alpha_2 = K ln\left(\frac{N_2^{\infty}}{N_2(0)}\right) + \omega_2 K t_e = K ln\left(\frac{N_2^{\infty}}{N_2(0)}\right) + K t_e\left(\omega_1 - \beta\left(\frac{1 - exp\left(-Kt_e\right)}{Kt_e}\right)\right)$$

$$\tag{47}$$

where eqn.[45] has been used. By a simple calculation

$$\alpha_2(t_e) = \alpha_1(t_e) + \beta exp(-Kt_e)$$
^[48]

At least for short time intervals after the end of chemotherapy, the survival patient probability should be proportional to the regrowth rate at the end of the therapy: larger regrowth rate corresponds to smaller survival probability. Therefore one can write

$$P_{i}(t) = 1 - \alpha_{i}(t_{e})(t - t_{e})$$
[49]

For not too large $t - t_e$, the value of the parameter α_1 can be fitted by approximating the curve ER- (ER/PgR absent) in fig. 1 of Colleoni et al.2(3). This gives $\alpha_1 \approx 0.071$ per year. To evaluate the survival probability with ER+ we use the corresponding formula with α_2 (t_e) (note that a different t_e is simply a rescaling of β). By comparison with the corresponding clinical data, the parameter β can be estimated by trial and error methods and turns out to be $\beta \sim 0.08$ per year for times in the range of 1-4 years. On the basis of such values, the survival probabilities can be estimated and the comparison with the corresponding clinical data is given in Table 1. These results are in good agreement with the data reported in 2β) for the curve ER+ (ER/PgR present) and suggest a generalization of eqn.[49] in exponential form, that is

$$P_i(t) = exp(-\alpha_i t)$$
^[50]

which gives the results reported in Table 1 by using $\beta \sim 0.13$ per year. One should note that the previous comparisons have been done by considering ω_1 time independent and without any optimization of the parameter β . This preliminary analysis does not show the saturation of the patient survival probability observed at long time intervals, however it suggests a simple relationship between the drug dose required to obtain a given regression rate and the survival probability of the patient which could be generalized in analogy to the so called "linear-quadratic model" used in radiotherapy. We shall discuss this point in detail in a forthcoming paper.

Discussion

In here we propose a model aimed at gaining more insights into how tumors respond to therapy, taking into account some features which are normally disregarded, and in particular:

- 1. the well known cancer characteristics of clonal heterogeneity and the growth instability it produces;
- 2. the analysis of the effects of such instabilities when occurring before and after the administration of chemotherapy;
- 3. their potential impact on the overall survival probability of the patient.

We specifically investigated the case of an initially monoclonal tumor in which, at time t_{δ} , a secondary, faster replicating strain emerges from the parent tumor cell population. The resulting "two population dynamics" is described in a way that is somewhat different from the usual interspecific competition of the two populations for survival and access to resources (prey-predator model, see, for example (24)) since the details about the dynamical evolution of the two distinct populations in the tumor required by such a more "traditional" approach would be unavailable, at least until the tumor is surgically approached and a biopsy taken for histological evaluation. In our case, therefore, the competition between the two cell populations

is encoded in the ratio $\frac{N_1^{\infty}}{N_2^{\infty}}$ and in the boundary condition for long intervals: formally the difference in specific growth is related to the saturation value of the cell number, $N(t \rightarrow \delta) =$ N^{∞} , given by eqn. [10].

Since the maximum number of individuals N^{∞} , the ratio and the delay in the onset of growth between the two subpopulations results in different specific growth rates, a variety of distinct realistic situations can be described without necessity of other "ad-hoc" assumptions. For instance, the occurrence of "dormancy" phases during tumor development, which has been simulated by Speer et al. (25) assuming that some parameters describing the overall Gompertzian growth may experience stepwise variations (responsible for the appearance of "plateaus" in the tumor growth curve), maybe generated by proper choices of the delay time t_{δ} (see Fig 3b). We also investigated how the conventional "late-intensity" chemotherapy schedule proposed by Norton and Simon, based on the underlying assumption of singlepopulation dynamics (with the parameters of a unique homogeneous Gompertzian growth), has to be modified when tumor time evolution is non uniform. We showed that in both case, i.e. if the instability starts before or after the onset of treatment the chemotherapeutic dose strongly depends on the balance of the two populations and on their specific growth rates.

Finally, by introducing a drug concentration non linear in time, the model was evaluated with some available clinical data, comparing its predictions for the subpopulation ER+ that emerges from the original breast cancer cells. Comparing the predicted survival probability following therapy, we conclude that the model is capable to prefigure the different responses on the basis of a small number of parameters, whose values can be easily fitted from clinical data. Most importantly, our simplified analysis indicates already that a global, average chemotherapeutic approach will not be successful if growth instabilities were to occur. It further argues that a better phenomenological and theoretical understanding of such therapeutic effects requires a more accurate description of these instabilities.

Acknowledgments

CG gratefully acknowledges the support from Regione Piemonte (PSF 2007, 2008) and University of Torino (60% 2007, 2008). This work has been supported by NIH grant CA 113004 (TSD) and by the Harvard-MIT (HST) Athinoula A. Martinos Center for Biomedical Imaging and the Department of Radiology at Massachusetts General Hospital.

References

- (1). Marusic M, Bajzer Z, Freyer JP, Vuc-Pavlovic S. Analysis of growth of multicellular tumour spheroids by mathematical models. Cell Prolif 1994;27:73-94. [PubMed: 10465028]
- (2). Retzky MW, Swartzendruber DE, Wardwell RH, Bame PD. Is Gompertzian or exponential kinetics a valid description of individual human cancer growth? Med Hypotheses 1990;33:95–106. [PubMed: 2259298]
- (3). Guiot C, Degiorgis PD, Delsanto PP, Gabriele P, Deisboeck TS. Does tumor growth follow a "universal law"? J Theor Biol 2003;225:147-151. [PubMed: 14575649]
- (4). Castorina P, Deisboeck TS, Gabriele P, Guiot C. Growth laws in cancer: implications for radiotherapy. Radiat Res 2007;168:349-356. [PubMed: 17705631]
- (5). Norton L, Simon R. Tumor size, sensitivity to therapy and the design of treatment protocols. Cancer Treat Rep 1976;61:1307–1317. [PubMed: 589597]
- (6). Skipper HE. Laboratory models: some historical perspectives. Cancer Treat Rep 1986;70:3-7. [PubMed: 3943114]
- (7). Norton L. Conceptual and practical implications of breast cancer geometry: Toward a more effective, less toxic therapy. The Oncologist 2005;10:370–381. [PubMed: 15967831]

- (8). Norton L, Simon R. The Norton-Simon hypothesis: Designing more effective and less toxic chemotherapeutic regimens. Nat. Clin Pract Oncol 2006;3:406–407. [PubMed: 16894366]
- (9). Fornier M, Norton L. Dose-dense adjuvant chemotherapy for primary breast cancer. Breast Cancer Res 2005;7:64–69. [PubMed: 15743513]
- (10). Traina AT, Theodoulou M, Feigin K, et al. Phase I Study of a Novel Capecitabine Schedule Based on the Norton-Simon Mathematical Model in Patients With Metastatic Breast Cancer. J Clin Oncol 2008;26:1797–1802. [PubMed: 18398145]
- (11). Goldie JH, Coldman AJ. Quantitative model for multiple levels of drug resistance in clinical tumours. Cancer Treat Rep 1983;67:923–931. [PubMed: 6627236]
- (12). Goldie JH, Coldman AJ. Application of theoretical models to chemotherapy protocol design. Cancer Treat Rep 1986;70:127–131. [PubMed: 3510731]
- (13). Coldman AJ, Murray JM. Optimal control for a stochastic model of cancer chemotherapy. Math Biosci 2000;168:187–200. [PubMed: 11121565]
- (14). Gardner SN. Modeling Multi-drug Chemotherapy: Tailoring Treatment to Individuals. J Theor Biol 2002;214:181–207. [PubMed: 11812172]
- (15). Monro HC, Gaffney EA. Modelling chemotherapy resistance in palliation and failed cure. J Theor Biol 2009;257:292–302. [PubMed: 19135065]
- (16). Luria SE, Delbruck M. Mutations of bacteria from virus sensibility to virus resistance. Genetics 1943;28:491–511. [PubMed: 17247100]
- (17). Norton L, Simon R. The Norton-Simon hypothesis revisited. Cancer Treat Rep 1986;70:163–169. [PubMed: 3510732]
- (18). Castorina P, Delsanto PP, Guiot C. Classification scheme for phenomenological universalities in growth problems in physics and other sciences. Phys Rev Lett 2006;96:188701–5. [PubMed: 16712405]
- (19). West GB, Brown JH, Enquist BJ. A general model for ontogenetic growth. Nature 2001;413:628–631. [PubMed: 11675785]
- (20). West GB, Woodruff WH, Brown JH. Allometric scaling of metabolic rate from molecules and mitochondria to cells and mammals. PNAS 2002;99(suppl 1):2473–2478. [PubMed: 11875197]
- (21). Clarke RB, Howell A, Potten CS, Anderson E. Dissociation between Steroid Receptor Expression and cell proliferation in the human breast. Cancer Res 1997;57:4987–4991. [PubMed: 9371488]
- (22). Viale G, Regan MM, Maiorano E, et al. Prognostic and predictive value of centrally reviewed expression of estrogen and progesterone receptors in a randomized trial comparing letrozole and tamoxifen adjuvant therapy for postmenopausal early breas cancer: BIG 1-98. J Clin Oncol 2007;25:3846–3852. [PubMed: 17679725]
- (23). Colleoni M, Viale G, Zahrieh D, et al. Expression of ER, PgR, HER1, HER2, and response: a study of preoperative chemotherapy. Annals Oncol 2008;19:465–472.
- (24). Royama, T. Analytic Population Dynamics. Chapman & Hall; London: 1992.
- (25). Speer JF, Petrosky VE, Retsky MW, Hardwell RH. A stochastic numerical model of breast cancer growth that simulates clinical data. Cancer Res 1984;44:4124–4130. [PubMed: 6744323]

Castorina et al.



Fig. 1.

Tumor mass predicted for each clone $(M_1 \text{ and } M_2)$, their sum $(M = M_1 + M_2)$ and the overall tumor according to the Gompertz Law (GL) assuming a delay t_{δ} between the two populations of 100 days (a) and 500 days (b). The other parameters values are: $M^{\infty} = 640$ g, M(0) = 6.4 g, $M_1^{\infty} = M_2^{\infty}$ and $M_2(t) = M_1(t\delta) 10^{-2}$.



Fig. 2.

Tumor mass predicted for each clone $(M_1 \text{ and } M_2)$, their sum $(M = M_1 + M_2)$ and the overall tumor according to the Gompertz Law (GL) assuming a delay t_{δ} between the two populations of 500 days. The ratio between the asymptotic masses are $M^{\infty}_1 = 0.4 M^{\infty}$ (a) and $\infty = 0.2 M^{\infty}$ (b). The other parameters values are: $M^{\infty} = 640 \text{ g}$, M(0) = 6.4 g and M^2 ($t^{\infty} = M_1$ ($t\delta$) 10⁻².



Fig. 3.

Tumor mass predicted for each clone $(M_1 \text{ and } M_2)$, their sum $(M = M_1 + M_2))$ and the overall tumor according to the Gompertz Law (GL). The time delay between the two populations is 500 days (a) and 2000 days (b). The other parameters values are: $M^{\infty} = 640$ g, M(0) = 6.4 g, $M_{21}(t\delta) = M_1(t\delta) 10^{-6}$ and $M_1 = 0.4 M^{\infty}$.





Ratio when the instability starts $at_{\delta} = 60$ days after the beginning of treatment. The parameters are the asymptotic mass, $M^{\infty} = 640$ g, M(0) = 6.4 g, $M_2(t^{\infty} = M_1(t\delta)10^{-1}, M_1^{\infty} M^{\infty}, \omega = 0.1$ per month and $K_g = 0.055$ per month. Different values for the ratio between K_1 and K_2 are depicted.





Ratio $D = \frac{C}{C_g}$ when the instability starts at $t \delta = 60$ days prior to the beginning of treatment. (a) $M_1^{\infty} = 0.1(M^{\infty} - M(t^{\infty})), M_2^{\infty} = 0.5 M_1^{\infty}, M_2 t^{\infty}) 10^{-1}$; (b) $M_1^{\infty} = 0.5 M^{\infty}, M_2 (t^{\infty}) = M_2 t^{\infty})$ 10^{-1} . The other parameters values are: $M^{\infty} = 640$ g, m(0) = 6.4 g, $\omega = 0.1$ per month and K_g = 0.055 per month. Different values for the ratio between K_1 and K_2 are depicted.

Table 1

Comparison with the experimental data for the linear interpolation as discussed in the text, eqns.[49,50]. The experimental error is not included.

Time (vrs)	P (Eqn. (49))	P (Eqn. (50))	Exp. value
1	0.85	0.82	0.87
2	0.70	0.67	0.65
3	0.55	0.54	0.5
4	0.40	0.45	0.45
5	0.25	0.38	0.45