

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**RhoA activity is required for fibronectin assembly and counteracts beta1B integrin inhibitory effect in FRT epithelial cells.**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/43378> since

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

## RhoA activity is required for fibronectin assembly and counteracts $\beta$ 1B integrin inhibitory effect in FRT epithelial cells

Gaetano Cali<sup>1</sup>, Claudia Mazzarella<sup>1</sup>, Marianna Chiacchio<sup>1</sup>, Rosa Negri<sup>1</sup>, Saverio Francesco Retta<sup>2</sup>, Mariastella Zannini<sup>1</sup>, Flaviana Gentile<sup>1</sup>, Guido Tarone<sup>3</sup>, Lucio Nitsch<sup>1</sup> and Corrado Garbi<sup>1,\*</sup>

<sup>1</sup>Centro di Endocrinologia ed Oncologia Sperimentale del CNR – Dipartimento di Biologia e Patologia Cellulare e Molecolare, Università degli Studi di Napoli 'Federico II', Napoli, Italy

<sup>2</sup>Istituto di Biologia, Facoltà di Medicina e Chirurgia, Università degli Studi di Palermo, Palermo, Italy

<sup>3</sup>Dipartimento di Genetica, Biologia e Chimica Medica, Università degli Studi di Torino, Torino, Italy

\*Author for correspondence (e-mail: garbi@unina.it)

Accepted 4 January; published on WWW 25 February 1999

### SUMMARY

FRT thyroid epithelial cells synthesize fibronectin and organize a network of fibronectin fibrils at the basal surface of the cells. Fibronectin fibril formation is enhanced by the overexpression of the ubiquitous  $\beta$ 1A integrin and is inhibited by the expression of the dominant-negative  $\beta$ 1B subunit. We tested the hypotheses that RhoA activity might mediate the integrin-dependent fibronectin fibrillogenesis and might counteract  $\beta$ 1B integrin inhibitory effect. FRT- $\beta$ 1A cells were transfected with a vector carrying a dominant negative form of RhoA (RhoAN19) or treated with the C3 transferase exoenzyme. Both treatments inhibited fibronectin assembly and caused loss of actin microfilaments and adhesion plaques. On the other hand, FRT- $\beta$ 1B cells

were transfected with the constitutively activated form of RhoA (RhoAV14) or treated with the *E. coli* cytotoxic necrotizing factor 1, which directly activates RhoA. Either treatment restored microfilament and adhesion plaque assembly and promoted fibronectin fibril organization. A great increase in fibronectin fibril assembly was also obtained by treatment of FRT- $\beta$ 1B cells with TGF- $\beta$ .

Our data indicate that RhoA is required to promote fibronectin matrix assembly in FRT cells and that the activation of the signal transduction pathway downstream of RhoA can overcome the inhibitory effect of  $\beta$ 1B integrin.

Key words: Integrin, Fibronectin, Epithelial cell, RhoA, TGF- $\beta$

### INTRODUCTION

Fibronectin (FN) matrix assembly is a dynamic process, in which soluble dimeric FN molecules are assembled into an insoluble, disulfide-bond stabilized, fibrillar matrix (Aguirre et al., 1994; Chernousov et al., 1991; Ichihara-Tanaka et al., 1995; Morla and Ruoslahti, 1992; Mosher et al., 1991, 1992; Schwarzbauer, 1991; Sottile and Wiley, 1994). Several insights into the mechanism of formation of FN fibrils have recently been provided. It has been found that the expression on the plasma membrane of specific FN binding integrins (Dzamba et al., 1994; Wu et al., 1993) is required. Among those,  $\alpha$ 5 $\beta$ 1,  $\alpha$ v $\beta$ 3 and  $\alpha$ IIb $\beta$ 3 integrins have been identified (Giancotti and Ruoslahti, 1990; Wennerberg et al., 1996; Wu et al., 1993, 1995). The activation state of integrins, i.e. the acquisition of a high affinity state for FN, is needed for fibrillogenesis in CHO cells (Wu et al., 1995). However, post-occupancy events following integrin activation and involving the  $\beta$  subunit cytoplasmic domain and an intact cytoskeleton are also required (Faull et al., 1993; Wu et al., 1995). Newly assembled FN fibrils align with bundles of actin filaments and with focal adhesions (Heggeness et al., 1978; Hynes and Destree, 1978; Singer and Paradiso, 1981; Wu et al., 1995), suggesting that actin stress fiber, adhesion plaque and FN fibril formation

are coordinately regulated (Ali and Hynes, 1977; Christopher et al., 1997; Sechler and Schwarzbauer, 1997).

Upon interaction with the substrate the integrins cluster and associate with a variety of cytoplasmic proteins to form focal complexes. Integrin-mediated adhesion induces cytoskeletal organization, leading to actin stress fiber formation (Burridge et al., 1988). Binding of integrins to their extracellular matrix also results in the transduction of tyrosine kinase-mediated signals (Clark and Brugge, 1995; Parsons, 1996; Schwartz et al., 1995). Several observations indicate the requirement of the small GTP binding protein RhoA in integrin signaling (Defilippi et al., 1997; Parsons, 1996; Schwartz et al., 1996). It is known that RhoA activation is required for the assembly of microfilament bundles and of adhesion plaques (Ridley and Hall, 1992), such as that induced by growth factors and by LPA (Ridley, 1994). However, the formation of stress fibers and focal adhesions can be obtained in the absence of added growth factors by plating cells on extracellular matrix proteins or by addition of the GRGDS peptide, which is recognized by several integrins (Barry et al., 1997). It has been demonstrated that microinjecting the cells with the RhoA inhibitor C3 transferase exoenzyme can block these effects (Barry et al., 1997), indicating that integrin-mediated actin cytoskeleton assembly may require RhoA activation. LPA has also been shown to

promote FN matrix assembly (Zhang et al., 1994). This effect appears to be correlated to actin stress fiber formation, cell contraction and increased FN binding (Zhang et al., 1997).

We recently demonstrated that  $\beta 1$  integrins are involved in the control of FN matrix assembly in FRT thyroid epithelial cells (Cali et al., 1998). In subconfluent cultures the FN matrix is almost exclusively organized at the basal cell surface. Stable expression of the human  $\beta 1A$  integrin in FRT cells (FRT- $\beta 1A$  cells) causes an increase in FN assembly that is paralleled by an increase of microfilaments and adhesion plaques formation. Expression of the  $\beta 1B$  (Altruda et al., 1990) dominant-negative variant (FRT- $\beta 1B$  cells) determines their inability to assemble FN fibers with concomitant loss of actin filaments and adhesion plaques (Cali et al., 1998).

The aim of this study was to investigate the role of RhoA in promoting FN matrix organization. We inhibited RhoA function in FRT- $\beta 1A$  cells, either by expressing a dominant-negative form of RhoA (RhoAN19) or by treating cells with the C3-transferase exoenzyme (Chardin et al., 1989; Nemoto et al., 1991), and we observed a dramatic reduction of FN assembly. The ability to organize FN fibrils was instead restored in FRT- $\beta 1B$  cells by transfecting a constitutively activated form of RhoA (RhoAV14), or by treatment with the cytotoxic necrotizing factor (CNF1) (Fiorentini et al., 1997). We further demonstrated that treatment with TGF- $\beta$  counteracted the dominant-negative effect of the  $\beta 1B$  integrin subunit.

These results indicate that RhoA has a role in the integrin-mediated FN assembly in epithelial cells and suggest a possible mechanism by which  $\beta 1B$  interferes with this process.

## MATERIALS AND METHODS

### Antibodies and reagents

Rabbit antiserum to rat FN was from Chemicon International (Temecula, CA); anti-paxillin mAb was from Zimed Lab. Inc. (San Francisco, CA); anti-phosphotyrosine (PT 66) mAb was from Sigma Chemical Co. (St Louis, MO); anti-c-myc (9E10) and anti-RhoA mAbs were from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA); mAb against  $\alpha 5\beta 1$  (PB1) was a gift of R. L. Juliano (University of North Carolina-Chapel Hill, Chapel Hill, NC; Brown and Juliano, 1985); horse radish peroxidase-conjugated anti-mouse antibody was from Amersham (Amersham, Buckinghamshire, UK); fluoresceinated avidin was from Oncor (Galthensburg, MD). CNF1 (cytotoxic necrotizing factor 1) was a kind gift of C. Fiorentini (Istituto Superiore di Sanità, Roma, Italy). It was purified from *E. coli* BM2 strain as previously described (Falzano et al., 1993). TGF- $\beta$ , type 1, was from R & D System Inc., (Minneapolis, USA). C3 ADP-ribosyltransferase was from Sigma. It was dissolved at a concentration of 100  $\mu\text{g/ml}$  in 120 mM KCl, 10 mM Tris/HCl, pH 7.4.

### Cell cultures

FRT cells and stable clones of FRT- $\beta 1A$  and FRT- $\beta 1B$  cells were cultured in Falcon tissue culture plastic dishes (Becton Dickinson Labware, Lincoln Park, NJ) in Coon's modified Ham's F12 medium (Sigma) containing 5% FBS (Gibco, Paisley, UK), penicillin, streptomycin and 500  $\mu\text{g/ml}$  of G418 (Sigma). For cell transfections and for all immunofluorescence assays  $1.5 \times 10^5$  cells were plated onto 12 mm diameter glass coverslips.

### Constructs and transfections

pEXVmyctag V14Rho was kindly provided by A. Hall (MRC,

University College of London); RhoAN19 was a gift of R. A. Cerione (Cornell University, Ithaca, New York).

The pEFmyctag RhoAN19 was generated by PCR amplification of the RhoAN19 coding region from the original plasmid (provided by A. Cerione) with a 5' primer and a 3' primer containing at their ends the *Bam*HI and *Eco*RI sites, respectively. The fragment was then subcloned in the corresponding sites of the expression vector EFpLink that was previously modified with the addition of a myc tag by R. Treisman. The construct was sequenced to control the fidelity of the frame between the myc tag and the inserted fragment. The DNA of all the plasmids was prepared by Qiagen cartridges (QIAGEN GmbH, Germany) and used for cell transfections.

FRT, FRT- $\beta 1A$  or FRT- $\beta 1B$  cells were transfected directly on 12 mm diameter glass coverslips with 0.5–2  $\mu\text{g}$  of pEXVmyctag V14Rho or pEXVmyctag RhoAN19 plasmid and 1.25–5  $\mu\text{g}$  of Lipofectin (Gibco BRL Life Technologies Paisley, UK). Transfections were performed as suggested by the factory with minor modifications. Briefly, cells were seeded onto 12 mm glass coverslips in a 24-well cell culture cluster (Costar, Cambridge, MA) in regular medium and incubated overnight. The culture medium was replaced with OPTIMEM 1 (Gibco BRL) 3 hours before transfection. The DNA/liposomes ratio was as suggested (1:2.5) and the final volume on the coverslips was 200  $\mu\text{l}$ .

### Immunofluorescence

Subconfluent cells on glass coverslips were fixed for 20 minutes with 4% paraformaldehyde (Sigma) in PBS containing 0.9 mM calcium and 0.5 mM magnesium (PBS CM) at room temperature, washed twice in 50 mM  $\text{NH}_4\text{Cl}$  in PBS CM and twice in PBS CM. Cells were permeabilized for 5 minutes in 0.5% Triton-X 100 (Bio-Rad) in PBS CM and washed twice, for 10 minutes, in 0.2% gelatin (Sigma) in PBS CM.

Cells were then incubated for 1 hour with the primary antibodies diluted in 0.5% BSA (Sigma) in PBS. After three washes with 0.2% gelatin in PBS CM cells were incubated for 20 minutes with the appropriate rhodamine- or fluorescein-tagged goat anti-mouse or anti-rabbit secondary antibody (Jackson ImmunoResearch, West Grove, PA), diluted 1:50 in 0.5% BSA in PBS. To visualize actin filaments, permeabilized cells were incubated with a 1:70 dilution of rhodamine-conjugated phalloidin (Sigma) for 20 minutes. After final washes with PBS, the coverslips were mounted on a microscope slide using a 50% solution of glycerol in PBS and examined with a Zeiss Axiophot microscope.

Samples were observed by three investigators independently, without knowledge of the experimental conditions. At least 100 fluorescence-positive and negative cells were counted for each experimental condition and representative fields were photographed using Ektachrome P1600 colour reversal film. Images were also acquired with a CCD camera (Sensys, Photometrics, Tucson, AZ). In either case the exposure length was fixed using the control as the index.

### Western blot

FRT and FRT- $\beta 1B$  cells were seeded at low confluence onto 100 mm diameter plastic dishes (Falcon). After 1 day the cells were washed three times, incubated overnight in medium without serum and then treated for 3 hours with 0.2 ng/ml CNF1. Cells were washed twice with PBS and lysed for 20 minutes in ice-cold lysis buffer (150 mM NaCl, 50 mM Tris-HCl, pH 8, 5 mM EDTA, pH 8, 10 mM NaF, 10 mM  $\text{Na}_2\text{P}_2\text{O}_7$ , 0.4 mM  $\text{Na}_3\text{VO}_4$ , 1% Nonidet NP-40) containing protease inhibitors. Cells were then scraped and centrifuged for 10 minutes at 4°C at 14,000 rpm (21,000 g). The pellets were discarded and protein concentration was determined using the Bio-rad protein assay (Bio-rad). 40  $\mu\text{g}$  of protein were solubilized in Laemmli sample buffer, boiled for 5 minutes, and analyzed by SDS/6–15%PAGE.

Gels were blotted onto nitrocellulose filters (Amersham) in a Bio-Rad apparatus (Bio-Rad). The filters were washed extensively with TTBS (150 mM NaCl, 20 mM Tris-HCl, pH 7.5, 0.05% Tween 20), blocked at room temperature for 2 hours with 1% non-fat dry milk

(NFDm; Bio-Rad) in TBS (150 mM NaCl, 20 mM Tris-HCl, pH 7.5), washed twice with TTBS and once with TBS, and incubated for 1 hour at room temperature with the anti-RhoA mAb (Santa Cruz) diluted 1:1000 in TBS containing 0.5% NFDm. Subsequently, the filters were washed extensively with TTBS and with TBS and incubated for 1 hour at room temperature with horse radish peroxidase-conjugated anti-mouse antibody (Amersham) diluted 1:1000 in 0.5% TBS. After extensive washing with TTBS and with TBS, the filters were developed using an ECL detection method (Amersham), according to the manufacturer's directions.

### Biotinylation of FN

80 µg (80 µl of 1 mg/ml) bovine serum FN (Sigma) was allowed to react with 400 µg (2 µl) of NHS-LC biotin (Pierce, Rockford, IL) in 420 µl PBS CM, for 20 minutes, at 4°C. The solution was dialyzed against PBS CM for 24 hours at 4°C.  $1.5 \times 10^5$  FRT-β1A cells were seeded onto 12 mm diameter glass coverslips and treated with 4 µg (125 µl) of biotinylated FN. Immunofluorescence was performed as described. Biotinylated FN was detected with fluorescein-tagged avidin (Oncor, Galthensburg, MD).

## RESULTS

### FRT-β1A cells organize an extracellular matrix containing both the cell-secreted FN and the serum FN

Subconfluent FRT cells cultured in the presence of 5% foetal calf serum were shown to assemble FN fibrils at the basal surface of the cells (Cali et al., 1998). In most of the experiments shown in this paper we have used two FRT-derived cell lines, FRT-β1A and FRT-β1B, which express the human β1A and β1B integrin subunit, respectively (Cali et al., 1998). FRT-β1A cells were able to deposit a much more abundant FN matrix (Fig. 1a) compared to wild-type FRT cells. In serum-free medium, a reduced amount of extracellular FN was still assembled (Fig. 1b), indicating that the cells can organize into fibrils the FN molecules that they synthesize and secrete. FRT-β1B cells, on the contrary, did not assemble FN fibrils in significant amount either in the presence or in the absence of serum (Fig. 1c-d). To demonstrate that FRT-β1A cells were indeed able to organize the FN present in the medium into fibrils, subconfluent cultures were grown in serum-free conditions in the presence of soluble biotinylated FN. The FN matrix deposited was detected by an anti-rat FN antibody (Fig. 1e) while the presence of biotinylated FN in the insoluble matrix was revealed by rhodamine-conjugated streptavidin (Fig. 1f). FN fibrils were labeled with fluorescent markers (Fig. 1e-f), indicating that newly synthesized as well as exogenous FN molecules participate in the assembly of the same extracellular FN matrix.

### α5β1 integrin is involved in the process of FN assembly

The involvement of the α5β1 integrin dimer in the fibrillogenesis process has been tested by interfering with the endogenous α5β1 activity. This was achieved by including in the culture medium the blocking antibody PB1 (Brown and Juliano, 1985; Wu et al., 1995) that is directed against the α5β1 FN receptor. FRT-β1A cells were cultured for 18 hours in the presence of PB1 and then analyzed by immunofluorescence for the presence of assembled FN. A

great reduction in the amount of cell-associated fibrils was seen in PB1-treated cells (Fig. 1g) as compared to control cells cultured in regular medium (Fig. 1h) or in the presence of non-specific mAbs (data not shown). This result indicates that α5β1 is the main integrin involved in the process of FN fibril organization in FRT cells.

### The expression of RhoAN19 inhibits FN matrix assembly


We have previously reported that in FRT-β1A cells the FN fibrillar network is paralleled by the organization of a well developed actin cytoskeleton and by the formation of a great number of adhesion plaques (Cali et al., 1998). To assess the role of RhoA in the process of extracellular FN deposition we investigated the effect of the expression of the dominant-negative form of RhoA, RhoAN19, in FRT-β1A cells. Cells were transiently transfected with an expression vector carrying the dominant-negative form of RhoA fused to a myc tag. After 24 hours cells were fixed and double-stained for immunofluorescence detection with the anti-myc antibody (Fig. 2a,c,e) and with rhodamine-conjugated phalloidin (Fig. 2b), anti-paxillin antibody (Fig. 2d) or anti-FN antibody (Fig. 2f). A significant reduction in microfilament and adhesion plaque assembly, and FN matrix organization, was found. It was observed that  $73 \pm 13\%$  of RhoAN19-expressing cells had reduced amounts of actin filaments and of FN matrix with respect to  $29 \pm 9\%$  of mock transfected cells. These data indicate that RhoA activity is required not only to assemble microfilaments and adhesion plaques but also to organize FN fibrils.

### Inactivation of endogenous RhoA by C3 transferase exoenzyme inhibits FN fibril formation

C3 ADP-ribosyltransferase of *C. botulinum* is a useful tool for investigating Rho-protein-mediated cellular responses because it specifically inactivates Rho protein (Aktories et al., 1992; Quilliam et al., 1989; Sekine et al., 1989). Although in other experimental systems the C3 transferase exoenzyme has been shown to penetrate the cells with some difficulty, this was not the case with FRT cells, which appeared to be quite sensitive to the action of the toxin. 1.5 µg/ml C3 transferase was added to FRT-β1A cells that had been cultured for 48 hours in standard conditions. After 8 or 18 hours cells were fixed and double-stained with rhodamine conjugated-phalloidin and with the anti-FN antibody (Fig. 3). After 8 hours of C3 transferase we observed that some cells showed a loss of microfilaments and a reduction in FN fibrils. These cells were frequently in small clusters (Fig. 3a-b) that were easily detected within the culture. By 18 hours of treatment most of the cells were found to be without microfilaments (Fig. 3c) and only residual deposits of FN, mostly in the form of irregular cables, were present (Fig. 3d). A similar result was obtained when wild-type FRT cells were treated with C3 transferase (data not shown). It appears that C3 inhibition of Rho activity leads to progressive disruption of the actin cytoskeleton, inhibition of organization of FN fibrils, and also removal of previously organized FN matrix.

### The expression of RhoAV14 in FRT-β1B cells restores the ability to assemble the FN matrix

FRT-β1B cells do not organize actin filaments and adhesion



**Fig. 1.** Analysis of FN deposition by FRT- $\beta$ 1A and FRT- $\beta$ 1B cells in different culture conditions. Subconfluent FRT- $\beta$ 1A cells (a,b) and FRT- $\beta$ 1B cells (c,d) were cultured for 18 hours in 5% FBS (a,c) or 0% FBS (b,d). FN was detected by immunofluorescence with an antiserum to rat FN. (e,f) FRT- $\beta$ 1A cells were cultured for 18 hours in the presence of 4  $\mu$ g biotinylated FN. FN fibrils were detected with an antiserum to rat FN (e) and biotinylated FN was detected with fluorescein-tagged avidin (f). (g,h) FRT- $\beta$ 1A cells were cultured for 18 hours in complete medium with (g) or without (h) the addition of the  $\alpha$ 5 $\beta$ 1 FN receptor blocking antibody PB1 (1:100 dilution). The experiments were performed at least three times. Bar, 25  $\mu$ m.


plaques, and assemble very few FN fibrils with respect to wild-type FRT cells (Cali et al., 1998). We speculated that FRT- $\beta$ 1B cells might be defective in one or more integrin-mediated signaling event(s) that might activate RhoA, and that lack of RhoA activity could be responsible for FRT- $\beta$ 1B cell phenotype. To prove this, FRT- $\beta$ 1B cells were transiently transfected with a vector carrying the constitutively activated form of RhoA, RhoAV14, fused to a myc tag. After 24 hours cells were fixed and double-stained for immunofluorescence detection with the anti-myc antibody (Fig. 4a,c,e) and with rhodamine-conjugated phalloidin (Fig. 4b), anti-paxillin antibody (Fig. 4d) or anti-FN antibody (Fig. 4f). The ability to assemble actin microfilaments (Fig. 4b), adhesion plaques (Fig. 4d) and FN fibrils (Fig. 4f) was rescued in a high number of transfected cells. The cells expressing RhoAV14 that organized FN fibrils and actin filaments were  $83 \pm 7\%$ , and  $24 \pm 10\%$  in mock transfected cells. Some of the myc-positive cells,

although expressing microfilaments, had a round shape and lacked FN fibrils.

#### Treatment of FRT- $\beta$ 1B cells with CNF1 activates RhoA and induces the organization of fibrillar FN

In order to increase the amount of active RhoA in FRT cells expressing the  $\beta$ 1B integrin, we treated the cells with CNF1, a toxin produced by pathogenic strains of *E. coli* that has been shown to directly bind and activate Rho (Fiorentini et al., 1997; Flatau et al., 1997; Schmidt et al., 1997). CNF1 has been shown to induce an increase in the content of actin stress fibers and focal contacts in cultured cells (Lacerda et al., 1997; Oswald et al., 1994). The activation of RhoA by CNF1 is correlated with a decrease in its electrophoretic mobility (Oswald et al., 1994). To determine if CNF1 was indeed causing a shift in the electrophoretic mobility of RhoA protein, FRT and FRT- $\beta$ 1B cells were treated with CNF1 and cell extracts were analyzed






**Fig. 2.** Effect of dominant-negative RhoAN19 on FN assembly in FRT- $\beta$ 1A cells. FRT- $\beta$ 1A cells grown on glass coverslips were transfected with 2  $\mu$ g pEXVmyctag RhoAN19 (see Materials and methods). Cells were stained and detected by immunofluorescence using the anti-myc antibody diluted 1:50 (a,c,e) to detect RhoAN19 expression and double stained with rhodamine-conjugated phalloidin (b), with anti-paxillin antibody diluted 1:100 (d) or with anti-FN antibody diluted 1:100 (f). Cells that were stained by the anti-myc antibody and show a reduction in actin microfilaments (a,b), adhesion plaques (c,d) and FN fibrils (e,f) are indicated by arrows. The experiments were performed at least five times. Bar, 10  $\mu$ m.

by western blot using antibodies against RhoA. A reduction in the migratory pattern of RhoA in CNF1-treated cells was observed (Fig. 5). To determine if CNF1 was promoting FN assembly, FRT- $\beta$ 1B cells were grown in standard culture conditions for 48 hours, treated with CNF1 for 18 hours and examined by immunofluorescence using rhodamine-conjugated phalloidin, anti-paxillin antibody or anti-FN antibody. A great increase in actin polymerization (Fig. 6a-b), adhesion plaque organization (Fig. 6c-d) and FN assembly (Fig. 6e-f) was observed. These effects were already


manifested by 6-8 hours of incubation, while changes in cell morphology due to CNF1 toxicity started to be seen after 24 hours. Similar results were observed when wild-type FRT cells were treated with CNF1 (data not shown).

### TGF- $\beta$ promotes the coordinated organization of FN matrix, actin microfilaments and adhesion plaques

TGF- $\beta$  is a multifunctional cytokine capable of inducing a great number of effects, depending on the cell type (Clark et al., 1998). In thyroid cells TGF- $\beta$  controls cell proliferation,



**Fig. 3.** Treatment of FRT- $\beta$ 1A cells with C3 transferase inhibits FN fibril formation. FRT- $\beta$ 1A cells were cultured on glass coverslips for 48 hours and treated with 1.5  $\mu$ g/ml C3 transferase for 8 hours (a,b) or 18 hours (c,d). Cells were double-stained with rhodamine-conjugated phalloidin (a,c) and with the anti-FN antibody (b,d). Lack of microfilaments (a) and a reduction in FN fibrils (b) is observed in localized areas (arrows) at 8 hours. An almost complete loss of microfilaments (c) and of FN fibrils (d) is observed at 18 hours. The experiments were performed at least three times. Bar, 15  $\mu$ m.




**Fig. 4.** Effect of the constitutively active form of RhoA (RhoAV14) on FN fibril assembly in FRT- $\beta$ 1B cells. FRT- $\beta$ 1B cells grown on glass coverslips were transfected with 2  $\mu$ g pEXVmyctag V14Rho (see Materials and methods). Cells were stained for immunofluorescence detection using the anti-myc antibody (a,c,e) to detect RhoAV14 expression and double-stained with rhodamine-conjugated phalloidin (b), with anti-paxillin antibody (d) or with anti-FN antibody (f). Cells that are stained by the anti-myc antibody and show an increase in actin microfilaments (a,b), adhesion plaques (c,d) and FN fibrils (e,f) are indicated by arrows. Note that in all panels the fields are almost entirely filled with cells. The experiments were performed at least five times. Bar, 20  $\mu$ m.

modulates some differentiated functions and acts as a potent inducer of stress fiber formation (Garbi et al., 1990). The exact mechanism by which this molecule exerts its intracellular effects is not known, though a recent report indicates that Rho activity might be required (Atfi et al., 1997). We treated FRT- $\beta$ 1B cells with 5 ng/ml of TGF- $\beta$ . Cells were fixed and stained with rhodamine-conjugated phalloidin, anti-phosphotyrosine or anti-FN antibodies. TGF- $\beta$  treatment induced within 24 hours the formation of a rich FN extracellular matrix paralleled by the formation of a great number of microfilaments and adhesion plaques (Fig. 7). However, at variance to what observed with CNF1, longer periods of incubation were needed to see these effects.

## DISCUSSION

We report in this paper the involvement of RhoA in the process of FN fibrillogenesis in FRT thyroid epithelial cells. These cells are able to synthesize and secrete FN (Canipari et al., 1992)




**Fig. 5.** CNF1 treatment induces a shift in the electrophoretic mobility of RhoA. Subconfluent FRT and FRT- $\beta$ 1B cells were treated for 3 hours with 0.2 ng/ml CNF1. Soluble proteins were separated by SDS/6-15%PAGE and analyzed by western blot using the anti-RhoA mAb diluted 1:1000. Molecular masses are indicated on the left side of the figure.

and to organize it into an insoluble fibrillar matrix with a polarized distribution. Unlike fibroblasts, where FN matrix is also present on the cell free surface (Wennerberg et al., 1996), FRT cells deposit FN only underneath the cells, at cell-substrate contact sites (Cali et al., 1998). We have generated two FRT-derived cell lines that constitutively express the human  $\beta$ 1A or  $\beta$ 1B integrin subunit and have demonstrated that  $\beta$ 1 integrins are involved in FN assembly (Cali et al., 1998). We observed that both the exogenous FN supplied with the serum and the cellular newly synthesized FN can be incorporated into the same fibrils, indicating that they are recognized by the cells with similar efficiency. Thus the different ability to organize FN matrix observed in FRT- $\beta$ 1A and FRT- $\beta$ 1B cells is likely not due to differences in FN synthetic potential.

Using the specific function-blocking antibody PB1 we were able to demonstrate that the  $\alpha$ 5 $\beta$ 1 dimers are essential for the process of FN fibrillogenesis. Since FRT cells are capable of sorting and retaining the  $\beta$ 1 integrin receptors almost exclusively on the basolateral surface (Cali et al., 1998), it is conceivable that polarized distribution of  $\alpha$ 5 $\beta$ 1 is responsible for the polarized assembly of FN. PB1 inhibition also suggests that  $\alpha$ 5 $\beta$ 1 cannot be substituted by other integrins in the FN assembly process, although FRT cells express the  $\alpha$ v $\beta$ 3 integrin that can replace  $\alpha$ 5 $\beta$ 1 in other cell model systems (Wennerberg et al., 1996).

To investigate the role of RhoA in FN assembly we have transiently expressed the dominant-negative RhoAN19 in FRT- $\beta$ 1A cells. A significant reduction in the amount of FN matrix assembly, paralleled by loss of microfilaments and reduction of focal adhesions, was observed in RhoAN19-expressing cells. A similar inhibition of FN assembly was also obtained upon exposure of FRT- $\beta$ 1A cells to the C3 transferase toxin. In quiescent Swiss 3T3 cells C3 transferase treatment also




**Fig. 6.** Treatment of FRT- $\beta$ 1B cells with CNF1 promotes FN fibril assembly. FRT- $\beta$ 1B cells were cultured on glass coverslips for 48 hours and treated with 0.2 ng/ml CNF1 for 8 hours. Untreated cells (a,c,e) and CNF1-treated cells (b,d,f) were stained with rhodamine-conjugated phalloidin (a,b), with anti-paxillin antibody (c,d) and with anti-FN antibody (e,f). An increase in actin filaments (b), adhesion plaques (d) and FN fibrils (f) is evident. The experiments were performed at least three times. Bar, 15  $\mu$ m.

determines a decrease in FN assembly (Zhong et al., 1998). In FRT- $\beta$ 1A cells, however, C3 transferase inhibited the formation of new fibrils and caused the removal of the FN matrix already deposited. Extracellular FN assembly should possibly be regarded as a dynamic process in which there is a

continuous balance between FN fibril formation and degradation. RhoA activity might be a permanent requirement in the interplay between the intracellular actin cytoskeleton and the extracellular FN matrix that is needed in these processes.

A major finding of this report is the demonstration that FN



**Fig. 7.** Treatment of FRT- $\beta$ 1B cells with TGF- $\beta$  induces FN matrix organization. FRT- $\beta$ 1B cells were cultured on glass coverslips for 48 hours and treated with 5 ng/ml TGF- $\beta$  for 24 hours. Untreated cells (a,c,e) and TGF- $\beta$  treated cells (b,d,f) were stained with rhodamine-conjugated phalloidin (a,b), with anti-phosphotyrosine antibody (c,d) and with anti-FN antibody (e,f). A prominent increase in actin filaments (b), adhesion plaques (d) and FN fibrils (f) is observed. The experiments were performed at least three times. Bar, 10  $\mu$ m.



fibrillogenesis can be restored in FRT- $\beta$ 1B cells by transient expression of the constitutively activated RhoA V14 or by treatment with CNF1. In both experiments microfilament assembly and adhesion plaques were concomitantly stimulated. These results indicate that  $\beta$ 1B inhibition is not permanent and that it can be overcome. Moreover they suggest that in FRT- $\beta$ 1B cells the signal transduction pathway downstream of RhoA operates as in wild-type FRT cells and that  $\beta$ 1B expression affects some other step of the integrin signaling pathway. FRT- $\beta$ 1B cells do not organize microfilaments and have a reduced number of very small adhesion plaques. These phenotypic properties correspond to what is observed in fibroblasts where RhoA activity is inhibited by the expression of RhoA N19 (Clark et al., 1998). It is therefore conceivable that FRT- $\beta$ 1B cells have impairment in the process that leads to RhoA activation.

The  $\beta$ 1B human integrin is an alternatively spliced variant of the  $\beta$ 1 subunit, where the cytosolic tail of the molecule is shorter than in  $\beta$ 1A and carries a specific COOH-terminal sequence 12 amino acids long (Altruda et al., 1990; Balzac et al., 1994). It has been recently demonstrated that  $\beta$ 1B expression in GD25  $\beta$ 1 null cells interferes with the  $\alpha$ v $\beta$ 3-dependent FN matrix assembly and that  $\beta$ 1B is present at the cell surface in an inactive conformation (Retta et al., 1998). It has been proposed that  $\beta$ 1B could bind and sequester molecules necessary for integrin signal transduction. A similar mechanism could be acting in FRT cells, although other possibilities, such as association to  $\alpha$  subunits, cannot be excluded. To date no direct interactions between integrins and RhoA have been reported. It has been proposed that in LOX melanoma cells the  $\alpha$ 6 $\beta$ 1 integrin signaling regulates the tyrosine phosphorylation state of p190 GAP and therefore influences the downstream signaling pathway through Rho (Nakahara et al., 1998).

How does the constitutive activation of RhoA promote FN assembly? Since RhoA is able to induce integrin clustering (Hotchin and Hall, 1995; Machesky and Hall, 1997), it is plausible that FN assembly can occur by recruitment of the endogenous  $\alpha$ 5 $\beta$ 1 integrins. An increase in FN assembly also occurs in CNF1-treated wild-type FRT cells. RhoA could also determine the acquisition of a high-affinity state of the  $\beta$ 1B integrin subunit. However, only small variations in the state of integrin activation have been observed as a consequence of RhoA activity changes (Zhong et al., 1998). A critical role in FN fibrillogenesis appears to be played by Rho-mediated cell contractility, which could generate tension in the FN molecules and expose cryptic self-assembly sites (Zhong et al., 1998).

The family of transforming growth factors beta has profound regulatory effects on cell growth and differentiation (Clark et al., 1998; Roberts et al., 1990). TGF- $\beta$  modulates the synthesis and accumulation of ECM components and the expression of cell surface receptors for ECM components (Ignatz and Massague, 1987). In many cell types TGF- $\beta$  promotes actin cytoskeleton organization. We have shown that, in FRTL5 rat thyroid epithelial cells, TGF- $\beta$  induces stress fiber formation and changes in the extracellular matrix organization (Garbi et al., 1990). Although the mechanism of receptor activation is reasonably understood, there is little information about the potential downstream targets of the receptor complex. Recent observations suggest that Rac (Mucsi et al., 1996) or Rho-like GTPases (Atfi et al., 1997) mediate some effects of TGF- $\beta$

signaling. Since TGF- $\beta$  effects on cytoskeleton and ECM mimic Rho activation, we speculated that there might be a possible link between TGF- $\beta$  activity and Rho in the FRT cell model system. Indeed, TGF- $\beta$  promoted assembly of FN and organization of microfilaments in FRT- $\beta$ 1B cells to the same extent as CNF1 induced RhoA activation.

In conclusion, RhoA activity is needed to determine  $\alpha$ 5 $\beta$ 1-dependent FN assembly in FRT epithelial cells.  $\beta$ 1B possibly inhibits FN assembly by interfering with some step of the integrin signal transduction pathway.

This work was partially supported by grants from Progetto Finalizzato Applicazioni Cliniche della Ricerca Oncologica (ACRO) of the CNR, from Associazione Italiana per la Ricerca sul Cancro (AIRC), from MURST-CNR Biotechnology Program L.95/95 and from MURST project 'La trasduzione dei segnali nei processi adesivi, migratori e proliferativi'. We thank Franco D'Agnello and Mario Berardone for the art work.

## REFERENCES

- Aguirre, K. M., McCormick, R. J. and Schwarzbauer, J. E. (1994). Fibronectin self-association is mediated by complementary sites within the amino-terminal one-third of the molecule. *J. Biol. Chem.* **269**, 27863-27868.
- Aktories, K., Mohr, C. and Koch, G. (1992). *Clostridium botulinum* C3 ADP-ribosyltransferase. *Curr. Top. Microbiol. Immunol.* **175**, 115-131.
- Ali, I. U. and Hynes, R. O. (1977). Effects of cytochalasin B and colchicine on attachment of a major surface protein of fibroblasts. *Biochim. Biophys. Acta* **471**, 16-24.
- Altruda, F., Cervella, P., Tarone, G., Botta, C., Balzac, F., Stefanuto, G. and Silengo, L. (1990). A human integrin beta 1 subunit with a unique cytoplasmic domain generated by alternative mRNA processing. *Gene* **95**, 261-266.
- Atfi, A., Djelloul, S., Chastre, E., Davis, R. and Gaspach, C. (1997). Evidence for a role of Rho-like GTPases and stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) in transforming growth factor beta-mediated signaling. *J. Biol. Chem.* **272**, 1429-1432.
- Balzac, F., Retta, S. F., Albin, A., Melchiorri, A., Kotliansky, V. E., Geuna, M., Silengo, L. and Tarone, G. (1994). Expression of beta 1B integrin isoform in CHO cells results in a dominant negative effect on cell adhesion and motility. *J. Cell Biol.* **127**, 557-565.
- Barry, S. T., Flinn, H. M., Humphries, M. J., Critchley, D. R. and Ridley, A. J. (1997). Requirement for Rho in integrin signalling. *Cell Adhes. Commun.* **4**, 387-398.
- Brown, P. J. and Juliano, R. L. (1985). Selective inhibition of fibronectin-mediated cell adhesion by monoclonal antibodies to a cell-surface glycoprotein. *Science* **228**, 1448-1451.
- Burridge, K., Fath, K., Kelly, T., Nuckolls, G. and Turner, C. (1988). Focal adhesions: transmembrane junctions between the extracellular matrix and the cytoskeleton. *Annu. Rev. Cell Biol.* **4**, 487-525.
- Cali, G., Retta, S. F., Negri, R., Damiano, I., Gentile, R., Tarone, G., Nitsch, L. and Garbi, C. (1998). Beta1B integrin interferes with matrix assembly but not with confluent monolayer polarity and alters some morphogenetic properties of FRT epithelial cells. *Eur. J. Cell Biol.* **75**, 107-117.
- Canipari, R., Zurzolo, C., Polistina, C., Garbi, C., Aloj, L., Cali, G., Gentile, R. and Nitsch, L. (1992). Polarized secretion of plasminogen activators by epithelial cell monolayers. *Biochim. Biophys. Acta* **1175**, 1-6.
- Chardin, P., Boquet, P., Madaule, P., Popoff, M. R., Rubin, E. J. and Gill, D. M. (1989). The mammalian G protein rhoC is ADP-ribosylated by *Clostridium botulinum* exoenzyme C3 and affects actin microfilaments in Vero cells. *EMBO J.* **8**, 1087-1092.
- Chernousov, M. A., Fogerty, F. J., Kotliansky, V. E. and Mosher, D. F. (1991). Role of the I-9 and III-1 modules of fibronectin in formation of an extracellular fibronectin matrix. *J. Biol. Chem.* **266**, 10851-10858.
- Christopher, R. A., Kowalczyk, A. P. and McKeown-Longo, P. J. (1997). Localization of fibronectin matrix assembly sites on fibroblasts and endothelial cells. *J. Cell Sci.* **110**, 569-581.
- Clark, E. A. and Brugge, J. S. (1995). Integrins and signal transduction pathways: the road taken. *Science* **268**, 233-239.

- Clark, E. A., King, W. G., Brugge, J. S., Symons, M. and Hynes, R. O. (1998). Integrin-mediated signals regulated by members of the rho family of GTPases. *J. Cell Biol.* **142**, 573-586.
- Defilippi, P., Venturino, M., Gulino, D., Duperray, A., Boquet, P., Fiorentini, C., Volpe, G., Palmieri, M., Silengo, L. and Tarone, G. (1997). Dissection of pathways implicated in integrin-mediated actin cytoskeleton assembly. Involvement of protein kinase C, Rho GTPase and tyrosine phosphorylation. *J. Biol. Chem.* **272**, 21726-21734.
- Dzamba, B. J., Bultmann, H., Akiyama, S. K. and Peters, D. M. (1994). Substrate-specific binding of the amino terminus of fibronectin to an integrin complex in focal adhesions. *J. Biol. Chem.* **269**, 19646-19652.
- Falzano, L., Fiorentini, C., Donelli, G., Michel, E., Kocks, C., Cossart, P., Cabanie, L., Oswald, E. and Boquet, P. (1993). Induction of phagocytic behaviour in human epithelial cells by *Escherichia coli* cytotoxic necrotizing factor type 1. *Mol. Microbiol.* **9**, 1247-1254.
- Faull, R. J., Kovach, N. L., Harlan, J. M. and Ginsberg, M. H. (1993). Affinity modulation of integrin alpha 5 beta 1: regulation of the functional response by soluble fibronectin. *J. Cell Biol.* **121**, 155-162.
- Fiorentini, C., Fabbri, A., Flatau, G., Donelli, G., Matarrese, P., Lemichez, E., Falzano, L. and Boquet, P. (1997). *Escherichia coli* cytotoxic necrotizing factor 1 (CNF1), a toxin that activates the Rho GTPase. *J. Biol. Chem.* **272**, 19532-19537.
- Flatau, G., Lemichez, E., Gauthier, M., Chardin, P., Paris, S., Fiorentini, C. and Boquet, P. (1997). Toxin-induced activation of the G protein p21 Rho by deamidation of glutamine. *Nature* **387**, 729-733.
- Garbi, C., Colletta, G., Cirafici, A. M., Marchisio, P. C. and Nitsch, L. (1990). Transforming growth factor-beta induces cytoskeleton and extracellular matrix modifications in FRTL-5 thyroid epithelial cells. *Eur. J. Cell Biol.* **53**, 281-289.
- Giancotti, F. G. and Ruoslahti, E. (1990). Elevated levels of the alpha 5 beta 1 fibronectin receptor suppress the transformed phenotype of Chinese hamster ovary cells. *Cell* **60**, 849-859.
- Heggeness, M. H., Ash, J. F. and Singer, S. J. (1978). Transmembrane linkage of fibronectin to intracellular actin-containing filaments in cultured human fibroblasts. *Ann. NY Acad. Sci.* **312**, 414-417.
- Hotchin, N. A. and Hall, A. (1995). The assembly of integrin adhesion complexes requires both extracellular matrix and intracellular rho/rac GTPases. *J. Cell Biol.* **131**, 1857-1865.
- Hynes, R. O. and Destree, A. T. (1978). Relationships between fibronectin (LETS protein) and actin. *Cell* **15**, 875-86.
- Ichihara-Tanaka, K., Titani, K. and Sekiguchi, K. (1995). Role of the carboxyl-terminal Fib2 domain in fibronectin matrix assembly. *J. Cell Sci.* **108**, 907-915.
- Ignatz, R. A. and Massague, J. (1987). Cell adhesion protein receptors as targets for transforming growth factor-beta action. *Cell* **51**, 189-197.
- Lacerda, H. M., Pullinger, G. D., Lax, A. J. and Rozengurt, E. (1997). Cytotoxic necrotizing factor 1 from *Escherichia coli* and dermonecrotic toxin from *Bordetella bronchiseptica* induce p21(rho)-dependent tyrosine phosphorylation of focal adhesion kinase and paxillin in Swiss 3T3 cells. *J. Biol. Chem.* **272**, 9587-9596.
- Machesky, L. M. and Hall, A. (1997). Role of actin polymerization and adhesion to extracellular matrix in Rac- and Rho-induced cytoskeletal reorganization. *J. Cell Biol.* **138**, 913-926.
- Morla, A. and Ruoslahti, E. (1992). A fibronectin self-assembly site involved in fibronectin matrix assembly: reconstruction in a synthetic peptide. *J. Cell Biol.* **118**, 421-429.
- Mosher, D. F., Fogerty, F. J., Chernousov, M. A. and Barry, E. L. (1991). Assembly of fibronectin into extracellular matrix. *Ann. NY Acad. Sci.* **614**, 167-180.
- Mosher, D. F., Sottile, J., Wu, C. and McDonald, J. A. (1992). Assembly of extracellular matrix. *Curr. Opin. Cell Biol.* **4**, 810-818.
- Mucsi, I., Skorecki, K. L. and Goldberg, H. J. (1996). Extracellular signal-regulated kinase and the small GTP-binding protein, Rac, contribute to the effects of transforming growth factor-beta1 on gene expression. *J. Biol. Chem.* **271**, 16567-16572.
- Nakahara, H., Mueller, S. C., Nomizu, M., Yamada, Y., Yeh, Y. and Chen, W. T. (1998). Activation of beta1 integrin signaling stimulates tyrosine phosphorylation of p190RhoGAP and membrane-protrusive activities at invadopodia. *J. Biol. Chem.* **273**, 9-12.
- Nemoto, Y., Namba, T., Kozaki, S. and Narumiya, S. (1991). Clostridium botulinum C3 ADP-ribosyltransferase gene. Cloning, sequencing and expression of a functional protein in *Escherichia coli*. *J. Biol. Chem.* **266**, 19312-19319.
- Oswald, E., Sugai, M., Labigne, A., Wu, H. C., Fiorentini, C., Boquet, P. and O'Brien, A. D. (1994). Cytotoxic necrotizing factor type 2 produced by virulent *Escherichia coli* modifies the small GTP-binding proteins Rho involved in assembly of actin stress fibers. *Proc. Natl Acad. Sci. USA* **91**, 3814-3818.
- Parsons, J. T. (1996). Integrin-mediated signalling: regulation by protein tyrosine kinases and small GTP-binding proteins. *Curr. Opin. Cell Biol.* **8**, 146-152.
- Quilliam, L. A., Lacal, J. C. and Bokoch, G. M. (1989). Identification of rho as a substrate for botulinum toxin C3-catalyzed ADP-ribosylation. *FEBS Lett.* **247**, 221-226.
- Retta, S. F., Balzac, F., Ferraris, P., Belkin, A. M., Fassler, R., Humphries, M. J., De Leo, G., Silengo, L. and Tarone, G. (1998). beta1-integrin cytoplasmic subdomains involved in dominant negative function. *Mol. Biol. Cell* **9**, 715-731.
- Ridley, A. J. (1994). Signal transduction through the GTP-binding proteins Rac and Rho. *J. Cell Sci. Suppl.* **18**, 127-131.
- Ridley, A. J. and Hall, A. (1992). The small GTP-binding protein rho regulates the assembly of focal adhesions and actin stress fibers in response to growth factors. *Cell* **70**, 389-399.
- Roberts, A. B., Flanders, K. C., Heine, U. I., Jakowlew, S., Kondaiah, P., Kim, S. J. and Sporn, M. B. (1990). Transforming growth factor-beta: multifunctional regulator of differentiation and development. *Phil. Trans. R. Soc. Lond. B* **327**, 145-154.
- Schmidt, G., Sehr, P., Wilm, M., Selzer, J., Mann, M. and Aktories, K. (1997). Gln 63 of Rho is deamidated by *Escherichia coli* cytotoxic necrotizing factor-1. *Nature* **387**, 725-729.
- Schwartz, M. A., Schaller, M. D. and Ginsberg, M. H. (1995). Integrins: emerging paradigms of signal transduction. *Annu. Rev. Cell. Dev. Biol.* **11**, 549-599.
- Schwartz, M. A., Toksoz, D. and Khosravi-Far, R. (1996). Transformation by Rho exchange factor oncogenes is mediated by activation of an integrin-dependent pathway. *EMBO J.* **15**, 6525-6530.
- Schwarzbauer, J. E. (1991). Identification of the fibronectin sequences required for assembly of a fibrillar matrix. *J. Cell Biol.* **113**, 1463-1473.
- Sechler, J. L. and Schwarzbauer, J. E. (1997). Coordinated regulation of fibronectin fibril assembly and actin stress fiber formation. *Cell Adhes. Commun.* **4**, 413-424.
- Sekine, A., Fujiwara, M. and Narumiya, S. (1989). Asparagine residue in the rho gene product is the modification site for botulinum ADP-ribosyltransferase. *J. Biol. Chem.* **264**, 8602-8605.
- Singer, II and Paradiso, P. R. (1981). A transmembrane relationship between fibronectin and vinculin (130 kd protein): serum modulation in normal and transformed hamster fibroblasts. *Cell* **24**, 481-492.
- Sottile, J. and Wiley, S. (1994). Assembly of amino-terminal fibronectin dimers into the extracellular matrix. *J. Biol. Chem.* **269**, 17192-17198.
- Wennerberg, K., Lohikangas, L., Gullberg, D., Pfaff, M., Johansson, S. and Fassler, R. (1996). Beta 1 integrin-dependent and -independent polymerization of fibronectin. *J. Cell Biol.* **132**, 227-238.
- Wu, C., Bauer, J. S., Juliano, R. L. and McDonald, J. A. (1993). The alpha 5 beta 1 integrin fibronectin receptor, but not the alpha 5 cytoplasmic domain, functions in an early and essential step in fibronectin matrix assembly. *J. Biol. Chem.* **268**, 21883-21888.
- Wu, C., Keivens, V. M., O'Toole, T. E., McDonald, J. A. and Ginsberg, M. H. (1995). Integrin activation and cytoskeletal interaction are essential for the assembly of a fibronectin matrix. *Cell* **83**, 715-724.
- Zhang, Q., Checovich, W. J., Peters, D. M., Albrecht, R. M. and Mosher, D. F. (1994). Modulation of cell surface fibronectin assembly sites by lysophosphatidic acid. *J. Cell Biol.* **127**, 1447-1459.
- Zhang, Q., Magnusson, M. K. and Mosher, D. F. (1997). Lysophosphatidic acid and microtubule-destabilizing agents stimulate fibronectin matrix assembly through Rho-dependent actin stress fiber formation and cell contraction. *Mol. Biol. Cell* **8**, 1415-1425.
- Zhong, C., Chrzanowska-Wodnicka, M., Brown, J., Shaub, A., Belkin, A. M. and Burridge, K. (1998). Rho-mediated contractility exposes a cryptic site in fibronectin and induces fibronectin matrix assembly. *J. Cell Biol.* **141**, 539-551.