

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## QTL analysis of plant development and fruit traits in pepper and performance of selective phenotyping

### This is the author's manuscript

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/100784> since

*Terms of use:*

#### Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*Theoretical and Applied Genetics 118 (6), 2009, DOI 10.1007/s00122-009-0970-0*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*<http://link.springer.com/article/10.1007/s00122-009-0970-0/fulltext.html>*

**Lorenzo Barchi<sup>1,+</sup> Véronique Lefebvre<sup>1</sup> Anne Marie Sage-Palloix<sup>1</sup> Sergio Lanteri<sup>2</sup>  
and Alain Palloix<sup>1</sup>**

**QTL analysis of plant development and fruit traits in pepper using selective phenotyping.**

<sup>1</sup> INRA-Avignon, Unité de Génétique et Amélioration des Fruits et Légumes, BP 94, F-84143 Montfavet Cedex, France.

<sup>2</sup> Di.Va.P.R.A., Plant Genetics and Breeding, University of Turin, Via L. da Vinci 44, I-10095 Grugliasco (Turin), Italy.

<sup>+</sup> Present adress : Di.Va.P.R.A., Plant Genetics and Breeding, University of Turin, Via L. da Vinci 44, I-10095 Grugliasco (Turin), Italy.

Corresponding author: Alain Palloix

alain.palloix@avignon.inra.fr

tel 33 (0)432 72 27 41

fax 33 (0)432 72 27 02

Manuscript with 3 tables and 2 figures

## Abstract

A QTL analysis was performed to determine the genetic basis of 13 horticultural traits conditioning yield in pepper (*Capsicum annuum*). The mapping population was a large population of 297 recombinant inbred lines (RIL) originating from a cross between the large-fruited bell pepper cultivar ‘Yolo Wonder’ and the small-fruited chilli pepper ‘Criollo de Morelos 334’. A total of 76 QTLs were detected for 13 fruit and plant traits, grouped in 28 chromosome regions. These QTLs explained together between 7% (internode growth time) and 91% (fruit diameter) of the phenotypic variation. The QTL analysis was also performed on two subsets of 141 and 93 RILs sampled using the MapPop software. The smaller populations allowed for the detection of a reduced set of QTLs and reduced the overall percentage of trait variation explained by QTLs. The frequency of false positives as well as the individual effect of QTLs increased in reduced population sets as a result of reduced sampling. The results from the QTL analysis permitted an overall glance over the genetic architecture of traits considered by breeders for selection. Colinearities between clusters of QTLs controlling fruit traits and/or plant development in distinct pepper species and in related solanaceous crop species (tomato and eggplant) suggests that shared mechanisms control the shape and growth of different organs throughout these species.

Key words: *Capsicum annuum*, QTL analysis, yield, fruit traits, plant growth, selective phenotyping

## Introduction

All commercial cultivars result from long and intensive breeding efforts to improve multiple traits including resistance to diseases, fruit or grain characteristics and plant development, which determine adaptation to agro-climatic as well as market conditions. The favourable alleles for these traits are primarily dispersed in distinct individuals in plant germplasm and the selection of elite cultivars gathering these traits requires multiple crossovers within the genetic background. This is particularly true in pepper (*Capsicum annuum*) where disease resistance and adaptation to agroclimatic conditions have to be introgressed into large and sweet fruited cultivars from small-fruited and pungent accessions, which form the large majority of genetic resources. Multi-trait selection is often hampered by unfavourable genetic linkages. However, the genetic mapping of valuable traits should shed light on the genomic regions of economic interest and the loci where genetic recombination should result in more genetic gain. Two questions related to this stake will be addressed in this article. One generic question is: can we reduce the efforts developed in genetic mapping for multiple traits that requires repeated experiments of large progenies? The second and specific question is: what is the architecture of genes and QTLs of interest in the pepper chromosomes, and can the comparison with genomes from other species help in their genome localization and characterization?

The power of QTL detection is determined by many parameters, including the choice of the parental lines (the more the parental lines are genetically distant, the easier is the detection of QTLs) previously reported by Crepieux et al. (2004), the size of the segregating population and the magnitude of the experimental error (Hackett 2002). Since large populations do allow for the identification of QTLs with weak effects (Tanksley 1993; Haley and Andersson 1997), the effort involved in the assessment of the phenotypes is large. Selective phenotyping has been proposed as a trade-off strategy to decrease phenotyping costs, and Brown and Vision (2000) developed the MapPop software which proposes to select a reduced subset of the most informative individuals, chosen from their genotype data. This reduced subset is further submitted to phenotyping, reducing the field experiments while losing as little information as possible. Such strategies were evaluated by Vales et al. (2005) and Birolleau-Touchard et al. (2007). They showed that the MapPop sampling method permits to select

population samples with balanced allele frequencies, and was superior to random sampling, particularly when QTL analysis has to be performed from unbalanced populations (doubled haploid or advanced backcross populations). However, performance in QTL detection remained affected by the population size, particularly for QTLs with low individual effects which are frequent for horticultural traits like fruit and plant growth traits in vegetables.

Considering pepper, gene and QTL mapping has been largely developed for disease resistance traits (Lefebvre 2005; Djian-Caporalino et al. 2006) and fruit traits, using both intraspecific (Ben Chaim et al. 2001,2006) and interspecific crosses (Rao et al. 2003; Zygier et al. 2005; Ben Chaim et al. 2006). Little information is available on QTLs controlling plant development traits: plant height and flowering time were investigated by Ben Chaim et al. (2001) and Rao et al. (2003), but primary components that determine plant growth and fruiting earliness, like the number of internodes on the primary stem, their length and growth speed, were not investigated. Considering fruit traits, results from Ben Chaim et al. (2001, 2006), Rao et al. (2003) and Zygier et al. (2005) are strongly coherent and revealed clusters of QTLs on pepper chromosomes with tight linkage or pleiotropy between QTLs for fruit size, diameter, length and shape. Presence of QTL clusters on chromosomes P2, P3, P4 and P10 between distinct crosses attested the conservation of QTL effects for fruit traits. Moreover, relationships with other solanaceous crops were explored, particularly for genetics of fruit size and shape. In tomato, numerous QTL mapping studies on horticultural traits were carried out, mainly in interspecific crosses (De Vicente and Tanksley 1993; Bernacchi et al. 1998; Grandillo et al. 1999; Ku et al.1999; Frary et al. 2004; Barrero and Tanksley 2004; Jiménez-Gómez et al. 2007). QTL mapping research on horticultural traits was also performed to a more limited extent in eggplant and potato (Van Eck et al. 1994; Doganlar et al. 2002; Frary et al. 2003). The large body of research concerned with trait inheritance in tomato has opened the way to the establishment of synteny (conservation of gene repertoire) and colinearity (conserved gene orders) among the *Solanaceae* (Tanksley et al. 1992; Livingstone et al. 1999; Ben Chaim et al. 2001, 2003a, b; Doganlar et al. 2002; Rao et al. 2003; Zygier et al. 2005; Ben Chaim et al. 2006; Paran and Van der Knaap 2007).

In the current paper, we report the analysis of 13 horticultural traits including fruit traits

and plant development traits, in a large intraspecific recombinant inbred progeny (RILs) of pepper that was used by Barchi et al. (2007) to develop a high-resolution genetic linkage map. QTL detection was performed in the whole population including 297 RILs and compared with informative subsets of 141 and 93 RILs to test the advantage of selective phenotyping in this progeny. QTL analysis for multiple traits informed the relative position of QTL clusters and genetic linkages between loci determining traits of interest. It also delivered target QTLs that suggest syntenic relationships with known horticultural QTLs in other *Solanaceae* species.

## Materials and methods

### *Plant material and trait evaluation*

The plant material was developed from an intraspecific cross between the large-fruited inbred line ‘Yolo Wonder’ (YW) and a small-fruited pungent inbred line ‘Criollo de Morelos 334’ (CM334) issued from the Mexican landrace ‘Criollo de Morelos’. The recombinant inbred line (RIL) population was based on a set of  $F_6$  derivatives, produced by self-pollination of the  $F_5$  RILs used to construct the genetic map (Barchi et al. 2007). The horticultural traits of the parental lines, their  $F_1$  hybrid and the RILs were measured in a single year experiment arranged in a randomized complete block design with three blocks of three individual plants (repeats) per genotype and block. The horticultural traits measured are included among the standard morphological descriptors for pepper (IPGRI 1995). The fruit traits were measured from two individual fruits per plant, except the fruit weight measured as an average of 5–10 fruits per plant. These fruit traits were the log-transformed mean of the fruit weight (Lfw), the fruit length (Frl) measured as the distance (in mm) between the pedicel attachment and the fruit apex, the fruit diameter (Frd) measured as the maximum fruit width (in mm), the fruit shape (Frs) defined as the Frl:Frd ratio, the pericarp thickness (Pet) in mm, the pedicel length (Pel) in mm, the number of fruit locules (Nlo). The plant traits were measured from each individual plant: the flowering earliness (Flw), i.e., the number of days from sowing to the anthesis of the first flower, the primary axis length (Axl) defined as the length (in cm) of the primary axis from the cotyledons to the first branch, the number of leaves on the primary axis (Nle), the mean internode length (Inl) given by the ratio Axl:Nle in cm, the axis growth speed (Axs) given by the ratio Axl:(Flw - 15 days) in which the 15 days corresponding to the time of hypocotyl and cotyledons emergence after sowing were deduced to obtain the growth time of the axis = epicotyl, and the mean internode growth time (Int) given by the ratio (Flw - 15 days):Nle.

### *Data and QTL analyses*

All statistical analyses were performed using R software (R Development Core Team 2006). Analyses of variance (ANOVA) were applied to estimate genotypic/environmental effects according to the model  $Y_{ij} = i + b_j + g_i + e_{ij}$ , where  $i$ ,  $b_j$ ,  $g_i$  and  $e_{ij}$  represent, respectively, the overall mean, block effect, genotypic effect



and error effect. Broad-sense heritability ( $h_{BS}^2$ ) values were calculated as  $\sigma_G^2 / (\sigma_G^2 + \sigma_E^2 / n)$  where  $\sigma_G^2$  is the genetic variance,  $\sigma_E^2$  the environmental variance (including block, interaction and error effects) and  $n$  the number of blocks. Correlations between traits were estimated using Pearson's coefficient. The normal distribution of traits was assessed by using the Shapiro–Wilks test ( $\alpha = 0.05$ ). Lines were declared transgressive when their value exceeded the highest parental value or were lower than the lowest parental value by more than the least significant difference (LSD) at  $P < 0.05$ .

QTL detection was based on the genetic map developed by Barchi et al. (2007), and made with QTL Cartographer software (Basten et al. 2002), employing both interval (IM) (Lander and Botstein 1989) and composite interval (CIM) (Zeng 1994) mapping. For each population size, LOD thresholds at the 5% probability level were empirically established by applying 1,000 permutations for each trait and for both methods (Churchill and Doerge 1994). The proportion of observed phenotypic variation attributable to an individual QTL was estimated by the coefficient of determination ( $R^2$ ). The total phenotypic variation explained was estimated by fitting a linear model (multiple regression) including all the putative QTLs for the respective trait simultaneously. For each QTL, the marker with the highest LOD was chosen as the QTL-representative marker variable in the multiple regression. Epistatic interactions between all the 250 markers of the core map were tested using a two-way ANOVA model with interaction effect between all possible two-locus combinations of core map marker genotypes, as described by Lefebvre and Palloix (1996). With 250 core markers, 32,125 marker combinations per trait were tested (404,625 for 13 traits) and epistasis was inferred to be significant at  $P < 10^{-5}$ . Individual QTLs were named using the three-first letters to indicate the trait, followed by a number indicating the pepper chromosome (e.g. 1 for P1) or by the linkage group involved (e.g. LG13), and a second number defining the position within the chromosome/LG to which the QTL mapped (e.g. 1.1). IM or CIM were added when a QTL was detected with only one of the two methods. MapChart software (Voorrips 2002) was used to produce visualisations of chromosomes carrying QTLs.

### *Choice of informative individuals*

26 of the full set of 297 RILs were removed from the population because of their poor fertility resulting in a lack of seeds for further experiments. From the remaining 271 RILs, two subpopulations ['A' (141 RILs) and 'B' (93 RILs)] were selected using MapPop software (Brown and Vision 2000), with the full linkage map as the input file. The selection criterion applied to identify the most informative individuals was the expected maximum bin length (eMBL), i.e. the expected maximum distance between two points subjected to recombination. As previously reported by Birolleau-Touchard et al. (2007), the best sample is the one with the most similar eMBL value compared to the original population obtained with a defined computational time and a chosen sample size. As a consequence, several iterations were run, until no further improvement in eMBL was achieved (data not shown).

## Results

### *Phenotypic variation and trait correlations*

The mean phenotypic values and estimated  $h_{BS}^2$  for each trait are listed in Table 1. The parental lines displayed significant contrasted phenotypes (Table 1) for fruit traits, with a larger, heavier fruit, a thicker pericarp and a higher number of locules for YW compare to the small and elongated fruit of CM334 which showed a higher fruit shape ratio. The pedicel length was similar for the two cultivars. Considering the plant traits, the two parental lines also displayed contrasted phenotypes (Table 1), YW presenting a later flowering date, a shorter primary axis and internode length and a slower growth speed of the axis than CM334. However, the two parental lines were very close for the number of leaves on the axis and internode growth time. In the full RIL set, all the traits were normally distributed. The RIL progeny and F<sub>1</sub> hybrid displayed intermediate values between the parental lines for all the fruit traits, except the pedicel length (Pel) and the fruit length (Frl) where about 69% and 11.7% of the progeny displayed significant transgressive phenotypes with higher or lower values than the parental lines ( $\alpha = 0.05$ ). Considering plant traits, for almost of them (except the flowering earliness) a large proportion (from 9.7% for Axs to 56% for Nle) of RILs displayed transgressive phenotypes; at the same time also the F<sub>1</sub> hybrid displayed higher (Axl, Inl, Axs) or lower (Flw, Nle) values than the parental lines.

Heritabilities were high for all the traits, ranging from 0.84 (Nlo) to 0.97 (Frd and Frs) (Table 1). Highly significant correlations were detected between fruit traits or between plant traits (Table 2). As a general feature, fruit weight, fruit diameter, locule number and pericarp thickness were positively correlated together but negatively correlated with fruit shape. Considering plant traits, the axis length increased with the number of leaves and the internode length. However, the growth speed of the axis (and of internodes) also increased with the axis length, so that flowering time (earliness) displayed a significant but weak correlation with Nle. Correlations between fruit and plant traits were either non-significant ( $P > 0.05$ ) or weak ( $< 0.3$ ).

Trait means in the two subpopulations are listed in Table 1. In no case, a subpopulation mean differed significantly ( $P < 0.05$ ) from the full population mean. The distribution of trait values was also very similar in both subpopulations and the full population for all the traits and the trait variances remained constant across subpopulations. Correlations

between traits were also comparable with those calculated for the full population (data not shown).

#### *QTL detection in the full and partial RIL populations*

Across the 13 traits, 76 QTLs were identified and grouped onto 28 genomic intervals distributed over 11 chromosomes and 14 small unassigned linkage groups (Fig. 1; Table 3). No significant epistatic interactions were identified between markers, even at higher  $P$  value ( $P < 10^{-2}$ ). Clusters of QTLs governing 4–6 distinct fruit traits were mapped on chromosomes P2, P3, P4, P10, P11, P12 and small linkage groups LG 17, 24, 25 and 27. Considering plant developmental traits, only four tightly linked QTLs affecting four different traits were mapped on chromosome P2, with smaller clusters on chromosomes P1, P4 and P9. Depending on the traits, 2–12 QTLs were detected, with individual  $R^2$  values ranging from 0.21 to 0.04. The total phenotypic variation explained by QTLs (global  $R^2$ ) ranged from 0.07 to 0.91. Traits with the highest global  $R^2$  were the fruit diameter (12 QTLs,  $R^2 = 0.91$ ) followed by the fruit weight (7 QTLs,  $R^2 = 0.48$ ), the number of locules (9 QTLs,  $R^2 = 0.46$ ), the axis growth speed (4 QTLs,  $R^2 = 0.44$ ) and the axis length (4 QTLs,  $R^2 = 0.42$ ). The traits which variation was poorly explained by QTLs were the internode growth time (2 QTLs,  $R^2 = 0.07$ ), the internode length (3 QTLs,  $R^2 = 0.10$ ) and the flowering earliness (5 QTLs,  $R^2 = 0.15$ ). For the fruit traits Lfw, Frd, Pet and Nlo, all the positive alleles were derived from the large-fruited parental line YW. For the fruit length (Frl) and shape (Frs), both parents contributed alleles at different QTLs that increased the trait value, that was expected since CM334 harbours long-shaped fruits. For the pedicel length (Pel) and all the plant traits, both parental alleles contributed to increase the trait.

In the subpopulations A (141 RILs) and B (93 RILs), 26 and 15 QTLs were detected respectively (Table 3). No significant epistatic interactions were identified in either subpopulation ( $P < 10^{-6}$ ). The LOD scores associated to the QTLs were always much lower in the reduced subpopulations than in the full RIL population, and only the QTLs with the highest LOD scores remained significant. For example, *Frd11.1* QTL showing a LOD score greater than 11 was common to all three populations, while other Frd QTLs, with lower LOD scores were progressively lost in the A and B subpopulations. A contrario, their individual  $R^2$  values were equal or higher in the subpopulations than in

the full RIL population. For example, the  $R^2$  value for *LfwLG24.1* increased from 0.08 in the original population to 0.10 and 0.14 in the subpopulations A and B respectively, while for *Flw1.1*, the  $R^2$  value increased from 0.10 (297 RILs) to 0.12 (A) and 0.17 (B). The total phenotypic variation explained by QTLs globally decreased with the population size, ranging from 0.91 to 0.07 in the whole population, from 0.36 to 0.08 in the A subpopulation and from 0.34 to 0.12 in the B subpopulation. This was spectacular for complex traits like Frd, which global  $R^2$  decreased from 0.91 (297 RILs) to 0.36 (subpopulation A) and 0.21 (subpopulation B). Finally, the use of subpopulations led to detect new QTLs for Pel, Nle and Int in A subpopulation and two new QTLs for Frd and Flw in B subpopulation. These QTLs were not detected in the whole population even as putative (close to the LOD threshold) QTLs.

## Discussion

The QTLs involved in the inheritance of 13 horticultural traits analysed provide a picture of the genetic complexity of these traits, and the location of the major loci should be informative for the genetic analysis and breeding of pepper. The mapping population was derived from an intraspecific cross between parental types widely used by pepper breeders. This choice serves both to maximise mapping precision by reducing the frequency of the skewed segregations commonly experienced in interspecific crosses, and to generate information with respect to QTL of relevance for genetic improvement within advanced germplasm.

### *General features of the data and QTL detection*

Phenotypes were assessed in a single environment, that may affect portability of QTLs to different agro-climatic conditions. The plant environment may poorly affect the relative values for fruit shape and weight within a segregating progeny, and this is confirmed by the fact that the main QTLs for fruit weight and shape appeared conserved in these distinct studies. No previous experiments permit this comparison as regards growth and development traits. However, the axis length, the number of leaves on primary axis and the flowering date are known by breeders to be affected by environment  $\times$  genotype interactions, despite these traits are very stable (genetically determined) in a constant environment (unpublished data). Thus, the results obtained here for plant traits may be relevant mainly in open field conditions in Mediterranean region. Nevertheless the high level of reproducibility between replicates resulted in high heritabilities of the traits in the tested environment. The heritability values were generally higher than those reported by Rao et al. (2003), but comparable to those estimated by Ben Chaim et al. (2001). The percentage of genetic variation explained by QTLs was highly trait-specific but not dependent on trait heritability. Frd was associated with the highest  $R^2$  value (0.91) and Int with the lowest one (0.07) despite they displayed similar heritabilities (0.97 and 0.91 respectively). The highest percentage of variation explained by QTLs occurred when many QTLs with minor effects (e.g. Frd) were detected. Global  $R^2$  depends on the accuracy of the linkage map as well as on the inter-marker distances (Jansen et al. 1995). The genetic structure of the trait, the size of the mapping population (Melchinger et al. 1998; Utz et al. 2000), the genetic

background, the environment, and interactions between QTLs can all contribute to a failure to detect all the QTLs segregating in a population. In the present study, the limited percentage of variation explained for some traits, may indicate that the heritabilities were partially overestimated due to a single year and single environment experiment (the variation between lines would not reflect exclusively the genetic variations), but also suggests that some QTLs remain undetected, possibly because of their low individual effects.

#### *Transgressive lines and contribution of parental alleles at the QTLs*

Transgressive segregations in the RIL progeny were observed for many traits, particularly when parental values were similar (pedicel length, number of leaves). In every case, alleles from both parental lines were shown to contribute to increase (or decrease) the trait value. Transgressive genotypes were already known to result from the combination of alleles from both parents that have effect on the same direction (De Vicente and Tanksley 1993). The effect of such allele combinations was tested regarding to the graphical genotypes at the detected QTLs of the three individuals with the highest and three individuals with the lowest values for the pedicel length (Fig. 2). The three transgressive individuals with the highest pedicel length were found to possess all the six alleles responsible for pedicel elongation (YW alleles at P1, P4 and LG42 and CM334 alleles at P2, P3 and P12), with the exclusion of one genotype, lacking one allele at *Pel12.1*. On the contrary, the three individuals with the shortest pedicel length possessed all the alleles which decreased the pedicel length, with the exclusion of one individual at *Pel2.1*. This result indicates that these six QTLs explain a major part (if not all) of the observed trait variation, and that the global  $R^2$  of 0.37 is an underestimation, probably resulting from incomplete linkage (recombination rate) between the QTLs and the markers inferred in the multiple regression model.

#### *Effect of selective sampling and reduced population size on QTL detection*

The detection power increases as the population size is maximized (Charcosset and Gallais 1996), an effect which is most severe for traits displaying lower levels of heritability (Gallais and Rives 1993). However practical considerations, particularly with regard to phenotyping, dictate that population sizes should be minimised as far as

possible. Thus Charmet (2000) has shown that 100 doubled haploid (DH) lines appear to be the critical limit for detecting QTLs of traits with a heritability above 0.3, while about 1,000 individuals are needed to detect QTLs for a trait with a heritability of 0.25. The precision of QTL detection also depends on the sampling method, which can be either random or based on either selective genotyping or selective phenotyping. We have chosen the latter strategy to generate the two subpopulations, one of which is about half the size of the full set, and the other about one-third.

The use of subpopulations brought to the identification of fewer QTLs, characterised by a lower LOD score than those detected in the original population. In most cases, only QTLs with the highest LOD scores in the full population remained significant in the subpopulations. The reduction of the number of QTLs detected when reducing the number of individuals used was also observed by Vales et al. (2005) and Birolleau-Touchard et al. (2007) who compared different sampling procedures, including the MapPop approach. As an expected consequence of the reduction of the number of detected QTLs, the use of small subsets of individuals led in our analyses to the reduction of the global  $R^2$  value for each trait. This result is in contrast with that reported by Vales et al. (2005) who obtained an increasing of global  $R^2$  values as the population size decreased. This probably resulted from the decrease of the whole phenotypic variance in their reduced subpopulations, instead of the phenotypic variances of our subpopulations A and B remained similar to that of the whole population. On the other side, Vales et al. (2005) also found that the use of reduced populations overestimates the individual  $R^2$  value of each QTL. The use of subpopulations also led to the detection of new QTLs, which were significant only in the smallest subpopulation B. The reduction in population size seemed to have generated unbalanced (skewed) samples which were responsible for detecting spurious QTLs (Birolleau-Touchard et al. 2007). We previously showed in this progeny that selective genotyping was useful for the fast mapping of additional markers to enhance a pre-existing genetic map (Barchi et al. 2007). In contrast, according to the results obtained, it seems that the selective phenotyping strategy performs poorly for traits governed by multiple genes with weak effects.



### *Genetic architecture of the QTLs determining fruit and plant traits*

The analyses performed led to detect a total of 76 QTLs distributed over all the pepper chromosomes (with the exclusion of chromosome P5) and most unassigned linkage groups. All the traits are polygenically controlled, mainly by small effect QTLs as the  $R^2$  values never overpassed 0.21 (Frl4.1). The fruit diameter was found to be controlled by 12 QTLs with individual  $R^2$  ranging from 0.05 to 0.12 and distributed in 12 independent genomic regions.

QTLs controlling fruit traits were mainly concentrated on P2, P3, P4, P10, P11 and P12, with main effects on P4 for fruit weight and length, on P3 for fruit shape and length (upper arm) and for diameter and pericarp thickness (lower arm). Many fruit trait QTLs landed near the same chromosomal location, as a result of linkage, but also pleiotropy since alleles that affect fruit length or diameter will also affect fruit shape, and alleles affecting any fruit dimension (including pericarp thickness) will affect fruit weight. Clusters of QTLs for fruit traits were previously detected in colinear positions on P2, P3, P4, P10 in intra as well as interspecific crosses (Zygier et al. 2005; Ben Chaim et al. 2001, 2003a), thus suggesting a conservation of alleles controlling these traits in both intra and interspecific germplasm. In the present study, we also detected a cluster on P11 of QTLs with minor effects for pericarp thickness, fruit weight, diameter and shape, linked to markers e41/m61\_137c and e41/m61\_270c that was also detected in interspecific crosses by Rao et al. (2003) but never reported in intraspecific crosses. Similarly, two QTLs for fruit shape were detected on P4, linked to markers p14/m39\_417y and p25/m45\_109y; fruit shape QTL clusters linked to these same markers were previously mapped in two separate interspecific crosses by Zygier et al. (2005). In addition, we detected a cluster of QTLs controlling the fruit weight, diameter, the number of locules and the pericarp thickness on chromosome P12 and linked to marker e44/m51\_263c which was not previously detected. This probably resulted from allelic differences specific to the (YW  $\times$  CM334) cross, displaying new alleles controlling the fruit parameters which could be potentially exploited in breeding programs. Co-localizations between fruit traits explain the correlations detected in the progeny, and suggest shared mechanisms between fruit components, i.e. pericarp thickness, fruit diameter and length that contribute to fruit weight of pepper.

Concerning the QTLs controlling plant growth, a tight co-localisation of four QTLs

affecting flowering earliness, axis length, axis growth speed and internode length was detected on chromosome P2. Additional clusters on chromosome P4 (flowering earliness, axis growth speed and internode growth time), on chromosome P9 (axis length and axis growth speed) and on LG47 (axis length, internode length and number of leaves) were also found. This clustering/co-localization of QTLs is in agreement with the correlations found between these traits. It may indicate linkage effects but also pleiotropic effects of genes on the morphology (internode length, number of leaves) or growth speed of vegetative organs. Co-localization between plant and fruit trait QTLs were detected on P2. A special interest should be brought on this chromosome where QTLs affecting the shape of several organs, i.e. the number of fruit locules, the fruit shape, the fruit diameter, the axis, internode and pedicel lengths, were tightly co-localized with overlapping LOD peaks. Only fine mapping would bring additional information on the linkage/pleiotropic determinism of these traits which all depend on cell division, elongation and regulation of the cell cycle. Several gene families involved in the cell cycle regulation were characterized for their functional role in plant growth and development (Goda et al. 2004; Inzé 2005). They may provide good candidates for further fine mapping these QTLs and looking for synteny relationships with *Arabidopsis* and solanaceous crops.

#### *Potential orthologous QTLs with other Solanaceae species*

Several studies revealed potential syntenies and colinearities among the *Solanaceae* species, in particular for fruit size and shape between pepper, tomato and eggplant (Tanksley et al. 1992; Livingstone et al. 1999; Ben Chaim et al. 2001; Doganlar et al. 2002; Rao et al. 2003; Ben Chaim et al. 2003a, b; Zygier et al. 2005; Ben Chaim et al. 2006; Paran and Van der Knaap 2007). However, the lack of common markers used to map these loci in our progeny prevents further evidence of synteny between these species. The detection of QTL clusters for pepper fruit traits on chromosomes P2, P3, P4 and P10 only consolidates the previous analyses performed by the authors cited above. Some additional information was obtained regarding the fruit QTL cluster on P2. In tomato, several QTLs controlling fruit size were mapped in the colinear genomic region of chromosome T2. Ben Chaim et al. (2001) and Zygier et al. (2005) detected QTLs for fruit weight, shape and diameter in different crosses in pepper that were

suggested to correspond to the tomato genomic region of the QTL *fw2.1*, due to their tight linkage with the *ovate* sequence in both pepper and tomato. In our study, we detected again fruit diameter and fruit shape QTLs (*Frd2.1*, *Frs2.1*) in this same genomic region, but also a QTL for the locule number (*Nlo2.1*) that might correspond to the *lcn2.1* which also affects the number of locules in the tomato fruit (Lippman and Tanksley 2001; Barrero and Tanksley 2004). This last QTL in tomato, depending on the cross, explains 19 or 32% of the phenotypic variability, while in pepper, its effect seems lower with a  $R^2$  of 5%. In pepper, fruit weight only depends on pericarp development, and fruit shape is mostly determined after fertilization, that suggests different mechanisms from tomato. However, co-localization of QTLs for the number of locules, the fruit diameter and shape and of the *ovate* gene strongly suggests synteny relationships of genes affecting cell division, elongation and polarity with colinearity relationships in tomato and eggplant. Additional candidate QTLs for synteny relationships include the QTL *Ax12.1* for height of the primary stem which has putative orthologues in both tomato chromosome T2 (Paran et al. 1997) and eggplant chromosome E2 (Frary et al. 2003.)

Further new QTLs which we identified and showing a putative synteny with other *Solanaceae* species are *pet10.1* (linked to marker e36/m47\_145y with a confidence interval of 21 cM) affecting the pericarp thickness, which is a possible orthologue to tomato *pcp10.1* (Frary et al. 2004) linked to TG52 marker. Finally, several QTLs were found in unassigned LGs and some of them (like *NloLG25.1* and *IntLG47.1*) explain a relatively high percentage of variation with individuals  $R^2$  values of 0.13 and 0.11 respectively.

Results obtained from our and previous studies show that the main constraint in synteny analysis is due to the complex rearrangements between *Capsicum* and *Solanum* genomes. Fine mapping with shared markers will be required for identifying orthologous regions among these *Solanaceae* species. Moreover, Zygier et al. (2005) pointed out that the level of orthology between pepper and tomato chromosome T2 and T4 for QTL controlling fruit shape was limited as a consequence of the divergent selection that occurred during the domestication of these species. The divergent selection for horticultural traits in *Solanaceae* species may have conserved only a limited number of orthologous QTL thus allowing an explanation to the limited number

of shared QTLs found between pepper, tomato and eggplant.

## **Conclusions**

Our results showed that fruit traits are mostly inherited independently from plant growth characteristics, except linkage relationships in chromosomes P2 and P4. This offers large degrees of freedom in breeding for plant and fruit ideotypes. Moreover, allelic diversity and contribution of alleles from distinct genotypes should increase the range of potential phenotypes. However, the clusters of QTLs for horticultural traits detected in chromosomes P4, P9 and P10 revealed linkage with previously mapped QTLs or major genes for resistance to major pepper diseases including *P. capsici*, tospoviruses and potyviruses on chromosomes P4 and P10, nematodes on chromosome P9 (for review see Djian-Caporalino et al. 2006). Selection of recombination events in these chromosomes have to be facilitated by a marker-assisted strategy, and markers localized in the chromosome regions where recombinant individuals have to be selected need to be further developed. The linkage between markers and QTLs for horticultural important traits may therefore represent a valuable tool for multi-trait marker-based breeding strategy in pepper.

## **Acknowledgments**

The salary of L. Barchi was supported by a grant of the C.I.P.E. (Resolution 17/2003) from the Italian Ministry of Agricultural Alimentary and Forest Politics. The authors thank P. Signoret, G. Nemouchi and T. Phaly for technical assistance.

## References

- Barchi L, Bonnet J, Boudet C, Signoret P, Nagy I, Lanteri S, Palloix A, Lefebvre V (2007) A high-resolution intraspecific linkage map of pepper (*Capsicum annuum* L.) and selection of reduced RIL subsets for fast mapping. *Genome* 50:51–60.
- Barrero L-S, Tanksley S-D (2004) Evaluating the genetic basis of multiple-locule fruit in a broad cross-section of tomato cultivars. *Theor Appl Genet* 109:669–679.
- Basten C-J, Weir B-S, Zeng ZB (2002) QTL cartographer, version 1.16. Department of Statistics. North Carolina State University, Raleigh.
- Ben Chaim A, Paran I, Grube R-C, Jahn M, van Wijk R, Peleman J (2001) QTL mapping of fruit-related traits in pepper (*Capsicum annuum*). *Theor Appl Genet* 102:1016–1028.
- Ben Chaim A, Borovsk E, Rao G-U, Tanyolac B, Paran I (2003a) Fs3.1: a major fruit shape QTL conserved in *Capsicum*. *Genome* 46:1–9.
- Ben Chaim A, Borovsky E, De Jong W, Paran I (2003b) Linkage of the A locus for the presence of anthocyanin and fs10.1, a major fruit-shape QTL in pepper. *Theor Appl Genet* 106:889–894.
- Ben Chaim A, Borovsky J, Rao G, Gur A, Zamir D, Paran I (2006) Comparative QTL mapping of fruit size and shape in tomato and pepper. *Isr J Plant Sci* 54:191–203.
- Bernacchi D, Beck-Bunn T, Emmatty D, Eshed Y, Inai S, Lopez J, Petiard V, Sayama H, Uhlig J, Zamir D, Tanksley S (1998) Advanced backcross QTL analysis of tomato. II. Evaluation of near-isogenic lines carrying single donor introgressions for desirable wild QTL alleles derived from *Lycopersicon hirsutum* and *L. pimpinellifolium*. *Theor Appl Genet* 97:381–397.
- Birolleau-Touchard C, Hanocq E, Bouchez A, Bauland C, Dourlen I, Seret J-P, Rabier D, Hervet S, Allienne J-F, Lucas P-H, Jaminon O, Etienne R, Baudhuin G, Giauffret C (2007) The use of MapPop1.0 for choosing a QTL mapping sample from an advanced backcross population. *Theor Appl Genet* 114:1019–1028.
- Brown D, Vision T (2000) MapPop 1.0: software for selective mapping and bin mapping. <http://www.bio.unc.edu/faculty/vision/lab/mappop/>.
- Charcosset A, Gallais A (1996) Estimation of the contribution of quantitative trait loci (QTL) to the variance of a quantitative trait by means of genetic markers. *Theor Appl Genet* 93:1193–1201.
- Charmet G (2000) Power and accuracy of QTL detection: simulation studies of one-QTL models. *Agronomie* 20:309–323.
- Churchill G-A, Doerge R-V (1994) Empirical threshold values for quantitative trait mapping. *Genetics* 138:963–971.
- Crepieux S, Lebreton C, Servin B, Charmet G (2004) Quantitative trait loci (QTL) detection in multicross inbred designs. Recovering QTL identical-by-descent status information from marker data. *Genetics* 168:1737–1749.
- De Vicente M-C, Tanksley S-D (1993) QTL analysis of transgressive segregation in an interspecific tomato cross. *Genetics* 134:585–596.
- Doganlar S, Frary A, Daunay M-C, Lester R-N, Tanksley S-D (2002) Conservation of gene function in the Solanaceae as revealed by comparative mapping of domestication traits in eggplant. *Genetics* 161:1713–1726.
- Djian-Caporalino C, Lefebvre V, Sage-Daubèze, A-M, Palloix A (2006) *Capsicum*. In: Singh RJ (ed) Genetic resources, chromosome engineering, and crop improvement, vol 3. Vegetable

- crops, CRC Press, Boca Raton, pp 185–243.
- Frary A, Doganlar S, Daunay M-C, Tanksley S-D (2003) QTL analysis of morphological traits in eggplant and implications for conservation of gene function during evolution of solanaceous species. *Theor Appl Genet* 107:359–370.
- Frary A, Fulton T-M, Zamir D, Tanksley S-D (2004) Advanced backcross QTL analysis of a *Lycopersicon esculentum* × *L. pennellii* cross and identification of possible orthologs in the Solanaceae. *Theor Appl Genet* 108:485–496.
- Gallais A, Rives M (1993) Detection, number and effects of QTL for a complex character. *Agronomie* 13:723–738.
- Goda H, Sawa S, Asami T, Fujioka S, Shimada Y, Yoshida S (2004) Comprehensive comparison of auxin-regulated and brassinosteroid-regulated genes in *Arabidopsis*. *Plant Physiol* 134:1–19.
- Grandillo S, Ku H-M, Tanksley S-D (1999) Identifying the loci responsible for natural variation in fruit size and shape in tomato. *Theor Appl Genet* 99:978–987
- Hackett C-A (2002) Statistical methods for QTL mapping in cereals. *Plant Mol Biol* 48:585–599.
- Haley C-S, Andersson L (1997) Linkage mapping of quantitative trait loci in plants and animals. In: Dear PH (ed) *Genome mapping. A practical approach*. Oxford University Press, Oxford, pp 49–71.
- Inzé D (2005) Green light for the cell cycle. *EMBO J* 24:657–662.
- IPGRI, AVRDC, and CATIE (1995) Descriptors for Capsicum, International Plant Genetic Resources Institute. Rome, Italy; the Asian Research and Development Center, Taipei, Taiwan and The Centro Agronomico Tropical de Investigacion y Ensenanza, Turrialba, Costa Rica, p 49. <http://www.ipgri.cgiar.org>.
- Jansen R-C, Van Ooijen J-M, Stam P, Lister C, Dean C (1995) Genotype-by-environment interaction in genetic mapping of multiple quantitative trait loci. *Theor Appl Genet* 91:33–37.
- Jiménez-Gómez J-M, Alonso-Blanco C, Borja A, Anastasio G, Angosto T, Lozano R, Martínez-Zapater J-M (2007) Quantitative genetic analysis of flowering time in tomato. *Genome* 50:303–315.
- Ku H-M, Doganlar S, Chen K-Y, Tanksley S-D (1999) The genetic basis of pear-shaped tomato fruit. *Theor Appl Genet* 9:844–850.
- Lander E-S, Botstein D (1989) Mapping mendelian factors underlying quantitative traits using RFLP linkage maps. *Genetics* 121:185–199.
- Lefebvre V (2005) Molecular markers for genetics and breeding: development and use in pepper (*Capsicum* spp.). In: Lörz H and Wenzel G (eds) *Molecular marker systems in plant breeding and crop improvement*. Biotechnology in Agriculture and Forestry, vol 55. Springer, Berlin, pp 189–214.
- Lefebvre V, Palloix A (1996) Both epistatic and additive effects of QTLs are involved in polygenic induced resistance to disease, a case study, the interaction pepper—*Phytophthora capsici* Leonian. *Theor Appl Genet* 93:503–511.
- Lippman Z, Tanksley S-D (2001) Dissecting the genetic pathway to extreme fruit size in tomato using a cross between the small-fruited wild species *L. pimpinellifolium* and *L. esculentum* var. Giant Heirloom. *Genetics* 158:413–422.
- Livingstone K-D, Lackney V-K, Blauth J-R, van Wijk R, Jahn M-K (1999) Genome mapping in

- capsicum and the evolution of genome structure in the Solanaceae. *Genetics* 152:1183–1202.
- Melchinger A-E, Utz H-F, Schoon C-C (1998) Quantitative trait locus (QTL) mapping using different testers and independent population samples in maize reveals low power of QTL detection and large bias in estimates of QTL effects. *Genetics* 149:383–403.
- Paran I, Van der Knaap E (2007) Genetic and molecular regulation of fruit and plant domestication traits in tomato and pepper. *J Exp Bot* 58(14):3841–3852.
- Paran I, Goldman I, Zamir D (1997) QTL analysis of morphological traits in a tomato recombinant inbred line population. *Genome* 40:242–248.
- Rao G-U, Ben Chaim A, Borovsky Y, Paran I (2003) Mapping of yield-related QTL in pepper in an interspecific cross of *Capsicum annuum* and *C. frutescens*. *Theor Appl Genet* 106:1457–1466.
- Tanksley S-D (1993) Mapping polygenes. *Ann Rev Genet* 27:205–233.
- Tanksley S-D, Ganai M-W, Prince J-P, de Vicente M-C, Bonierbale M-W, Broun P, Fulton T-M, Giovannoni J-J, Grandillo S, Martin J-B, Messeguer R, Miller J-C, Miller L, Paterson A-H, Pineda O, Roder M-S, Wing R-A, Wu W, Young N-D (1992) High density molecular linkage maps of the tomato and potato genomes. *Genetics* 132:1141–1160.
- R Development Core Team (2006) R: a language and environment for statistical computing. <http://www.R-project.org>
- Utz H-F, Melchinger A-E, Schön CC (2000) Bias and sampling error of the estimated proportion of genotypic variance explained by quantitative trait loci determined from experimental data in maize using cross validation and validation with independent samples. *Genetics* 154:1839–1849.
- Vales M, Schon C, Capettini F, Chen X, Corey A, Mather D, Mundt C, Richardson K, Sandoval-Islas J, Utz H, Hayes P (2005) Effect of population size on the estimation of QTL: a test using resistance to barley stripe rust. *Theor Appl Genet* 111:1260–1270.
- Van Eck H-J, Jacobs J-M, Stam P, Ton J, Stiekema W-J, Jacobsen E (1994) Multiple alleles for tuber shape in diploid potato detected by qualitative and quantitative genetic analysis using RFLPs. *Genetics* 137:303–309.
- Voorrips R-E (2002) MapChart: software for the graphical presentation of linkage maps and QTL. *J Hered* 93:77–78.
- Zeng Z-B (1994) Precision mapping of quantitative trait loci. *Genetics* 136:1457–1468
- Zygier S, Chaim A-B, Efrati A, Kaluzky G, Borovsky Y, Paran I (2005) QTL mapping for fruit size and shape in chromosomes 2 and 4 in pepper and a comparison of the pepper QTL map with that of tomato. *Theor Appl Genet* 111:437–445.

Table 1: Trait means, standard deviation (SD) and heritabilities for the horticultural traits measured in the parental lines, the F1 hybrid, the RIL population of pepper (n=297) and the sub-populations A (n=141) and B (n=93) issued from selective genotyping.

Trait	code	Yolo Wonder	Criollo de Morelos 334	Significant mean difference among parental values (t test)	F1	F6	Sub-population A	Sub-population B	Heritability
Logarithm of fruit weight	Lfw	5.38±0.12	1.83±0.08	Yes: $p < 10^{-6}$	3.27±0.11	3.33±0.37	3.33±0.41	3.27±0.39	0.96
Fruit length (mm)	Frl	91.47±12.70	58.83±5.22	Yes: $p < 10^{-6}$	81.88±12.63	77.8±14.70	78.05±14.25	76.53±12.985	0.96
Fruit diameter (mm)	Frd	90.43±8.44	18.54±1.56	Yes: $p < 10^{-6}$	37.57±3.36	36.98±6.52	36.98±7.34	36.36±7.18	0.97
Fruit shape	Frs	1.02±0.19	3.17±0.27	Yes: $p < 10^{-6}$	2.20±0.38	2.19±0.57	2.21±0.58	2.55±0.58	0.97
Pericarp thickness (mm)	Pet	7.45±0.66	2.48±0.25	Yes: $p < 10^{-6}$	3.64±0.63	4.06±0.65	4.12±0.66	4.06±0.65	0.92
Number of locules	Nle	4.02±0.98	2.29±0.46	Yes: $p < 10^{-6}$	2.46±0.51	2.85±0.37	2.83±0.40	2.82±0.38	0.84
Pedicle length (mm)	Pel	40.67±6.41	38.23±4.24	No: $p=0.67$	49.94±5.63	47.63±8.17	46.75±6.94	46.37±6.53	0.95
Flowering (days)	Flw	60.96±3.19	53.07±3.07	Yes: $p < 10^{-6}$	52.73±2.51	56.62±2.26	56.61±2.23	56.48±2.21	0.88
Axis length (cm)	Axl	18.01±2.42	22.92±3.67	Yes: $p < 10^{-6}$	25.1±2.7	22.06±3.66	21.96±3.75	21.85±3.37	0.94
Number of leaves	Nle	12.12±1.56	12.50±1.91	No: $p=0.47$	11.26±1.42	11.56±1.47	11.59±1.52	11.44±1.57	0.92
Internode length (cm)	Inl	1.49±0.16	1.85±0.24	Yes: $p < 10^{-6}$	2.25±0.32	1.93±0.28	1.91±0.27	1.93±0.26	0.91
Internode growth time (days)	Int	3.86±0.59	3.09±0.42	Yes: $p < 10^{-6}$	3.39±0.37	3.68±0.48	3.67±0.44	3.71±0.45	0.90
Axis growth speed (mm/day)	Axs	3.93±0.55	6.01±1.00	Yes: $p < 10^{-6}$	6.66±0.68	5.32±0.84	5.82±0.84	5.27±0.74	0.93



Table 2. Pearson correlations between traits assessed in the F6YC RIL population. The traits analysed were Lfw (logarithm of the mean fruit weight), Frl (fruit length), Frd (fruit diameter), Frs (fruit shape), Pet (pericarp thickness), Nlo (number of fruit locules), Pel (pedicel length), Flw (earliness), Axl (axis length), Nle (number of leaves), Inl (mean internode length), Int (mean internode growth time), Axs (axis growth speed). Coloured boxes indicate significant correlations ( $p < 0.05$ ).

Trait	Lfw	Frl	Frd	Frs	Pet	Nlo	Pdl	Flw	Axl	Nle	Inl	Int
Frl	0.41											
Frd	0.87	0.06										
Frs	-0.31	0.69	-0.64									
Pet	0.73	0.06	0.68	-0.43								
Nlo	0.23	-0.29	0.36	-0.46	0.17							
Pdl	0.14	0.40	-0.02	0.30	-0.04	-0.22						
Flw	0.23	0.03	0.26	-0.14	0.20	0.03	0.25					
Axl	-0.09	0.03	-0.11	0.08	-0.03	0.02	0.08	0.26				
Nle	-0.16	-0.11	-0.13	0.01	-0.10	0.06	-0.13	0.34	0.50			
Inl	0.05	0.14	0.00	0.09	0.06	-0.05	0.21	0.01	0.66	-0.30		
Int	0.28	0.12	0.27	-0.09	0.20	-0.05	0.26	0.12	-0.39	-0.88	0.34	
Axs	-0.17	0.03	-0.20	0.14	-0.10	0.01	-0.01	-0.08	0.94	0.40	0.68	-0.45

Table 3: QTLs detected in the whole F5YC RIL population, the sub-populations A and B obtained by selective genotyping. For each trait the LOD significance threshold at  $p=0.05$  determined after 1000 permutations is indicated. The closest marker to the QTL and the direction of the QTL (i.e., which parent contributes positively to the trait) are indicated, along with the LOD value of the QTL, the determination coefficient of the individual QTL and for the whole trait. IM, CIM: QTLs detected only with Interval mapping or Composite interval mapping.

	QTL	Marker	Direction	LOD	Variation explained	Heritability	Additive
					Locus	Trait	
Logarithm of fruit weight Lod>3.08	Lfw3.1	e40/m49_198y	Yolo Wonder	5.14	0.11	0.48	0.12
	Lfw4.1	p17/m32_240c	Yolo Wonder	9.63	0.15		0.15
	Lfw11.1	e41/m61_137c	Yolo Wonder	4.61	0.07		0.1
	Lfw12.1 CIM	e44/m51_263c	Yolo Wonder	4.22	0.05		0.09
	LfwLG15.1	e38/m60_224y	Yolo Wonder	5.03	0.08		0.1
	LfwLG24.1 IM	e41/m54_184y	Yolo Wonder	4.49	0.08		0.11
	LfwLG45.1	Epms_402	Yolo Wonder	4.01	0.08		0.1
Fruit length Lod>3.12	Frl3.1	e36/m52_158y	Criollo de Morelos	7.26	0.1	0.39	-3.9
	Frl4.1	e38/m60_109y	Yolo Wonder	17.16	0.21		6.5
	Frl7.1	e41/m61_140c	Yolo Wonder	5.53	0.1		3.43
	FrlLG22.1	e34/m53_181c	Yolo Wonder	5.99	0.1		3.95
Fruit diameter Lod>3.15	Frd2.1 CIM	e36/m52_116c	Yolo Wonder	4.73	0.12	0.91	2.23
	Frd3.1	p14/m39_221y	Yolo Wonder	3.72	0.11		2.17
	Frd4.1	e41/m48_078y	Yolo Wonder	4.63	0.05		1.46
	Frd8.1 CIM	p11/m49_274c	Yolo Wonder	3.23	0.06		1.38
	Frd10.1	e36/m47_145y	Yolo Wonder	3.32	0.07		1.73
	Frd11.1	e41/m61_137c	Yolo Wonder	11.27	0.13		2.34
	Frd12.1 IM	e44/m51_263c	Yolo Wonder	3.43	0.06		1.65
	FrdLG15.1	e38/m60_224y	Yolo Wonder	7.64	0.1		2.06
	FrdLG17.1	e38/m61_144y	Yolo Wonder	4.74	0.06		1.57
	FrdLG24.1 IM	e41/m54_184y	Yolo Wonder	4.57	0.08		1.85
	FrdLG25.1	p15/m40_091c	Yolo Wonder	6.09	0.07		1.7
	FrdLG37.1	p15/m40_319c	Yolo Wonder	3.94	0.06		1.56
Fruit shape Lod>3.12	Frs2.1 CIM	e36/m52_116c	Criollo de Morelos	3.24	0.04	0.31	-0.11
	Frs3.1	e43/m54_256y	Criollo de Morelos	11.75	0.15		-0.22
	Frs4.1	p14/m39_417y	Yolo Wonder	4.01	0.05		0.11
	Frs4.2CIM	p25/m45_109y	Yolo Wonder	3.83	0.08		0.16
	Frs10.1	e38/m60_117c	Criollo de Morelos	4.68	0.06		-0.14
	Frs11.1	e41/m61_270c	Criollo de Morelos	4.41	0.07		-0.13
	FrsLG17.1	e40/m47_239y	Criollo de Morelos	5.21	0.07		-0.14
	FrsLG25.1	e41/m54_351c	Criollo de Morelos	3.79	0.06		-0.12
Pericarp thickness Lod>3.06	Pet3.1	PG101	Yolo Wonder	5.15	0.14	0.31	0.24
	Pet6.1 IM	p25/m45_185y	Yolo Wonder	3.59	0.05		0.15
	Pet10.1	e36/m47_145y	Yolo Wonder	6.87	0.09		0.19
	Pet11.1 IM	e41/m61_270c	Yolo Wonder	3.69	0.06		0.15
	Pet12.1CIM	e44/m51_263c	Yolo Wonder	3.31	0.04		0.13
	PetLG15.1	e38/m60_224y	Yolo Wonder	7.82	0.1		0.2
	PetLG24.1IM	e41/m54_184y	Yolo Wonder	4.27	0.07		0.18
	PetLG27.1	e40/m49_305c	Yolo Wonder	5.75	0.08		0.18
Number of locules Lod>3.00	Nlo2.1 CIM	e36/m47_146c	Yolo Wonder	3.42	0.05	0.46	0.08
	Nlo8.1	Hpms1_214	Yolo Wonder	3.83	0.06		0.09
	Nlo12.1	e44/m51_263c	Yolo Wonder	4.08	0.05		0.08
	Nlo12.2	e36/m47_237c	Yolo Wonder	4.88	0.1		0.08
	NloLG17.1 IM	e38/m61_144y	Yolo Wonder	4.06	0.08		0.1
	NloLG22.1CIM	e34/m53_181c	Criollo de Morelos	3.13	0.04		-0.07
	NloLG25.1	p15/m40_091c	Yolo Wonder	8.12	0.13		0.12
	NloLG30.1	e44/m51_376c	Yolo Wonder	8.12	0.11		0.12
	NloLG39.1	TntC07y	Yolo Wonder	5.72	0.1		0.1
Pedicel length Lod>3.03	Pel1.1 CIM	e41/m61_199y	Yolo Wonder	3.48	0.04	0.37	1.58
	Pel2.1	e36/m52_116c	Criollo de Morelos	9.04	0.1		-2.61
	Pel3.1 IM	e34/m53_077c	Criollo de Morelos	5.48	0.12		-2.79
	Pel4.1	e41/m48_078y	Yolo Wonder	5.68	0.07		2.05
	Pel12.1	p25/m45_087c	Criollo de Morelos	7.2	0.12		-2.83
	PelLG42.1	p25/m45_335c	Yolo Wonder	6.14	0.08		2.15
Flowering earliness Lod>3.07	Flw1.1	e34/M53_233c	Yolo Wonder	8.04	0.1	0.15	0.72
	Flw2.1	Epms409	Criollo de Morelos	6.64	0.1		-0.7
	Flw4.1 CIM	e38/m61_168c	Yolo Wonder	3.34	0.05		0.49
	FlwLG15.1	e38/m60_224y	Yolo Wonder	4.83	0.07		0.58
	FlwLG17.1	p14/m33_851c	Yolo Wonder	4.19	0.07		0.59
Axis length Lod>3.02	Axl2.1	e36/m47_146c	Criollo de Morelos	4.65	0.07	0.42	-0.96
	Axl6.1 IM	p14/m41_060y	Yolo Wonder	3.29	0.05		0.83
	Axl9.1	e37/m54_92c	Criollo de Morelos	3.83	0.08		-0.98
	AxlLG24.1 IM	Epms376	Yolo Wonder	3.53	0.06		0.88
	AxlLG47.1	p14/m33_311c	Yolo Wonder	6.15	0.1		1.18

Number of leaves Lod>3.17	Nle3.1 IM	e40/m49_198y	Criollo de Morelos	3.29	0.06	0.25	0.92	-0.36
	NleLG38.1 CIM	p17/m32_344c	Criollo de Morelos	3.3	0.04			-0.31
	NleLG45.1 IM	Epms_402	Criollo de Morelos	3.08	0.05			-0.33
	NleLG47.1	p14/m33_311c	Yolo Wonder	6.31	0.11			0.49
Internode length Lod>3.13	Inl1.1	e36/m52_190y	Yolo Wonder	4.22	0.06	0.1	0.91	0.07
	Inl2.1	e36/m47_146c	Criollo de Morelos	6.25	0.11			-0.08
	InlLG28.1 IM	e41/m54_221c	Yolo Wonder	5.1	0.08			0.08
Internode growth time Lod>3.00	Int4.1IM	e42/M48_116y	Yolo Wonder	3.47	0.05	0.07	0.9	0.1
	IntLG47.1	p14/m33_311c	Criollo de Morelos	4.86	0.11			-0.14
Axis growth speed Lod>3.05	Axs2.1 CIM	e36/m47_146c	Criollo de Morelos	6.27	0.1	0.44	0.93	-0.26
	Axs2.2 CIM	p25/m42_268c	Yolo Wonder	3.69	0.05			0.18
	Axs4.1	e38/m61_168c	Criollo de Morelos	4.48	0.07			-0.2
	Axs9.1	CT145	Criollo de Morelos	4.46	0.09			-0.24

## Subpopulation A

QTL		Marker	Direction	LOD	Variation explained	Additive	
					Locus	Trait	
Logarithm of fruit weight Lod>3.03	Lfw3.1	P14/m39_221y	Yolo Wonder	5.14	0.13	0.31	0.12
	Lfw4.1	p17/m32_240c	Yolo Wonder	4.56	0.15		0.16
	LfwLG24.1 CIM	EPMS_376	Yolo Wonder	3.57	0.1		0.13
Fruit length Lod>3.06	Fr14.1	e38/m60_109y	Yolo Wonder	3.86	0.12	0.24	5.7
	Fr1LG22.1	p17/m32_155c	Yolo Wonder	3.86	0.1		5.07
Fruit diameter Lod>3.21	Frd11.1	e41/m61_137c	Yolo Wonder	5.96	0.14	0.36	2.89
	FrdLG15.1	e38/m60_224y	Yolo Wonder	4.55	0.12		2.60
	FrdLG17.1	p14/m33_851c	Yolo Wonder	4.55	0.12		2.57
	FrdLG24.1 IM	p11/m49_197y	Yolo Wonder	3.86	0.12		2.66
Fruit shape Lod>3.18	Frs3.1	E34/m53_077c	Criollo de Morelos	4.1	0.13	0.36	-0.21
	Frs11.1 CIM	e41/m61_270c	Criollo de Morelos	3.3	0.06		-0.15
	FrsLG17.1	e40/m47_239y	Criollo de Morelos	3.37	0.11		-0.19
Pericarp thickness Lod>3.10	Pet3.1 IM	PG101	Yolo Wonder	3.33	0.14	0.16	0.25
	Pet10.1	e36/m47_145y	Yolo Wonder	5.42	0.15		0.24
	PetLG15.1	e38/m60_224y	Yolo Wonder	3.44	0.11		0.21
Number of loges Lod>3.21	NloLG17.1 IM	e38/m61_144y	Yolo Wonder	3.86	0.15	0.22	0.15
	NloLG25.1	p15/m40_091c	Yolo Wonder	8.42	0.21		0.21
Pedicel length Lod>3.18	Pel2.1 CIM	e36/m52_116c	Criollo de Morelos	4.40	0.1	0.19	-2.29
	Pel4.1 IM	p17/m32_240c	Yolo Wonder	3.18	0.1		2.31
	Pel4.2 CIM	E42/m48_116hy	Yolo Wonder	3.32	0.07		1.94
Flowering earliness Lod>3.16	Flw1.1	e34/M53_233c	Yolo Wonder	5.42	0.12	0.29	0.79
	FlwLG15.1	e38/m60_224y	Yolo Wonder	5.61	0.13		0.85
Axis length Lod>3.09	Axl9.1	e37/m54_92c	Criollo de Morelos	3.43	0.12	0.12	-1.3
Number of leaves Lod>3.00	Nle11.1 CIM	EPMS_410	Yolo Wonder	3.20	0.08	0.08	-0.44
Internode growth time Lod>3.18	Int9.1CIM	E44/m61_515c	Criollo de Morelos	4.06	0.09	0.09	-0.14
Axis growth speed Lod>3.12	Axs9.1 CIM	CT145	Criollo de Morelos	3.66	0.13	0.13	-0.31

## Subpopulation B

QTL		Marker	Direction	LOD	Variation explained	Additive	
					Locus	Trait	
Logarithm of fruit weight Lod>3.24	Lfw4.1	P14/m41_648c	Yolo Wonder	4.74	0.17	0.13	0.18
	LfwLG24.1 CIM	EPMS_376	Yolo Wonder	3.27	0.14		0.15
Fruit length Lod>3.16	Fr14.1	p17/m32_240c	Yolo Wonder	7.16	0.2	0.2	6.7
Fruit diameter Lod>3.36	Frd1.1 CIM	E38/m60_221y	Yolo Wonder	4.13	0.1	0.21	2.43
	Frd4.1 CIM	E38/m60_109y	Yolo Wonder	5.74	0.16		3.03
	Frd11.1	e41/m61_137c	Yolo Wonder	6.72	0.17		2.96
Pericarp thickness Lod>3.29	Pet10.1	E38/m60_117c	Yolo Wonder	3.3	0.16	0.16	0.27
Pedicel length Lod>3.28	Pel12.1 IM	E36/m47_237c	Criollo de Morelos	3.51	0.18	0.18	-2.83
Flowering earliness Lod>3.26	Flw1.1CIM	TntC09y	Yolo Wonder	4.31	0.17	0.34	0.93
	FlwLG15.1IM	e38/m60_224y	Yolo Wonder	3.60	0.17		0.93
	FlwLG24.1IM	E41/m54_184y	Yolo Wonder	3.88	0.19		0.97
Internode length Lod>3.27	Inl1.1	C33/m54_327y	Yolo Wonder	4.99	0.12	0.12	0.11
	Inl2.1IM	Ee44/m51_258c	Criollo de Morelos	3.28	0.17		-0.11
Axis growth speed Lod>3.19	Axs2.1	EPMS_497	Criollo de Morelos	3.22	0.12	0.32	-0.25
	Axs2.2CIM	P15/m43_153	Yolo Wonder	3.95	0.14		0.29

Figure 1. Map positions of the significant QTLs on each chromosome or LG.





