

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**MUC1 gene polymorphism in three Nelore lines selected for growth and its association with growth and carcass traits**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/85325> since 2016-10-10T11:19:20Z

*Published version:*

DOI:10.1007/s11033-011-0893-0

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

***MUC1 gene polymorphism in three Nelore lines selected for growth and its association with growth and carcass traits,***

*F.R.P. de Souza, S. Maione, S. Sartore, D. Soglia, V. Spalenza, E. Cauvin, L.R. Martelli,  
M.E. Zerlotti Mercadante, P. Sacchi, L. Galvão de Albuquerque, R. Rasero*

*MOLECULAR BIOLOGY REPORTS (ISSN:1573-4978), pp. 1541-1549. Vol. 39. (2012)*

*DOI: 10.1007/s11033-011-0893-0*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*<http://dx.doi.org/10.1007/s11033-011-0893-0>*

Editorial Manager(tm) for Molecular Biology Reports  
Manuscript Draft

Manuscript Number: MOLE-3236R1

Title: MUC1 gene polymorphism in three Nelore lines selected for growth and its association with growth and carcass traits

Article Type: Manuscript

Keywords: beef cattle; QTL; VNTR; marker assisted selection; zebu

Corresponding Author: Fabio Souza

Corresponding Author's Institution: Universidade Estadual Paulista

First Author: Fabio Souza

Order of Authors: Fabio Souza; Sandra Maione, Dr.; Stefano Sartore, Dr.; Dominga Soglia, Dr.; Veronica Spalenza, Dr.; Elsa Cauvin, Dr.; Lucia Regina Martelli, Dr.; Maria Eugênia Zerlotti Mercadante, Dr.; Paola Sacchi, Dr.; Lucia Galvão Albuquerque, Dr.; Roberto Rasero, Dr.

1 Authors: Fabio Ricardo Pablos de Souza<sup>\*1</sup>, Sandra Maione<sup>§</sup>, Stefano Sartore<sup>§</sup>, Dominga Soglia<sup>§</sup>,  
2 Veronica Spalenza<sup>§</sup>, Elsa Cauvin<sup>§</sup>, Lucia Regina Martelli<sup>\*\*</sup>, Maria Eugênia Zerlotti Mercadante<sup>†</sup>, Paola  
3 Sacchi<sup>§</sup>, Lucia Galvão de Albuquerque<sup>\*</sup>, and Roberto Rasero<sup>§</sup>  
4  
5  
6  
7

8 Title: MUC1 gene polymorphism in three Nelore lines selected for growth and its association  
9 with growth and carcass traits  
10  
11  
12

13 Affiliation:  
14  
15  
16  
17

18 \* Departamento de Zootecnia, Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual  
19 Paulista. Via de Acesso Prof. Paulo Donato Castellane - 14884-900 Jaboticabal (SP), Brazil.  
20  
21

22 \*\* Departamento de Genética, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo. Av.  
23 Bandeirantes 3900 - 14049-900 Ribeirão Preto (SP), Brazil.  
24  
25

26 § Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia, University of Torino. Via Leonardo da  
27 Vinci 44 - 10095 Grugliasco (TO) - Italy  
28  
29

30 † Instituto de Zootecnia, Centro de Pesquisa em Pecuária de Corte. Cx.63. 14160-000 Sertãozinho (SP),  
31 Brazil.  
32  
33  
34  
35

36 Address for correspondence:  
37

38 F. R. P. Souza  
39

40 E-mail: [fabiopablos@hotmail.com](mailto:fabiopablos@hotmail.com)  
41

42 Tel/Fax: 5501632092678  
43

44 <sup>1</sup>Present address:  
45

46 Via de acesso Paulo Donato Castellane s/n. Departamento de Zootecnia, Prédio 2. CEP: 14884-900,  
47 Jaboticabal, SP, Brazil  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 **Summary**  
3  
4  
5

6           The objective of this study was to describe the VNTR polymorphism of the mucin 1 gene  
7 (MUC1) in three Nelore lines selected for yearling weight to determine whether allele and genotype  
8 frequencies of this polymorphism were affected by selection for growth. In addition, the effects of the  
9 polymorphism on growth and carcass traits were evaluated. Birth, weaning and yearling weights, rump  
10 height, Longissimus muscle area, backfat thickness, and rump fat thickness, were analyzed. A total of 295  
11 Nelore heifers from the Beef Cattle Research Center, Instituto de Zootecnia de Sertãozinho, were used,  
12 including 41 of the control line, 102 of the selection line and 152 of the traditional. The selection and  
13 traditional lines comprise animals selected for higher yearling weight, whereas control line animals are  
14 selected for yearling weight close to the average. Five alleles were identified, with allele 1 being the most  
15 frequent in the three lines, especially in the lines selected for higher means for yearling weight.  
16 Heterozygosity was significantly higher in the control line. Association analyses showed significant  
17 effects of allele 1 on birth weight and weaning weight while the allele 3 exert significant effects on  
18 yearling weight and back fat thickness. Despite these findings, application of this marker to marker-  
19 assisted selection requires more consistent results based on the genotyping of a larger number of animals  
20 in order to increase the accuracy of the statistical analyses.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 **Keywords:** beef cattle, QTL, VNTR, marker assisted selection, zebu  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

1  
2  
3  
4 In beef cattle, growth and carcass traits are important economical traits. With improved selection  
5 schemes by marker assisted selection in mind, the approach of candidate gene has been applied by  
6 molecular biologists and animal breeders for identification of molecular markers in livestock. Several  
7 genes with significant associations for these traits has been identified including *AMPD1* and *PPAR $\gamma$*  for  
8 carcass traits, and *NUCB2*, *GAD1* and *SDC1* for growth traits [ 1,2,3,4,5].  
9

10  
11  
12  
13  
14 Epithelial mucins are heavily O-glycosylated proteins found in the mucus layer or at the cell  
15 surface of many epitheliums. They are responsible for the physical properties of mucus gels and are  
16 involved in epithelial cell protection. Thus, the central role played by mucins lies in accommodating the  
17 resident commensal flora and limiting infectious disease of mucosal tissues that presents an enormous  
18 surface area to the exterior environment [6].  
19  
20  
21  
22  
23

24 Like other mucins, mucin 1 (MUC1), whose main function is the protection and lubrication of  
25 epithelial surfaces, is a transmembrane glycoprotein that is co-dominantly expressed on the apical surface  
26 of secretory epithelial cells of the respiratory, digestive, urinary and reproductive tracts of mammals [7, 8,  
27 9]. Additionally, MUC1 has been reported to play a role in cell growth, fetal development, epithelial  
28 regeneration and differentiation, epithelial integrity, tumor genesis, and metastasis [10].  
29  
30  
31  
32  
33

34 Bovine MUC1 is well characterized and is also known as PAS due to its affinity for periodic  
35 acid Schiff reagent. In Danish Holstein cattle, the complete molecule consists of 580 amino acids, with  
36 the 22-amino acid N-terminal sequence comprising the signal peptide. After cleavage of the signal  
37 peptide, the sequence can be divided into three domains: a phosphorylated intracellular domain consisting  
38 of 70 amino acids, a hydrophobic transmembrane domain, and a long extracellular domain. The SEA  
39 domain, comprising 120 amino acids, and the polymorphic VNTR, consisting of tandem repeats of 20  
40 amino acids, are found in the extracellular domain [11]. The VNTR is an important region of the  
41 molecule due to its dense O-glycosylation and its relationship with the adhesive or anti-adhesive role of  
42 MUC1 in the organism.  
43  
44  
45  
46  
47  
48  
49  
50  
51

52 In domestic ruminants, due to the abundant expression of MUC1 in the mammary glands [12],  
53 the VNTR polymorphism of the gene has been characterized mainly in species reared for milk production  
54 such as cattle (*Bos primigenius taurus*), goats (*Capra hircus*) and sheep (*Ovies aries*) [13-16]. In Holstein  
55 cattle, associations of the polymorphism with milk production and composition traits, as well as with  
56  
57  
58  
59  
60

1 traits related to fertility and disease resistance, have been reported [17]. The polymorphism was also  
2 characterized in Zebu beef cattle (*Bos primigenius indicus*) and its effect on growth, fertility and carcass  
3 traits was investigated, but no evidence of an association was found [18-21].  
4

5  
6 In view of the scarcity of association studies between the MUC1 gene polymorphism and traits  
7 of economic interest in beef cattle, the objective of the present investigation was to analyze the VNTR  
8 polymorphism of the MUC1 gene in three Nelore lines selected for yearling weight in order to determine  
9 whether allele and genotype frequencies were affected by selection for growth. In addition, the effects of  
10 the polymorphism on growth and carcass traits of the animals were evaluated.  
11  
12  
13  
14  
15  
16  
17

## 18 **Methods**

### 19 **Animals**

20  
21  
22  
23  
24  
25  
26 A total of 295 Nelore heifers born between 2003 and 2005, which belonged to three lines  
27 selected for growth at the Estação Experimental de Zootecnia de Sertãozinho (EEZS), a research facility  
28 of the Instituto de Zootecnia, located in the northern region of the State of São Paulo, Brazil (21°10' south  
29 latitude and 48°5' west longitude), were used. This region is characterized by a wet tropical climate, with  
30 average annual temperature and rainfall of 24°C and 1,312 mm, respectively. Pastures mainly consist of  
31 *Panicum maximum* and *Brachiaria brizantha*, which are common tropical grasses in Brazil.  
32  
33  
34  
35  
36  
37

38 The animals are part of the Zebu Selection Program started in 1978 and have been allocated to  
39 the following three selection lines since 1980: Nelore selection line (NeS), traditional Nelore line (NeT),  
40 and control Nelore line (NeC). In the NeS and NeT lines, males are selected for increasing weight  
41 adjusted to 378 days of age ( $W_{378}$ ) after performance testing in confinement. Females are selected for  
42 weight adjusted to 550 days of age ( $W_{550}$ ) on pasture. In the NeC line, the animals are selected based on  
43  $W_{378}$  and  $W_{550}$  close to the average (selection differential of zero). The NeC and NeS lines are closed,  
44 whereas the NeT line differs from the NeS line because it eventually received sires from other herds,  
45 including commercial ones. The mean breeding values for yearling weight of animals born in the last four  
46 years (2004 to 2007), corresponding to 5.5 generations selected for growth, are 0.2, 49.5 and 55.3 kg for  
47 the NeC, NeS and NeT lines, respectively [22]. Forty-one heifers of the NeC line, 102 of the NeS line,  
48 and 152 of the NeT line were genotyped.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 Traits studied  
3  
4

5  
6 Birthweight (BW), weaning weight (adjusted to 210 days of age,  $W_{210}$ ), and yearling weight  
7 (adjusted to 550 days of age,  $W_{550}$ ) are obtained as part of the selection program. Yearling height ( $YH_{550}$ )  
8 is obtained at the time of weight recording at 550 days.  
9

10  
11 Longissimus muscle area (LMA), subcutaneous backfat thickness (BF), and rump fat thickness  
12 (RF) were measured by ultrasound at an average age of  $22 \pm 2.5$  months. BF and LMA were measured on  
13 transverse ultrasound images taken between the 12th and 13th ribs over the Longissimus dorsi muscle. RF  
14 was measured at the intersection between the Gluteus medius and Biceps femoris muscles located between  
15 the hook and pin bone. Two ultrasound devices were used depending on the occasion of measurement:  
16 Aloka 500V (Corometrics Medical Systems, Inc., Wallingford, CT, USA) equipped with a linear 3.5-  
17 MHz probe (17.5 cm) (transducer: Aloka Co. Ltd., Tokyo, Japan), and Pie Medical 401347 – Aquila  
18 (Esaote Europe B.V., The Netherlands) equipped with a linear 3.5-MHz probe (18 cm). Vegetable oil and  
19 a standoff were used for capture of the images to guarantee acoustic contact between the linear probe and  
20 body of the animal. The images were saved and subsequently interpreted using the Echo Image Viewer  
21 1.0 software (Pie Medical Equipment B.V., 1996).  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 Genotyping  
37  
38  
39

40 Blood (5 mL) was collected from each animal by puncture of the jugular vein into Vacutainer  
41 tubes containing 7.5 mg EDTA. Genomic DNA was extracted according to the saline standard protocol.  
42

43  
44 Genotyping was performed by the polymerase chain reaction (PCR) using the Qiagen Hot Start  
45 Taq enzyme (Qiagen GmbH, Hilden, Germany). The reaction mixture contained 100 ng genomic DNA,  
46 1X buffer, 1X solution Q, 1.5 mM  $MgCl_2$ , 1 mM dNTPs, 0.5  $\mu$ M of each primer, and 2.5 units of the  
47 enzyme in a volume of 50  $\mu$ L. The P4 - P5 primer pair described by Rasero [13] were used. PCR was  
48 carried out in a GeneAmp PCR System 2400 (Applied Biosystems, Foster City, CA, USA) under the  
49 following conditions: denaturation at 95° C for 15 min, followed by 37 cycles of denaturation at 94° C for  
50 30 s, primer annealing at 60° C for 30 s, and extension at 72° C for 1 min and 30 s. A final extension was  
51 carried out at 72° C for 10 min.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 The amplified fragments were separated by electrophoresis on 1.5% agarose gel. The length of  
2 the alleles was estimated using the GeneRuler™ 100-bp DNA Ladder Plus molecular weight marker  
3  
4 (Fermentas International, Inc., Burlington, Ontario, Canada).  
5  
6

#### 7 8 Statistical analysis 9

10  
11 Allele and genotype frequencies were compared between the three lines by Fisher's exact test  
12 using the population differentiation module of the GENEPOP program, version 3.4 [23]  
13  
14 (<http://genepop.curtin.edu.au/>). A P value  $\leq 0.05$  was considered to be significant.  
15  
16

17  
18 The association analyses were based on the methods described by Byun, Forrest, Lin, and  
19  
20 Hickford [24-27].  
21

22 Association analysis between MUC1 genotypes and the traits studied was performed using the  
23 PROC MIXED procedure of the SAS/STAT 9.1.3 program (SAS Institute, Inc., Cary, NC,USA). The  
24 analysis model included marker genotype (1, ..., 10), contemporary group (selection line and year of birth,  
25  
26 1, ..., 9) and month of birth (September, October, November) as fixed effects, age of dam and age at  
27  
28 recording (only for YH<sub>50</sub> and the carcass traits LMA, BF and RF) as linear covariates, and the random  
29  
30 effect of sire (1, ..., 41). All genotypes were included in the analysis.  
31  
32

33  
34 A second set of analyses was performed in order to evaluate the effect of the presence/absence of  
35  
36 the MUC1 alleles on each trait. The allele presence or absence (coded as 1 or 0, respectively) was  
37  
38 considered in place of the marker genotype. Initially, a "single-allele" model was performed testing the  
39  
40 presence/absence of each allele in a trait. Subsequently, for the alleles that showed  $P < 0.20$  in the "single  
41  
42 allele" model, a "multi-allele" model was performed which included all the alleles.  
43

44 Finally, to analyze the effect of the number of copies of each allele on each trait the number of  
45  
46 copies (0, 1, and 2 copies) of each allele was considered. Only alleles that had homozygous forms  $> 1\%$   
47  
48 of all genotypes were included. Using this approach, each allele was tested separately and, subsequently,  
49  
50 alleles with  $P < 0.20$  were included in a "multi-allele" model with all alleles. To ascertain whether  
51  
52 additive, dominant or recessive effects were present, the significant effects were followed by multiple  
53  
54 pair-wise comparisons (Least Significant Difference tests).  
55

56 For the association analyses all P-values were considered statistically significant when  $P < 0.05$   
57  
58 and trends were considered when  $0.05 < P < 0.10$ .  
59  
60

## Results

The means and standard deviations obtained for each trait according to selection line are shown in [Online resource 1](#). In general, the NeS and NeT lines presented higher means for all traits, except for BF and RF.

Five alleles of different lengths were amplified and named allele 1, allele 2, allele 3, allele 4, and allele 5. Allele 1 was the predominant allele in the three selection lines, followed by allele 4 with a frequency of 0.25 in the NeC line, 0.08 in the NeS line, and 0.07 in the NeT line. In general, alleles 2 and 5 were the least frequent, with allele 2 not being detected in the NeC line. Compared to the selection lines NeS and NeT, the NeC line was characterized by a high frequency of alleles 4 and 5. Genotype 1/1 was the most frequent in all selection lines, followed by genotype 1/4 (Table 1). Although the NeC line presented only five genotype classes, heterozygosity (0.46) was higher than in the NeS line with seven genotype classes (0.16) and in the NeT that presented nine genotype classes (0.25).

Fisher's exact test showed a significant difference in allele and genotype frequencies between the lines selected for higher growth (NeT and NeS) and the control line. Regarding allele frequency, P values of less than 0.001 were observed for comparison between the NeC line and the NeS and NeT lines, whereas no significant difference ( $P = 0.38$ ) was found between the two lines selected for higher growth (NeS and NeT). With respect to genotype frequency, the P values for comparison between the NeC line and the NeS and NeT lines were less than 0.001, whereas no significant difference ( $P = 0.46$ ) was observed between the two lines selected for higher growth (NeS and NeT).

The statistical analysis most frequently applied in association studies in animal breeding revealed no evidence of a significant association between the genotypes and the traits analyzed ([Online Resource 2](#)). The statistical analysis considering the presence/absence of the MUC1 alleles revealed that the alleles 1 and 3 were associated with some economic traits in Nelore cattle (Table 2). Using the single-allele model, the absence of allele 3 was significantly associated with higher mean for  $W_{550}$  ( $P < 0.05$ ). A trend was also observed between the absence of allele 3 and higher mean for BF ( $P < 0.10$ ). Considering the multi-allele model, which included the alleles with  $P < 0.20$  in the single allele model, a significant effect ( $P < 0.05$ ) and a trend ( $P < 0.10$ ) for the allele 1 were found for higher means on BW and  $W_{210}$ , respectively.

1  
2 For the allele 3, changing from the single-allele to the multi-allele model, its effect in  $W_{550}$  became non-  
3 significant whereas the trend between higher means for BF remained.

4 In the third set of analyses the effects of the number of copies of each allele in each trait was  
5 investigated (Table 3). The analysis was performed for those alleles with sufficiently common  
6 homozygous forms (>1% of all genotypes), so only the alleles 1, 3 and 4 were selected. In the single  
7 allele-model the significant effects of allele 1 on BW and of allele 3 on  $W_{550}$  were observed ( $P<0.05$ ).  
8 Considering BW, animals with just one copy of allele 1 presented means significantly higher than animals  
9 with any copy or two copies of this allele. For  $W_{550}$  was observed that animals with no or one copy of  
10 allele 3 presented means significantly higher than animals with two copies of this allele. In the multi-  
11 allele model, which included the alleles with  $P<0.20$  in the single allele model, just the effect of allele 3  
12 on  $W_{550}$  persisted ( $P\leq 0.05$ ).  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

## 24 **Discussion**

25  
26  
27  
28 In humans, MUC1 protein have several isoforms originating from alternative splicing,  
29 including MUC1/transmembrane, MUC1/secreted and MUC1/Y. MUC1/secreted lacks the hydrophobic  
30 transmembrane and cytoplasmic domains required for anchorage to the cell membrane, and MUC1/Y  
31 lacks the tandem repeat domains. The MUC1/transmembrane is the complete peptide expressed as a  
32 continuous polypeptide, however, the molecule is cleaved between the transmembrane and VNTR  
33 domains during its processing in complex Golgi complex although the two parts of the molecule remain  
34 associated via noncovalent forces [6].  
35  
36  
37  
38  
39  
40  
41

42 Like all mucins, the best-described functions of MUC1/transmembrane are those of mucus  
43 layers: hydration, lubrication, protection from proteases, and defense against pathogens. The glycosilation  
44 pattern provides a hydrophilic environment ideal for hydration and lubrication of epithelia. In addition,  
45 the large protein core and dense sugar chains prevent access to the epithelium below. Thus, MUC1 extend  
46 considerably further from the cell surface than many cellular receptors, providing a mechanism that can  
47 disrupt the adhesion of cells and pathogens [28]  
48  
49  
50  
51  
52  
53

54 The gene MUC1 bovine lies in BTA3 and like in humans is formed by seven exons and six  
55 introns. In humans, the promoter of MUC1 gene contains numerous potential binding sites for  
56 transcriptional regulators, among them Sp1, AP1-4, NF-1, NF-B, an E-box, GC boxes, and estrogen and  
57  
58  
59

1 progesterone receptor sites. The VNTR in exon 2 of MUC1 gene is an important region because of the  
2 abundant presence of codons coding for serine and threonine residues in all mammalian species studied to  
3 date. These are the main amino acids that undergo dense O-glycosylation during processing of the protein  
4 in the Golgi complex, permitting steric hindrance that confers a broad functional role to the molecule in  
5 the organism of mammals [6].  
6  
7

8  
9 The VNTR polymorphism of the MUC1 gene has been extensively studied in humans and  
10 domestic animals. In humans, the number of repeats at the VNTR locus ranges from 25 to 125 and this  
11 polymorphism has been associated with severe acne [29], conditions that precede gastric carcinoma [30],  
12 lung adenocarcinoma [31], and infertility due to failure in embryo implantation [32, 33]. In ruminants, the  
13 number of repeats ranges from 10 to 27 in cattle [13, 18], from 30 to 50 in goats [14], and from 19 to 22  
14 bp in sheep [15]. An association between the polymorphism and milk production, health and fertility has  
15 been reported for the Holstein dairy breed (*Bos primigenius taurus*) [17]. The present study described the  
16 pattern of this polymorphism in three lines selected for yearling weight and its association with growth  
17 and carcass traits in Nelore beef heifers (*Bos primigenius indicus*).  
18  
19

20 As reported in previous studies, Nelore is characterized by 5 alleles of the MUC1 gene VNTR  
21 [18, 20, 21]. The allele 1 was the predominant allele in the Nelore breed in both the sample as a whole  
22 and in each selection line, especially in the NeS and NeT lines. However, in contrast to the results  
23 reported so far for Nelore animals, the second most frequent allele in all selection lines was allele 4,  
24 especially in the NeC line. The frequencies of allele 2 and allele 3 were lower than the frequencies  
25 reported in previous studies and these alleles showed a lower frequency than allele 5 in the NeC line.  
26 Alleles 4 and 5 have so far been considered the rarest alleles in the Nelore breed [18,20,21].  
27  
28

29 This is the first study reporting significant associations between the MUC1 gene polymorphism  
30 and economic traits in beef cattle. In previous studies using samples from a commercial Nelore herd  
31 [20,21], this polymorphism was characterized in this breed and its effects on early puberty and expected  
32 progeny differences for growth, carcass and fertility traits were investigated. No evidence of an  
33 association was found.  
34  
35

36 In the present study, the statistical methods commonly used in association studies between  
37 genetic polymorphisms and economic traits showed no significant associations. Therefore, the approach  
38 reported by Hickford [27] and Forrest [25] was used as an alternative for a more complete analysis.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 The presence of allele 1 seems to be favorable for BW and  $W_{210}$  since it showed significant  
2 effects and trends on the multi allele model, respectively. For BW, the analysis of the number of copies  
3 also showed significant effects of this allele, being the heterozygous animals the heavier animals. Weight  
4 at different ages in Nelore and Japanese Black cattle have positive correlation that decreases with the  
5 increasing of the interval between the ages [34,35,36]. This allele showed higher frequencies in the two  
6 lines selected for yearling weight (NeS and NeT - 378 days for sires and 550 days for females),  
7 suggesting that this selection scheme is favoring the allele 1 in this lines and indirectly increasing the  
8 means for BW and  $W_{210}$  in the animals.  
9

10 The allele 3 seems unfavorable for  $W_{550}$  and BF since its presence was associated with low  
11 means for these traits in the single allele model. In the multi allele model just the trend for BF remained  
12 for this allele. Smith [37] related low correlation (0.10) between slaughter weight and BF in Brahman  
13 cattle. Similarly, Yokoo [38], described a correlation close to zero (0.06) between  $W_{550}$  and BF in Nelore  
14 cattle, thus, these effects on  $W_{550}$  and BF cannot be attributed to genetic correlation. Considering the  
15 analysis of the number of copies, the allele 3 is significative in both analysis (single and multi allele  
16 models) for  $W_{550}$ , being animals with any or one copy heavier than animals with two copies. These results  
17 suggests that this allele have recessive effects for  $W_{550}$ , although caution is needed in drawing this  
18 conclusion, as the number of heifers carrying allele 3, especially in a homozygous genotype, is low.  
19 Although the frequency of the allele 3 in the line NeC is low its effect on  $W_{550}$  may explain the lower  
20 frequencies of the allele 3 in the lines NeS and NeT, which the females are selected considering higher  
21 means for  $W_{550}$ , but it is also necessary to consider the low frequency of this allele in other studies with  
22 Nelore cattle and the probability of its low frequency be a characteristic of the herd that initiated this three  
23 selection lines in the EEZS at 1978 [18, 20, 21].  
24

25 The main explanation between these effects and MUC1 variation could lie in the protection and  
26 lubrication of epithelial tissues. [The ability of bacteria to colonise the intestinal tract and initiate  
27 pathogenesis is intimately associated with their ability to bind receptors displayed on the surface of  
28 digestive epithelial cells \[39\].](#) The variations of MUC1 may influence the health of digestive tract of  
29 calves and mature animals permitting weight gain and to deposit fat. Thus, MUC1 acts as a barrier against  
30 opportunistic infections by some bacterial strains [9,16,40]. Another explanation is that the alleles 1 and 3  
31 are in linkage disequilibrium with another DNA variation that influences directly the traits under study.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Indeed, there is evidence suggesting the existence of five QTLs in *Bos primigenius taurus* X *Bos primigenius indicus* crosses with additive effects on BW on chromosome 3, where the MUC1 gene is located [41]. Beever [42] also reported the existence of a QTL for weaning weight on this chromosome in nine half-sib families of sires originating from the European breeds Simmental, Gelbvieh, or South Devon. In addition to the QTL for BW, Casas [43] also found QTLs for subcutaneous fat thickness and for marbling on chromosome 3.

In conclusion, the results showed a high frequency of allele 1 in three selection lines, especially in the lines NeS and NeT selected for higher yearling weight. The allele 1 was associated with higher means for BW and  $W_{210}$ , while the allele 3 exert significant unfavorable on  $W_{550}$  and BF. However, application of this marker to marker-assisted selection requires more consistent results based on the genotyping of a larger number of animals in order to increase the number of representatives in each genotype class and, consequently, the accuracy of the statistical analyses.

## References

1. He H, Liu X, Gu Y, Liu Y (2010) A novel 18-bp deletion mutation of the AMPD1 gene affects carcass traits in Qinchuan cattle. *Mol Biol Rep* 37:3945-3949. doi: 10.1007/s11033-010-0052-z
2. Fan YY, Zan LS, Fu CZ, Tian WQ, Wang HB, Liu YY, Xin YP (2011) Three novel SNPs in the coding region of PPAR $\gamma$  gene and their associations with meat quality traits in cattle. *Mol Biol Rep.* 38:131-137. doi: 10.1007/s11033-010-0086-2
3. Li F, Chen H, Lei CZ, Ren G, Wang J, Li ZJ, Wang JQ (2010) Novel SNPs of the bovine NUCB2 gene and their association with growth traits in three native Chinese cattle breeds. *Mol Biol Rep* 37:541-546. doi: 10.1007/s11033-009-9732-y
4. Li F, Chen H, Lei CZ, Ren G, Wang J, Li ZJ, Wang JQ (2010) Novel SNPs of the bovine GAD1/gad67 gene and their association with growth traits in three native Chinese cattle breeds. *Mol Biol Rep* 37:501-505. doi: 10.1007/s11033-009-9699-8
5. Sun J, Jin Q, Zhang C, Fang X, Gu C, Lei C, Wang J, Chen H (2011) Polymorphisms in the bovine ghrelin precursor (GHRL) and Syndecan-1 (SDC1) genes that are associated with growth traits in cattle. *Mol Biol Rep* 38:3153-3160. doi: 10.1007/s11033-010-9986-4

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
6. Linden SK, Sutton P, Karlsson NG, Korolik V, McGuckin MA (2008) Mucins in the mucosal barrier to infection. *Mucosal Immunol* 1:183-197. doi: 10.1038/mi.2008.5
  7. Gendler SJ (2001) MUC1, the renaissance molecule. *J Mammary Gland Biol Neoplasia* 6: 339–353. doi: 10.1023/A:1011379725811
  8. Zhang J, Zhang X, Zhu Y, Chen Z, Xu Z, Miao Y (2010) Transcriptional regulation of human mucin gene MUC4 in pancreatic cancer cells. *Mol Biol Rep* 6:2797-2802. doi: 10.1007/s11033-009-9826-6
  9. Ji H, Ren J, Yan X, Huang X, Zhang B, Zhang Z, Huang L (2011) The porcine MUC20 gene: molecular characterization and its association with susceptibility to enterotoxigenic *Escherichia coli* F4ab/ac. *Mol Biol Rep* 3:1593-1601. doi: 10.1007/s11033-010-0268-y
  10. Moniaux N, Escande F, Porchet N, Aubert JP, Batra SK (2001) Structural organization and classification of the human mucin genes. *Front Biosci* 6: 1192–1206.
  11. Pallesen LT, Andersen MH, Nielsen RL, Berglund L, Petersen TE, Rasmussen LK, Rasmussen JT (2001) Purification of MUC1 from bovine milk-fat globules and characterization of a corresponding full-length cDNA clone. *J Dairy Sci* 12: 2591–2598. doi: 10.3168/jds.S0022-0302(01)74713-3
  12. Patton S, Gendler SJ, Spicer AP (1995) The epithelial mucin, MUC1, of milk, mammary gland and other tissues. *Biochim Biophys Acta* 1241: 407–424.
  13. Rasero R, Sacchi P, Rosati S, Cauvin E, Maione S (2002) Molecular analysis of the length polymorphic MUC1 gene in cattle. *J Anim Breed Genet* 119: 342–349. doi: 10.1046/j.1439-0388.2002.00351.x
  14. Sacchi P, Caroli A, Cauvin E, Maione S, Sartore S, Soglia D, Rasero R (2004) Analysis of the MUC1 gene and its polymorphism in *Capra hircus*. *J Dairy Sci* 87: 3017–3021. doi: 10.3168/jds.S0022-0302(04)73433-5
  15. Rasero R, Bianchi L, Cauvin E, Maione S, Sartore S, Soglia D, Sacchi P (2007) Analysis of the sheep MUC1 gene: Structure of the repetitive region and polymorphism. *J Dairy Sci* 90: 1024-1028. doi: 10.3168/jds.S0022-0302(07)71588-6
  16. Sando L, Pearson R, Gray C, Parker P, Hawken R, Thomson PC, Meadows JR, Kongsuwan K, Smith S, Tellam RL (2009) Bovine Muc1 is a highly polymorphic gene encoding an extensively glycosylated mucin that binds bacteria. *J Dairy Sci* 92: 5276-5291. doi: 10.3168/jds.2009-2216

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
17. Hens JR, Rogers GW, Huott ML, Patton S (1995) Association of the epithelial mucin, PAS-1, with yield, health and reproductive traits in Holstein dairy cows. *J Dairy Sci* 78: 2473–2480. doi: 10.3168/jds.S0022-0302(95)76875-8
  18. Souza FR, Dentillo DB, Meola J, Biase FH, Andréa MV, Vozzi PA, Lôbo RB, Martelli LR (2007) The polymorphism in MUC1 gene in Nelore cattle. *J Anim Breed Genet* 124: 42-46. doi: 10.1111/j.1439-0388.2007.00628.x
  19. Hiendleder S, Lewalski H, Janke A (2008) Complete mitochondrial genomes of *Bos taurus* and *Bos indicus* provide new insights into intra-species variation, taxonomy and domestication. *Cytogenet Genome Res* 120: 150-156. doi: 10.1159/000118756
  20. Souza FRP, Boligon AA, Baldi F, Mercadante, MEZ, Vila RA, Galerani MAV, Lôbo RB, Martelli LR (2009) MUC1 VNTR polymorphism is not associated with early puberty in Nelore cattle (*Bos primigenius indicus*). *Anim Reprod* 6: 545-549.
  21. Souza FR, Vozzi PA, Vila RA, Boligon AA, Galerani MAV, Lôbo RB, Martelli LR (2010) Association between MUC1 gene polymorphism and expected progeny differences in Nelore cattle (*Bos primigenius indicus*). *Genet Mol Biol* 33: 68-70. doi: 10.1590/S1415-47572009005000098
  22. Razook A G, Mercadante M E Z (2007). Ganhos de produtividade com o uso de touros provados. In: F. A. P. Santos (ed) *Requisitos de qualidade na bovinocultura de corte*, ed. FEALQ, Piracicaba, SP, 93-114.
  23. Raymond M, Rousset F (1995) An exact test for population differentiation. *Evolution* 49: 1280-1283.
  24. Byun S O, Zhou H, Frampton C M, Hickford J G H (2010) No association between variation in the ovine calpastatin gene and either longevity or fertility in sheep. *Anim Genet* 41: 223-224. doi: 10.1111/j.1365-2052.2009.01982.x
  25. Forrest RH, Itenge-Mweza TO, McKenzie GW, Zhou H, Frampton CM, Hickford JG (2009) Polymorphism of the ovine beta3-adrenergic receptor gene (ADRB3) and its association with wool mean staple strength and yield. *Anim Genet* 40: 958-962. doi: 10.1111/j.1365-2052.2009.01926.x
  26. Lin L, Flisikowski K, Schwarzenbacher H, Scharfe M, Severitt S, Blöcker H, Fries R (2009) Characterization of the porcine AMPK alpha 2 catalytic subunit gene (PRKAA2): genomic



- structure, polymorphism detection and association study. *Anim Genet* 41: 203–207. doi:  
10.1111/j.1365-2052.2009.01971.x
27. Hickford JG, Forrest RH, Zhou H, Fang Q, Han J, Frampton CM, Horrell AL (2010) Polymorphisms in the ovine myostatin gene (MSTN) and their association with growth and carcass traits in New Zealand Romney sheep. *Anim Genet* 41: 64-72. doi: 10.1111/j.1365-2052.2009.01965.x
28. Hatstrup CL, Gendler SJ (2008) Structure and function of the cell surface (tethered) mucins. *Annu Rev Physiol* 70:431-457. doi: 10.1146/annurev.physiol.70.113006.100659
29. Ando I, Kukita A, Soma G, Hino H (1998) A large number of tandem repeats in the polymorphic epithelial mucin gene is associated with severe acne. *J Dermatol* 25: 150-152.
30. Carvalho F, Seruca R, David L, Amorim A, Seixas M, Bennett E, Clausen H, Sobrinho-Simões M (1997) MUC1 gene polymorphism and gastric cancer - An epidemiological study. *Glycoconj J* 14: 107-111. doi: 10.1023/A:1018573201347
31. Mitsuta K, Yokoyama A, Kondo K, Nakajima M, Arita K, Kohno N (2005) Polymorphism of the MUC1 mucin gene is associated with susceptibility to lung adenocarcinoma and poor prognosis. *Oncol Rep* 14: 185-189.
32. Goulart LR, Vieira GS, Martelli L, Inácio J, Goulart IM, Franco JG Jr (2004) Is MUC1 polymorphism associated with female infertility? *Reprod Biomed Online* 8: 477-482.
33. Horne AW, White JO, Margara RA, Williams R, Winston RM, Lalani E (2001) MUC1: a genetic susceptibility to infertility? *The Lancet* 357: 1336-1337. doi: 10.1016/S0140-6736(00)04502-5
34. Nobre PR, Misztal I, Tsuruta S, Bertrand JK, Silva LO, Lopes PS (2003) Analyses of growth curves of nellore cattle by multiple-trait and random regression models. *J Anim Sci* 81: 918-26.
35. Aziz MA, Nishida S, Suzuki K, Nishida A (2005) Estimation of direct and maternal genetic and permanent environmental effects for weights from birth to 356 days of age in a herd of Japanese Black cattle using random regression. *J Anim Sci* 83: 519-530.
36. Boligon AA, Mercadante ME, Forni S, Lôbo RB, Albuquerque LG (2010) Covariance functions for body weight from birth to maturity in Nellore cows. *J Anim Sci* 88: 849-59. doi:10.2527/jas.2008-1511

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42
37. Smith T, Domingue JD, Paschal JC, Franke DE, Bidner TD, Whipple G (2007) Genetic parameters for growth and carcass traits of Brahman steers. *J Anim Sci* 85: 1377–1384. doi: 10.2527/jas.2006-653
38. Yokoo MJ, Lobo RB, Araujo FR, Bezerra LA, Sainz RD, Albuquerque LG (2010) Genetic associations between carcass traits measured by real-time ultrasound and scrotal circumference and growth traits in Nelore cattle. *J Anim Sci* 88: 52-58. doi: 10.2527/jas.2008-1028
39. Parker P, Sando L, Pearson R, Kongsuwan K, Tellam RL, Smith S (2010) Bovine Muc1 inhibits binding of enteric bacteria to Caco-2 cells. *Glycoconj J.* 27:89-97. doi: 10.1007/s10719-009-9269-2
40. Lindén SK, Sheng YH, Every AL, Miles KM, Skoog EC, Florin TH, Sutton P, McGuckin MA (2009) MUC1 limits *Helicobacter pylori* infection both by steric hindrance and by acting as a releasable decoy. *PLOS Pathog* 10: e1000617. doi: 10.1371/journal.ppat.1000617
41. Kim JJ, Farnir F, Savell J, Taylor JF (2003) Detection of quantitative trait loci for growth and beef carcass fatness traits in a cross between *Bos taurus* (Angus) and *Bos indicus* (Brahman) cattle. *J Anim Sci* 81: 1933-1942.
42. Beever J E, Da Y, Lewin H A (1997) A genome scan for QTL affecting growth traits in beef cattle. *Plant & Animal Genome V Conference*, San Diego, CA, p 295.
43. Casas E, Shackelford SD, Keele JW, Koohmaraie M, Smith TP, Stone RT (2003) Detection of quantitative trait loci for growth and carcass composition in cattle. *J Animal Sci* 81: 2976-2983.

#### 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65

**Acknowledgements**

This work was supported by Fondazione CRT – World Wide Style, Torino, Italy, and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). We are indebted to the Instituto de Zootecnia for providing the cattle blood samples and the data set used in this study.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table 1. Allele and genotype frequencies of the MUC1 polymorphism in the total sample and each Nelore selection line.

	Total		NeC <sup>1</sup> (N = 41)		NeS <sup>1</sup> (N = 102)		NeT <sup>1</sup> (N = 152)								
	Allele frequency	Genotype frequency	Allele frequency	Genotype frequency	Allele frequency	Genotype frequency	Allele frequency	Genotype frequency							
1	0.82	1/1	0.71	1	0.68	1/1	0.48	1	0.87	1/1	0.80	1	0.83	1/1	0.71
2	0.02	1/2	0.02	2	0	1/3	0.05	2	0.01	2/2	0.01	2	0.03	1/2	0.04
3	0.05	2/2	0.01	3	0.03	1/4	0.34	3	0.03	1/3	0.05	3	0.06	2/2	0.01
4	0.10	1/3	0.06	4	0.25	4/4	0.06	4	0.08	3/3	0.01	4	0.07	1/3	0.09
5	0.01	2/3	0.01	5	0.04	4/5	0.07	5	0.01	1/4	0.10	5	0.01	2/3	0.01
		3/3	0.02							4/4	0.02			3/3	0.01
		1/4	0.12							4/5	0.01			1/4	0.09
		4/4	0.03											4/4	0.02
		1/5	0.02											1/5	0.02
		4/5	0.01												

<sup>1</sup>NeC: control line; NeS: selection line; NeT: traditional line.

Table 2. Associations of MUC1 alleles with growth and carcass traits in Nellore cattle.

Trait <sup>1</sup>	Allele	Other alleles in the Model	Mean±Standard Error				Pvalue
			Allele absent	N	Allele present	N	
BW	1	None	27.77±0.90	18	29.13±0.36	277	0.13
	2	None	29.04±0.36	286	29.11±1.25	9	0.95
	3	None	29.02±0.36	271	29.32±0.80	24	0.69
	4	None	28.88±0.38	246	29.59±0.59	49	0.24
	5	None	29.05±0.36	288	28.40±1.43	7	0.64
	<b>1</b>	<b>2,3,4 and 5</b>	<b>28.11±1.15</b>	<b>18</b>	<b>30.41±1.21</b>	<b>277</b>	<b>0.02</b>
W <sub>210</sub>	1	None	169.86±5.16	18	176.94±1.61	276	0.18
	2	None	176.33±1.57	285	179.09±7.44	9	0.71
	3	None	176.54±1.59	270	174.11±4.54	24	0.59
	4	None	176.58±1.81	245	175.78±3.13	49	0.82
	5	None	176.23±1.58	287	181.74±8.37	7	0.51
	<b>1</b>	<b>2,3,4 and 5</b>	<b>174.60±6.6</b>	<b>18</b>	<b>185.22±7.19</b>	<b>277</b>	<b>0.09</b>
W <sub>550</sub>	1	None	287.64±6.12	18	293.51±2.05	276	0.34
	2	None	292.86±2.02	285	303.12±8.67	9	0.23
	<b>3</b>	<b>None</b>	<b>293.92±1.95</b>	<b>270</b>	<b>282.47±5.31</b>	<b>24</b>	<b>0.03</b>
	4	None	292.34±2.24	245	295.35±3.79	49	0.48
	5	None	292.85±2.00	288	302.68±10.69	6	0.36
	3	1,2,4 and 5	303.49±6.91	270	294.69±8.90	24	0.11
H <sub>550</sub>	1	None	130.61±0.85	18	131.77±0.31	276	0.17
	2	None	131.67±0.31	285	132.39±1.18	9	0.53
	3	None	131.75±0.31	270	130.85±0.75	24	0.22
	4	None	131.70±0.34	245	131.63±0.54	49	0.90
	5	None	131.70±0.31	288	131.06±1.47	6	0.66
	1	2,3,4 and 5	130.73±1.11	18	132.05±1.20	276	0.19
LMA	1	None	45.53±1.29	18	46.55±0.53	267	0.43
	2	None	46.42±0.52	276	48.23±1.82	9	0.31
	3	None	46.51±0.52	261	45.70±1.16	24	0.47
	4	None	46.47±0.57	236	46.42±0.82	49	0.95
	5	None	46.45±0.52	279	46.54±2.22	6	0.96
BF	1	None	2.08±0.28	18	1.89±0.11	267	0.50
	2	None	1.90±0.11	276	2.33±0.40	9	0.26
	<b>3</b>	<b>None</b>	<b>1.94±0.11</b>	<b>261</b>	<b>1.53±0.25</b>	<b>24</b>	<b>0.09</b>

	4	None	1.87±0.12	236	2.00±0.18	49	0.52
	5	None	1.90±0.11	279	2.04±0.48	6	0.77
	<b>3</b>	<b>1,2,4 and 5</b>	<b>2.20±0.32</b>	<b>261</b>	<b>1.78±0.41</b>	<b>24</b>	<b>0.10</b>
RF	1	None	3.79±0.37	18	3.73±0.16	268	0.86
	2	None	3.72±0.15	277	4.53±0.52	9	0.11
	3	None	3.75±0.15	262	3.56±0.33	24	0.56
	4	None	3.74±0.17	237	3.71±0.24	49	0.92
	5	None	3.74±0.15	280	3.34±0.64	6	0.53
	2	1,3,4 and 5	3.45±0.37	277	4.24±0.65	9	0.14

<sup>1</sup>BW: birthweight; W<sub>210</sub>: weaning weight; W<sub>550</sub>: yearling weight; H<sub>550</sub>: yearling height; LMA: Longissimus muscle area; BF: backfat thickness; RF: rump fat thickness.

Table 3. Nominal adjusted P values, least square means and standard errors obtained for the effects of the number of copies of each allele of the MUC1 gene polymorphism on growth and carcass traits.

Trait <sup>1</sup>	Allele	Other alleles in the Model	Mean ± Std Error						Pvalue
			0 copy	N	1 copy	N	2 copies	N	
BW	<b>1</b>	None	<b>27.52<sup>A</sup>±0.90</b>	<b>18</b>	<b>29.72<sup>B</sup>±0.40</b>	<b>66</b>	<b>28.75<sup>A</sup>±0.37</b>	<b>211</b>	<b>0.01</b>
	3	None	29.02±0.36	271	29.64±0.85	21	27.22±2.06	3	0.50
	4	None	28.87±0.38	246	30.00±0.64	41	27.77±1.27	8	0.14
	1	2,3,4 and 5	27.80±1.11	18	29.48±1.35	66	28.77±1.82	211	0.49
	4	1,2,3 and 5	28.53 <sup>A</sup> ±0.79	246	29.15 <sup>A</sup> ±1.44	41	28.36 <sup>A</sup> ±2.16	8	0.84
W <sub>210</sub>	1	None	169.87±5.01	18	177.07±1.94	66	176.41±1.72	211	0.40
	3	None	176.54±1.59	270	176.48±4.85	21	158.14±12.42	3	0.33
	4	None	176.51±1.81	245	177.54±3.47	41	167.65±7.61	8	0.49
W <sub>550</sub>	1	None	287.40±6.03	18	294.73±2.53	65	292.74±2.27	211	0.46
	<b>3</b>	None	<b>293.90<sup>A</sup>±1.96</b>	<b>270</b>	<b>285.26<sup>A</sup>±5.67</b>	<b>21</b>	<b>264.01<sup>B</sup>±14.32</b>	<b>3</b>	<b>0.03</b>
	4	None	292.30±2.25	245	296.22±4.18	41	291.28±8.91	8	0.68
	<b>3</b>	1,2,4 and 5	<b>293.57<sup>A</sup>± 3.41</b>	<b>270</b>	<b>275.57<sup>A</sup>±10.23</b>	<b>21</b>	<b>257.25<sup>B</sup>±18.70</b>	<b>3</b>	<b>0.05</b>
H <sub>550</sub>	1	None	130.42±0.87	18	131.92±0.39	65	131.53±0.36	211	0.21
	3	None	131.75±0.31	270	131.06±0.80	21	129.44±1.96	3	0.35
	4	None	131.70±0.34	245	131.73±0.59	41	131.18±1.22	8	0.91
LMA	1	None	45.35±1.25	18	46.31±0.54	65	46.14±0.51	202	0.76
	3	None	46.53±0.52	261	46.30±1.24	21	42.06±3.00	3	0.32
	4	None	46.46±0.57	236	46.16±0.91	41	47.51±1.85	8	0.80
BF	1	None	2.08±0.27	18	1.86±0.11	65	1.87±0.11	202	0.75
	3	None	1.94±0.11	261	1.51±0.27	21	1.67±0.65	3	0.24
	4	None	1.87±0.12	236	1.92±0.19	41	2.35±0.40	8	0.51
RF	1	None	3.78±0.36	18	3.66±0.16	65	3.71±0.15	203	0.93
	3	None	3.75±0.15	262	3.54±0.36	22	3.70±0.86	3	0.83
	4	None	3.74±0.17	237	3.65±0.26	41	3.99±0.53	8	0.84

<sup>1</sup>BW: birthweight; W<sub>210</sub>: weaning weight; W<sub>550</sub>: yearling weight; H<sub>550</sub>: yearling height; LMA: Longissimus muscle area; BF: backfat thickness; RF: rump fat thickness. For the significant effects, values followed by different letters in the same row differ significantly (p<0.05)

## Supplementary Material

[Click here to download Supplementary Material: Souza online resource MBR.doc](#)