



since

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effect of culture media and pH on the biomass production and biocontrol efficacy of a Metschnikowia pulcherrima strain to be used as a biofungicide for postharvest disease control

This is the author's manuscript

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/75208

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

1	
2	
5 4	
5	This is an author version of the contribution published on:
6	Questa è la versione dell'autore dell'opera:
7	[Spadaro D., Ciavorella A., Zhang D., Garibaldi A., Gullino M.L. (2010) –
8	Effect of culture media and pH on the biomass production and biocontrol
9	efficacy of a Metschnikowia pulcherrima strain to be used as a
10	biofungicide for postharvest disease control. Canadian Journal of
11	Microbiology, 56 (2), 128-137. DOI: 10.1139/W09-117]
12	
13	The definitive version is available at:
14	La versione definitiva è disponibile alla URL:
15	[http://www.nrcresearchpress.com/toc/cjm/56/2]
16	
17	

18	Effect of culture media and pH on the biomass production and biocontrol efficacy
19	of a Metschnikowia pulcherrima strain to be used as a biofungicide for postharvest
20	disease control
21	
22	Spadaro D. ¹² , Ciavorella A. ² , Dianpeng Z. ² , Garibaldi A. ² , Gullino M.L. ²
23	
24	¹ DiVaPRA – Plant Pathology, University of Turin, Grugliasco (TO), Italy. ² Centre of
25	Competence for the Innovation in the Agro-environmental Sector (AGROINNOVA),
26	University of Turin, Grugliasco (TO), Italy.
27	
28	Correspondence
29	Davide Spadaro, Di.Va.P.R.A Plant Pathology and AGROINNOVA - Centre of
30	Competence for the Innovation in the Agro-environmental Sector, University of Turin,
31	Via L. da Vinci 44, I-10095 Grugliasco (TO), Italy.
32	E-mail: <u>davide.spadaro@unito.it</u>
33	Telephone: +39-011-6708942
34	Fax: +39-011-6709307

36 Abstract

37

38 Few strains of Metschnikowia pulcherrima are under development for control of 39 postharvest pathogens on fruit. A substrate was developed to optimize the biomass 40 production M. pulcherrima strain BIO126. Different complex nutrient sources, with or 41 without pH control, were tested. Growth in Yeast Extract provided at concentrations equal to or higher than 30 g l⁻¹ resulted in the highest biomass. The addition of two 42 carbon sources, D-Mannitol and L-Sorbose at 5 g l⁻¹ each, significantly improved the 43 44 yeast growth. Initial pH values of the medium ranging from 5.0 to 7.5 permitted the 45 highest growth of the yeast. A combination of Yeast Extract, D-Mannitol and L-Sorbose 46 (YEMS), probably with diauxic utilization, showed synergistic effect, widening the exponential phase (the maximum specific growth rate was 0.45 h⁻¹), and increasing the 47 final cell number (1.5 x 10^9 cells ml⁻¹) and dry biomass (6.0 g l⁻¹) in well controlled 48 49 batch fermentation. In efficacy trials on 'Golden Delicious' apples, the microorganism 50 grown in YEMS effectively reduced incidence and severity of B. cinerea (51.1% and 51 70.8%) and P. expansum (41.7% and 14.0%). Also on 'Gala' apples, the best reduction 52 of grey and blue mould incidence was obtained with cells grown in YEMS (58.1% and 50.5%, respectively). 53

54

55 Keywords: batch fermentation, biological control, pH effect, postharvest, yeast.

- 56
- 57

58 **1. Introduction**

59 Several microorganisms have been evaluated to obtain microbial based products useful 60 in agriculture, such as biofertilisers or biopesticides. Among the biofungicides, during 61 the last twenty years, several yeast have been widely investigated against postharvest 62 fungal pathogens of different host species (Janisiewicz and Korsten 2002; Spadaro and 63 Gullino 2004; Wilson and Wisnieswki 1994).

64 Among the microorganisms under development, there are few strains of Metschnikowia 65 pulcherrima antagonistic against fungi causing postharvest decay of fruit. Some strains 66 are effective against Botrytis cinerea, Penicillium expansum, or Alternaria alternata of 67 apples (Janisiewicz et al. 2001; Piano et al. 1997; Spadaro et al. 2002; 2008), other 68 strains were selected against *Penicillium digitatum* on grapefruit, B. cinerea, Rhizopus 69 stolonifer, and Aspergillus niger on table grape, or B. cinerea and R. stolonifer on 70 cherry tomato (Schena et al. 2000). Epiphytic isolates reduced A. carbonarius and A. 71 niger colonization on grapes (Bleve et al. 2006). Moreover, a strain of M. pulcherrima proved effective in preventing the growth or survival of food-borne human pathogens, 72 73 such as Listeria monocytogenes or Salmonella enterica, on fresh-cut apple tissue 74 (Leverentz et al. 2006). M. pulcherrima could act through competition for iron 75 (Saravanakumar et al. 2008) or production of hydrolases, such as chitinases and 76 glucanases (Saravanakumar et al. 2009).

The efficacy of many antagonists of wound pathogens is directly related to the number of antagonist propagules applied (Hofstein et al. 1994). Sinigaglia et al. (1998) found that the antagonistic effects of isolates of four yeast species, including *M. pulcherrima*, against *Penicillium glabrum* were more pronounced at high cell concentrations. Moreover, an increasing efficacy in reducing the germination of *B. cinerea* was

demonstrated by increasing the number of yeast cells applied (Spadaro et al. 2002). A simple way to increase the effectiveness of such biofungicides is the application of a higher number of cells. Mass production of yeast cells is an essential step in the commercialization of a biocontrol agent. A rapid, efficient and cheap mass production of yeast antagonists, generally by liquid fermentation, is one of the key issues to achieve the commercial use of the biofungicide (Wraight et al. 2001).

88 Due to the recent interest over the use of the yeast species *M. pulcherrima* as biocontrol 89 agent, efforts must be intensified to produce M. pulcherrima in a laboratory scale 90 fermenter to provide relevant information for the scale-up production. Operating 91 conditions (aeration, agitation, pH and temperature) as well as medium constituents may 92 affect the quality and quantity of the tested microorganisms. To increase the biomass 93 production of an antagonistic yeast on a laboratory scale, the optimization of the growth 94 conditions, using different complex nutrient sources, is essential. The culture media can 95 greatly influence the efficacy of the biocontrol agents (Wraight et al. 2001). The aim of 96 developing a substrate for laboratory purposes is to optimize the biomass production, to 97 find optimal conditions for stabilization and formulation, and to develop a quality 98 control system. To scale-up a laboratory fermentation process to an industrial level, it is fundamental to find cheap nutrient sources, generally industrial by-products, with 99 100 nutritional values similar to the laboratory standardized media.

In this work, the influence of different complex nutrient media on the growth of *M*. *pulcherrima* strain BIO126 in 5 L batch fermentation was considered. The aim of the research was to find which sources provided the highest biomass production (as number of living cells and as dry biomass) of the antagonistic yeast and what was the optimal concentration for the identified sources. The experiments were carried out first in shake 106 flasks as a preliminary screening, and then in a 5-liter fermenter, optimizing aeration, 107 temperature and pH. Finally, biological control assays were used to test the efficacy of 108 the yeast cells produced through well controlled batch fermentation.

109

110 **2. Materials and Methods**

111 2.1. Microorganism

Metschnikowia pulcherrima (Pitt) M.W. Miller strain BIO126 was isolated from the carposphere of a 'Golden Delicious' apple harvested from an unsprayed orchard located in Piedmont, Northern Italy (Spadaro et al. 2002). The strain was stored as a cell suspension in 20% V/V glycerol at -80°C at the Microorganism Culture Collection of the Centre of Competence for the Innovation in the Agro-environmental Sector of the University of Torino (Italy). The strain was deposited within the American Type Culture Collection on June 19, 2007 with deposit designation PTA-8486.

119

120 2.2. Inoculum preparation

121 The yeast inoculum was prepared by subculturing in Yeast extract-Peptone-Dextrose 122 (YPD) [10 g Γ^1 granulated yeast extract (Merck, Darmstadt, Germany); 20 g Γ^1 triptone-123 peptone of casein (Difco, Detroit, MI, USA); 20 g Γ^1 D(+)-glucose monohydrate 124 (Merck)] on a rotary shaker (100 rpm) at 25°C for 48 h. Yeast cells were collected by 125 centrifugation at 2500 x g for 7 minutes, washed, resuspended in sterilized Ringer 126 solution (pH 6.9±0.1; Merck) and used as inoculum for the different liquid substrates 127 evaluated in shake flask and batch fermentation experiments.

128

129 2.3. Shake flask experiments

Cell suspensions of *M. pulcherrima* strain BIO126 (3 ml; 5 x 10^8 cfu ml⁻¹) were 130 131 inoculated in 1 litre Erlenmeyer flasks containing 300 ml of liquid media and grown on 132 a rotary shaker (150 rpm) at 25°C for 48 h. Three flasks were prepared per each medium 133 and two samples were collected from each flask. The final number of viable cells (cfu 134 ml⁻¹) was determined by plating on NYDA (Nutrient broth-Yeast extract-Dextrose-Agar 135 as in Droby et al., 1989): 10-fold dilutions of each suspension were prepared in 136 sterilized Ringer solution (pH 6.9+0.1; Merck) and spread-plated in order to calculate 137 the cell number. Plates were incubated at 25°C for 48 h and the number of colony 138 forming units per millilitre (cfu ml⁻¹) was determined. The shake flask experiments were 139 repeated two times.

140

141 2.4. Culture media

142 Complex nutrient media selection. The complex sources selected for the experiments 143 were rich organic sources (yeast extract, nutrient broth, malt extract, meat peptone, 144 casein peptone, bacto-peptone and casein hydrolyzed). Every source was tested at 10 g 1⁻¹. The pH values were registered at the beginning of the experiment, after 24 and 48 h 145 146 of culture. In a second experiment, the initial pH of the seven complex nutrient sources 147 was adjusted to 7.00 \pm 0.05, using a 1.0 M phosphate buffer (Na₂HPO₄ and NaH₂PO₄) 148 solution, in order to evaluate the pH effect on the yeast growth. Every source was tested at 10 g l⁻¹. The pH values were registered after growth for 24 and 48 h. 149

150

151 *Concentration of the complex nutrient source.* Yeast extract medium was tested at 152 different concentrations (5, 10, 15, 20, 30, 40, 50 and 60 g l^{-1}). The pH values were 153 registered after growth for 48 h.

155 *Initial pH.* Yeast extract (30 g l^{-1}) was put in all the flasks. The pH of the media 156 prepared was adjusted using either 0.1 N HCl or NaOH to obtain initial pH ranging 157 from 1.0 to 11.0. The pH values were registered after growth for 24 and 48h.

158

159 *Carbon source addition.* In a first assay, the carbon sources were tested in a medium 160 containing 30 g Γ^1 of yeast extract. Three monosaccharide sugars (D-Glucose, D-161 Fructose and L-Sorbose), two disaccharide sugars (Maltose and Sucrose, Sigma 162 Chemical Co.) and two sugar alcohols (L-Sorbitol and D-Mannitol) were added to the 163 yeast extract, at 10 and 20 g Γ^1 . The pH values were registered after growth for 48h.

In a second assay, D-Mannitol and L-Sorbose were tested either individually at different concentrations (from 2.5 g Γ^1 to 20 g Γ^1) or mixed at concentrations of 5+5 g Γ^1 or 6+6 g Γ^1 in a medium containing 30 g Γ^1 of yeast extract. The pH values were registered after growth for 48h. The results of the two experiments were similar, so they could be analyzed together and combined (Table 3).

169

170 2.5. Fermentation experiments

Well-controlled fermentations of 4.0 L working volume (nominal volume, 5 L) were carried out in Applikon BioConsole ADI 1025 glass stirred tank vessels (AppliconTM Biotechnology, Schiedam, The Netherlands), integrated with the software Bioexpert Lite for data acquisition. Operating conditions included temperature controlled at $25\pm$ 0.2° C, dissolved oxygen permitted to float and monitored using a polarographic probe, agitation with two equally spaced Rushton impellers controlled at a constant speed of 450 rpm, and air sparging through a submerged ring sparger controlled at 4.0L/min or 1 178 vvm (volume of air per volume of medium).

179 The tested substrates were YE (Yeast Extract 30 g l^{-1}), YEM (Yeast Extract 30 g l^{-1} ; D-

180 Mannitol 10 g l⁻¹), YES (Yeast Extract 30 g l⁻¹; L-Sorbose 10 gl⁻¹) and YEMS (Yeast

181 Extract 30 g l^{-1} ; D-Mannitol 5 g l^{-1} ; L-Sorbose 5 g l^{-1}). 0.05 ml/L of silicone antifoam

(Sigma antifoam 204) were added. Every 2 h, starting from the inoculation of the fermenter to the end of the experiment, a 5 ml sample was harvested in order to measure the microorganism cell concentration. The concentration of viable cells (cfu ml⁻¹) was

185 determined by serial dilutions and plating, as indicated in the shake flask experiments.

186 The fermenter experiments were carried out twice.

In order to know the dry biomass produced by fermentation, the dry weight was determined. After 36 h, the liquid cultures were collected and centrifuged (7500 rpm) at 4°C for 10 min (Beckman J21-2 centrifuge, Palo Alto, CA, USA) and the supernatant was discarded. The cell pellet was dried at 105°C for 30h and the dry mass was weighed.

192

193 2.6. Biocontrol assay

194 To evaluate the effect of the growth of *M. pulcherrima* strain BIO126 in four substrates (YE, YEM, YES and YEMS) on the biocontrol efficacy, four trials were carried out 195 against Botrytis cinerea and Penicillium expansum on 'Golden Delicious' and 'Gala' 196 197 apples. Yeast cells were grown in the four substrates for 36h, centrifuged and suspended to 10⁷ cfu ml⁻¹in 100 l tanks. Five strains per each pathogen were isolated from rotted 198 199 apples and selected for their virulence. Each strain was stored in slant on Potato Dextrose Agar (PDA; Merck) with 50 mg l⁻¹ streptomycin Merck at 4°C. Spore 200 201 suspensions were prepared by growing the fungal pathogens on Petri dishes for two

weeks on PDA with 50 mg l^{-1} of streptomycin. Spores from the five strains were 202 203 collected, suspended in sterile Ringer's solution, filtered through 8 layers of sterile cheese-cloth and brought to a final concentration of 10⁵ spores ml⁻¹ per strain. Apples 204 were artificially wounded at the equatorial region (3 mm diameter; 6 mm depth; 3 205 206 wounds per fruit). The fruits were artificially inoculated by dipping for 60 seconds in a 100 l tank containing a conidial suspension (10^5 spores ml⁻¹ per pathogen) of *B. cinerea* 207 or *P. expansum*. After 3 hours, biocontrol isolates were applied at 10^7 cells ml⁻¹ by 208 209 completely dipping the boxes of fruits for 60 sec in a 100 l tank containing the yeast cell 210 suspensions prepared as described. Fifty apples per replicate and three replicates per 211 treatment were used. After incubation at 1°C for 28 days, the incidence of rotten fruits 212 and the lesion diameters were measured. A chemical control treatment consisted of 213 fruits treated with thiabendazole (Tecto 20 S, Elf Atochem Agri Italy, 19.7 % a.i., 20 g a.i. $100 l^{-1}$). The experiment was carried out twice. 214

215

216 2.7. Statistical analysis

217 The fermentation experiments in bioreactor were performed twice, while the growth 218 experiments in shake flasks and the biocontrol experiments were repeated twice. No 219 significant differences were found among corresponding experiments so that the trials 220 were pooled and statistical analysis was performed by using the SPSS software (SPSS 221 Inc., version 13.0, Chicago, IL, USA). Statistical significance was generally judged at 222 the level of P<0.05 for the shake flasks growth and biocontrol experiments, but at 223 P<0.01 for the assay of concentration of the complex nutrient source. When the analysis 224 of variance was statistically significant either in the shake flask growth or in the 225 biocontrol experiments, Duncan's multiple range test was used for the separation of 226 means.

227

- 228 **3. Results**
- 229
- 230 3.1. Growth in complex nutrient media

Maximum exponential growth rate and biomass production of the strain BIO126 of *M*. *pulcherrima* varied with the complex nutrient source (Table 1a). The total nitrogen content of the tested media ranged from 8.0% to 15.4%, but it was very low for Malt Extract (1.1%). The highest yield was obtained with Yeast Extract (1.2 x 10^8 cfu ml⁻¹) and Nutrient Broth (8.5 x 10^7 cfu ml⁻¹), followed by Malt Extract and Meat Peptone. The Ringer solution, used as control, permitted to keep alive the initial inoculum.

237 Five out of seven complex nutrient sources, with an initial pH almost neutral, favoured the growth of *M. pulcherrima*, and the two substrates providing the highest biomass 238 239 resulted in an increased pH value after 24 and 48 h from the inoculum. In the case of 240 Malt Extract and Casein Hydrolyzed, the pH, initially acidic, decreased further at the 241 end of the microorganism growth. For this reason, the experiment was repeated, 242 adjusting the initial pH to 7.00±0.05 with a phosphate buffer (Table 1b). Yeast Extract 243 and Nutrient Broth confirmed the highest viable biomass. The pH values after 24 and 48 244 h were higher and slightly acidic when the microorganism was grown in buffered Malt 245 Extract, but the final biomass obtained was lower. On the opposite, the viable cells 246 obtained in buffered Casein Hydrolyzed were higher compared to the not buffered 247 substrate, and the pH value after 48 h was basic (8.04).

An increase in initial Yeast Extract concentration from 5 to 30 g l⁻¹ gave a proportional increase in the biomass produced at the stationary phase (Fig.1). No significant increase in biomass was observed from 40 to 60 g I^{-1} of Yeast Extract, which is likely due to the Crabtree effect (Crabtree 1928; Boulton et al. 1998). At the stationary phase, pH values of the culture substrate ranged from 8.07 to 8.70.

To assess the effect of the initial pH value on the final biomass produced, *M. pulcherrima* was grown in Yeast Extract (30 g Γ^1) whose initial pH was adjusted at values ranging from 1.0 to 11.0 (Table 2). The pH values lower than 3.0 and higher than 10.0 did not permit the growth of *M pulcherrima*. A growth of at least 10⁸ cfu ml⁻¹ was possible at initial pH values ranging from 4.0 to 8.5. When the initial pH ranged between 5.0 and 7.5, a viable population higher than 3.0 x 10⁸ cfu ml⁻¹ was achieved and the final pH ranged between 8.05 and 8.34.

260

261 *3.2. Effect of carbon addition*

The effect of different carbon sources on the growth of M. pulcherrima was assessed in 262 presence of Yeast Extract at two different concentrations: 10 and 20 g l^{-1} (Table 3). All 263 the carbon sources tested increased the yeast biomass production when used at 10 g l^{-1} . 264 At 20 g l⁻¹ of carbon source, only D-Fructose did not provide a statistically significant 265 increase in the biomass of *M. pulcherrima*. In general, 20 g l^{-1} of carbon source did not 266 improve the yeast biomass compared to 10 g l^{-1} suggesting that high external carbon 267 268 source concentration are not beneficial to growth of this yeast. Only D-Glucose, applied at 20 g l^{-1} , provided a *M. pulcherrima* biomass higher than at 10 g l^{-1} . 269

D-Mannitol and L-Sorbose at the concentration of 10 g Γ^1 provided the highest biomass, 1.5 x 10⁹ cfu ml⁻¹ and 8.0 x 10⁸ cfu ml⁻¹, respectively. Also Sucrose, either at 10 or 20 g 1⁻¹, was a good carbon source, resulting in 6.7 and 6.9 x 10⁸ cfu ml⁻¹ of *M. pulcherrima*, respectively.

The addition of different carbon sources resulted in a lower pH after 48 h. For D-274 Fructose, D-Mannitol, D-Sorbitol, and Sucrose, the pH was below 7.0 even after 48 h. 275 276 D-Mannitol and L-Sorbose were used alone or combined at different concentrations in a second assay to evaluate the effect on the final growth of the yeast strain (Table 3). The 277 maximum growth was obtained using 5.0 g l^{-1} of D- Mannitol + 5.0 g l^{-1} of L-Sorbose, 278 that caused increase in the final number of cells to 1.7×10^9 cfu ml⁻¹. At equal 279 280 concentrations, D-Mannitol provided more yeast growth than L-Sorbose. In particular, the addition of 7.5 and 10.0 g Γ^1 of D-Mannitol were the most effective concentrations 281 resulting in a final cell number of 1.5×10^9 cfu ml⁻¹. The highest growth with L-Sorbose 282 was achieved at concentration of 12.5 g l^{-1} (1.2 x 10^9 cfu m l^{-1}). Addition of D-Mannitol 283 and the mixture of D-Mannitol and L-Sorbose reduced pH after 48h growth. Increasing 284 285 concentrations of D-Mannitol contributed to increase in the final pH value, while 286 increasing concentrations of L-Sorbose had the opposite effect.

287

288 3.3. Fermentation experiments

The biomass production process was scaled-up from shaking flasks to a 5-1 fermenter. The yeast biomass resulted significantly higher $(1.6 \times 10^9 \text{ cfu ml}^{-1})$ in YEMS medium compared to simple YE, YE with L-Sorbose, or YE with D-Mannitol (Fig.2*a*). At 5 1 min⁻¹ of aeration and 450 rpm, the stationary phase was achieved after 32 h of batch culture in YEMS medium, while in the other substrates it was reached from 2 to 4 h later. The maximum specific growth rate during the exponential phase was 0.45 h^{-1} in YEMS, while it was 0.33 h^{-1} in YES and 0.34 h^{-1} in YE and YEM.

Initial pH was 6.9 in YE and in YEM (Fig.2*b*). The presence of L-Sorbose contributedto lower the initial pH to 6.5 in YES and YEMS. As the growth approached the

stationary phase, pH tended to increase. The metabolism of D-Mannitol contributed to
lower the pH of 0.4 units: the minimum value achieved was 6.40 for YEM and 6.03 for
YEMS.

301 The consumption of oxygen is an indication of exponential growth (Fig.2c). The 302 reduction in the dissolved oxygen became visible when the viable population reached around 10^7 cfu ml⁻¹. In YE, the length of the exponential phase and the final number of 303 304 cells obtained were reduced. In YES and YEM, the dissolved oxygen started to decrease 305 earlier than in YE. In YE, moreover, even during the exponential phase, the dissolved 306 oxygen was never reduced to 0%, but in YEMS it declined to 0% from the hour 24 to 307 the hour 31, for the longest period, indicating a long exponential phase. The evolution 308 of the dissolved oxygen indicated the sequential metabolism of D-Mannitol, followed 309 by L-Sorbose and finally by the amino acids and proteins contained in Yeast Extract.

At the end of the fermentation experiments, 1000 ml of cultural broth were harvested, centrifuged and dried to measure the wet and the dry biomass produced. Using YE, YES, YEM and YEMS, the wet biomass was 11.1, 12.0, 19.8, and 26.4 g l^{-1} , respectively, and the dry biomass was 1.4, 2.1, 4.2, and 6.0 g l^{-1} , respectively. These results confirmed the values obtained by plate counting.

315

316 *3.4. Efficacy trials*

In the efficacy trials carried out on apples by treating with *M. pulcherrima* strain BIO126 grown on four substrates, yeast cells grown in YEMS were more effective. Generally, yeast concentrations being equal, the efficacy of the antagonistic cell suspension was influenced by the growth culture substrate (Fig. 3). On 'Golden Delicious' apples (Fig.3*a*), grey mould incidence and severity on the fruits treated with 322 BIO126 grown on YEMS were 48.9% and 29.2% compared to the inoculated control 323 (whose incidence and lesion diameter were 91.3% and 51.9 mm, respectively). Among 324 the fruits treated with the four yeast cell suspensions, grey mould incidence was 325 significantly lower when fruit were treated with cells grown in YEMS, but the mean 326 lesion diameters observed were not significantly different. Considering the efficacy 327 against P. expansum on 'Golden Delicious' apples, the cell suspension of M. 328 pulcherrima BIO126 grown in YEMS reduced the incidence from 86.3% (inoculated 329 control) to 50.3% and the mean lesion diameter from 37.9 mm (inoculated control) to 330 32.6 mm. On the opposite, when the yeast was grown in YE, it was ineffective in 331 reducing blue mould severity compared to the control.

332 On 'Gala' apples (Fig. 3*b*), the grey mould incidence was significantly reduced by 333 BIO126 grown on YEMS, YES, or YEM, but not when the yeast was grown on YE. 334 The lesion diameter of grey rots was not significantly reduced by any yeast application. 335 Similarly, against *Penicillium expansum* on 'Gala' apples, none of the biological 336 treatments could significantly reduce the lesion diameter, but cell suspensions of the 337 yeast grown in YES or YEMS significantly reduced blue mould incidence.

The fungicide thiabendazole reduced the incidence of grey mould and blue mould on
both apple cultivars, but its efficacy in reducing the lesion diameter was higher against *B. cinerea* on 'Golden Delicious' apples, lower, although significant, against *P. expansum* on 'Golden Delicious' apples, and not significant against both pathogens on
'Gala' apples.

343

344 **4. Discussion**

345 The cell production is an essential step in the commercialization of a yeast with

industrial application as a biopesticide. To our knowledge, there are not studies that
have addressed the production process of *M. pulcherrima*, considering the biomass
viable count and the biocontrol efficacy as an objective function of the process.

349 The growth parameters were initially optimized in flask experiments: the complex 350 nutrient source that provided the maximum biomass of the antagonistic yeast M. 351 pulcherrima was Yeast Extract. Although other sources tested (nutrient broth and the 352 three peptones) contained higher nitrogen content, probably Yeast Extract possesses a 353 more balanced equilibrium of amino acids and peptides (it contained 11.4% nitrogen), 354 together with vitamins and carbohydrates, able to promote and sustain a rapid growth of 355 a yeast microorganism (Peppler 1982; Perez et al. 1992). Moreover Yeast Extract shows 356 a buffer ability (Gaudreau et al. 1997) and this could contribute to reach the highest 357 viable cells concentration. The concentrations of Yeast Extract providing the highest growth were 30 g l^{-1} or more. 358

The trial carried out using the complex source media adjusted to pH 7.00 ± 0.05 allowed to determine the effect of the pH control on the biomass production by the strain BIO126. The buffering permitted to significantly improve the final yeast biomass obtained using Casein Hydrolyzed and Nutrient Broth, but it did not affect the result with the other complex source media. For this reason, the trials continued using Yeast Extract without phosphate buffer.

A wide range of pH permitted the growth of *M. pulcherrima* (from 3.0 to 10.0),
although initial pH values ranging from 5.0 to 7.5 provided the highest culture growth.
Thus the subsequent trials were carried out measuring the initial pH values.

368 The results obtained in flask experiments were confirmed in well-controlled batch 369 fermentations. The higher viable cells count obtained in the top bench fermenter could

370 be linked to a better oxygenation as compared to the shake flasks. Maximum 371 exponential growth rate, maximum culture density and maximum wet and dry 372 biomasses were reached using YEMS medium. Moreover, the stationary phase was 373 reached in YEMS from 2 to 4 h earlier than in the other substrates tested. Our results on 374 M. pulcherrima are in accordance or higher than previously obtained results. Abadias et al. (2003) obtained 8 x 10^8 cfu ml⁻¹ of *Candida sake* after 30 h growth in a 5 l lab-scale 375 376 fermenter. By growing Rhodotorula minuta in a shake flasks for 48 h, the best results in terms of viable microorganisms (over 10^9 cells ml⁻¹) were obtained with a PYD, a 377 378 medium containing soluble potato starch, dextrose and yeast extract (Patiño-Vera et al. 379 2005).

380 In the fermentation experiments performed, L-Sorbose lowered the initial pH when it 381 was added to YES and YEMS, and the metabolism of D-Mannitol contributed to lower 382 the pH of the medium during the exponential growth phase. For this reason the 383 combination of the two carbon sources, probably with diauxic utilization, showed a 384 synergistic effect, improving the final yeast biomass over those obtained with the single 385 carbohydrates, and lowering the pH up to 6.08 during the exponential phase. The 386 sequential metabolism of D-Mannitol, followed by L-Sorbose, and finally by the amino 387 acids and proteins contained in Yeast Extract, suggests that the growth response is 388 diauxic, reflecting a sequential rather than simultaneous utilization of the carbon sources 389 (Collier et al. 1996).

Actively growing yeast acidifies the growth medium through a differential ion uptake and direct secretion of organic acids and carbon dioxide (Walker 1998). An increase of the pH was related to the beginning of the stationary phase. During the experiments, when the microorganism was approaching the stationary phase of growth, the

respiration growth decreased, carbon dioxide production decreased and, consequently, pH tended to increase. An experiment was carried out in the top bench fermenter using YEMS as a substrate and with pH control kept at 6.00 (data not shown). The final cell concentration obtained was not significantly different from the experiment without pH buffering, as already shown in the shake flask experiments, so that pH control was not considered as an essential factor to increase the yeast biomass.

400 Oxygen consumption could be considered an indicator of exponential growth (Abadias
401 et al. 2003). The dissolved oxygen rapidly decreased during the exponential phase
402 because of cell respiration and, on the opposite, it started back to increase because of a
403 decrease in the respiration rate of the cells (Meesters et al. 2003).

404 Cells were grown for 48 h in shake flask experiments and 40 h in well-controlled batch 405 experiments, periods largely sufficient for the microorganism to reach the stationary 406 phase of growth. Harvesting stationary phase cells is desirable to enhance cell survival 407 under stress conditions such as low water potential or drying (Abadias et al. 2001). This 408 could be an advantage as, after production, M. pulcherrima cells have to be formulated, 409 probably, through a drying process, such as freeze drying (Melin et al. 2007) or fluid 410 bed drying (Bayrock and Ingledew 1997). In both cases, yeast cells harvested at the 411 stationary phase are more resistant to the osmotic stress caused by the drying process 412 (Wraight et al. 2001). The efficacy trials conducted on two commercial varieties of apple 413 permitted to test the influence of the growing media on the biocontrol capability of M. 414 pulcherrima BIO126. The growth of both pathogens and the biocontrol efficacy of 415 BIO126 were affected by the host cultivar. The two different cultivars chosen for the 416 test, "Gala" and "Golden Delicious", have different pH and titrable acidity, able to 417 modify the growth and fitness of postharvest pathogens (Morales et al. 2008). On 'Gala,

418 the incidence of fruits with grey or blue mould was lower than on 'Golden Delicious' 419 apples. On 'Golden Delicious' apples, the microorganism cells grown in YEMS 420 provided a higher reduction of incidence and severity of B. cinerea and P. expansum. 421 Results were more promising against grey mould than blue mould, probably because P. 422 expansum is growing at a faster rate. On 'Gala' apples, the best results on the reduction 423 of the disease incidence were obtained with cells grown on YEMS or YES. Previous 424 studies, carried out to test the influence of the growing media on the biocontrol efficacy 425 of other antagonists, such as *Pantoea agglomerans*, did not show any statistically 426 significant effect of the substrates (Costa et al. 2001).

427 As a chemical control, thiabendazole was used. Benzimidazoles can effectively control 428 grey mould on apples, but they are almost ineffective against *P. expansum*, due to the 429 high level of resistance developed by most of the Italian strains of this pathogen 430 (Bertetti et al. 2003). To represent conditions more similar to the postharvest 431 environment, both pathogen mixtures used in the efficacy trials were formed by four 432 benzimidazole-sensitive isolates and one benzimidazole-resistant isolate. Penicillium 433 expansum can cause economical losses due to blue mould, but also produces patulin, a 434 mycotoxin often found in apple juices (Spadaro et al. 2007). For this reason, effective 435 control strategies against blue mould are necessary, and strain BIO126 could constitute 436 an effective alternative for blue mold control after harvest.

The results obtained were encouraging and a good substrate was developed for laboratory purposes (i.e. to find optimal conditions for biomass production, stabilization, formulation, and to develop a quality control system), but the following step will be the scaling up of the production process to the level of pilot plant and the subsequent stabilization and formulation. To scale-up a laboratory fermentation process

442 to an industrial level, it is fundamental to find nitrogen and carbon sources that provide 443 maximum biomass production and minimum cost of media, whilst maintaining 444 biocontrol efficacy (Mousdale et al. 1999). Yeast extract was the best complex 445 nutritional source for BIO126 but it is expensive for an industrial process. The use of 446 commercial by-products with the same nutritional qualities can result in a cheap 447 alternative for the yeast biomass production (Ghribi et al. 2006). Anyway, often by-448 products are not standardized as purified products and they may contain impurities that 449 need to be removed before fermentation (Stanbury et al., 1995). Moreover, their 450 composition may vary according to season and origin. For these reasons, appropriate 451 procedures should be employed to standardize the industrial growth media (Thomsen 452 2005). Commercial dry beer yeast could replace synthetic yeast extract, but it should be 453 filtered before autoclaving, to discard the insoluble fraction (Reed and Nagodawithana 454 1991).

The knowledge gained about the addition of D-Mannitol and L-Sorbose will be useful to develop a cheap substrate containing a complex nutritional source similar to yeast extract together to the sugars selected in this study. After producing the biomass of the antagonistic yeast, the next step will be the cell stabilization by freeze-drying or fluid bed drying (Brian and Etzel 1997).

460 Acknowledgements

This research was carried out with a grant from the Piedmont Region for the project "Selection, study of the efficacy and of the mechanism of action, characterization and development of antagonistic yeast against post-harvest fruit pathogens". The authors gratefully acknowledge Dr. Pilar Bustos, Dr. Piergiuseppe Nestola and Ms. Incoronata Luongo for their help in the realization of the trials and Dr. Jeanne Griffin for her

466 linguistic advice. A special thank to the anonymous reviewers of the manuscript.

467

468 **References**

- 469 Abadias, M., Teixidó, N., Usall, J., and Viñas, I. 2003. Optimization of growth
- 470 conditions of the postharvest biocontrol agent Candida sake CPA-1 in a lab-scale
- 471 fermenter. J. Appl. Microbiol. **95**: 301-309.
- 472 Abadias, M., Teixidó, N., Usall, J., Viñas, I., and Magan, N. 2001. Improving water
- 473 stress tolerance of the biocontrol yeast *Candida sake* grown in molasses-based media by
- 474 physiological manipulation. Can. J. Microbiol. **4**: 23-129.
- 475 Bayrock, D. and Ingledew, W.M. 1997. Fluidized bed drying of baker's yeast: moisture
- 476 levels, drying rates, and viability changes during drying. Food Res. Int. **30**: 407-415.
- 477 Bertetti, D., Kejji, S., Garibaldi, A., and Gullino, M.L., 2003. Valutazione della
- 478 sensibilità di alcuni agenti di marciume su pomacee in post-raccolta nei confronti di
- 479 diversi fungicidi. Inf.tore Fitopatol. 53 (6): 57-60.
- 480 Bleve, G., Grieco, F., Cozzi, G., Logrieco, A., and Visconti, A. 2006. Isolation of
- 481 epiphytic yeasts with potential for biocontrol of Aspergillus carbonarius and A. niger on
- 482 grape. Int. J. Food Microbiol. 108: 204-209.
- 483 Boulton, R.B., Singleton, V.L., Bisson, L.F., and Kunkee, R.E. 1998. Principles and
- 484 Practices of Winemaking. Aspen Publishers, New York.
- 485 Brian, C.S. and Etzel, M.R. 1997. Spray drying, freeze drying, or freezing of three
- 486 different lactic acid bacteria species. J. Food Sci. **62:** 576-578.
- 487 Collier, D.N., Hager, P.W., and Phibbs, P.V. 1996. Catabolite repression control in the
- 488 Pseudomonads. Res. Microbiol. 147: 551–561.
- 489 Costa, E., Teixidó, N., Usall, J., Atarés, E., and Viñas, I. 2001. Production of the

- 490 biocontrol agent Pantoea agglomerans strain CPA-2 using commercial products and
- 491 by-products. Appl. Microbiol. Biotechnol. 56: 367-371.
- 492 Crabtree, H.G. 1928. The carbohydrate metabolism of certain pathological overgrowths.
- 493 Biochem. J. 22 (5): 1289-1298.
- 494 Droby, S., Chalutz, E., Wilson, C.L., and Wisniewski, M.E. 1989. Characterization of
- 495 the biocontrol activity of *Debaryomices hansenii* in the control of *Penicillium digitatum*
- 496 on grapefruit. Can. J. Microbiol. **35**: 794-800.
- 497 Gaudreau, H., Champagne, C.P., Goulet, J., and Conway, J. 1997. Lactic fermentation
- 498 of media containing high concentrations of yeast extract. J. Food Sci. 62: 1072-1075.
- 499 Ghribi, D., Zouari, N., Trigui, W., Jaoua, S. 2006. Use of sea water as salts source in
- starch- and soya bean-based media, for the production of *Bacillus thuringiensis*bioinsecticides. Process Biochem. 42: 374–378.
- 502 Hofstein, R., Friedlender, B., Chalutz, E., and Droby, S. 1994. Large scale production
- 503 and pilot testing of biocontrol agents of postharvest diseases. In Biological Control of
- 504 Postharvest Diseases Theory and Practice. *Edited by* C.L. Wilson and M. Wisniewski.
- 505 CRC Press Inc, Boca Raton, pp. 89-100.
- 506 Janisiewicz, W.J., and Korsten, L. 2002. Biological control of postharvest diseases of
- 507 fruits. Annu. Rev. Phytopathol. 40: 411-441.
- 508 Janisiewicz, W.J., Tworkoski, T.J., and Kurtzman, C.P. 2001. Biocontrol potential of
- 509 Metschnikowia pulcherrima strains against blue mold of apple. Phytopathology 91:
- 510 1098-1108.
- 511 Leverentz, B., Conway, W.S., Janisiewicz, W., Abadias, M., Kurtzman, C.P., and
- 512 Camp, M.J. 2006. Biocontrol of the food-borne pathogens Listeria monocytogenes and

- *Salmonella enterica* serovar Poona on fresh-cut apples with naturally occurring bacterial
 and yeast antagonists. Appl. Environ. Microbiol. **72**:1135–40.
- 515 Meesters, P.A.E.P., Huijberts, G.N.M., and Eggink, G. 1996. High-cell density 516 cultivation of the lipid accumulating yeast *Cryptococcus curvatus* using glycerol as
- 517 carbon source. Appl. Microbiol. Biotechnol. **45:** 575-579.
- 518 Miller, M.W., and Phaff, H.J. 1998. *Metschnikowia* Kamienski. *In* The Yeasts. *Edited*
- 519 by C.P. Kurtzmann and J.W. Fell. Elsevier, Amsterdam, pp. 256-257.
- 520 Morales, H., Barros, G., Marín, S., Chulze, S., , Ramos, A.J., and Sanchis, V. 2008.
- 521 Effects of apple and pear varieties and pH on patulin accumulation by Penicillium
- 522 *expansum.* J. Sci. Food Agric. **88**: 2738-2743.
- 523 Mousdale, D.M., Melville, J.C., and Fischer, M. 1999. Optimization of fermentation
- 524 process by quantitative analysis: from analytical biochemistry to chemical engineering.
- 525 In Fermentation Microbiology and Biotechnology. Edited by E.M.T. El-Mansi and
- 526 C.F.A. Bryce. Taylor and Francis, London, pp. 147-178.
- 527 Patiño-Vera, M., Jiménez, B., Balderas, K., Ortiz, M., Allende, R., Carrillo, A. and
- 528 Galindo, E., 2005. Pilot-scale production and liquid formulation of *Rhodotorula minuta*,
- 529 a potential biocontrol agent of mango anthracnose. J. Appl. Microbiol. **99**: 540-550.
- 530 Peppler, H.J. 1982. Yeast extracts. In Fermented Foods. Edited by A.H. Rose. Academic
- 531 Press, London, pp. 293-312.
- 532 Perez, F., Riba, J.P., and Strehaiano, P. 1992. Effect of yeast extract concentration on
- 533 growth of *Schizoccharomyces pombe*. Biotechnol. Lett. 14: 123-126.
- 534 Piano, S., Neyrotti, V., Migheli, Q., and Gullino, M.L. 1997. Characterization of the
- 535 biocontrol capability of *Metschnikowia pulcherrima* against Botrytis postharvest rot of
- 536 apple. Postharvest Biol. Technol. **11**: 131-140.

- 537 Reed, G. and Nagodawithana, T.W. 1991. Yeast Technology, 2nd edition, Van Nostrand
 538 Rheinold, New York.
- 539 Saravanakumar, D., Ciavorella, A., Spadaro, D., Garibaldi, A., and Gullino, M.L. 2008.
- 540 Metschnikowia pulcherrima strain MACH1 outcompetes Botrytis cinerea, Alternaria
- 541 *alternata* and *Penicillium expansum* in apples through iron depletion. Postharvest Biol.
- 542 Technol. **49**: 121-128.
- 543 Saravanakumar D., Spadaro D., Garibaldi A., and Gullino M.L. 2009. Detection of
- 544 enzymatic activity and partial sequence of a chitinase in *Metschnikowia pulcherrima*
- 545 strain MACH1 used as postharvest biocontrol agent. Eur. J. Plant Pathol. **123**: 183-193.
- 546 Schena, L., Ippolito, A., Zahavi, T., Cohen, L., and Droby, S. 2000. Molecular 547 approaches to assist the screening and monitoring of postharvest biological yeasts. Eur.
- 548 J. Plant Pathol. **106**: 681-691.
- 549 Sinigaglia, M., Corbo, M.R., and Ciccarone, C. 1998. Influence of temperature, pH and
- water activity on "in vitro" inhibition of *Penicillium glabrum* (Wehmer) Westling by
 yeasts. Microbiol Res. 153:137-43.
- Spadaro, D., and Gullino, M.L. 2004. State of art and future perspectives of biological
 control of postharvest fruit diseases. Int. J. Food Microbiol. 91: 185-194.
- Spadaro, D., Sabetta, W., Acquadro, A., Portis, E., Garibaldi, A., and Gullino, M.L.
 2008. Efficacy and genetic diversity of *Metschnikowia pulcherrima* strains isolated
 from different food matrices against postharvest diseases in apple. Microbiol. Res. 163:
 523-530.
- Spadaro, D., Vola, R., Piano, S., and Gullino, M.L. 2002. Mechanisms of action,
 efficacy and possibility of integration with chemicals of four isolates of the yeast *Metschnikowia pulcherrima* active against postharvest pathogens on apples. Postharvest

- 561 Biol. Technol. 24: 123-134.
- 562 Spadaro, D., Ciavorella, A., Frati, S., Garibaldi, A., and Gullino, M.L. 2007. Incidence
- 563 and level of patulin contamination in pure and mixed apple juices marketed in Italy.
- 564 Food Cont. **18**: 1098-1102.
- 565 Stanbury, P.F., Whitaker, A., and Hall, S.J. 1995. Principles of Fermentation
- 566 Technology. Pergamon Press, Oxford.
- 567 Thomsen, M.H. 2005. Complex media from processing of agricultural crops for
 568 microbial fermentation. Appl. Microbiol. Biotechnol. 68: 598-606.
- 569 Walker, G.M. 1998. Yeast Physiology and Biotechnology. John Wiley and Sons,
- 570 Chichester.
- 571 Wilson, C.L., and Wisniewski, M.E. 1994. Biological Control of Postharvest Diseases.
- 572 Theory and Practice. CRC Press, Boca Raton.
- 573 Wraight, S.P., Jackson, M.A., and de Kock, S.L. 2001. Production, Stabilization and
- 574 Formulation of Fungal Biocontrol Agents. In Fungi as Biocontrol Agents. Edited by
- 575 T.M. Butts, C. Jackson and N. Magan. CAB International, Wallingford, Great Britain,
- 576 pp. 253-287.
- 577

Table 1 Growth of *Metschnikowia pulcherrima* strain BIO126 (cfu m⁻¹) in different complex nutrient media (10 g Γ^1 each), without (a, left) or with (b, right) adjustment of initial pH₀ 7.00±0.05 with a phosphate buffer. Values of pH (±0.05) at initial time (pH₀), after 24h (pH₂₄) and 48h culture (pH₄₈) are shown. Cultures (5 x 10⁵ cfu ml⁻¹) were inoculated in 300 ml of liquid media and grown on a rotary shaker (150 rpm) at 25°C for 48 h.

	Total	Complex nutrient media (a)			Buffe	Buffered complex nutrient media (b)			
	nitrogen content	pH ₀	pH ₂₄	pH ₄₈	Mean (cfu ml ⁻¹)	pH ₂₄	pH48	Mean (cfu ml ⁻¹)	
Yeast Extract	11.4	7.13	7.91	8.56	1.2 x 10 ⁸ a*	8.05	8.61	1.1 x 10 ⁸ a*	
Nutrient Broth	13.0	7.11	7.72	8.57	8.5×10^7 a-b	8.19	8.73	1.2×10^8 a	
Malt Extract	1.1	5.17	4.94	3.84	5.3×10^7 b	6.77	6.70	1.8×10^7 c	
Meat Peptone	13.9	7.36	7.30	7.20	2.3×10^{7} b-c	7.10	7.34	1.8×10^{7} c	
Casein Peptone	13.5	7.04	7.35	7.50	1.5×10^{7} c	7.14	7.22	2.7×10^{7} c	
Bacto-peptone	15.4	7.09	7.56	7.53	1.3×10^{7} c	7.26	7.35	1.4×10^{7} c	
Casein Hydrolyzed	8.0	5.42	5.53	5.67	1.3×10^{7} c	7.45	8.04	8.0 x 10 ⁷ b	
Ringer solution		6.35	6.30	6.54	4.9 x 10 ⁵ d	7.42	7.32	$5.6 \times 10^{5} d$	

^{\$585} * Values in the same column followed by the same letter are not statistically different

586

by

587 Duncan's Multiple Range Test (*P*<0.05).

Table 2 Effect of the initial pH value on the growth of *Metschnikowia pulcherrima* strain BIO126 (inoculum: 5×10^5 cfu ml⁻¹) produced in liquid medium containing yeast extract (30 g l⁻¹) at 25°C for 48 h. Values of pH (±0.05) at initial time (pH₀), after 24 (pH₂₄) and 48h culture (pH₄₈) are shown.

pH_0	pH 24h	Mean _{24h} (cfu ml ⁻¹)	$pH_{\rm \ 48h}$	Mean _{48h} (cfu ml ⁻¹)
1.0	1.06	no living cells	0.94	no living cells
1.5	1.58	no living cells	1.42	no living cells
2.0	2.08	no living cells	1.95	no living cells
2.5	2.53	no living cells	2.45	no living cells
3.0	3.22	5.2 x 10 ⁵	7.35	9.5 x 10 ⁶ b*
3.5	6.03	1.2 x 10 ⁶	7.67	$1.6 ext{ x } 10^{7} ext{ b-c}$
4.0	7.72	1.2 x 10 ⁸	7.32	2.6 x 10 ⁸ d-f
4.5	7.70	1.1 x 10 ⁸	6.59	2.1 x 10 ⁸ d-e
5.0	8.02	1.3 x 10 ⁸	8.12	4.9 x 10 ⁸ g
5.5	7.02	1.6 x 10 ⁸	8.27	3.8 x 10 ⁸ f-g
6.0	6.80	1.5 x 10 ⁸	8.16	3.7 x 10 ⁸ f-g
6.5	8.23	1.4 x 10 ⁸	8.33	3.2 x 10 ⁸ e-g
7.0	7.98	7.3×10^{7}	8.34	3.1 x 10 ⁸ e-g
7.5	7.68	5.8 x 10 ⁷	8.05	3.4 x 10 ⁸ e-g
8.0	7.91	5.5×10^{7}	6.70	$2.2 \times 10^{8} \text{ d-f}$
8.5	8.30	6.3 x 10 ⁷	8.28	1.6 x 10 ⁸ d
9.0	7.05	3.3×10^{7}	7.62	$3.6 \times 10^{-7} c$
9.5	8.53	1.4×10^{7}	8.23	$2.3 \times 10^{-7} \text{ b-c}$
10.0	8.86	3.9 x 10 ⁵	8.90	1.5 x 10 ⁵ a
11.0	9.30	3.9×10^{5}	9.43	no living cells

* See Table 1.

Table 3 Growth of *Metschnikowia pulcherrima* strain BIO126 (cfu ml⁻¹) in media containing yeast extract (30 g l⁻¹) and different carbon sources. Cultures (5 x 10⁵ cfu ml⁻¹) were inoculated in 300 ml of liquid media and grown on a rotary shaker (150 rpm) at 25°C for 48 h. Values of pH (± 0.05) at initial time (pH₀) and 48h culture (pH₄₈) are shown.

Nitrogen source	Carbo	pH_{0h}	$pH_{\rm \ 48h}$	Mean _{48h} (cfu ml ⁻¹)		
	D-Glucose	10 g l^{-1}	6.88	7.38	3.7×10^8	b-c*
		$20 \text{ g} \text{ l}^{-1}$	6.80	8.18	5.2×10^8	b-c
	D-Fructose	$10 \text{ g } \text{ l}^{-1}$	6.89	8.54	3.8×10^8	b-c
		$20 \text{ g } \text{ l}^{-1}$	6.90	6.78	$3.0 \ge 10^8$	a-b
	Sucrose	10 g l ⁻¹	6.90	7.64	6.7 x 10 ⁸	c-e
		20 g l^{-1}	6.89	6.85	6.9 x 10 ⁸	c-e
	Maltose	10 g l ⁻¹	6.89	8.20	6.2×10^8	b-d
		20 g l ⁻¹	6.87	7.70	$5.7 \ge 10^8$	b-d
	D-Sorbitol	10 g l ⁻¹	6.92	7.18	5.9 x 10 ⁸	b-d
		20 g l ⁻¹	6.90	6.94	$5.6 \ge 10^8$	b-d
	D-Mannitol	2.5 g l ⁻¹	6.90	6.40	$2.4 \text{ x} 10^8$	a-b
		5.0 g l ⁻¹	6.92	6.40	6.5 x 10 ⁸	c-e
		7.5 g l ⁻¹	6.92	6.50	$1.5 \ge 10^9$	g
Yeast		10.0 g l ⁻¹	6.90	6.60	$1.5 \ge 10^9$	g
extract		12.5 g l ⁻¹	6.90	7.00	$1.2 \ge 10^9$	f-g
$(30 \text{ g } \text{l}^{-1})$		15.0 g l ⁻¹	6.90	7.20	$1.3 \ge 10^9$	f-g
		17.5 g l ⁻¹	6.91	7.20	8.2×10^8	d-e
		20.0 g l ⁻¹	6.91	7.50	6.6 x 10 ⁸	c-e
	L-Sorbose	2.5 g l ⁻¹	6.90	8.40	$1.5 \ge 10^8$	а
		5.0 g l ⁻¹	6.90	8.40	2.4×10^8	a-b
		7.5 g l ⁻¹	6.91	8.30	7.3×10^8	d-e
		10.0 g l^{-1}	6.89	8.20	$8.0 \ge 10^8$	d-e
		12.5 g l ⁻¹	6.90	8.20	$1.2 \ge 10^9$	f-g
		15.0 g l ⁻¹	6.89	8.00	9.0 x 10 ⁸	e-f
		17.5 g l ⁻¹	6.90	7.95	$7.0 \ge 10^8$	c-e
		20.0 g l ⁻¹	6.90	7.95	$7.0 \ge 10^8$	c-e
	D-Mannitol +	$5.0 \text{ g l}^{-1} + 5.0 \text{ g l}^{-1}$	6.97	6.69	1.7×10^9	g
	L-Sorbose	$6.0 \text{ g } \text{l}^{-1} + 6.0 \text{ g } \text{l}^{-1}$	6.93	7.23	$1.3 \ge 10^9$	f-g
	None		6.91	8.40	1.5×10^8	а

604 * See Table 1.



606 **Fig. 1** Effect of the Yeast Extract concentration on the growth of *Metschnikowia* 607 *pulcherrima* strain BIO126 (cfu ml⁻¹) produced in shake flasks at 25°C for 48 h. Values 608 of pH after 48h culture are shown between parentheses.

Fig 2 (*a*) Evolution of biomass production (cfu ml⁻¹) of *Metschnikowia pulcherrima* strain BIO126, (*b*) pH and (*c*) dissolved oxygen using YE (Yeast Extract; \blacktriangle), YES (Yeast Extract + D-Sorbitol; x), YEM (Yeast Extract + L-Mannitol; \blacklozenge) and YEMS (Yeast Extract + D-Sorbitol + L-Mannitol; \blacksquare) media in a 5 1 fermenter, maintaining temperature at 25°C, stirring at 450 rpm and oxygen flow at 5 1 min⁻¹ for 40 h. (*a*)





Fig 3 Influence of the cultivation substrate on the efficacy of M. pulcherrima strain 640 BIO126 (10⁷ cfu ml⁻¹) against *Botrytis cinerea* (left) and *Penicillium expansum* (right) 641 on apples 'Golden Delicious' (a) and 'Gala' (b). Fruits were artificially inoculated with 642 643 the pathogen $(10^5 \text{ conidia ml}^{-1})$ 3 hours before treatment with the biocontrol agent, and 644 then stored at 1°C for 28 days. Disease incidence was expressed as percentage of rotten 645 fruits and disease severity was assessed by measuring the lesion diameter of the rots 646 (mm). Thiabendazole was used as chemical control (Tecto 20 S, Elf Atochem Agri Italy, 19.7 % a.i., 20 g a.i. 100 l⁻¹). 647





Values in columns of the same colour followed by the same letter are not statistically

653 different by Duncan's Multiple Range Test (P < 0.05).