

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Defective neurogenesis in the absence of Dlx5.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/79203> since

Publisher:

Attuale: ACADEMIC PRESS INC ELSEVIER SCIENCE, 525 B ST, STE 1900, SAN DIEGO, USA, CA, 92101-4495

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Defective neuronogenesis in the absence of *Dlx5*

Marzia Perera,^a Giorgio R. Merlo,^b Sara Verardo,^{a,c} Laura Paleari,^a
Giorgio Corte,^{a,c} and Giovanni Levi^{d,*}

^aLaboratorio di Biologia dello Sviluppo, Istituto Nazionale per la Ricerca sul Cancro-IST, 16132 Genova, Italy

^bDulbecco Telethon Institute CNR-ITB, 20090 Segrate, Milano, Italy

^cDipartimento di Oncologia, Biologia e Genetica, Università di Genova, 16132 Genova, Italy

^dLaboratoire de Physiologie Generale et Comparee, Museum National d'Histoire Naturelle, UMR5166 CNRS-MNH Evolution des Régulations Endocriniennes, Paris Cedex 5, France

Received 2 July 2003; revised 23 September 2003; accepted 7 October 2003

Dlx genes play an important role in the control of the development of the central nervous system (CNS). Single or compound inactivation of *Dlx1*, *Dlx2*, or *Dlx5* in the mouse causes defects of neuronal migration and differentiation. *Dlx5*, in particular, is essential for the correct development of the olfactory system. Targeted inactivation of *Dlx1* and *Dlx2* in the mouse results in abnormal neuronal differentiation in the embryonic subcortical forebrain and is associated to the loss of *Dlx5* and *Dlx6* expression. So far, however, it has been impossible to investigate the role of *Dlx* genes on late neurogenesis, as their inactivation leads to perinatal death. We have now generated cultures of neural stem cells (NSCs) derived from embryonic and newborn *Dlx5*-null mice, and we have compared their capacity to differentiate *in vitro* to that of equivalent cells derived from normal littermates. We show here that in the absence of *Dlx5*, NSCs derived from newborn animals have a severely reduced capacity to generate neurons. This is not the case for cells derived from E12.5 embryos. Forced expression of *Dlx5* in cultures of newborn mutant NSCs fully restores their neuronogenic potential. Our data suggest that *Dlx5* is essential for secondary (postnatal) neuronogenesis.

© 2003 Elsevier Inc. All rights reserved.

Introduction

The fate of neural progenitor cells (NPCs) results from the integration of multiple environmental signals with intrinsic genetic programs (Okano, 2002). Some of these molecular mechanisms, such as, for example, those underlying the generation of spinal motoneurons (Wichterle et al., 2002) or of dopaminergic mesen-

cephalic neurons (Wagner et al., 1999), are beginning to be revealed (Schuermans and Guillemot, 2002). Despite these important results, most of the intrinsic and extrinsic determinants of cell fate in the developing central nervous system (CNS) remain to be elucidated. In this respect, it is essential to identify which genes are involved in the initiation of neuronal differentiation of uncommitted NPCs (see for example Nieto et al., 2001; Sun et al., 2001). One approach is to analyze the properties of NPCs derived from mice carrying targeted mutations of genes potentially important for the control of neural differentiation.

Dlx genes encode for homeobox-containing transcription factors related to *Drosophila Distal-less (Dll)*. In mammals, there are six *Dlx* genes arranged in three pairs of closely linked transcription units (Simeone et al., 1994; Porteus et al., 1991; Price et al., 1991; Robinson et al., 1991; Robinson and Mahon, 1994; Quinn et al., 1997; McGuinness et al., 1996). *Dlx1*, *Dlx2*, *Dlx5*, and *Dlx6* mRNAs are expressed in the subcortical forebrain mainly in those territories where GABAergic differentiation occurs and in precursors of the GABAergic lineage (Panganiban and Rubenstein, 2002).

As a direct evidence of a role of *Dlx* genes in the initiation of the GABAergic differentiation program, exogenous expression of *Dlx2* or *Dlx5* in cortical neurons leads to the appearance of molecular markers specific for the GABAergic phenotype (Stuhmer et al., 2002). Furthermore, *Dlx1* and *Dlx2* genes are required for the differentiation of projection neurons in the basal ganglia since the maturation of striatal cells that synthesize GABA, acetylcholine, and dopamine is impaired in mice lacking both *Dlx1* and *Dlx2* (Anderson et al., 1997a; Marin et al., 2000; Yun et al., 2002). Notably in single *Dlx1*-, *Dlx2*-, or *Dlx5*-null mice, defects in basal ganglia formation have not been reported.

In the ganglionic eminences (GE), *Dlx* genes appear to be expressed in a hierarchical order of transcriptional regulation (Zerucha et al., 2000), which is reflected in a patterned expression in the neurogenic ventricular–subventricular zone (VZ–SVZ): *Dlx2* is found in very early progenitors, capable of initiating the neuronal, astrocytic, and oligodendrocytic lineages (He et al., 2001; Marshall and Goldman, 2002), while *Dlx1*, *Dlx5*, and *Dlx6* are found in progressively restricted subset of *Dlx2*-positive cells.

Abbreviations: GE, ganglionic eminence; LGE, lateral ganglionic eminence; MGE, medial ganglionic eminence; NPC, neural progenitor cells; NSC, neural stem cells; OB, olfactory bulb; RMS, rostral migratory stream; SVZ, subventricular zone; VZ, ventricular zone.

* Corresponding author. Laboratoire de Physiologie Generale et Comparee, Museum National d'Histoire Naturelle, UMR5166 CNRS-MNH Evolution des Régulations Endocriniennes, 7, rue Cuvier, 75231 Paris Cedex 5, France. Fax: +33-1-40-79-36-07.

E-mail address: glevi@mnhn.fr (G. Levi).

Available online on ScienceDirect (www.sciencedirect.com.)




Fig. 1. Expression of *Dlx5* and *Dlx6* in the neonatal mouse brain. Coronal (frontal) sections were used for X-gal staining of the β -galactosidase reporter expressed in *lacZ-Dlx5*^{+/-} and *lacZ-Dlx5*^{-/-} mice (C, D, H, and J) for in situ hybridization with *Dlx6* probe (in E, F, and G) or for immunohistochemical staining (A, anti- β -galactosidase; B, anti-Emx2). In A and B, sections correspond to the area indicated in C. Note the high level of expression of *Dlx5* (A, C) but not of *Dlx6* (red asterisk in E) in a region just under the SVZ. Inactivation of *Dlx5* did not eliminate the population of cells in which the *Dlx5* and *Dlx6* promoter is active in the CP and in the RMS (C and D; F–J). Abbreviations: CP, caudate-putamen; RMS, rostral migratory stream (red arrows).




Fig. 2. *lacZ* expression in cultured NSCs. (Top) Undifferentiated NPCs after 3 h culture; β -gal staining is visible in a subpopulation of both *Dlx5*^{+/-} (B) and *Dlx5*^{-/-} (C) cells. (Bottom) NSCs after 24 h culture in differentiating conditions, no staining is seen in astrocytes (black arrows) independently from their genotype. In cultures of *Dlx5*^{+/-} cells, expression of the *lacZ-Dlx5* allele is clearly visible in differentiated neurons (arrowheads in E) that can be recognized by their distinctive morphology. No neuronal differentiation and not β -gal positive cells could be found after 24 h in cultures of *Dlx5*^{-/-} NSCs (F). The weak blue staining seen in F is an artefact due to the conditions of illumination of the sample.




Fig. 3. Defective neurogenesis in NPCs from P0 brains. Micrographs of cultured NPCs after 7 days of differentiation, stained with anti-GFAP (red), and TuJ1 anti- β -tubulin (green). NPCs obtained from E12.5 (top) and P0 (bottom) animals are compared. The genotype at the *Dlx5* locus is indicated on the top right of each panel. Inserts in A and B show an example of differentiated neuron (TuJ1 positive cells) from the corresponding culture. Note the strongly reduced number of TuJ1-positive cells in cultures of *Dlx5*^{-/-} neonatal NSCs.

Thus, in the VZ–SVZ of the embryonic GE *Dlx* gene expression follows a temporal, positional, and functional sequence: *Dlx2*, *Dlx1*, and then *Dlx6* (Porteus et al., 1994; Liu et al., 1997; Eisenstat et al., 1999; Panganiban and Rubenstein, 2002). In the embryonic GE, *Dlx2* and *Dlx1* are mainly found in neuroepithelial cells of the VZ, while *Dlx5* is mostly expressed in cells of the SVZ, in cells of the rostral migratory stream (RMS), and of the olfactory bulb (OB) (Bulfone et al., 1993; Eisenstat et al., 1999; Levi et al., 2003). Whether this functional and histological organization of *Dlx* expressing cells is retained in the neonatal SVZ is not entirely known, nor we have evidence of a function of *Dlx* genes during perinatal and postnatal neurogenesis in the SVZ–RMS–OB system.

The analysis of the brain phenotypes of mice with targeted disruption of individual *Dlx* genes (*Dlx1*^{-/-}, *Dlx2*^{-/-}, and *Dlx5*^{-/-}) or combinations of them (*Dlx1*^{-/-}/*Dlx2*^{-/-}) reveals partially overlapping functions of *Dlx* genes in the development of the basal forebrain. *Dlx* genes play fundamental roles in the control of migration of NPCs along the RMS toward the OBs, in the tangential migration from the GE into the dorsal cortex, and the differentiation of immature precursors into GABAergic neurons (Anderson et al., 1997b, 2001; Marin and Rubenstein, 2001; Panganiban and Rubenstein, 2002; Stuhmer et al., 2002; Levi et al., 2003). The brain of *Dlx1*^{-/-}/*Dlx2*^{-/-} mice has been highly informative: in addition to a block in tangential migration from the GE to the cortex, these mice show a block in neurogenesis in the basal telencephalon and a block in the differentiation of GABAer-

gic and dopaminergic interneurons in the OB and other brain areas (Marin et al., 2000; Marin and Rubenstein, 2001; Anderson et al., 1997a,b; Bulfone et al., 1998). Notably, both *Dlx2*^{-/-} and *Dlx5*^{-/-} mice show a reduction in the number of GABAergic and dopaminergic neurons in the OB (Qiu et al., 1995; Bulfone et al., 1998; Long et al., 2003; Levi et al., 2003).

In *Dlx5*-null mice (Acampora et al., 1999; Depew et al., 1999), a unique olfactory phenotype is observed, with defects affecting both the olfactory epithelium (OE) and the OBs. In these mice, we (Levi et al., 2003) and others (Long et al., 2003) have observed a profound disorganization of the OB cell layers and a severe impairment of dopaminergic and GABAergic interneuron differentiation. We could also demonstrate that the reduction of differentiated interneurons is a direct consequence of the mutation in a cell-autonomous fashion rather than the result of the lack of axonal connectivity between the OE and the OB (Levi et al., 2003). OB interneurons derive mainly from precursors that reside in VZ–SVZ of the forebrain, although neural stem cells (NSCs) have been isolated from all the rostral extension regions of the olfactory areas, including the OBs (Gritti et al., 2002). At later stages of development and in postnatal life, proliferating cells in the SVZ migrate rostrally along the RMS and home specifically to the OBs (Weiss et al., 1996; Lois and Alvarez-Buylla, 1994; Alvarez-Buylla et al., 2002; Alvarez-Buylla and Garcia-Verdugo, 2002). Based on these observations, we raise the hypothesis that the impaired OB interneuronal differentiation seen in *Dlx5*-null brains reflects an

intrinsic defect of differentiation of *Dlx5*-deficient NPCs. This would necessarily imply a function of *Dlx5* in the regulation of the intrinsic differentiation capacity of NPCs of the olfactory neurogenic system. Embryonic neurogenesis is severely impaired in the double *Dlx1–Dlx2*-null brains (Anderson et al., 1997a,b); this effect is associated to the loss of *Dlx5* expression via the *Dlx*-dependent sub-pallium-specific enhancer element in the intergenic *Dlx5–Dlx6* region (Zerucha et al., 2000). The disruption of *Dlx5*, in turn, leads to a less dramatic reduction of interneuronal cell differentiation in the OB with an expansion of the progenitor cell compartment (Long et al., 2003).

The SVZ–OB system represents an ideal and well-characterized model to unravel mechanisms regulating the genesis, migration, and differentiation of NPCs in the postnatal mammalian brain (Peretto et al., 1999). However, efforts to define the role of *Dlx5* in the biology of NPCs have been hampered by the facts that (a) *Dlx* genes are expressed in CNS territories simultaneously involved in neurogenesis, migration, and differentiation, making *in vivo* histological analyses difficult, and (b) *Dlx5*-null mice die soon after birth, making the analysis of postnatal neurogenesis impossible.

In this study, we directly investigate growth and differentiation of *Dlx5*^{−/−} NSCs by employing an *in vitro* approach based on neurosphere assay (Stemple and Anderson, 1992; Reynold and Weiss, 1992; Gritti et al., 1996, 1999). NSCs obtained from *Dlx5*-null brains displayed a normal proliferation rate but exhibited a markedly reduced neuronal differentiation. Therefore, *Dlx5* controls key steps necessary for neuronal differentiation of NSCs after cell proliferation.

Results

Expression of *Dlx5* in the SVZ of the mouse brain

The pattern of expression of *Dlx* genes in the embryonic mouse brain has been reported (Bulfone et al., 1993; Simeone et al., 1994; Merlo et al., 2000; Panganiban and Rubenstein, 2002), but still little is known on the expression of *Dlx5* and *Dlx6* in the newborn and adult brain.

To study the pattern of expression of *Dlx5* and *Dlx6*, we used the knock-in reporter *lacZ* as well as *in situ* hybridization; we have already shown that the expression of the *lacZ* reporter reproduces faithfully that of *Dlx5* in *lacZ–Dlx5* knock-in (Acampora et al., 1999), and we confirmed this observation in this study (data not shown). In the newborn brain, very high levels of expression of *Dlx5*, but not of *Dlx6* (Figs. 1A and C) are observed adjacent to the lateral and ventral border of the telencephalic ventricles, a region corresponding to a subependymal layer where the presence of NPCs has been clearly documented. To clarify whether *Dlx5*-expressing cells coincide with the NPCs, we stained adjacent sections with anti-Emx2, a marker of NPCs in the SVZ (Gangemi et al., 2001), and anti- β -galactosidase antibodies. The anti-Emx2 antibody stains a population of cells more adjacent to the ventricle with respect to the location of *Dlx5*/ β -gal positive cells. Although we cannot exclude that a fraction of NPCs are *Dlx5* positive, the large majority of these cells are in a territory not compatible with an NPC identity where usually more differentiated cells are found.

No qualitative difference was observed in the distribution of β -galactosidase positive cells, indicating that the cell lineage normally expressing *Dlx5* is not lost in the mutant animal. Analysis of

lacZ–Dlx5 and *Dlx6* positive cells along the RMS of normal and *Dlx5*^{−/−} brains (Figs. 1G–J) indicates that the cell lineage normally expressing *Dlx5* and *Dlx6* is still present in the absence of *Dlx5* (see also Long et al., 2003).

Expression of *Dlx5* in cultured NSCs

The distribution of *Dlx5* in the SVZ and the defective differentiation and lamination of the OB in *Dlx5*^{−/−} mice could suggest that *Dlx5* plays a role in the differentiation of NPCs. First, we examined if *Dlx5* is expressed in cultured NSCs obtained from the SVZ of wild-type, *Dlx5*^{+/-}, or *Dlx5*^{−/−} newborn brains. NSCs were maintained in culture as neurospheres under non-differentiating conditions (Gritti et al., 1996). We could not detect any significant difference either in the number, in the appearance, or in the self-renewal capacity of neurospheres generated from *Dlx5*-null mice compared to those from normal (wild type or *Dlx5*^{+/-}) brains. Similarly, neurosphere formation was unaffected when cells were derived from E12.5 brains.

Neurospheres were then dissociated, the resulting cell suspensions were plated, and the adhering cells were stained for the expression of *lacZ* before or after induction of differentiation. Few hours after plating, 50–60% of the undifferentiated *Dlx5*^{+/-} or *Dlx5*^{−/−} cells were positive (Figs. 2A–C), indicating that the *Dlx5* promoter is active. When *lacZ* expression was analyzed 24 h after induction of differentiation, the situation changed radically. We never observed *lacZ* expression in cell characterized by an astrocyte-like morphology (independently of their genotype) (Figs. 2E and F, black arrows). Cells with a phenotype characterized by long bipolar processes reminiscent of early neurons were clearly visible in cultures of *Dlx5*^{+/-} NSCs (indicated by arrowheads in Figs. 2D and E), these cells express the *lacZ* reporter in *Dlx5*^{+/-} cultures. We did not observe any of these cells in cultures of *Dlx5*^{−/−} cells after 1 day of differentiation. It must be pointed out, however, that at these early stages of differentiation in culture, none of the classical “neuronal” marker is yet expressed; these markers appear much later during differentiation. It is therefore difficult to call these cells “neurons” as they are not yet completely differentiated. This observation and the defects in OB formation observed *in vivo* in *Dlx5*^{−/−} mice (Long et al., 2003; Levi et al., 2003) suggested that the lack of *Dlx5* impairs the proliferation and/or the proper neuronal differentiation of NSCs.

Proliferation and differentiation of cultured NPCs

Cell proliferation was assayed by incorporation of tritiated thymidine on culture neurospheres obtained from normal and *Dlx5*^{−/−} brains. The same experiment was carried out on cells derived from either newborn or E12.5 brains. We did not observe any significant difference in the rate of thymidine incorporation of NSCs from normal, *Dlx5*^{+/-}, and *Dlx5*^{−/−} animals of the same gestational age (either E12.5 or P0), suggesting that *Dlx5* does not play a major role in the control of NSCs proliferation (data not shown).

Single cell suspensions derived from neurospheres of normal, *Dlx5*^{+/-}, and *Dlx5*^{−/−} mutant mice at E12.5 and at birth were replated in differentiating medium for 7 days. Under these conditions, wild type NSCs differentiate and cause astrocytes, oligodendrocytes, and neurons in a reproducible distribution. To identify and quantify these three cell populations, differentiated cultures




Fig. 4. Efficiency of neuronal differentiation in cultures of *Dlx* mutant NPCs. Cells obtained from embryonic brain (E12.5, left) and neonatal brains (P0, right) are compared. In both cases, the percentage of Tuj1-positive cells over the total number of cells examined is shown for the three *Dlx5* genotypes (indicated on the bottom). The means (six independent experiments per point) and standard deviations are reported. Cultures of neonatal *Dlx5*^{-/-} NSCs generate a significantly (double asterisk = $P < 0.002$) lower number of neurons.

were stained with antibodies that recognize population-specific markers (galactocerebrosidase for oligodendrocytes; β -tubulin for neurons; glial fibrillary acidic protein for astrocytes) (Fig. 3). To estimate the percentage of each cell type present, a minimum of 4,000 cells were counted in each individual experiment. The pattern of differentiation of cultures derived from *Dlx5*^{+/-} brains did not differ significantly from that of wild type cells at any gestational age. In all experiments, a very low percentage of oligodendrocytes (1–2%) was found. This makes a quantitative comparison rather difficult. In spite of this, we could not observe any significant disturbance of the differentiation rate of this cell type. The absence of *Dlx5* was associated with a significant and consistent reduction of the percentage of neurons formed in cultures of newborn *Dlx*^{-/-} NSCs (Figs. 3 and 4). Fig. 4 reports as an histogram the percentage of neurons obtained from differentiating normal and mutant NSCs.

The reduced number of neurons observed in the absence of *Dlx5* could be explained either by impaired ability of NSCs to differentiate or by the activation of a cell death program after the initiation of differentiation. If the latter explanation were true, one should expect a higher number of neurons at early stages of differentiation, which would subsequently decline with time. To rule out this possibility, we examined differentiation following short induction times. No obvious neuronal differentiation could be observed at Day 1 (Fig. 3E and data not shown), and the number of β -tubulin positive cells remained low when cells were examined at intermediate times (starting from Day 3 when the first cells displaying this differentiation marker begin to appear). Moreover, we also examined apoptotic cell death either by estimating the number of morphologically apoptotic nuclei or by staining the cells with an antibody that reacts with activated caspase-3 (data not shown). No increased apoptotic cell death could be observed in *Dlx5*^{-/-} NSCs cultures compared to the control cells.

When the same *in vitro* differentiation experiments were performed with NSCs derived from *Dlx5*^{+/-} or *Dlx*^{-/-} E12.5 embryos, we did not observe any significant difference in the

number of differentiated β -tubulin-immunoreactive cells or of any other cell type in the absence of *Dlx5* (Figs. 3 and 4).

Dlx5 transfection restores normal differentiation of *Dlx5*^{-/-} NSCs

To confirm that the reduced capacity to differentiate into neurons of newborn *Dlx5*^{-/-} NSCs was a consequence of the absence of the gene, we evaluated if forced expression of *Dlx5* could rescue the neuronal differentiation pathway. NSCs derived from *Dlx5*^{-/-} mice at birth were transduced *in vitro* with the pLXIN/*Dlx5* retroviral vector (see Experimental Methods) and selected for 7 days in G418-containing medium. As a control, *Dlx5*^{-/-} NSCs were mock transduced with an empty pLXIN retroviral vector and similarly selected. The transduced cells were then assayed by RT-PCR for *Dlx5* mRNA expression. As shown in Fig. 5, a PCR product corresponding to *Dlx5* mRNA was only present in transduced NSCs.

Mock transduced cells, nontransduced cells, and cells transduced with *Dlx5* were then plated in the differentiating medium along with control cells, as described above. The relative number of the three different cell lineages was assessed by immunostaining as indicated above. Transfection of exogenous *Dlx5* in *Dlx5*^{-/-}




Fig. 5. Transduction with wild-type *Dlx5* cDNA restores neuronal differentiation in *Dlx5*^{-/-} NPCs. Top: RT-PCR for the detection of murine *Dlx5* and β -actin mRNAs in samples from wild-type (left), *Dlx5*^{-/-} NSCs transduced with empty vector (middle), and *Dlx5*^{-/-} NSCs transduced with pLXIN/*Dlx5*. Bottom: Efficiency of neuronal differentiation *in vitro* of wild-type (left), *Dlx5*^{-/-} NSCs transduced with empty vector (middle), and *Dlx5*^{-/-} NSCs transduced with pLXIN/*Dlx5* (right). The percentage of Tuj1-positive cells over the total number of cells examined is shown. The means (three independent experiments per point) and standard deviations are reported. Cultures of neonatal *Dlx5*^{-/-} NSCs transduced with empty vector generate a significantly ($P < 0.02$) lower number of neurons compared both to wild type and pLXIN/*Dlx5*-transduced mutant cultures. The difference in neurogenic potential between wild type NSCs and pLXIN/*Dlx5*-transduced mutant NSCs is not statistically significant.

NSCs restored the capacity of these cells to produce β -tubulin positive neurons (Fig. 5), indicating that the absence of this gene is directly associated with the reduced neurogenic potential of NSCs.

Discussion

Mammalian brain development and patterning depend on the concerted proliferation, migration, differentiation, and death of NPCs (Gage, 2000; Van der Kooy and Weiss, 2000; Anderson, 2001) and in the establishment and selection of appropriate connectivity. All these processes are under the control of gene regulatory networks that are beginning to be unravelled thanks to the generation of targeted mutations in vivo. Indeed, targeted mutations of genes expressed in NPCs can profoundly subvert their migration and homing of to specific target territories (Marin et al., 2000; Marin and Rubenstein, 2001). In this study, we used cultures of NPCs to gain insight on the role of *Dlx5*, a member of the *Dlx* homeobox gene family on the neurogenic process. Our results indicate that *Dlx5* is essential for neuronal differentiation of perinatal, but not embryonic, NPCs.

Around E12.5 of mouse development migratory progenitors from the medial and lateral ganglionic eminence (MGE and LGE) express members of the *Dlx* gene family, including *Dlx1* and *Dlx2* (Anderson et al., 1997a,b) and *Dlx5* (Porteus et al., 1994; Long et al., 2003; Levi et al., 2003), and contribute to the generation of GABAergic interneurons both in the cortex and in the OB. *Dlx* genes are expressed in a partially overlapping pattern: *Dlx2* is expressed in cells throughout the VZ, while *Dlx5* is expressed in cells of the SVZ and in committed cells of the mantle zone. *Dlx2* expression begins before that of *Dlx1* and marks the earliest progenitors; instead expression of *Dlx5* appears later in *Dlx1*- and *Dlx2*-positive postmitotic neuronal progenitors. *Dlx6* expression is minimal in the SVZ and appears strongly in the mantle zone of the GE (Eisenstat et al., 1999). Moreover, progenitors from the GE have been shown to have the potential to differentiate into neurons and glia in vitro (He et al., 2001), and *Dlx2*-positive migratory NPCs can differentiate into astrocytes and oligodendrocytes in the dorsal telencephalon in vivo (Marshall and Goldman, 2002). Hence, while *Dlx2* is expressed in the most uncommitted NPCs in the GE (embryonic) and in the SVZ (neonatal), *Dlx5* and *Dlx6* are likely to be involved in later steps of NPCs maturation; however, there is no information on the postnatal function of *Dlx5* in the differentiation of SVZ cells, as *Dlx5*-null mice die at birth. The culture assay employed here reveals that *Dlx5* does not play any major role in the genesis of astrocytes and oligodendrocytes. This result is not entirely unexpected as only *Dlx2*-positive uncommitted progenitors from the GE have the potential to differentiate into neurons and glia (He et al., 2001; Marshall and Goldman, 2002) while *Dlx5* marks more committed NPCs.

In the embryonic and neonatal mouse brain, *Dlx5* is expressed in both the MGE and the LGE. There are significant differences in the fate of NPCs in these adjacent structures. The LGE hosts progenitors that either generate striatal projection neurons (Deacon et al., 1994; Olsson et al., 1995, 1997, 1998; Anderson et al., 1997a; Wichterle et al., 2001) or migrates along the RMS and contributes granule cells and interneurons to the OBs. The phenotype of mice with a combined loss-of-function mutation of *Dlx1* and *Dlx2* is consistent with the presence of a common precursor for striatal projection neurons and OB interneurons in the LGE since

differentiation defects are observed in both of these compartments. The MGE contributes to some precommitted interneuronal precursor populations that migrate tangentially to various areas of the dorsal telencephalon, hippocampus, and projection neurons in the basal ganglia (Anderson et al., 1997a, 2001; Ladvas et al., 1999; Sussel et al., 1999; Pleasure et al., 2000; Wichterle et al., 1999, 2001; Yun et al., 2002; reviewed in Marin and Rubenstein, 2001).

The most evident lesion observed in the *Dlx5*^{-/-} brain is the overall size reduction of the OB (about 60–70% of its normal size) accompanied by an enlargement of its VZ–SVZ. This defect has been shown by examining the expression pattern of *Dlx1*, *Dlx2*, *Dlx5*^{lacZ}, and *Dlx6* in the mutant OB (Long et al., 2003; Levi et al., 2003). This observation suggests an expansion of the NPCs and/or precommitted cells compartment linked to a reduction of the number of committed cells. In fact, a detailed analysis of OB differentiated interneurons indicates a severe reduction of both TH- and GAD67-positive cells. Our in vitro results indicate that *Dlx5* does not have a major role in controlling NSCs proliferation. This is in line with in vivo studies on *Dlx5*^{-/-} mice: BrdU incorporation analysis and in situ hybridization with *Hes5* have not shown any significant difference in the rate of mitosis and in the size of the proliferative area of the VZ–SVZ in *Dlx5*^{-/-} brains (Long et al., 2003; Levi et al., 2003). The fact that *Dlx5* does not play any major role in control of NSCs cell cycle is also in line with the fact that expression of *Dlx5* in the embryonic VZ–SVZ appears to be mostly confined to postmitotic precursors (Liu et al., 1997; Eisenstat et al., 1999; Stuhmer et al., 2002). Conversely, it is interesting to note that the absence of *Dlx5* does not increase the rate of cell death in the OB (Long et al., 2003; Merlo, unpublished observation). The question remains as to what is the cellular basis for the reduced size of the OB. Most likely, *Dlx5* plays a significant role in the secondary neuronogenesis that occurs in the postnatal brain, while embryonal neuronogenesis is minimally or not affected. The results presented here, based on neurosphere assays, warrant a detailed in vivo analysis of mice bearing a conditional *Dlx5* mutation when these will be available.

If proliferation and apoptosis are not affected, the simpler way to explain the severe reduction of the neurogenic capacity of perinatal *Dlx5*^{-/-} NSCs is to assume that these cells have a preferential capacity to differentiate into glial cells and have lost, in great part, the capacity to differentiate into neurons. This is in agreement with two other findings. First, cultures of *Dlx5*^{+/-} NSCs differentiate in neurons and glia, but *Dlx5* is expressed only in differentiating neurons suggesting that this transcription factor is essential for the neuronal lineage. The second finding derives from our in vivo analyses of *Dlx5*^{-/-} olfactory system: in the *Dlx5*^{-/-} mice, the axons of OE receptor neurons are present but fail to produce functional connections with the OBs. This defect appears to be intrinsic to the OE (Levi et al., 2003). Conversely, in the OB, immature interneurons fail to a large extent to express the differentiation markers TH or GAD67 (Long et al., 2003; Levi et al., 2003). On the basis of in vivo grafting experiments, we could also rule out the possibility that the lack of GABAergic and dopaminergic interneurons in the OB is a mere consequence of the lack of innervation (Levi et al., 2003). This led to the hypothesis that the absence of *Dlx5* directly causes an intrinsic block in the ability of precursor cells to differentiate into neurons.

Lineage tracing using *Dlx5/6*-reporter transgenic mice have indicated that the SVZ of the LGE contains two distinct populations of progenitor cells, both expressing *Dlx* genes: one that expresses the LIM-homeobox protein Islet1 (ISL1) and one that

is negative for this protein but positive for the ETS transcription factor Er81 (Stenman et al., 2003). These two cell populations have a clearly distinct fate: the $Dlx^+/ISL1^+$ population composes mainly striatal progenitors, while the $Dlx^+/Er81^+$ cells contribute mainly to granule cells and interneurons of the OBs. It is interesting to note that defects in the striatal formation have not been reported in $Dlx5^{-/-}$ mice. In the $Dlx5$ mutant brain, cells in the striatum are properly organized and express major differentiation markers (Merlo, unpublished observation). This observation suggests that in spite of a much wider distribution in the LGE and MGE, $Dlx5$ controls the fate of only a subset of NPCs: those implicated in the development of the olfactory system. As the contribution of neurons to the OB from the SVZ via RMS and migration begins mostly perinatally, this could explain why, in our setting, we observe no significant difference in neuronal differentiation from embryonic NSCs.

Experimental methods

Generation of $Dlx5$ null mice

Mice with targeted disruption of $Dlx5$ have been reported (Acampora et al., 1999). The targeted allele had the first and second exons of the gene replaced with the *lacZ* reporter. The modified allele, denominated $Dlx5^{lacZ}$, allows for easy detection of the $Dlx5$ -expressing cell lineage. β -galactosidase expression in $Dlx5^{+/-}$ embryos recapitulates the known $Dlx5$ expression pattern (Simeone et al., 1994; Merlo et al., 2000). Genotypes were determined by PCR amplification of the wild-type and mutant alleles, as described (Acampora et al., 1999).

Brain histology and histochemistry

Embryos up to E14.5 were immersion fixed in 4% PAF in 0.1 M phosphate buffer (PB, pH 7.4) for 6 h, washed in PBS, and either paraffin embedded or cryprotected (20% sucrose). Animal E16.5 or older were perfused with 4% PAF, the brain was dissected, fixed for additional 2–3 h in 4% PAF, then paraffin embedded. For vibratome sectioning, brains were dissected following perfusion, postfixed for 30 min, and sectioned at 100 μ M. For the detection of Emx2 protein in histology sections, brains were perfused with NaCl 0.9%, dissected, snap frozen in isopentane/liquid N_2 for 20–30 s, then transferred in methanol at -70° and stored for 30 days at -70° . The brains were then transferred in 100% ethanol and paraffin embedded.

β -gal was detected by X-gal staining of either whole-mount embryos or vibrating microtome sections by incubating the samples for 16–24 h at 32° , as described (Acampora et al., 1999). Stained sections were fixed in PAF 4% for 2–4 h before mounting and microscopic examination. Immunohistochemistry was carried out in Tris buffer saline using standard protocols. The following primary antibodies were used: rabbit anti- β -gal (1:5000; ICN Biomedical), rabbit anticalretinin (1:1000; Chemicon), rabbit antityrosine hydroxylase (TH; 1:2000; Chemicon), rabbit antiglutamic acid decarboxylase 67 kDa (GAD67; 1:2000; Chemicon), and rabbit anti-Emx1 and anti-Emx2 (1:500). Rabbit and goat antisera were revealed with goat antirabbit (EnVision+, Dako) and rabbit antigoat (Rockland) peroxidase-conjugated secondary antibodies, respectively. Mouse monoclonal antibodies

were revealed with the ARK kit (Dako) according to the manufacturer's specifications. Peroxidase was developed with DAB (Dako).

Probes and in situ hybridization

In vitro transcribed (Promega) DIG-labeled $Dlx5$ cRNA probe was obtained from the entire murine cDNA ($Dlx5$), while the $Dlx6$ probe is obtained from a partial cDNA fragment of 260 bp comprising exons III and IV. These probes specifically recognize the correct mRNA on Northern blot; in situ hybridization of the $Dlx5$ probe on $Dlx5$ null embryo yielded negligible background. Hybridization was carried out on 100- μ m thick vibratome sections of newborn brains perfused with PAF 4% according to the published procedures (Wilkinson, 1992). Prehybridization and hybridization were done in 50% formamide, $5\times$ SSC pH 4.5, 1% SDS, 50 μ g/ml heparin, 50 μ g/ml of yeast tRNA, and 1 μ g/ml of DIG-labeled probe at 70° C. Signal was detected with AP-conjugated anti-DIG Fab fragments (1:5000; Roche-Boehringer) followed by color development with NBT/BCIP. Three brains of each genotype were analyzed.

Primary cultures of NSCs

Primary cultures of NSCs were established as described (Gangemi et al., 2001; Gritti et al., 1996, 1999). Briefly, brains of P0 normal, $Dlx5^{+/-}$, and $Dlx5^{-/-}$ mice were microdissected to obtain the rostral periventricular region after coronal sections; brains of E12.5 mice were microdissected to obtain the whole telencephalic vesicle. The tissue was dissociated with 0.1% trypsin in phosphate-buffered saline (PBS) for 10 min at 37° C in the presence of 10 mM glucose. After centrifugation, cells were carefully dissociated and resuspended in serum-free NSA medium (Euroclone) supplemented with N2 (Gibco/BRL), 2 mM glutamine, and 20 ng/ml EGF and 10 ng/ml bFGF (Peprotech). Neurospheres were mechanically dissociated every 3 days and replated in tissue culture flasks. To assess self-renewal, primary floating neurosphere colonies were subcloned by mechanically dissociating a single neurosphere colony in 0.2 ml of serum-free medium containing FGF2, heparin, and EGF and plated in uncoated 96-well (0.2 ml/well) plates (Nunc). Stem-cell self-renewal was assessed by identifying new neurospheres after a further 7 days in vitro.

For the examination of differentiation properties, dissociated neurospheres were plated at 3×10^4 cells/well in 24-well plates (Costar) on matrigel-coated glass coverslips (Becton Dickinson). After adhesion, cells were cultured for 5 days in serum-free medium containing bFGF and then in medium containing 1% serum for 3–7 days.

Immunocytochemistry on cultured NSCs

Cells were fixed with 4% PAF in PBS for 15 min at room temperature (RT), then incubated with the appropriate primary antibody for 1 h, washed twice with PBS, and incubated with the secondary antibody for 45 min. The following primary antibodies were used: monoclonal anti-galactocerebrosidase (Boehringer Mannheim) 1:10; monoclonal anti β -tubulin (Sigma) 1:100; rabbit anti cow glial fibrillary acidic protein (Dako) 1:300, and rabbit anticaspase-3 1:1000 (BD Pharmingen). Secondary anti-

bodies were: goat antimouse IgG (H + L) FITC and goat antirabbit TRITC (Jackson Immunoresearch) used at 1:200 and 1:250, respectively. For β -gal staining, cells were fixed in glutaraldehyde 0.5% for 15 min, rinsed with PBS, and stained as above.

Estimation of NSCs proliferation in culture

The number of cells in S-phase of the cell cycle was determined by thymidine incorporation. Ten thousand cells/well were plated on flat bottom 96-well plates in quadruplicate and grown for 24, 48, and 72 h. ^3H -methyl-thymidine, 1 μCi /well (Amersham) was added 12 h before harvesting. Cells were lysed at the desired time, blotted on filter paper, and following addition of 5 ml of scintillation fluid, ^3H emission was measured in a β -counter.

Retroviral vector construction and infection

A bicistronic retroviral vector (pLXIN2, Clontech) was constructed with the complete *Dlx5* cDNA and the *neo* gene under the control of the viral LTR (pLXIN/*Dlx5*). The same bicistronic vector with the *neo* gene alone was used as control. Retroviral plasmids were used to transfect the Phoenix packaging cell line. The retroviral titer was evaluated on 3T3 cells and ranged from 5×10^5 to 1×10^6 colony forming unit (CFUs). The efficiency of NSC infection ranged from 40% to 50%. The infection was repeated twice for 8 h, then NP cells were selected for 7 days with G418. Expression of exogenous *Dlx5* mRNA in the transduced cells was tested by RT-PCR. Total RNA was prepared from normal and *Dlx5*^{-/-} newborn mice and from pLXIN/*Dlx5*-infected NSCs using a commercial kit (RNeasy, Qiagen). PCR amplification of *Dlx5* was performed for 35 cycles with the following 5' and 3' oligonucleotides:

5' AGAGTCCCAAGCATCCGATCC 3'

5' CCAGCACAACTGTAGTCCC 3'

Acknowledgments

We thank Dr. Stefano Mantero, Dr. Sonya Trombino, and Ms. Olinda Alegria for their contribution and excellent technical help, Dr. Stefano Bertuzzi for critical reviewing of the manuscript, and Dr. Ottavia Barbieri for maintaining the mouse colony. G.C. was supported by grants from MIUR, Fondi di Ateneo Università di Genova and Ministero della Sanità. G.R.M. is recipient of a Telethon-Italy Career Award (TCP99003) and is supported by Ministero della Sanità and Cariplo Foundation. G.L. is supported by Telethon-Italy (project GP0218/01), ARSEP, the EU consortium "GENOSPORA" QLK6-1999-02108, and a grant from "Vaincre la Mucoviscidose".

References

Acampora, D., Merlo, G., Paleari, L., Zerega, B., Mantero, S., Barbieri, O., Postiglione, M.P., Simeone, A., Levi, G., 1999. Craniofacial, vestibular and bone defects in mice lacking the *Distal-less*-related gene *Dlx5*. *Development* 126, 3795–3809.

Alvarez-Buylla, A., Garcia-Verdugo, J.M., 2002. Neurogenesis in the adult subventricular zone. *J. Neurosci.* 22, 629–634.

Alvarez-Buylla, A., Seri, B., Doetsch, F., 2002. Identification of neural stem cells in the adult vertebrate brain. *Brain Res. Bull.* 57, 751–758.

Anderson, D.J., 2001. Stem cells and pattern formation in the nervous system: the possible versus the actual. *Neuron* 30, 19–35.

Anderson, S.A., Qiu, M., Bulfone, A., Eisenstat, D.D., Meneses, J., Pedersen, R., Rubenstein, J.L.R., 1997a. Mutations of the homeobox genes *Dlx-1* and *Dlx-2* disrupt the striatal subventricular zone and differentiation of late born striatal neurons. *Neuron* 19, 27–37.

Anderson, S.A., Eisenstat, D.D., Shi, L., Rubenstein, J.L., 1997b. Interneuron migration from basal forebrain to neocortex: dependence on *Dlx* genes. *Science* 278, 474–476.

Anderson, S.A., Marin, O., Horn, C., Jennings, K., Rubenstein, J.L.R., 2001. Distinct cortical migrations from the medial and lateral ganglionic eminences. *Development* 128, 353–363.

Bulfone, A., Puelles, L., Porteus, M.H., Frohman, M.A., Martin, G.R., Rubenstein, J.L.R., 1993. Spatially restricted expression of *Dlx-1*, *Dlx-2* (*Tes-1*), *Gbx-2*, and *Wnt3* in the embryonic day 12.5 mouse forebrain defines potentially transverse and longitudinal segmental boundaries. *J. Neurosci.* 13, 3155–3172.

Bulfone, A., Wang, F., Hevner, R., Anderson, S., Cutforth, T., Chen, S., Meneses, J., Pedersen, R., Axel, R., Rubenstein, J.L.R., 1998. An olfactory sensory map develops in the absence of normal projection neurons or GABAergic interneurons. *Neuron* 21, 1273–1282.

Deacon, T.W., Pakzaban, P., Isacson, O., 1994. The lateral ganglionic eminence is the origin of cells committed to striatal phenotypes: neural transplantation and developmental evidence. *Brain Res.* 668, 211–219.

Depew, M.J., Liu, J.K., Long, J.E., Presley, R., Meneses, J.J., Pedersen, R., Rubenstein, J.L.R., 1999. *Dlx5* regulates regional development of the branchial arches and sensory capsules. *Development* 126, 3831–3846.

Eisenstat, D.D., Liu, J.K., Mione, M., Zhong, W., Yu, G., Anderson, S.A., Ghattas, I., Puelles, L., Rubenstein, J.L.R., 1999. *DLX-1*, *DLX-2*, and *DLX-5* expression define distinct stages of basal forebrain differentiation. *J. Comp. Neurol.* 414, 217–237.

Gage, F.H., 2000. Mammalian neural stem cells. *Science* 287, 1433–1438.

Gangemi, R.M., Daga, A., Marubbi, D., Rosatto, N., Capra, M.C., Corte, G., 2001. *Emx2* in adult neural precursor cells. *Mech. Dev.* 109, 323–329.

Gritti, A., Parati, E.A., Cova, L., Frolichsthal, P., Galli, R., Wanke, E., Faravelli, L., Morassutti, D.J., Roisen, F., Nickel, D.D., Vescovi, A.L., 1996. Multipotential stem cells from the adult mouse brain proliferate and self-renew in response to basic fibroblast growth factor. *J. Neurosci.* 16, 1091–1100.

Gritti, A., Frolichsthal, P., Galli, R., Parati, E.A., Pagano, S.F., Bjornson, C.R., Vescovi, A.L., 1999. Epidermal and fibroblast growth factors behave as mitogenic regulators for a single, multipotent, stem-like cells population from the ventricular region of the adult mouse forebrain. *J. Neurosci.* 19, 3287–3297.

Gritti, A., Bonfanti, L., Doetsch, F., Caille, I., Alvarez-Buylla, A., Lim, D.A., Galli, R., Garcia-Verdugo, J.M., Herrera, D.G., Vescovi, A., 2002. Multipotent neural stem cells reside into the rostral extension and olfactory bulb of adult rodents. *J. Neurosci.* 22, 437–445.

He, W., Ingraham, C., Rising, L., Goderie, S., Temple, S., 2001. Multipotent stem cells from the mouse basal forebrain contribute GABAergic neurons and oligo-dendrocytes to the cerebral cortex during embryogenesis. *J. Neurosci.* 21, 8854–8862.

Ladvas, A.A., Grigoriou, M., Pachnis, V., Parnavelas, J.G., 1999. The medial ganglionic eminence gives rise to a population of early neurons in the developing cortex. *J. Neurosci.* 19, 7881–7888.

Levi, G., Puche, A.C., Mantero, S., Barbieri, O., Trombino, S., Paleari, L., Egeo, A., Merlo, G.R., 2003. The *Dlx5* homeodomain gene is essential for olfactory development and connectivity in the mouse. *Mol. Cell. Neurosci.* 22, 530–543.

Liu, J.K., Ghattas, I., Liu, S., Chen, S., Rubenstein, J.L.R., 1997. *Dlx* genes encode DNA-binding proteins that are expressed in an overlapping and

- sequential pattern during basal ganglia differentiation. *Dev. Dyn.* 210, 498–512.
- Lois, C., Alvarez-Buylla, A., 1994. Long-distance neuronal migration in the adult mammalian brain. *Science* 264, 1145–1148.
- Long, J.E., Garel, S., Depew, M.J., Tobet, S., Rubenstein, J.L.R., 2003. *DLX5* regulates development of peripheral and central components of the olfactory system. *J. Neurosci.* 23, 568–578.
- Marin, O., Rubenstein, J.L.R., 2001. A long remarkable journey: tangential migration in the telencephalon. *Nat. Rev., Neurosci.* 2, 780–790.
- Marin, O., Anderson, S.A., Rubenstein, J.L.R., 2000. Origin and molecular specification of striatal interneurons. *J. Neurosci.* 20, 6063–6076.
- Marshall, C.A.G., Goldman, J.E., 2002. Subpallial *Dlx2*-expressing cells give rise to astrocytes and oligodendrocytes in the cerebral cortex and white matter. *J. Neurosci.* 22, 9821–9830.
- McGuinness, T., Porteus, M.H., Smiga, S., Bulfone, A., Kingsley, C., Qiu, M., Liu, J.K., Long, J.E., Xu, D., Rubenstein, J.L., 1996. Sequence, organization, and transcription of the *Dlx-1* and *Dlx-2* locus. *Genomics* 35, 473–485.
- Merlo, G.R., Zerega, B., Paleari, L., Trombino, S., Mantero, S., Levi, G., 2000. Multiple functions of *Dlx* genes. *Int. J. Dev. Biol.* 44, 619–626.
- Nieto, M., Schuurmans, C., Britz, O., Guillemot, F., 2001. Neural bHLH genes control the neuronal versus glial fate decision in cortical progenitors. *Neuron* 29, 401–413.
- Okano, H., 2002. Stem cells biology of the central nervous system. *J. Neurosci. Res.* 69, 698–707.
- Olsson, M., Campbell, K., Wictorin, K., Bjorklund, A., 1995. Projection neurons in fetal striatal transplant are predominantly derived from the lateral ganglionic eminence. *Neuroscience* 69, 1169–1182.
- Olsson, M., Campbell, K., Turnbull, D.H., 1997. Specification of mouse telencephalic and mid-hindbrain progenitors following heterotopic, ultrasound-guided embryonic transplantation. *Neuron* 19, 761–772.
- Olsson, M., Bjorklund, A., Campbell, K., 1998. Early specification of striatal projection neurons and interneuronal subtypes in the lateral and medial ganglionic eminence. *Neuroscience* 84, 867–876.
- Panganiban, G., Rubenstein, J.L.R., 2002. Developmental functions of the *Distal-less/Dlx* homeobox genes. *Development* 129, 4371–4386.
- Peretto, P., Merighi, A., Fasolo, A., Bonfanti, L., 1999. The subependymal layer in rodents: a site of structural plasticity and cell migration in the adult mammalian brain. *Brain Res. Bull.* 49, 221–243.
- Pleasure, S.J., Anderson, S., Hevner, R., Bagri, A., Marin, O., Lowenstein, D.H., Rubenstein, J.L.R., 2000. Cell migration from the ganglionic eminences is required for the development of hippocampal GABAergic interneurons. *Neuron* 28, 727–740.
- Porteus, M.H., Bulfone, A., Ciaranello, R.D., Rubenstein, J.L., 1991. Isolation and characterization of a novel cDNA clone encoding a homeo-domain that is developmentally regulated in the ventral forebrain. *Neuron* 7, 221–229.
- Porteus, M.H., Bulfone, A., Liu, J.K., Puellas, L., Lo, L.C., Rubenstein, J.L.R., 1994. *DLX-2*, *MASH-1*, *MAP-2* expression and bromodeoxyuridine incorporation define molecularly distinct cell populations in the embryonic mouse forebrain. *J. Neurosci.* 14, 6370–6383.
- Price, M., Lemaistre, M., Pischetola, M., Di Lauro, R., Duboule, D., 1991. A mouse gene related to *Distal-less* shows a restricted expression in the developing forebrain. *Nature* 351, 748–751.
- Qiu, M., Bulfone, A., Martinez, S., Meneses, J.J., Shimamura, K., Pedersen, R.A., Rubenstein, J.L.R., 1995. Null mutation of *Dlx-2* results in abnormal morphogenesis of proximal first and second branchial arch derivatives and abnormal differentiation of the forebrain. *Genes Dev.* 9, 2523–2538.
- Quinn, L.M., Johnson, B.V., Nicholl, J., Sutherland, G.R., Kalionis, B., 1997. Isolation and identification of homeobox genes from the human placenta including a novel member of the *Distal-less* family, *DLX4*. *Gene* 187, 55–61.
- Reynold, B.A., Weiss, S., 1992. Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. *Science* 255, 1707–1710.
- Robinson, G.W., Mahon, K.A., 1994. Differential and overlapping expression domains of *Dlx2* and *Dlx3* suggest distinct roles for *Distal-less* homeobox genes in craniofacial development. *Mech. Dev.* 48, 199–215.
- Robinson, G.W., Wray, S., Mahon, K.A., 1991. Spatially restricted expression of a member of a new family of murine *Distal-less* homeobox genes in the developing forebrain. *New Biol.* 3, 1183–1194.
- Schuurmans, C., Guillemot, F., 2002. Molecular mechanisms underlying cell fate specification in the developing telencephalon. *Curr. Opin. Neurobiol.* 12, 26–34.
- Simeone, A., Acampora, D., Pannese, M., D'Esposito, M., Stornaiuolo, A., Gulisano, M., Mallamaci, A., Kastury, K., Druck, T., Huebner, K., Boncinelli, E., 1994. Cloning and characterization of two members of the vertebrate *Dlx* gene family. *Proc. Natl. Acad. Sci. U. S. A.* 91, 2250–2254.
- Stemple, D.L., Anderson, D.J., 1992. Isolation of a stem cell for neurons and glia from the mammalian neural crest. *Cell* 71, 973–985.
- Stenman, J., Toresson, H., Campbell, K., 2003. Identification of two distinct progenitor populations in the lateral ganglionic eminence: implication for striatal and olfactory bulb neurogenesis. *J. Neurosci.* 23, 167–174.
- Stuhmer, T., Anderson, S., Ekker, M., Rubenstein, J.L.R., 2002. Ectopic expression of *Dlx* gene induces glutamic acid decarboxylase and *Dlx* expression. *Development* 129, 245–252.
- Sun, Y., Nadal-Vicens, M., Misono, S., Lin, M.Z., Zubiaga, A., Hua, X., Fan, G., Greenberg, M.E., 2001. Neurogenin promotes neurogenesis and inhibits glial differentiation by independent mechanisms. *Cell* 104, 365–376.
- Sussel, L., Marin, O., Kimura, S., Rubenstein, J.L.R., 1999. Loss of *Nkx2.1* homeobox gene function results in a ventral to dorsal molecular re-specification within the basal telencephalon: evidence for a transformation of the pallidum into the striatum. *Development* 126, 3359–3370.
- Van der Kooy, D., Weiss, S., 2000. Why stem cells? *Science* 287, 1439–1441.
- Wagner, J., Akerud, P., Castro, D.S., Holm, P.C., Canals, J.M., Snyder, E.Y., Perlmann, T., Arenas, E., 1999. Induction of a midbrain dopaminergic phenotype in *Nurr1*-overexpressing neural stem cells by type 1 astrocytes. *Nat. Biotechnol.* 17, 653–659.
- Weiss, S., Dunne, C., Hewson, J., Wohl, C., Wheatley, M., Peterson, A.C., Reynolds, B.A., 1996. Multipotent CNS stem cells are present in the adult mammalian spinal cord and ventricular neuroaxis. *J. Neurosci.* 16, 7599–7606.
- Wichterle, H., Garcia-Verdugo, J.M., Herrera, D.G., Alvarez-Buylla, A., 1999. Young neurons from medial ganglionic eminence disperse in adult and embryonic brain. *Nat. Neurosci.* 2, 461–466.
- Wichterle, H., Turnbull, D.H., Nery, S., Fishell, G., Alvarez-Buylla, A., 2001. In utero fate mapping reveals distinct migratory pathways and fates of neurons born in the mammalian basal forebrain. *Development* 128, 3759–3771.
- Wichterle, H., Lieberman, I., Porter, J.A., Jessell, T.M., 2002. Directed differentiation of embryonic stem cells into motor neurons. *Cell* 110, 3985–3999.
- Wilkinson, D.G., 1992. Whole-mount in situ hybridization of vertebrate embryos. In: Wilkinson, D.G. (Ed.), *In Situ Hybridization: A Practical Approach*. IRL Press, Oxford, pp. 75–83.
- Yun, K., Fischman, S., Johnson, J., Hrabe de Angelis, M., Weinmaster, G., Rubenstein, J.L., 2002. Modulation of the notch signaling by *Mash1* and *Dlx1/2* regulates sequential specification and differentiation of progenitor cell types in the subcortical telencephalon. *Development* 129, 5029–5040.
- Zerucha, T., Stuhmer, T., Hatch, G., Park, B.K., Long, Q., Yu, G., Gamberotta, A., Schultz, J.R., Rubenstein, J.L.R., Ekker, M.A., 2000. Highly conserved enhancer in the *Dlx5/Dlx6* intragenic region is the site of cross-regulatory interactions between *Dlx* genes in the embryonic forebrain. *J. Neurosci.* 20, 709–721.