

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Molecular identification of *Fusarium* spp. associated with bakanae disease of rice in Italy and assessment of their pathogenicity

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/81274> since

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16

1. ***This is an author version of the contribution published on:***

Questa è la versione dell'autore dell'opera:

[Amatulli M.T., Spadaro D., Gullino M.L., Garibaldi A. (2010) - Molecular identification of *Fusarium* spp. associated with bakanae disease of rice in Italy and assessment of their pathogenicity. *Plant Pathology*, 59, 839-844.

] .

]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-3059.2010.02319.x/abstract>]

17
18
19 **Molecular identification of *Fusarium* spp. associated with bakanae disease of rice in Italy**
20 **and assessment of their pathogenicity**

21

22 M.T. Amatulli¹, D. Spadaro^{1,2}, M.L. Gullino¹, A. Garibaldi¹

23

24 ¹ Agroinnova, Centre of Competence for the Innovation in the Agro-Environmental Field,
25 University of Torino, Via Leonardo da Vinci 44, 10095 Grugliasco (To), Italy

26 ² DiVaPRA – Plant Pathology, University of Torino, Via Leonardo da Vinci 44, 10095
27 Grugliasco (To), Italy

28

29 *Abstract*

30 In the last 10 years the *Fusarium* disease of rice has spread through Italian rice fields becoming a
31 serious problem for seed production and for seed companies. Between 2006 and 2008, 146
32 isolates of *Fusarium* spp. were obtained from diseased rice plants and seeds coming from the
33 major rice-growing regions of Italy. These isolates were identified, based on the translation
34 elongation factor sequence and pathogenicity tests to assess their aggressiveness against the
35 susceptible rice cultivar Galileo. Use of the EF-1 α sequence gave reliable identification and
36 showed that *Fusarium fujikuroi*, the causal agent of bakanae disease, is the most abundant
37 *Fusarium* spp. isolated. These data were confirmed by inoculation of the isolates in the
38 greenhouse, showing that only *F. fujikuroi* strains are able to cause bakanae disease. We
39 identified pathogenic strains with different levels of virulence. Phylogenetic analysis based on
40 EF-1 α sequences generated a tree which separated the various *Fusarium* species into different

41 clusters with high bootstrap values.

42

43 *Key words: Fusarium fujikuroi, Oryza sativa, translation elongation factor, aggressiveness.*

44

45 **Introduction**

46

47 Northern Italy has the largest production area (232,500 ha in 2007) of rice in Europe (FAO data).

48 Several fungal diseases affect this crop; among these, *Fusarium* species are the agents of many

49 different plant and seed diseases, such as bakanae. In the last decades, bakanae has emerged as a

50 problem in Italian rice production (Desjardins *et al.*, 1997). The most evident symptom of this

51 disease is yellowing and abnormal elongation of infected seedlings, due to gibberellic acid

52 production, which led to the Japanese name *bakanae*, meaning “foolish seedling.” In older plants,

53 the roots, crowns, stems, leaf sheaths, and panicles can be infected. The disease is seed-borne and

54 primarily seed transmitted (Desjardins *et al.*, 1997).

55 Although bakanae disease was first described over 100 years ago in Japan, it is still not clear

56 which *Fusarium* species are associated with the different symptoms. Early work in Japan

57 identified the pathogen as “*Fusarium moniliforme*” in a broad sense (Ou, 1985); however, this

58 taxon comprises a number of distinct species, now collectively termed the *Gibberella fujikuroi*

59 species complex. The presence of the *Gibberella* sexual stage can distinguish mating populations,

60 or biological species, within this group (Leslie, 1995). Three mating populations of the *G.*

61 *fujikuroi* complex have been associated with bakanae diseased rice. Mating population C (MP-C)

62 (anamorph *Fusarium fujikuroi*, Nirenberg) was first identified in 1977 among strains isolates

63 from Taiwan rice (Hsieh, 1977). MP-A (anamorph *Fusarium verticillioides* (Sacc.) Nirenberg)

64 and MP-D (anamorph *Fusarium proliferatum* (Matsushima) Nirenberg) have been isolated from

65 rice in Asia, and MP-D has also been isolated from rice from Africa, Australia, and the United
66 States (Desjardins *et al.*, 1997; Voigt *et al.*, 1995). The names for the sexual stage of these
67 pathogens are *G. fujikuroi*, *G. intermedia* and *G. moniliforme*. *Fusarium* species within this
68 complex are also able to produce mycotoxins that can affect human and animal health (Leslie *et*
69 *al.*, 1992). In particular *F. fujikuroi* could produce moniliformin, beauvericin, and fumonisins; *F.*
70 *verticillioides* can produce fumonisins, and *F. proliferatum* are able to synthesize fumonisins,
71 beauvericin, and fusaproliferin. Recently molecular techniques to identify and group the species
72 are becoming important. Recent work has revealed great diversity among *Fusarium* species,
73 underestimated by earlier morphological criteria (O'Donnell *et al.*, 1998a, b; Aoki & O'Donnell,
74 1999; O'Donnell *et al.*, 2000; Ward *et al.*, 2002). Different techniques such as AFLP (amplified
75 fragment length polymorphism), RAPD (random amplification of polymorphic DNA), and RFLP
76 (restriction fragment length polymorphism) analysis have been used to characterize and identify
77 the different mating populations in the *G. fujikuroi* complex (Voigt *et al.*, 1995; Patiño *et al.*,
78 2006; Moretti *et al.*, 2004) as well as multiple gene phylogeny (O'Donnell *et al.*, 2000). Geiser
79 (2003) reported that intron-rich parts of protein-coding genes are excellent markers for
80 species-level phylogenetics in fungi. Gene introns seem better than the internal transcribed spacer
81 (ITS) regions of the nuclear ribosomal RNA gene repeat because they tend to evolve at a higher
82 rate (O'Donnell *et al.*, 2000; O'Donnell *et al.*, 1998a). Non-orthologous copies of the ITS2 may
83 also lead to incorrect phylogenetic inferences (O'Donnell *et al.*, 1997). In these circumstances, it
84 is better to identify regions with high phylogenetic utility. The translation elongation factor 1- α
85 gene (EF-1 α gene), which encodes an essential part of the protein translation machinery, is a
86 good single-locus identification tool in *Fusarium* because it shows high sequence polymorphism
87 among closely related species, even compared to the intron-rich portions of protein-coding genes
88 such as calmodulin, beta-tubulin and histone H3, and non-orthologous copies of the gene have

89 not been detected in the genus.

90 Since the identity of the bakanae causal agent seems to be still unclear, the objectives of this
91 study were: i) to assess the variability of *Fusarium* from bakanae-infected rice in Italy, ii) to
92 identify species by amplification and sequencing of the EF-1 α gene, and iii) to evaluate the
93 pathogenicity of the Italian isolates on the susceptible rice cultivar Galileo.

94

95 **Materials and methods**

96

97 *Collection and isolation of Fusarium strains*

98 Bakanae-infected rice samples were collected during the whole growing season for 3 years
99 (2006-2008) in the major rice production areas of north-western Italy (Novara, Vercelli and Pavia
100 provinces). Fungi were isolated from the stem just above the crown of diseased plants of different
101 varieties. Tissues were washed under running tap water, sterilized with 1% NaOCl for 30 s,
102 rinsed twice in sterile distillate water (SDW), placed in Petri plates containing Komada medium
103 (Komada, 1975), and kept at room temperature. Fungi were isolated from the seeds after washing
104 whole seeds under running tap water. The seed surface was treated with NaOCl for 45 s, rinsed
105 twice in SDW, and placed on PDA. After 4 days, mycelia grown from tissues and seeds were
106 transferred to fresh Petri dishes containing potato dextrose agar (PDA, Difco, USA) plus 25 mg/l
107 streptomycin (AppliChem, Germany) in order to induce microconidia. In addition four
108 representative strains (ITEM504 *F. verticillioides*, ITEM1720 *F. proliferatum*, ITEM231 *F.*
109 *verticillioides*, and ITEM1746 *F. verticillioides*) from the of Institute of Food Production
110 Sciences Culture Collection (Bari, Italy), and one (M-1149=ATCC 38939, *F. fujikuroi*) from the
111 Fusarium Research Center (FRC, Pennsylvania State University, USA) were used. The strains
112 were stored both at 4°C in Spezieller Nährstoffarmer agar (SNA: 1.0 g/L KH₂PO₄, 1.0 g/L KNO₃,

113 0.5 g/L MgSO₄·7H₂O, 0.5 g/L glucose, 0.2 g/L sucrose) and at -80°C in 50% glycerol.

114

115 *DNA extraction and amplification*

116 About 100 mg of mycelium was scraped from a Petri dish containing PDA, and DNA extracted
117 from it using the NucleoMag Kit (Macherey Nagel GmbH and Co., Duren, USA) according to the
118 manufacturer's instructions, adding 10 µL of proteinase K (10 mg/l) and 10 µL of RNase A (12
119 mg/l) to MC1 buffer in each tube. DNA was extracted using Kingfisher (Thermo, Waltham,
120 USA) according to the manufacturer's protocols. The PCR amplified the EF-1α gene using
121 specific primers (O'Donnell *et al.*, 1998b): ef1 (5'-ATGGGTAAGGA(A/G)GACAAGAC-3')
122 and ef2 (5'- GGA(G/A)GTACCAGT(G/C)ATCATGTT-3'). The reaction mixture of 20 µL
123 contained 1 µl of buffer 10X, 1 U of Taq DNA polymerase (QIAGEN, Chatsworth, USA), 0.5
124 mM each of dATP, dCTP, dGTP, and dTTP, 0.5 mM of each primer, and 1.5 mM of MgCl₂.
125 Amplification was carried out in a T-Gradient thermal cycler (Biometra, Göttingen, Germany)
126 using the following steps: 95°C for 3 min, 35 cycles with denaturation at 95°C for 1 min, 60°C
127 for 1 min, 72°C for 3 min and a final extension at 72°C for 5 min. Five µl of the PCR products
128 were separated by electrophoresis in 1.5 % agarose gel (Agarose D-1 LOW EEO Eppendorf,
129 Milan, Italy) containing 1 µl/100 ml of SYBR Safe DNA gel stain (Invitrogen, Eugene, USA) for
130 30 min at 3.3 V in 1 X TAE running buffer.

131

132 *Sequencing and phylogenetic analysis*

133 The PCR products were purified using QIAquick PCR purification kit (QIAGEN, Chatsworth,
134 USA) according to the manufacturer's instructions and were sequenced in both directions by
135 BMR Genomics Center (Padova, Italy). Sequencing data were aligned with the programme DNA
136 baser V2.71.0 (Heracle Software Lilienthal, Germany) and then blasted using both NCBI

137 (HYPERLINK "<http://www.ncbi.nih.gov/>" National Center for Biotechnology Information,
138 Rockville Pike, Bethesda, MS, USA) and FUSARIUM-ID (Geiser *et al.*, 2004) to confirm the
139 results. Sequence alignments and analysis were performed using MEGA version 4 (Tamura *et al.*,
140 2007). The dendrogram was obtained using bootstrap analyses (1000 replicates, removing gaps)
141 with UPGMA. Sequences from strains M1148, M1150, M3096, M1250 (*F.fujikuroi*), M5598 (*F.*
142 *thapsinum*), M7491 (*F. nygamai*), M5331 (*F. verticillioides*) from the Fusarium Research Center
143 (FRC, The Pennsylvania State University, USA) were included in the analysis.

144
145 *Pathogenicity tests*

146 The susceptible cultivar Galileo was used to assess the pathogenicity of 121 strains of *Fusarium*
147 spp. Fungal cultures were sub-cultivated in PDA at room temperature. After 15 days plates were
148 flooded with sterile water and scraped with a sterile spatula. The resulting suspensions were
149 filtered through 2 layers of sterile cotton lint and brought to a final concentration of 10⁶ spores/ml
150 in SDW. Before inoculation, rice seeds were treated with hot water for 15 minutes at 60°C, to
151 remove any surface contaminants. A hundred and fifty rice seeds were soaked in the spore
152 suspension and shaken for 10 minutes while control seeds were soaked in SDW for 10 minutes
153 then were air dried. The inoculated seeds were sown in plastic pots (50 seeds per pot) containing
154 sterile soil (60% peat, 40% sand). The plants were watered three times a day. The greenhouse
155 temperature was maintained at 24°-26°C during the day and 16-18°C during the night.
156 Fertilization with macro and micro nutrients was done twice during the 40-day period. Disease
157 symptoms were scored weekly starting from 15 days after inoculation using a 0-5 scale where 0
158 corresponded to asymptomatic plants, 1 to stunted plants with narrow and yellow leaves, 2 to thin
159 plants with narrow, pale yellow leaves, 3 to plants with narrow, pale yellow leaves, thin and long
160 internodes, short root system, 4 to plants with heavy bakanae symptoms and a starting crown

161 necrosis, and 5 corresponded to plant death due to bakanae. The score obtained was standardized
162 onto a 0–100 scale.

163

164 *Statistical analysis*

165 For each strain, the final disease index was calculated as the means of the 3 pots in the 2
166 repetitions. Data analysis was performed by using the SPSS software (SPSS Inc., version 17.0,
167 Chicago, USA). Statistical significance was judged at the level of /p/-value < 0.05. When the
168 analysis of variance was statistically significant, Tukey's test was used to compare the means of
169 pathogenicity level.

170

171 **Results and Discussion**

172

173 **Collection of *Fusarium* spp. from rice and molecular identification of the species**

174 *Fusarium* species were isolated from 98% of plant samples, including tissues and seeds of
175 different rice varieties, collected in different fields. One hundred and forty-six isolates were
176 confirmed as *Fusarium* spp. by their growth and morphological characteristics on PDA.
177 Sequencing was used for further identification. Amplification of the translation elongation factor
178 1 α gave a 660 bp product and its sequence was used for classification based on a BLAST analysis.
179 All the 79 strains obtained from diseased tissues belonged to the *G. fujikuroi* complex, in
180 particular 73 were *F. fujikuroi*. Of the 67 strains isolated from rice seeds, 26 proved to be *F.*
181 *fujikuroi*, 12 *F. proliferatum*, and 7 *F. verticillioides*. Other strains belonging to different species
182 (*F. equiseti*, *F. oxysporum*, *F. graminearum*, *F. brevicatenuatum*, *F. napiforme*, *F. avenaceum*
183 and *F. sporotrichioides*) were identified (Table 1). Molecular identification showed that 85% of
184 our isolates belonged to the *G. fujikuroi* species complex, confirming an earlier report from Italy

185 (Moretti *et al.*,2007). In particular, 72 out of 79 strains isolated from rice tissue showing bakanae
186 symptoms were *F. fujikuroi* confirming the species as the most abundant in rice fields (Anderson,
187 2005; Zainudin *et al.*, 2008). The other species were detected at very low levels and mainly from
188 seeds.

189

190 **Phylogenetic analysis**

191 The EF-1 α sequences obtained were used to create a dendrogram (Figure 1). The three clusters of
192 *F. fujikuroi*, *F. proliferatum* and *F. verticillioides*, although sharing a high level of homology,
193 grouped together with *F. oxysporum* strains and the other species of the *G. fujikuroi* complex in a
194 larger cluster. In another cluster were grouped the other species: *F. equiseti*, *F. graminearum*, *F.*
195 *avenaceum* and *F. sporotrichioides*. The isolates of *F. brevicatenuatum* and *F. napiforme*
196 diverged from both groups. Phylogenetic trees obtained from analyses of EF-1 α sequences
197 clearly grouped the different species of *Fusarium* with high bootstrap values confirming that
198 EF-1 α gene is a good marker for species identification inside *Fusarium* spp..

199

200 **Pathogenicity**

201 One hundred and twenty one isolates of *Fusarium* spp., obtained from naturally infected rice
202 plants, were tested for pathogenicity. The typical bakanae symptoms appeared 14 days after seed
203 sowing only from seedlings inoculated with *F. fujikuroi* strains. The most severely infected plants
204 died within 34 days of inoculation. Bakanae symptoms were not seen on healthy control plants.
205 The *F. fujikuroi* strains were ranked based on their pathogenicity. Sixty one were pathogenic
206 (rated between 52 and 97), 5 were slightly pathogenic (between 24 and 43), and 9 were not
207 pathogenic (between 6 and 0) (Table 2). No bakanae symptoms were induced by other species of
208 the *G. fujikuroi* complex or other *Fusarium* species. Reduction in germination were observed in

209 the plants inoculated with *F. graminearum* strains (15%). Pathogenicity tests showed that only *F.*
210 *fujikuroi* strains can cause bakanae symptoms. All the 61 pathogenic isolates showed abnormal
211 elongation of the seedling, thinning and yellowing, and most plants started to die from 25 dpi.
212 Some isolates did not show the classical elongation symptom, but only tanning and yellowing,
213 confirming the observations of Ou (1985) and Zainudin *et al.* (2008). Different disease indices
214 were observed also among isolates obtained from the same area or the same cultivar. Further
215 studies need to clarify if the variation in aggressiveness among the isolates is due to physiological
216 features of the strain or to environmental factors, such as temperature and humidity during the
217 experiment itself (Pasco *et al.*, 2005). All the pathogenic and not pathogenic *F. fujikuroi* strains
218 and all the strains of *F. verticillioides* and *F. proliferatum* were reisolated from diseased
219 seedlings. This shows that other species besides *F. fujikuroi* can colonise rice tissues but they are
220 not involved in bakanae disease. *F. graminearum* was never re-isolated from inoculated rice
221 plants, though in literature it is reported as a rice pathogen (Lee *et al.*, 2009). The other species,
222 all isolated from seeds, were not re-isolated, and they could probably be considered as epiphytic
223 or saprophytic species. *F. equiseti*, for instance, is known as a saprophytic coloniser that can
224 regularly be recovered as secondary invader of diseased tissues (Summerel *et al.*, 2003).

225

226 **Conclusions**

227

228 The introduction of *F. fujikuroi* in Italy seems to have followed the use of infected material
229 originating from endemic areas. Infected rice seed is the primary source of inoculum, so the best
230 management strategy is to use healthy seed (Dodan *et al.*, 1994). Tanning and thermotherapy (in
231 water at 60-61°C for 15 minutes or aerated steam at 74 ° C for 2 minutes) of seed can reduce the
232 disease by more than 90% (Titone *et al.*, 2003). Others have studied the use of bacteria and fungi

233 as agents of biological control (Adhikari *et al.*, 2001). Good management is also important to
234 avoid contamination with mycotoxins. In the last decade, the organization of *Fusarium* spp. into
235 well defined lineages and their mapping to specific geographic locations have led to the
236 employment of genes involved in mycotoxin biosynthesis or other metabolic processes to study
237 the pathogen populations prevalent in those regions (Brown *et al.*, 2007; Teichert *et al.*, 2006).
238 For this reason, knowledge of the distribution and aggressiveness pattern of *Fusarium* spp. in
239 northern Italian rice fields is very useful for understanding and developing strategies to control
240 the occurrence of bakanae disease.

241

242 **Acknowledgements**

243

244 Work carried out with a grant from the Piedmont region "Control of *Fusarium* diseases on rice:
245 identification of the causal agents, monitoring of mycotoxins in rice-derived products, and
246 selection of resistant varieties". The authors thank Dr. Patrizia Titone and Dr. Veluswamy
247 Karthikeyan for providing the isolates, and Dr. Robert Milne for critically revising the paper.

248

249 **References**

250

251 Adhikari TB, Joseph CM, Yang G, Phillips DA, Nelson LM, 2001. Evaluation of bacteria
252 isolated from rice for plant growth promotion and biological control of seedling disease of rice
253 *Canadian Journal of Microbiology* **47**, 916–924.

254 Anderson L, 2005. *Bakanae disease of rice in California: investigations of disease incidence,*
255 *spread, and pathogen population structure.* Davis, California: University of California, Ph.D
256 thesis.

257 Aoki T, O'Donnell K, 1999. Morphological and molecular characterization of *Fusarium*
258 *pseudograminearum* sp nov., formerly recognized as the Group 1 population of *F. graminearum*.
259 *Mycologia* **91**, 597–609.

260 Brown DW, Butchko RA, Busman M, Proctor R, 2007. The *Fusarium verticillioides* *FUM* gene
261 cluster encodes a Zn(II)₂Cys₆ protein that affects *FUM* gene expression and fumonisin
262 production. *Eukaryot Cell* **6**, 1210–1218.

263 Desjardins AE, Plattner RD, Nelson P, 1997. Production of Fumonisin B1 and Moniliformin by
264 *Gibberella fujikuroi* from Rice from Various Geographic Areas. *Applied and environmental*
265 *microbiology* **63**, 1838-1842.

266 Dodan D, Ram S, Sunder S, 1994. Survival of *Fusarium moniliforme* in infected rice grains and
267 its chemical control. *Indian Journal of Mycology and Plant Pathology* **24**, 135-138.

268 FAO, 2008. FAOSTAT *ProdSTAT. Updated 11 June 2008. *Internet Resource:
269 <http://faostat.fao.org/site/567/default.aspx#ancor>

270 Geiser DM, Gasco MJ, Kang S, Makalowska I, Veeraraghavan N, Ward TJ, Zhang N, Kuldau
271 GA, O'Donnell K, 2004. FUSARIUM-ID v. 1.0: A DNA sequence database for identifying
272 *Fusarium*. *European Journal of Plant Pathology* **110**, 473–479.

273 Geiser DM, 2003. Practical molecular taxonomy of fungi. In: Large L and Tkacz J, eds. *Advances*
274 *in Fungal Biotechnology for Industry, Medicine and Agriculture*. Kluwer, Academic Publishers,
275 Dordrecht, The Netherlands.

276 Komada H, 1975. Development of a selective medium for quantitative isolation of *Fusarium*
277 *oxysporum* from natural soils. *Review of Plant Protection Research* **8**, 114-124.

278 Lee J, Chang IY, Kim H, Yun SH, Leslie JF, Lee YW, 2009 Genetic diversity and fitness of
279 *Fusarium graminearum* populations from rice in Korea. *Applied and Environmental*
280 *Microbiology* **75**, 3289-3295.

281 Leslie, JF, 1995. *Gibberella fujikuroi*: available populations and variable traits. *Canadian*
282 *Journal of Botany* **73**, S282–91.

283 Leslie JF, Plattner RD, Desjardins, AE, Klittich CJR, 1992. Fumonisin B1 production by strains
284 from different mating populations of *Giberella fujikuroi* (*Fusarium moniliforme*).
285 *Phytopathology* **82**, 341-345.

286 Moretti A, Somma S, Picco AM, Rodolfi M, Ritieni A, 2007. Biological, molecular and toxin
287 characterization of *Fusarium* species isolated from bakanae diseased rice plants. *Proceedings of*
288 *4th International Rice conference*, 164-165.

289 Moretti A, Mulè G, Susca A, González-Jaén MT, Logrieco A, 2004. Toxin Profile, Fertility and
290 AFLP Analysis of *Fusarium verticillioides* from Banana Fruits. *European Journal of Plant*
291 *Pathology*, **110**, 601-609.

292 O'Donnell K, Nirenberg HI, Aoki T, Cigelnik E, 2000. A multigene phylogeny of the *Gibberella*
293 *fujikuroi* species complex: Detection of additional phylogenetically distinct species. *Mycoscience*
294 **41**, 61-78.

295 O'Donnell K, Cigelnik E, Nirenberg HI, 1998a. Molecular systematics and phylogeography of
296 the *Gibberella fujikuroi* species complex. *Mycologia* **90**, 465–493.

297 O'Donnell K, Kistler HC, Cigelnik E, Ploetz RC, 1998b. Multiple evolutionary origins of the
298 fungus causing Panama disease of banana: Concordant evidence from nuclear and mitochondrial
299 gene genealogies. *Applied biological Sciences* **95**, 2044-2049.

300 O'Donnell K, Cigelnik E, Weber NS, Trappe JM, 1997. Phylogenetic relationships among
301 ascomycetous truffles and the true and false morels from 18S and 28S ribosomal DNA sequence
302 analysis. *Mycologia* **89**, 48–65.

303 Ou SH, 1985. Rice diseases. Commonwealth Mycological Institute, Kew, England.

304 Patiño B, Mirete S, Vázquez C, Jiménez M, Rodríguez MT, González-Jaén MT, 2006.

305 Characterization of *Fusarium verticillioides* strains by PCR-RFLP analysis of the intergenic
306 spacer region of the rDNA. *Journal of the Science of Food and Agriculture* **86**, 429–435

307 Pasco C, Jouan B, Andrivon D, 2005. Resistance of potato genotypes to common and netted
308 scab-causing species of *Streptomyces*. *Plant Pathology* **54**, 383-392

309 Summerell BA, Salleh B, Leslie JF, 2003. A utilitarian approach to *Fusarium* identification.
310 *Plant Disease* **87**, 117-128.

311 Tamura K, Dudley J, Nei M, Kumar S, 2007. *MEGA4*: Molecular evolutionary genetics analysis
312 (MEGA) software version 4.0. *Molecular Biology and Evolution* **24**,1596-1599.

313 Teichert S, Wottawa M, Schönig B, Tudzynski B, 2006. Role of the *Fusarium fujikuroi* TOR
314 Kinase in Nitrogen Regulation and Secondary Metabolism *Eukaryotic Cell* **5**, 1807-1819.

315 Titone P, Tamborini L, Polenghi G, Garibaldi A, 2003. Efficacia della concia chimica e fisica nei
316 confronti della fusariosi del riso. *Sementi Elette* **49(4)**, 23-28.

317 Voigt K, Schleier S, Bruckner B, 1995. Genetic variability in *Gibberella fujikuroi* and some
318 related species of the genus *Fusarium* based on random amplification of polymorphic DNA
319 (RAPD). *Current Genetetics* **27**, 528–535.

320 Ward TJ, Bielawski JP, Kistler HC, Sullivan E, O’Donnell K, 2002. Ancestral polymorphism and
321 adaptive evolution in the trichothecene mycotoxin gene cluster of phytopathogenic *Fusarium*.
322 *PNAS* **99**, 9278–9283.

323 Zainudin NAIZ, Razak AA, Salleh B, 2008. Bakanae disease of rice in Malaysia and Indonesia:
324 etiology of the causal agent based on morphological, physiological and pathogenicity
325 characteristics. *Journal of Plant Protection and Research* **48**, 475-485.

326

327 **Table 1.** *Fusarium* spp. isolates from diseased rice tissues and seeds identified by sequences in
 328 the EF-1 α gene.

329

<i>Fusarium</i> species	N° of isolates identified for each species				Total
	From seeds		From rice tissues		
	2006-2007	2008	2006-2007	2008	
<i>F. fujikuroi</i> *	11	15	19	53	98
<i>F. proliferatum</i> *	10	2	6	0	18
<i>F. verticillioides</i> *	6	1	1	0	8
<i>F. equiseti</i> *	6	2	0	0	8
<i>F. graminearum</i>	3	1	0	0	4
<i>F. oxysporum</i>	4	0	0	0	4
<i>F. avenaceum</i>	0	1	0	0	1
<i>F. brevicatenuatum</i>	1	0	0	0	1
<i>F. napiforme</i>	1	0	0	0	1
<i>F. sporotrichioides</i>	0	1	0	0	1
<i>Fusarium</i> spp.	2	0	0	0	2
Total	67		79		146

330

331 * The number of samples isolated from seed and tissue cannot be considered identical, based on
 332 the χ^2 test (p -value < 0.01).

333

334

335

336 **Table 2.** List of *F. fujikuroi* isolates tested for pathogenicity on rice susceptible cv. Galileo, their
337 degree of pathogenicity and Disease Severity Index (DSI) on inoculated rice seedling at 34 dpi.

Isolate Code	1st R	2nd R	M*	Isolate Code	1st R	2nd R	M*	Isolate Code	1st R	2nd R	M*
pathogenic				CSA 44	98	81	90 h	SG12	91	85	88 gh
11-3	96	88	92 h	D4	98	79	89 h	VAE2	78	95	87 fgh
11-80	79	87	83 gh	D8	85	56	71 e-h	VAE5	91	84	88 fgh
11-93	90	86	88 h	Da3	55	63	59 d-h	VAE7	65	62	64 d-h
14-22	96	98	97 h	Da5	71	67	69 e-h	VAL1	74	78	76 e-h
15-66	70	68	69 d-h	G1S	87	83	85 gh	Ve13	79	77	78 e-h
19-77	79	70	75 e-h	G3S	65	73	69 e-h	Ve9	81	83	82 gh
2-179	78	98	88 h	G6S	47	64	56 d-h	VN4	95	81	88 fgh
2-42	77	95	86 gh	GI7S	81	79	80 gh	VPG2	90	77	84 gh
2-76	89	99	94 h	I 1,1	47	60	54 d-h	slightly pathogenic			
2-82	62	66	64 d-h	I 1,3	93	90	92 h	2-10	27	20	24 abc
A 2-44	91	77	84 gh	I 1,5	89	74	82 gh	2-147	24	35	30 a-d
A10	69	81	75 fgh	I 2,1	54	50	52 d-h	11-85	27	12	20 abc
A4-27	45	67	56 d-h	I 2,4	49	59	54 d-h	I 1,6	39	46	43 b-e
A5	97	91	94 h	I 2,6	75	83	79 fgh	S7	39	47	43 b-e
A6	69	79	74 e-h	I 3,1	64	62	63 d-h	not pathogenic			
A7	60	64	62 d-h	L5	98	80	89 h	11-23	6	5	6 ab
AC6	71	59	65 e-h	LS3	45	64	55 d-h	11-50	1	4	3 a
AG8	71	79	75 e-h	M1149a	73	66	70 e-h	19-53	1	8	5 ab
C1S	84	84	84 fgh	S10	85	71	78 fgh	AG2	8	0	4 a
C3S	73	82	78 fgh	S2	68	55	62 d-h	C2S	2	2	2a
C4S	73	76	75 e-h	S2S	96	92	94 h	LS6	5	1	3 a
C5S	68	65	67 e-h	S3	77	89	83 gh	SG4	1	0	1 a
C6S	98	79	89 fgh	S4	69	65	67 e-h	VAG6	1	0	1 a
Ca1	79	81	80 fgh	S5S	85	91	88 h	VPG6	0	2	1 a
Ca2	67	79	73 e-h	S6	63	75	69 e-h				
CFG	67	54	61 d-h	S6S	94	84	89 h	Control	0	0	0 a

339 ^a strains obtained from the Fusarium Research Center (FRC, The Pennsylvania State University, USA).

340 ^bDisease severity index . Statistical analysis were performed using Tukey test. Values with different letters are significantly different at

341 $p \leq 0.05$.

342 **Figure caption**

343

344 **Figure 1.** UPGMA analysis inferred from sequences of the elongation factor gene. Bootstrap
345 values are indicated as percentage above the nodes for maximum parsimony analysis (1,000
346 replications). Because of the high number of samples, 5 Italian isolates were used as
347 representative of *F. fujikuroi*, *F. proliferatum* and *F. verticillioides* (GenBank accession number
348 is given next to the strain code).