



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Infinitely many solutions for a class of semilinear elliptic equations in RN

This is a pre print version of the following article:
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/107183 since
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

## Infinitely many solutions for a class of semilinear elliptic equations in $\mathbb{R}^N$

Francesca Alessio<sup>1</sup>, Paolo Caldiroli<sup>2</sup> and Piero Montecchiari<sup>3</sup>

<sup>1</sup> Dipartimento di Matematica, Università degli Studi di Torino via Carlo Alberto, 10 – I 10129 Torino e-mail: alessio@dm.unito.it

<sup>2</sup> Scuola Internazionale Superiore di Studi Avanzati via Beirut, 2-4 – I 34013 Trieste e-mail: paolocal@sissa.it

<sup>3</sup> Dipartimento di Matematica, Università degli Studi di Ancona via Brecce Bianche – I 60131 Ancona e-mail: montecch@popcsi.unian.it

In this article we discuss some results obtained by the authors about the semilinear elliptic problem

$$\begin{cases} -\Delta u + u = a(x)f(u) & \text{in } \mathbb{R}^N\\ u \in H^1(\mathbb{R}^N) \end{cases}$$
(P<sub>a</sub>)

where  $a \in L^{\infty}(\mathbb{R}^N)$  is such that  $\liminf_{|x|\to\infty} a(x) > 0$ , and  $f \in C^1(\mathbb{R})$  satisfies:

- (f1) there exists C > 0 such that  $|f(s)| \leq C(1 + |s|^p)$  for any  $s \in \mathbb{R}$ , where  $p \in (1, \frac{N+2}{N-2})$  if  $N \geq 3$ , or p > 1 if N = 2,
- (f2) there exists  $\theta > 2$  such that  $0 < \theta F(s) \le f(s)s$  for any  $s \ne 0$ , where  $F(s) = \int_0^s f(r) dr$ ,

 $<sup>$^2$</sup>Comunicazione tenuta dal secondo autore il 15 settembre 1999 a Napoli in occasione del XVI Convegno U.M.I.$ 

(f3)  $\frac{f(s)}{s} < f'(s)$  for any  $s \neq 0$ .

Note that  $f(s) = |s|^{p-1}s$  verifies (f1)-(f3) for a subcritical exponent p, namely, for  $p \in (1, \frac{N+2}{N-2})$  if  $N \ge 3$ , or p > 1 if N = 1, 2.

Such kind of problem has been widely studied by variational methods. In particular, solutions to  $(P_a)$  can be obtained as critical points of the functional  $\varphi_a: H^1(\mathbb{R}^N) \to \mathbb{R}$  defined by

$$\varphi_a(u) = \frac{1}{2} ||u||^2 - \int_{\mathbb{R}^N} a(x) F(u(x)) dx$$

Here ||u|| denotes the standard norm on the Sobolev space  $H^1(\mathbb{R}^N)$ . Thanks to the assumptions made on a and f, the functional  $\varphi_a$  turns out to be of class  $C^1$ on  $H^1(\mathbb{R}^N)$ , and it satisfies the geometric properties of the mountain pass lemma. However, in general, the Palais-Smale condition fails, because of the noncompactness of the Sobolev embedding of  $H^1(\mathbb{R}^N)$  into  $L^2(\mathbb{R}^N)$ .

We point out that this lack of compactness is not just a technical difficulty for the existence problem, due to the variational approach, but it is an intrinsic feature of the problem itself, that may be responsible of nonexistence results, as it happens in some cases. Nevertheless, as we will discuss in the next section, one can take advantage of the lack of compactness in order to recover even existence of infinitely many solutions to  $(P_a)$  in a "generic" situation.

Before explaining this viewpoint in more detail, it is worth giving an overview of some results already known in the literature, which show quite clearly that existence/nonexistence or multiplicity of solutions to  $(P_a)$  is strongly affected by the behaviour of the coefficient a.

In the simplest case  $a(x) \equiv 1$ , discussed in [33], one can recover compactness, and then existence of a positive solution, by restricting to the subspace of radial functions  $H^1_{rad}(\mathbb{R}^N)$ , which is compactly embedded into  $L^q(\mathbb{R}^N)$  for q > 2, q subcritical. We also remark that in this case some renowned results by Gidas, Ni, and Nirenberg [21], based on the maximum principle, also guarantee the uniqueness, up to translations, of the positive solution to  $(P_a)$ .

An important progress in the study of the problem  $(P_a)$  has been made with the celebrated papers by P.L. Lions on the concentration-compactness principle [26]. This has supplied a better understanding of the possible ways of losing compactness, and has permitted to consider the case of a nonnegative coefficient a such that there exists  $\lim_{|x|\to\infty} a(x) \in (0, +\infty)$ . Under this additional assumption one can prove that the Palais-Smale condition holds true at some levels. This fact has been used to obtain many different existence results. Among many others, we mention, as examples, [19], [34], [9], [10], [13], [23] and [5].

On the other side, small  $L^{\infty}$  perturbations of the coefficient may change from a problem in which compactness holds to a problem with no nontrivial solution. For instance, in [20] it is showed that if *a* is nonconstant and monotone in one direction then the related problem  $(P_a)$  has only the solution  $u \equiv 0$ .

Another different kind of phenomena appear when the coefficient a has an oscillatory behaviour. For example, when a is periodic, the invariance under translations permits to prove existence, [30], and also multiplicity results, as in [8], [17], [1], [27], where infinitely many solutions (distinct up to translations) are found. In fact, in this case, the noncompactness of the problem can be exploited to set up a new minimax argument, in the spirit of the works [16] and [32], and then to exhibit a rich structure of the set of solutions.

Multiplicity results have been obtained also without periodicity or asymptotic assumptions on a, in some "perturbative" settings, where concentration phenomena occur and a localization procedure can be used to get some compactness in the problem. A first result in this direction is the paper [24] concerning the prescribed scalar curvature problem on  $\mathbb{S}^3$  and  $\mathbb{S}^4$ . We also mention [31], [6], [7], [18], [22], [25] and the references therein, for the case of a nonlinear stationary Schrödinger equation  $-\epsilon^2 \Delta u + V(x)u = f(u)$  with  $\epsilon > 0$  small and  $V \in C^1(\mathbb{R}^N)$ ,  $V(x) \ge V_0 > 0$  in  $\mathbb{R}^N$ , having local maxima or minima or other topologically stable critical points. Similar concentration phenomena occur also considering the equation  $-\Delta u + \lambda u = a(x)f(u)$ for  $\lambda > 0$  large enough (see [15]) or  $-\Delta u + u = a(x)|u|^{p-1}u$  with  $p = \frac{N+2}{N-2} - \epsilon$ ,  $\epsilon > 0$ small, and  $N \ge 3$ , where a blow-up analysis can be done (see [28]).

In the paper [4], we adopt a quite different viewpoint from the ones followed in the works quoted in the Introduction, and we show that the existence of infinitely many solutions for the problem  $(P_a)$  is a generic property with respect to  $a \in L^{\infty}(\mathbb{R}^N)$ , with  $\liminf_{|x|\to\infty} a(x) \geq 0$ . Precisely we prove

**Theorem 1.** Let  $f \in C^1(\mathbb{R})$  satisfy (f1)-(f3). Then there exists a set  $\mathcal{A}$  open and dense in  $\{a \in L^{\infty}(\mathbb{R}^N) : \liminf_{|x|\to\infty} a(x) \geq 0\}$  such that for every  $a \in \mathcal{A}$  the problem  $(P_a)$  admits infinitely many solutions.

In fact, given any  $a \in L^{\infty}(\mathbb{R}^N)$  with  $\liminf_{|x|\to\infty} a(x) > 0$ , for all  $\bar{\alpha} > 0$  we are able to construct a function  $\alpha \in C(\mathbb{R}^N)$ ,  $0 \leq \alpha(x) \leq \bar{\alpha}$  in  $\mathbb{R}^N$ , such that the problem  $(P_{a+\alpha})$  admits infinitely many solutions.

The function  $\alpha$  is obtained in a constructive way that can be roughly described as follows. First, we introduce the variational setting and we make a careful analysis of the hull of the functionals "at infinity"  $\{\varphi_b : b \in H_{\infty}(a)\}$ , where  $H_{\infty}(a)$  is the set of the  $w^*-L^{\infty}$  limits of the sequences  $a(\cdot + x_j)$  with  $(x_j) \subset \mathbb{R}^N$ ,  $|x_j| \to \infty$ . All the functionals at infinity have a mountain pass geometry and, called c(b) the mountain pass level of  $\varphi_b$ , we can show that there exists  $a_{\infty} \in H_{\infty}(a)$  such that  $c(a_{\infty}) \leq c(b)$  for any  $b \in H_{\infty}(a)$  and the corresponding problem  $(P_{a_{\infty}})$  admits a solution characterized as mountain pass critical point. Then, following a suitable sequence  $(x_j) \subset \mathbb{R}^N$  such that  $a(\cdot + x_j) \to a_{\infty} w^*-L^{\infty}$ , we construct a family  $\alpha_{\omega}$  $(\omega > 0 \text{ small})$  of perturbations of a by setting:

$$\alpha_{\omega}(x) = \begin{cases} \bar{\alpha}(1 - \frac{\omega^2}{4}|x - x_j|^2) & \text{for } |x - x_j| \le \frac{2}{\omega} , \ j \ge j(\omega) \\ 0 & \text{otherwise,} \end{cases}$$

where  $\bar{\alpha} > 0$  is fixed small enough, and  $j(\omega)$  is a suitable positive integer (in fact,  $j(\omega) \to +\infty$  as  $\omega \to 0$ ). Note that for every  $\omega > 0$  small,  $\|\alpha_{\omega}\|_{\infty} = \bar{\alpha}$  and the support of  $\alpha_{\omega}$  is the union of infinitely many disjoint balls of radius  $\frac{2}{\omega}$  and center  $x_j$ . Then, we focus on the family of functionals  $\varphi_{\omega} = \varphi_{a+\alpha_{\omega}}$  and we prove that they satisfy local compactness properties in some sets of the type

$$A_{j}(\omega) = \{ u \in H^{1}(\mathbb{R}^{N}) : \varphi_{\omega}(u) \leq c(a_{\infty} + \bar{\alpha}) + \epsilon, \\ \|\varphi'_{\omega}(u)\| \leq \epsilon, \\ \exists y \in B_{\frac{1}{\alpha}}(x_{j}) \ s.t. \ \|u\|_{H^{1}(B_{1}(y))} \geq \bar{\rho} \}$$

where  $\epsilon, \bar{\rho} > 0$  are some constant independent of  $\omega$  and j. Finally, the mountain pass structure of the limiting functional  $\varphi_{\infty} = \varphi_{a_{\infty}+\bar{\alpha}}$  permits to define local minimax classes for the perturbed functionals  $\varphi_{\omega}$  related to the sets  $A_j(\omega)$ . This fact, toghether with the compactness properties that hold in  $A_j(\omega)$ , yields the existence of a critical point in  $A_j(\omega)$ , for any  $\omega > 0$  small enough, and  $j \in \mathbb{N}$  sufficiently large. Such as critical point turns out to be "stable" under perturbations of the cofficient  $a + \alpha_{\omega}$  which are small in  $L^{\infty}$ . Hence, the following result holds true:

**Theorem 1.** There exists  $\hat{\omega} > 0$  such that for every  $\omega \in (0, \hat{\omega})$  the problem  $(P_{a+\alpha_{\omega}})$  admits infinitely many solutions. In addition, there exists  $\beta_0 > 0$  such that for all  $\omega \in (0, \hat{\omega})$  and  $\beta \in L^{\infty}(\mathbb{R}^N)$  with  $\|\beta\|_{L^{\infty}(\mathbb{R}^N)} \leq \beta_0$ , also the problem  $(P_{a+\alpha_{\omega}+\beta})$  admits infinitely many solutions.

We point out that the perturbation  $\alpha_{\omega}$  is explicitly known, up to the sequence  $(x_j)$  that depends on the behaviour of a at infinity. However, in some simple cases, also the sequence  $(x_j)$  can be prescribed a priori. For instance, if  $a(x) \to a_{\infty} \in (0, +\infty)$  as  $|x| \to \infty$ , then any sequence  $(x_j) \subset \mathbb{R}^N$  such that  $|x_{j+1}| - |x_j| \to +\infty$  works well. Similarly, if a is periodic in each variable, then one can take  $(x_j)$  on the period lattice, with the same divergence property as before.

We note that, by a standard argument (taking  $\bar{f}$  instead of f, defined by  $\bar{f}(t) = 0$ for  $t \leq 0$  and  $\bar{f}(t) = f(t)$  for t > 0), it is possible to show the existence of infinitely many positive classical solutions of the problem  $(P_a)$  for any  $a \in \mathcal{A}$ , a smooth.

Finally we want to point out some possible easy extensions of our result. We observe firstly that with minor change, our argument can be used to prove an analogous result for the class of the nonlinear Schrödinger equations  $-\Delta u + b(x)u = a(x)f(u)$  with  $b \in L^{\infty}(\mathbb{R}^N)$ ,  $b(x) \geq b_0 > 0$  for a. e.  $x \in \mathbb{R}^N$ , and a and f as above. Moreover, we point out that in proving Theorems 1 and 2 we never use comparison theorems based on the maximum principle. Then our argument can be repeated exactly in the same way to study systems of the form  $-\Delta u + u = a(x)\nabla F(u)$  where  $F \in C^2(\mathbb{R}^N, \mathbb{R}^M)$  satisfies properties analogous to (f1)-(f3). In particular the result can be established in the framework of the homoclinic problem for second order Hamiltonian systems in  $\mathbb{R}^M$  (see [3] and the references therein).

Secondly we remark that the solutions we find satisfy suitable stability properties. These can be used to prove that in fact the perturbed problem  $(P_{a+\alpha})$  admits multibump type solutions (see [32]) with bumps located around the points  $x_j$ . We refer in particular to [2] for a proof that can be adapted in this setting.

Finally we mention also the fact that if a is assumed to be positive and almost periodic (see [12]) then it is not known whether or not the problem  $(P_a)$  admits solutions. Following [3] it is possible to show that in this case one can construct a perturbation  $\alpha$  almost periodic and with  $L^{\infty}$  norm small as we want, in such a way that the problem  $(P_{a+\alpha})$  admits infinitely many (actually multibump type) solutions. Then we get a genericity result (with respect to the property of existence of infinitely many solutions) for the class of problems  $(P_a)$  with  $a \in C(\mathbb{R}^N)$  positive and almost periodic.

## References

- S. Alama and Y.Y. Li, Existence of solutions for semilinear elliptic equations with indefinite linear part, J. Diff. Eq. 96 (1992), 88-115.
- [2] F. Alessio and P. Montecchiari, Multibump solutions for a class of Lagrangian systems slowly oscillating at infinity, Ann. Inst. H. Poincaré, Anal. non linéaire, 16 (1999), 107-135.
- [3] F. Alessio, P. Caldiroli and P. Montecchiari, Genericity of the multibump dynamics for almost periodic Duffing-like systems, Proc. Roy. Soc. Edinburgh Sect. A 129 (1999), 885-901.
- [4] F. Alessio, P. Caldiroli and P. Montecchiari, Genericity of the existence of infinitely many solutions for a class of semilinear elliptic equations in ℝ<sup>N</sup>, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 27 (1998), 47-68.
- [5] F. Alessio, P. Caldiroli and P. Montecchiari, On the existence of homoclinics for the asymptotically periodic Duffing equation, Top. Meth. Nonlinear Anal. 12 (1998), 275-292.
- [6] A. Ambrosetti and M. Badiale, Homoclinics: Poincarè-Melnikov type results via a variational approach, Ann. Inst. H. Poincaré, Anal. Non Linéaire, 15 (1998), 233-252.
- [7] A. Ambrosetti, M. Badiale and S. Cingolani, Semiclassical states of nonlinear Schrödinger equation, Arch. Rat. Mech. Anal. 140 (1997), 285-300.
- [8] S. Angenent, The Shadowing Lemma for Elliptic PDE, Dynamics of Infinite Dimensional Systems (S.N. Chow and J.K. Hale eds.) F37 (1987).
- [9] A. Bahri and Y.Y. Li, On a Min-Max Procedure for the Existence of a Positive Solution for Certain Scalar Field Equation in  $\mathbb{R}^n$ , Rev. Mat. Iberoamericana 6 (1990), 1-15.
- [10] A. Bahri and P.L. Lions, On the existence of a positive solution of semilinear elliptic equations in unbounded domains, Ann. Inst. H. Poincaré, Anal. non linéaire 14 (1997), 365-413.

- [11] H. Berestycki and P.L. Lions, Nonlinear scalar field equations, Arch. Rat. Mech. Anal. 82 (1983), 313-345.
- [12] A.S. Besicovitch, Almost Periodic Functions, Dover Publications Inc. (1954).
- [13] D.M. Cao, Positive solutions and bifurcation from the essential spectrum of a semilinear elliptic equation in  $\mathbb{R}^n$ , Nonlinear Anal. T.M.A. **15** (1990), 1045-1052.
- [14] D.M. Cao, Multiple solutions of a semilinear elliptic equation in  $\mathbb{R}^n$ , Ann. Inst. H. Poincaré, Anal. Non Linéaire **10** (1993), 593-604.
- [15] D.M. Cao and E.S. Noussair, Multiplicity of positive and nodal solutions for nonlinear elliptic problems in  $\mathbb{R}^n$ , Ann. Inst. H. Poincaré, Anal. Non Linéaire **13** (1996), 567-588.
- [16] V. Coti Zelati, I. Ekeland and E. Séré, A variational approach to homoclinic orbits in Hamiltonian systems, Math. Ann. 288 (1990), 133-160.
- [17] V. Coti Zelati and P.H. Rabinowitz, Homoclinic type solutions for a semilinear elliptic PDE on  $\mathbb{R}^n$ , Comm. Pure Appl. Math. 45 (1992), 1217-1269.
- [18] M. Del Pino and P.L. Felmer, Multi-peak bound states for nonlinear Schrödinger equations, Ann. Inst. H. Poincaré, Anal. Non Linéaire, 15 (1998), 127-149.
- [19] W.Y. Ding and W.M. Ni, On the existence of a positive entire solution of a semilinear elliptic equation, Arch. Rat. Mech. Anal. 91 (1986), 283-308.
- [20] M.J. Esteban and P.L. Lions, Existence and nonexistence results for semilinear elliptic problems in unbounded domains, Proc. Roy. Soc. Edinburgh 93 (1982), 1-14.
- [21] B. Gidas, W.-M. Ni and L. Nirenberg, Symmetry of positive solutions of nonlinear elliptic equations in R<sup>n</sup>, Math. Anal. Appl. Part A, Adv. Math. Suppl. Studies 7A (1981), 369-402.
- [22] C. Gui, Existence of multi-bump solutions for nonlinear Schrödinger equations via variational methods, Comm. Part. Diff. Eq. 21 (1996), 787-820.
- [23] Y. Li, Remarks on a semilinear elliptic equations on  $\mathbb{R}^N$ , J. Diff. Eq. **74** (1988), 34-39.
- [24] Y.Y. Li, Prescribing scalar curvature on S<sup>3</sup>, S<sup>4</sup> and related problems, J. Funct. Anal. 118 (1993), 43-118.
- [25] Y.Y. Li, On a singularly perturbed elliptic equation, Adv. Diff. Eq. 2 (1997), 955-980.
- [26] P.-L. Lions, The concentration-compactness principle in the calculus of variations: the locally compact case, Part I and II, Ann. Inst. H. Poincaré, Anal. Non Linéaire 1 (1984), 109-145 and 223-283.

- [27] P. Montecchiari, Multiplicity results for a class of Semilinear Elliptic Equations on  $\mathbb{R}^m$ , Rend. Sem. Mat. Univ. Padova **95** (1996), 1-36.
- [28] R. Musina, Multiple positive solutions of a scalar field equation in R<sup>n</sup>, Top. Meth. Nonlinear Anal. 7 (1996), 171-185.
- [29] W.M. Ni, Some aspects of semilinear elliptic equations Nonlinear diffusion equations and their equilibrium states (W.M. Ni, L.A. Peletier and J. Serrin, eds.) Springer Verlag, Berlin (1988).
- [30] P.H. Rabinowitz, A note on a semilinear elliptic equation on  $\mathbb{R}^n$ , Nonlinear Analysis, a tribute in honour of Giovanni Prodi (A. Ambrosetti and A. Marino, eds., Quaderni della Scuola Normale Superiore, Pisa) (1991).
- [31] P.H. Rabinowitz, On a class of nonlinear Schrödinger equations, Z. Angew. Math. Phys. 43 (1992), 270-291.
- [32] E. Séré, Looking for the Bernoulli shift, Ann. Inst. H. Poincaré, Anal. Non Linéaire 10 (1993), 561-590.
- [33] W.A. Strauss, Existence of solitary waves in higher dimensions, Comm. Math. Phys. 55 (1979), 149-162.
- [34] C.A. Stuart, Bifurcation in  $L^p(\mathbf{R}^n)$  for a semilinear elliptic equation, Proc. London Math. Soc. (3) 57 (1988), 511-541.