

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Role of allogeneic transplantation in multiple myeloma in the era of new drugs.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/100198> since

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Perspectives

Role of Allogeneic Transplantation in Multiple Myeloma in the Era of New Drugs

Benedetto Bruno, Luisa Giaccone, Moreno Festuccia and Mario Boccadoro.

Division of Hematology at the S. Giovanni Battista Hospital, University of Torino, Via Genova 3, Torino, Italy.

Correspondence to: Benedetto Bruno, M.D., Ph.D., Divisione Universitaria di Ematologia, Azienda Ospedaliera San Giovanni Battista, Via Genova 3, 10126, TORINO, Italy, Phone +39-011-6334354 E-mail: benedetto.bruno@unito.it

Competing interests: The author have declared that no competing interests exist.

Published: June 1, 2010

Received: May 10, 2010

Accepted: May 31, 2010

Medit J Hemat Infect Dis 2010, 2(2): e2010013, DOI 10.4084/MJHID.2010.013

This article is available from: <http://www.mjhid.org/article/view/5944>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

High-dose melphalan with autologous stem cell rescue has been regarded as the standard of care for patients with newly diagnosed myeloma up to the age of 65-70 years. The recent development of agents with potent anti-tumor activity such as thalidomide, lenalidomide and bortezomib has further improved overall survival and response rates. However, relapse is a continuous risk.

Allografting is a potentially curative treatment for a subset of multiple myeloma patients for its well documented *graft-vs-myeloma* effects. However, its role has been hotly debated. Even though molecular remissions have been reported up to 50% after high-dose myeloablative conditionings, their applications, given the high toxicity, have been for long limited to younger relapsed/refractory patients. These limitations have greatly been reduced through the introduction of non-myeloablative/reduced-intensity conditionings.

The introduction of new drugs, characterised by low risks of early mortality, indeed requires to define role and timing of an allograft to capture the subset of patients who may most benefit from *graft-vs-myeloma* effects.

Ultimately, new drugs should not be viewed as mutually exclusive with an allograft. They may be employed to achieve profound cytoreduction before and enhance *graft-versus-myeloma* effects as consolidation/maintenance therapy after an allograft. However, this combination should be explored only in well-designed clinical trials.

Introduction: Multiple myeloma is a fatal plasma cell disorder, though recent advances in the understanding of its pathogenesis has identified peculiar mechanisms that have become targets of

agents with potent anti-myeloma activity such as lenalidomide and bortezomib. High-dose chemotherapy and autologous transplantation with/without these newer agents has been regarded

The introduction of reduced-intensity and non-myeloablative conditionings greatly renewed the interest in allografting and the incidence of this procedure peaked in the early 2000'. These regimens allowed to increase the eligible age for an allograft up to 70 years even in medically unfit

This manuscript aims at reviewing the current evidence of *graft-vs-myeloma* effects; the results obtained with conventional myeloablative and, more recently, with non-myeloablative conditionings; and the possible integration of so-called new drugs in the setting of allografting to improve clinical outcomes.

Myeloablative conditioning regimens (Table 1): The most commonly used myeloablative conditioning regimens included cyclophosphamide with total body irradiation or busulfan, or melphalan and total body irradiation.¹⁻¹³

1989 - 2008	➡	1086
Before 1989	➡	22




Table 1. Myeloablative Conditioning Regimens for Allografting in Multiple Myeloma

Author	Patients	Median Age (years)	Conditioning	Transplant-Related Mortality %	Complete Remission %	Overall Survival %
Bensinger et al.	136	43-48 (<60)	Bu, Cy, \pm Total Body Irradiation	48 (at day 100) 63 (at 1 year)	34	22 (at 5 years)
Barlogie et al.	36	≤ 55	Melphalan (100 mg/m ²), Total Body Irradiation (12Gy)	53 (at 1 year)	---	39 (at 7 years)
Reece et al.	26	43	Cy, Total Body Irradiation Bu,Cy Melphalan (100 mg/m ²), Total Body Irradiation	19 (at day 100)	62	47 (at 3 years)
Alyea et al.	24	46	Cy, Total Body Irradiation (14Gy) Bu,Cy	10	---	55 (at 2 years)
Kulkarni et al.	33	38	Melphalan (110 mg/m ²), Total Body Irradiation (10.5Gy) Cy, Total Body Irradiation (9.5Gy) Cy, Melphalan Bu,Cy	54	37	36 (at 3 years)
Le Blanc et al.	37	47	Cy, TBI (12Gy) Melphalan (140 mg/m ²), Total Body Irradiation (10.5Gy) Bu,Cy Others	22	57	32 (at 40 months)
Couban et al.	22	43	Melphalan (160 mg/m ²), Total Body Irradiation (12Gy) Cy, TBI (12Gy) Bu,Cy	59	50	32 (at 3 years)
Varterasian et al.	24	43	Cy, Total Body Irradiation Melphalan, Total Body Irradiation Bu,Cy, Total Marrow Irradiation Others	25	---	40 (at 3 years)

Abbreviations: Bu: Busulfan; Cy: cyclophosphamide)

The high transplant-related mortality up to 60% limited this approach to young, medically fit patients.¹⁻³ Causes of death comprised regimen-related, graft-vs-host disease (GvHD) and its transplant-related complications. Strong myeloma effects on baseline organ functions and severe immunodeficiency may be responsible for transplant-related mortality observed in other malignancies. Most representative experiences on the use of myeloablative conditioning regimens in multiple myeloma come from Seattle, the US Intergroup Trial S9321, and a European Bone Marrow Transplantation (EBMT) Registry study.^{2,4,14} The largest single-center experience comes from the Seattle group at the Fred Hutchinson Cancer Research Center.^{2,5} One-

hundred-thirty-six heavily pre-treated or disease refractory patients received an allograft between 1987 and 1999 from related (84%) or unrelated donors (16%). A day-100 transplant-related mortality of 48% was reported. The 5-year survival was 22% with disease-free survival of 14%. In 34% of patients who achieved complete remission, overall and disease-free survivals at 5 years were 48% and 37%. Subgroup analyses showed that early transplant-related mortality was approximately 20% for patients with chemo-sensitive disease who were transplanted within one year from diagnosis.

A North-American prospective trial compared autografting with myeloablative allografting.⁴ The US intergroup trial (S9321) of early vs late autografting included a third option that allowe

patients with HLA-identical siblings, under the age of 55, to undergo an allograft after melphalan and total body irradiation. This arm of the study was prematurely closed after the first 36 patients were enrolled given an excessively high transplant-related mortality of 53%. After a follow up of 7 years, however, the overall survivals were identical at 39% for both autologous and allogeneic recipients, while the progression-free survivals were 15% for autologous recipients as compared to 22% for allogeneic recipients, respectively. However, while the risk of relapse and death continues in the cohorts treated with an autograft, the overall survival curve for the allogeneic cohort reached a plateau with follow up extending to 10 years.

A large retrospective registry analysis by the EBMT group showed a remarkable improvement in overall survival in the late 90' due to a reduction in transplant-related mortality through improved supportive care and more careful patient selection.¹⁴ In this analysis, 690 patients, median age at transplant 44 years, who underwent a myeloablative allograft were divided into two cohorts: patients who received a bone marrow allograft between 1983-93 and those between 1994-98. In this latter cohort, some patients also received granulocyte-colony-stimulating factor (G-CSF) mobilized peripheral blood hematopoietic cells. Transplant-related mortality at 6 and 24 months was lower in the cohort transplanted between 1994-1998 than between 1983-1993, 21% versus 38% and 30% versus 46%. The reduced toxicity was associated with an increase in overall and progression-free survivals at 3 years from 35% to 55% and from 7 to 19 months for patients transplanted between 1994-1998. Furthermore, no differences in clinical outcomes were observed between patients who received marrow and those who received peripheral blood hematopoietic cells.

The interpretation of these studies to draw definitive conclusions is extremely difficult as the reported patients were not included in prospective control trials. Most patients were heavily pretreated, were chemo-resistant at the time of transplant and received a variety of conditionings and GvHD prophylaxes. The most consistent finding, however, was the high treatment-related mortality.

Despite selection bias, it was widely assumed that better clinical outcomes were associated with patients with chemo-sensitive myeloma at transplant. In most studies, only 10-25% of patients eventually became long-term disease-free survivors and were possibly cured.

Reduced-intensity and non-myeloablative conditioning regimens (Table 2): Though higher in multiple myeloma, the transplant-related morbidity and mortality associated with myeloablative conditioning regimens and allografting for the treatment of hematological malignancies have always been a matter of concern. These clinical observations prompted investigators, in the late 90', to explore highly immunosuppressive, though less myelosuppressive and less intense, conditionings which could possibly establish stable donor engraftment while reducing transplant-related organ toxicities. Pioneering studies were carried out in Seattle where it was shown that donor engraftment could be obtained with the sole combination of low dose non-myeloablative total body irradiation (200 cGy) and fludarabine, followed by peripheral blood stem cells and potent immunosuppression with cyclosporine and mycophenolate mofetil.¹⁵ Shortly thereafter, the tandem approach of an autologous transplant followed, 2-4 months later, by a non-myeloablative allograft was also designed for patients with newly diagnosed multiple myeloma.¹⁶ In 52 patients treated with this tandem modality the complete remission rate was 48% while progression free survival and overall survival were 48% and 69% respectively. The same "tandem concept" was also developed by Kroger et al using melphalan, fludarabine and anti-thymocyte globulin with related and unrelated donors.¹⁷

The tandem approach of an autologous transplant followed by a low dose non-myeloablative total body irradiation has become the most widely used conditioning for myeloma patients. The rationale for this tandem "autologous-allogeneic" approach was to separate in time the high-dose cytoreduction with melphalan at standard 200 mg/m² and the graft-vs-myeloma effect with the potential of drastically reducing treatment-related toxicity and mortality.

Two large series from Seattle and Italy have recently reported on more than 200 patients using the tandem auto/allo strategy. Long-term clinical outcomes of 102 patients treated with this approach, after a follow up of 6.3 years, were recently reported by Rotta et al.¹⁸ However, unlike the first report by the same group, patients were not uniformly in first line treatment. Overall, 42% of patients developed grade II-IV acute GvHD and 74% experienced chronic GvHD. Transplant-related mortality at 5 years was 18%, mostly due to GvHD and/or infections. Overall response rate was 94%, with 65% and 29% of patients achieving complete

Table 2. Non-myeloablative/Reduced Intensity Conditioning Regimens for Allografting in Multiple Myeloma

Author	Patients	Conditioning	Transplant-Related Mortality %	Chronic GVHD %	Complete Remission %	Overall Survival %
Mohty et al.	41	Bu, Fluda, ATG	17	41	24	62 (at 2 years)
Peggs et al.	20	Total Body Irradiation, Fluda, alemtuzumab	15	---	10	71 (at 2 years)
Einsele et al.	22	Total Body Irradiation (2Gy), Fluda, Cy	23	32	27	26 (at 2 years)
Giralt et al.	22	Fluda , Melphalan (90/140 mg/m ²)	41	27	32	30 (at 2 years)
Gerull et al.	52	Total Body Irradiation (2Gy), Fluda	17	70	27	41 (at 1.5 years)
Maloney et al.	54	Total Body Irradiation (2Gy)/ Total Body Irradiation (2Gy), Fluda	22	60	57	69 (at 5 years)
Lee et al.	45	Melphalan (100 mg/m ²), Total Body Irradiation (2Gy), Fluda	38	13	64	36 (at 3 years)
Kroger et al.	17	Melphalan (100 mg/m ²), Fluda, ATG	18	7	73	74 (at 2 years)
Kroger et al.	21	Melphalan (100-140 mg/m ²), Fluda, ATG	24	12	40	74 (at 2 years)

Abbreviations: Bu: Busulfan; Fluda: fludarabine; ATG anti-thymocyte globulin

and partial remissions respectively. Median overall survival was not reached and progression-free survival was 3 years. Estimated 5 year overall and progression free survivals were 64% and 36%. Results were recently reported also by the Gruppo Italiano Trapianti di Midollo.¹⁹ One-hundred newly diagnosed patients younger than 65 years were registered in a prospective multi-center study. Major strength of the study was the rigid enrolment of untreated myeloma patients who underwent the same vincristin, adriamycin and dexamethasone (VAD)-based induction before the autologous cytoreductive transplant. Primary objectives were overall and event-free survivals from diagnosis. After a median follow up of 5 years, overall survival was not reached and event free survival was 37 months. Incidences of acute and chronic GvHD were 38% and 50%, respectively. Complete remission, achieved in 53% of patients, or very good partial remission prior to allografting were significantly associated with achievement of post-transplant remission and longer event-free survival. Interestingly, in both studies from Seattle and from

the Italy graft-vs-myeloma effects were not associated with clinical GvHD.

In recent years, several reduce-intensity regimens have been designed including melphalan, 100-140 mg/m², with or without fludarabine, and intermediate-dose busulfan.²⁰⁻²⁷ Moreover, anti-thymocyte globulin or alemtuzumab have been employed in some trials to reduce GvHD.^{20,21} In a review of the EBMT registry, 26 different conditioning regimens, with/without T cell depletion, in 229 patients were reported.^{28,29} Almost 80% of patients received peripheral blood stem cells. Acute grade II-IV GVHD developed in 31% extensive chronic GVHD in 25%. Transplant-related mortality was rather low at 22%, however, 3 year overall survival and progression free survival were disappointing at 41% and 21%. Best clinical outcome was observed in those patients who were transplanted in first remission and did not receive more than one autograft. The use of alemtuzumab to prevent GVHD had a negative impact on transplant-related mortality, progression free survival and overall survival. Achievement of complete remission and occurrence of chronic GvHD were

associated with prolonged progression free survival. It is imperative to underline that fact the patients cohorts were highly heterogeneous and study designs greatly differ. No definite conclusions could be drawn.

More recently, studies comparing allografting after reduced-intensity conditionings and autografting have been published. The concept of Mendelian or genetic randomization has been applied to the assessment of outcomes in patients with hematological disorders who were treated with allografting or other therapies.³⁰⁻³³ This concept relies on the biological process through which offspring randomly inherit genetic traits half from each parent so that one in four siblings is expected to have a potential HLA-identical sibling donor. The comparison by the intention-to-treat principle between patients with HLA-identical siblings, who can be assigned to allografting, and those without such siblings, and who cannot receive an allograft, is used as a surrogate for an unbiased randomization.

The first such study was reported by the French group. The study compared two trials which included high risk myeloma patients carrying elevated serum β_2 -microglobulin and del(13).³⁴ All patients underwent an autograft after melphalan at 200 mg/m². Sixty-five patients with HLA-identical sibling donors then received an allograft after a conditioning with busulfan, fludarabine and high-dose anti-thymocyte globulin, 12.5 mg/kg. Outcomes were compared with 219 high risk patients who were treated with a second autograft after melphalan at 220 mg/m². Transplant-related mortality and response rates were not different. After a median follow-up of 2 years, overall and event free survivals were 35% and 25%, and 41% and 30% for the double autologous and the autologous-allogeneic cohorts, respectively. The Authors concluded that patients with high risk features may not benefit from a reduced-intensity allograft. This study was criticized for the inclusion of high dose anti thymocyte globulin, 12.5 mg/kg, in the conditioning regimen. As a matter of fact, though the incidence of chronic GvHD was 7%, the high dose of anti-thymocyte globulin may have highly prevented potentially curative graft-vs-myeloma effects. This study was also updated.³⁵ By intent-to-treat analysis on all 284 patients, after a median follow-up of 56 months, event-free survival did not significantly differ between tandem autologous and a single autograft followed by a reduced-intensity allograft (median 22 versus 19 months, *p* 0.58). There was a trend for a superior

overall survival in the tandem autologous cohort (median 48 versus 34 months, *p* 0.07).

Another study by Bruno et al. reported on 245 consecutive newly diagnosed myeloma patients, up to the age of 65 years, diagnosed between 1998-2004 where 162 out of 199 with at least one sibling were HLA-typed with their potential sibling donors.³⁶ The novelty of the study was the treatment assignment in function of the presence/absence of an HLA-identical sibling donor. Patients received induction with VAD-based regimens followed by a standard autograft with melphalan. Eighty patients with at least one HLA-identical sibling were offered total body irradiation -based non-myeloablative conditioning followed by an allograft with G-CSF mobilized peripheral blood stem cells. Eighty-two patients without an HLA-identical sibling were assigned to receive a second autograft after high-dose, 140-200 mg/m², or intermediate-dose, 100 mg/m², of melphalan. After a median follow up of 45 months, overall and event-free survivals were significantly longer in patients with donors: 80 versus 54 months and 35 versus 29 months. By multivariate analysis, having an HLA-identical sibling was an independent variable significantly associated with longer overall and event-free survivals. Overall, 58 and 46 patients completed the tandem autologous-allogeneic and the tandem autologous programs, with complete remission rates of 55% versus 26%. Transplant-related mortality was 10% and 2% respectively. Median overall survival was not reached in the tandem autologous-allogeneic cohort and was 58 months in the tandem autologous cohort. Event-free survival was 43 and 33 months, respectively. Criticisms to the study were that only 58 and 46 patients in the tandem autologous-allogeneic cohort and in the tandem autologous cohort, respectively, completed their assigned treatments and the relatively poor outcome of the patients assigned to the tandem autograft. This study was also updated after a median follow up of 6 years. Overall survival was not reached for the 80 patients with an HLA-identical sibling and was 52 months for those without, *p*=0.004; event free survival remained significantly longer in patients with HLA-identical siblings: 35 versus 29 months, *p*=0.009. Median overall survival was not reached in the 58 patients who completed the tandem autologous-allogeneic program and was 64 months in the 46 who completed the double autologous program, *p*=0.04. Event-free survival was 37 and 33 months *p*=0.06.

A third biologically randomized study was reported by the Spanish PETHEMA group.³⁷ One-

hundred-ten patients, after failing to reach at least near-complete remission after a first autograft, received either a second autograft (No. 85) or an allograft (No.25) after a reduced-intensity conditioning with melphalan and fludarabine. There was a higher complete remission rate, 40% versus 11%, $p=0.001$, and a trend towards a longer progression-free survival, median 31 months versus not reached, $p=0.08$, in the reduced-intensity group. Patients who underwent an allograft showed a trend towards a higher transplant-related mortality, 16% versus 5%, $p=0.07$, and no difference in overall and event-free survivals.

Finally, 4 large prospective randomized studies, the Blood and Marrow Transplant Clinical Trials Network (BMT-CTN) 0102 trial in the U.S.A.; the Dutch-Belgian Hemato-Oncology Cooperative Group (HOVON) trial,³⁸ the EBMT trial³⁹ and the study by the German DSMM group⁴⁰ in Europe, have recently been presented.

The large BMT-CTN 0102 trial comparing double autologous transplant versus tandem autologous/non-myeloablative allogeneic transplant completed the accrual in March 2007. More than 150 patients were biologically randomized to the latter cohort. The results from this study are eagerly awaited and should be released in 2010.

In the HOVON 54 study, newly diagnosed patients with an HLA-identical sibling donor included in the HOVON 50 study, a phase 3 study for the evaluation of thalidomide combined with high-dose melphalan, were allowed to proceed to a non-myeloablative allogeneic transplant from 2 and 6 months after a standard autograft, whereas patients without a suitable donor were randomized to thalidomide or interferon maintenance. By intent-to-treat analysis, no difference in progression free survival and overall survival were observed with an interim analysis that included 126 patients with a donor and 141 patients without.³⁸

In the EBMT trial, progression free survival at 60 months was 35% for the tandem auto/allo cohort as compared to 18% for double auto, and overall survival 65% and 57% respectively.³⁹ This trend was observed in both deletion 13 and non-deletion 13 patients. Final analyses of the Hovon and of the EBMT trials are expected in 2010.

Another prospective study comparing double autologous transplant versus tandem autologous/reduced-intensity allogeneic transplant, after a conditioning with fludarabine and melphalan, has been reported by the German DSMM.⁴⁰ This study only included patients with deletion 13q14. Transplants from HLA-matched

unrelated donors were allowed. Preliminary data showed a higher complete remission rate in patients with deletion 13q14 who received an allograft as compared to the autologous group (59% versus 32 % $p=0.003$). However, the projected overall survival at 3 years was 70% for the double autologous group and 60% for the allogeneic group ($P=0.22$). In the latter, transplant-related mortality at 2 years was only 12.7% even though 60% of patients received an allograft from an unrelated donor.

The potentially curative role of allografting:

graft-vs-myeloma: The potentially unique, curative role of allografting consist of the immune reaction of donor T cells against myeloma cells through the recognition of possibly disease-specific antigens. Evidence for the existence of such reactions was initially documented by the achievement of complete remissions after the discontinuation of immunosuppression or after the infusion of donor T lymphocytes in patients with recurrent disease post-transplant.⁴¹⁻⁴³ Some Authors, however, reported that the strongest predictors for response to donor lymphocyte infusions were acute and chronic GVHD⁴⁴⁻⁴⁷ indicating that GVHD and graft-vs.-myeloma may share the same antigenic targets. Chronic GVHD has been associated with longer response duration and prolonged overall survival. Recently, the Gruppo Italiano Trapianti di Midollo (GITMO), however, reported that the development of chronic GVHD did not correlate with the remission rates and response duration.¹⁹ Thus, subclinical graft-vs.-host reactions, especially after a non-mieloablative conditioning, may occur in the absence of detrimental GVHD. Finally, further evidence for graft-vs-myeloma are the molecular remissions, prelude to possible complete eradication, that have been reported up to 50% of patients following allografting.⁴⁸

Role of “new drugs”: So called “new drugs” have greatly changed the treatment options for multiple myeloma. Not only do they target malignant plasma cells but also affect their cross-talk with the marrow microenvironment due to several immunomodulatory properties. Interestingly, they modulate T cell subpopulations that may play a pivotal role in graft-vs-myeloma effects. Thus, their role in combination with allografting should be extensively investigated.

Thalidomide, lenalidomide and bortezomib have recently been included in a number of randomized clinical trials in both young and elderly

patients.⁴⁹⁻⁵² Response rates have significantly been improved even though longer follow-up is needed to evaluate the impact on long-term overall survival.

In the setting of allografting, these new drugs have first been employed in patients relapsing after allografting. The addition of thalidomide to donor lymphocyte infusions improved efficacy of salvage treatment without increasing GvHD.⁵³

Remarkable results have been obtained with lenalidomide in a cohort of patients with progressive disease after reduced-intensity allografting.⁵⁴⁻⁵⁵ In a study, 14/15 (93%) patients responded; however, a severe flare of GVHD in some patients was observed. Lenalidomide has also been employed as maintenance treatment to enhance graft-vs-myeloma in a prospective phase II study by the HOVON group.⁵⁶ After, an autologous/non-myeloablative tandem transplant, patients were given lenalidomide at the dose of 10 mg/day for 21 days and then 7 days of rest. Treatment was started between 1 and 6 months post-transplant in patients with no GvHD. Preliminary results showed a drop out rate of 41% primarily due to acute flare of severe GVHD that strongly correlated with the start of maintenance. Given this recently reported toxicity profile, the GITMO group is conducting a study where lenalidomide, employed as maintenance, is started at 6 months post non-myeloablative transplant in patients without signs and/or symptoms of chronic GVHD.

Bortezomib has also been shown to be effective in patients with relapsed disease.⁵⁷⁻⁶⁰ Interestingly, bortezomib may play a role in the immunomodulation of GVHD: in a preclinical murine model, it down-regulated cytokine synthesis, induced T cell apoptosis, prevented GvHD. Importantly, graft-vs-tumor effects were not affected.^{61,62} More recently, Blanco et al showed that bortezomib induced selective depletion of allo-reactive T lymphocytes, decreased the production of Th1 cytokines and allowed the emergence of a suppressor T cell subset.^{63,64} Of note, another study has shown that the combination of bortezomib with tacrolimus and methotrexate was very effective in the prevention of GvHD after reduced-intensity allografts from HLA-mismatched unrelated donors.⁶⁵ These findings appear attractive for studies in myeloma patients.

Conclusions: Overall, myeloablative allografts have cured a minority of patients who obtained complete clinical remission after transplant.

Reduced-intensity and non-myeloablative conditionings represent a clinical and biological breakthrough given that toxicity was greatly reduced and the existence of graft-vs-myeloma effects were indubitably shown. Long-term disease control and disabling chronic GVHD in a subset of patients represent important issues.

If an allograft should be part of first-line treatment plans or of salvage therapy for refractory/relapsed patients is still hotly debated. In newly diagnosed patients with chemosensitive disease, therefore in complete or very good remissions, a non-myeloablative conditioning would safely allow for donor engraftment with a reduced risk of toxicity and would potentially add a curative graft-vs.myeloma effect in a subset of patients. To support this, many reports show that better outcome is associated with chemosensitive disease at transplant and that allografting at an earlier disease phase is associated with stronger graft-vs-myeloma effects.^{66,67} This almost unanimously reported observation may be related to an antigen expression profile of potential targets for donor T cells that change through the disease phases. Siegel et al. reported the identification of HLA-A*0201-presented T cell epitopes, derived from the oncofetal antigen-immature laminin receptor protein, in many haematological malignancies.⁶⁸ However, it was interestingly observed that the expression of these antigens on plasma cells was lost over time. Even though very different in design, long-term results of donor-vs-no donor comparisons of the Blood and Marrow Transplant Clinical Trials Network (BMT CTN), the Dutch Hovon, the EBMT, and the German DSMM studies may allow valuable information on the use of up-front allografting.

Other Authors underline the fact that new treatment schemas may likely translate into longer overall survival and would be more inclined to offer an allograft at relapse. In this case, however, disease reduction prior to transplant and a more intense conditioning, rather than a non-myeloablative regimen, would be required despite a higher risk of toxicity.

In conclusion, future studies cannot be designed without the combination of new drugs that may enhance graft-versus-myeloma effects to allow long-term disease control and prolong survival even in patients with high risk disease. Profound cytoreduction before and enhanced graft-versus-myeloma effects after allografts through the immunomodulatory properties of lenalidomide and

bortezomib may be key factors to improve clinical outcomes.

Optimal timing of an allograft and dosage of new drugs remain to be determined and should be

explored prospectively only in the context of clinical trials and not routinely recommended.

References:

- Gahrton G, Tura S, Ljungman P, et al. Allogeneic bone marrow transplantation in multiple myeloma. *N Engl J Med* 1991;325:1267-1273.
- Bensinger WI, Buckner CD, Anasetti C, et al. Allogeneic marrow transplantation for multiple myeloma: An analysis of risk factors on outcome. *Blood* 1996;88:2787-2793.
- Gahrton G, Tura S, Ljungman P, et al. Prognostic factors in allogeneic bone marrow transplantation for multiple myeloma. *J Clin Oncol* 1995;13:1312-1322.
- Barlogie B, Kyle RA, Anderson KC, et al. Standard chemotherapy compared with high-dose chemoradiotherapy for multiple myeloma: final results of phase III US Intergroup Trial S9321. *J Clin Oncol* 2006;24(6): 929-936.
- Bensinger WI, Maloney D, Storb R. Allogeneic hematopoietic cell transplantation for multiple myeloma. *Semin Hematol* 2001;38: 243-249.
- Reece DE, Shepherd JD, Klingemann HG, et al. Treatment of myeloma using intensive therapy and allogeneic bone marrow transplantation. *Bone Marrow Transplant* 1995;15: 117-123.
- Alyea E, Weller E, Schlossman R, et al. T-cell-depleted allogeneic bone marrow transplantation followed by donor lymphocyte infusion in patients with multiple myeloma: induction of graft-versus-myeloma effect. *Blood* 2001;98: 934-939.
- Kulkarni S, Powles RL, Treleaven JG, et al. Impact of previous high-dose therapy on outcome after allografting for multiple myeloma. *Bone Marrow Transplant* 1999;23: 675-680.
- Le Blanc R, Montminy-Métivier S, Bélanger R, et al. Allogeneic transplantation for multiple myeloma: further evidence for a GVHD-associated graft-versus-myeloma effect. *Bone Marrow Transplant* 2001;28: 841-848.
- Couban S, Stewart AK, Loach D, et al. Autologous and allogeneic transplantation for multiple myeloma at a single centre. *Bone Marrow Transplant* 1997;19:783-789.
- Varterasian M, Janakiraman N, Karanes C, et al. Transplantation in patients with multiple myeloma: a multicenter comparative analysis of peripheral blood stem cell and allogeneic transplant. *Am J Clin Oncol* 1997;20: 462-466.
- Russell NH, Mifflin G, Stainer C, et al. Allogeneic bone marrow transplant for multiple myeloma. *Blood* 1997;89: 2610-2611. (Letter)
- Cavo M, Bandini G, Benni M, et al. High-dose busulfan and cyclophosphamide are an effective conditioning regimen for allogeneic bone marrow transplantation in chemosensitive multiple myeloma. *Bone Marrow Transplant* 1998;22:27-32.
- Gahrton G, Svensson H, Cavo M, et al. Progress in allogeneic bone marrow and peripheral blood stem cell transplantation for multiple myeloma: a comparison between transplants performed 1983-93 and 1994-8 at European Group for Blood and Marrow Transplantation centres. *Br J Haematol* 2001;113:209-216.
- McSweeney PA, Niederwieser D, Shizuru JA, et al. Hematopoietic cell transplantation in older patients with hematologic malignancies: replacing high-dose cytotoxic therapy with graft-versus-tumor effects. *Blood* 2001;97:3390-3400.
- Maloney DG, Molina AJ, Sahebi F, et al. Allografting with nonmyeloablative conditioning following cytoreductive autografts for the treatment of patients with multiple myeloma. *Blood* 2003;102:3447-3454.
- Kröger N, Schwerdtfeger R, Kiehl M, et al. Autologous stem cell transplantation followed by a dose-reduced allograft induces high complete remission rate in multiple myeloma. *Blood* 2002;100:755-760.
- Rotta M, Storer BE, Sahebi F, et al. Long-term outcome of patients with multiple myeloma after autologous hematopoietic cell transplantation and nonmyeloablative allografting. *Blood* 2009;113:3383-91.
- Bruno B, Rotta M, Patriarca F, et al. Non-myeloablative allografting for newly diagnosed multiple myeloma: the experience of the Gruppo Italiano Trapianti di Midollo. *Blood* 2008;113:3375-82.
- Mohty M, Boiron JM, Damaj G, et al. Graft-versus-myeloma effect following antithymocyte globulin-based reduced intensity conditioning allogeneic stem cell transplantation. *Bone Marrow Transplant* 2004;34:77-84.
- Peggs KS, Mackinnon S, Williams CD, et al. Reduced-intensity transplantation with in vivo T-cell depletion and adjuvant dose-escalating donor lymphocyte infusions for chemotherapy-sensitive myeloma: Limited efficacy of graft-versus-tumor activity. *Biol Blood Marrow Transplant* 2003;9: 257-265.
- Einsele H, Schäfer HJ, Hebart H, et al. Follow-up of patients with progressive multiple myeloma undergoing allografts after reduced-intensity conditioning. *Br J Haematol* 2003;121: 411-418.
- Giralt S, Aleman A, Anagnostopoulos A, et al. Fludarabine/melphalan conditioning for allogeneic transplantation in patients with multiple myeloma. *Bone Marrow Transplant* 2002;30:367-373.
- Gerull S, Goerner M, Benner A, et al. Long-term outcome of nonmyeloablative allogeneic transplantation in patients with high-risk multiple myeloma. *Bone Marrow Transplant* 2005;36:963-969.
- Bruno B, Sorasio R, Patriarca F, et al. Unrelated donor hematopoietic cell transplantation after non-myeloablative conditioning for patients with high-risk multiple myeloma. *Eur J Haematol* 2007;78:330-337.
- Lee CK, Badros A, Barlogie B, et al. Prognostic factors in allogeneic transplantation for patients with high-risk multiple myeloma after reduced intensity conditioning. *Exp Hematol* 2003;31:73-80.
- Kröger N, Sayer HG, Schwerdtfeger R, et al. Unrelated stem cell transplantation in multiple myeloma after a reduced-intensity conditioning with pretransplantation antithymocyte globulin is highly effective with low transplantation-related mortality. *Blood* 2002;100:3919-3924.
- Crawley C, Iacobelli S, Björkstrand B, et al. Reduced-intensity conditioning for myeloma: lower nonrelapse mortality but higher relapse rates compared with myeloablative conditioning. *Blood* 2007;109:3588-3594.
- Crawley C, Lallancette M, Szydlo R, et al. Outcomes for reduced-intensity allogeneic transplantation for multiple myeloma: an analysis of prognostic factors from the Chronic Leukaemia Working Party of the EBMT. *Blood* 2005; 105:4532-39.
- Wheatley K, Gray R. Commentary: Mendelian randomization--an update on its use to evaluate allogeneic stem cell transplantation in leukaemia. *Int J Epidemiol* 2004;33:15-7.
- Balduzzi A, Valsecchi MG, Uderzo C, et al. Chemotherapy versus allogeneic transplantation for very-high-risk childhood acute lymphoblastic leukaemia in first complete

- remission: comparison by genetic randomisation in an international prospective study. *Lancet* 2005;366:635-42.
32. Woods WG, Neudorf S, Gold S, et al. A comparison of allogeneic bone marrow transplantation, autologous bone marrow transplantation, and aggressive chemotherapy in children with acute myeloid leukemia in remission. *Blood* 2001;97:56-62.
33. Suci S, Mandelli F, de Witte T, et al. Allogeneic compared with autologous stem cell transplantation in the treatment of patients younger than 46 years with acute myeloid leukemia (AML) in first complete remission (CR1): an intention-to-treat analysis of the EORTC/GIMEMAAML-10 trial. *Blood* 2003;102:1232-40.
34. Garban F, Attal M, Michallet M, et al. Prospective comparison of autologous stem cell transplantation followed by dose-reduced allograft (IFM99-03 trial) with tandem autologous stem cell transplantation (IFM99-04 trial) in high-risk de novo multiple myeloma. *Blood* 2006;107:3474-3480.
35. Moreau P, Garban F, Attal M, et al. Long-term follow-up results of IFM99-03 and IFM99-04 trials comparing nonmyeloablative allotransplantation with autologous transplantation in high-risk de novo multiple myeloma. *Blood* 2008;112:3914-5.
36. Bruno B, Rotta M, Patriarca F, et al. A comparison of allografting with autografting for newly-diagnosed myeloma. *N Engl J Med* 2007;356:1110-1120.
37. Rosiñol L, Pérez-Simón JA, Sureda A, et al. A prospective PETHEMA study of tandem autologous transplantation versus autograft followed by reduced-intensity conditioning allogeneic transplantation in newly diagnosed multiple myeloma. *Blood* 2008;112:3591-3.
38. Lokhorst H, Sonneveld P, van der Holt B, et al. Donor Versus No Donor Analysis of Newly Diagnosed Myeloma Patients Included in the HOVON 50/54 Study. *Blood (ASH Annual Meeting Abstracts)* 2008; 112: 461.
39. Gahrton G, Björkstrand B, Iacobelli S, et al. Tandem Autologous(ASCT)/Allogeneic Reduced Intensity Conditioning Transplantation (RIC) with Identical Sibling Donor Versus ASCT in Previously Untreated Multiple Myeloma (MM): Long Term Follow up of a Prospective Controlled Trial by the EBMT. *Blood (ASH Annual Meeting Abstracts)* 2009; 114:52.
40. Knop S, Liebisch P, Hebart H, et al. Allogeneic Stem Cell Transplant versus Tandem High-Dose Melphalan for Front-Line Treatment of Deletion 13q14 Myeloma – An Interim Analysis of the German DSMM V Trial. *Blood (ASH Annual Meeting Abstracts)* 2009; 114:51.
41. Verdonck LF, Lokhorst HM, Dekker AW, et al. Graft-versus-myeloma effect in two cases. *Lancet* 1996;347:800-801.
42. Aschan J, Lönnqvist B, Ringdén O, et al. Graft-versus-myeloma effect. *Lancet* 1996;348:346. (Letter)
43. Libura J, Hoffmann T, Passweg J, et al. Graft-versus-myeloma after withdrawal of immunosuppression following allogeneic peripheral stem cell transplantation. *Bone Marrow Transplant* 1999;24:925-927.
44. Lokhorst HM, Schattenberg A, Cornelissen JJ, et al. Donor leukocyte infusions are effective in relapsed multiple myeloma after allogeneic bone marrow transplantation. *Blood* 1997;90: 4206-4211.
45. Lokhorst HM, Schattenberg A, Cornelissen JJ, et al. Donor lymphocyte infusions for relapsed multiple myeloma after allogeneic stem-cell transplantation: predictive factors for response and long-term outcome. *J Clin Oncol* 2000;18: 3031-3037.
46. Lokhorst HM, Wu K, Verdonck LF, et al. The occurrence of graft-versus-host disease is the major predictive factor for response to donor lymphocyte infusions in multiple myeloma. *Blood* 2004;103: 4362-4364.
47. Van de Donk NW, Kröger N, Hegenbart U, et al. Prognostic factors for donor lymphocyte infusions following non-myeloablative allogeneic stem cell transplantation in multiple myeloma. *Bone Marrow Transplant* 2006;37:1135-1141.
48. Corradini P, Voena C, Tarella C, et al. Molecular and clinical remissions in multiple myeloma: role of autologous and allogeneic transplantation of hematopoietic cells. *J Clin Oncol* 1999;17:208-215.
49. Richardson PG, Sonneveld P, Schuster MW, et al. Bortezomib or high-dose dexamethasone for relapsed multiple myeloma. *N Engl J Med*. 2005;352:2487-2498.
50. Dimopoulos M, Spencer A, Attal M, et al. Lenalidomide plus dexamethasone for relapsed or refractory multiple myeloma. *N Engl J Med*. 2007;357:2123-2132.
51. Weber DM, Chen C, Niesvizky R, et al. Lenalidomide plus dexamethasone for relapsed multiple myeloma in North America. *N Engl J Med*. 2007;357:2133-2142.
52. Palumbo A, Bringhen S, Caravita T, et al.: Italian Multiple Myeloma Network, GIMEMA. Oral melphalan and prednisone chemotherapy plus thalidomide compared with melphalan and prednisone alone in elderly patients with multiple myeloma: randomised controlled trial. *Lancet*. 2006;367:825-31.
53. Kröger N, Shimoni A, Zagrivnaja M, et al: Low-dose thalidomide and donor lymphocyte infusion as adoptive immunotherapy after allogeneic stem cell transplantation in patients with multiple myeloma. *Blood* 2004;104:3361-3363.
54. Minnema MC, van der Veer MS, Aarts T, et al: Lenalidomide alone or in combination with dexamethasone is highly effective in patients with relapsed multiple myeloma following allogeneic stem cell transplantation and increases the frequency of CD4+Foxp3+ T cells. *Leukemia* 2009; 23:605-607.
55. Lioznov M, El-Cheikh J Jr, Hoffmann F, et al: Lenalidomide as salvage therapy after allo-SCT for multiple myeloma is effective and leads to an increase of activated NK (NKp44(+)) and T (HLA-DR(+)) cells. *Bone Marrow Transplant* 2010; 45:349-353
56. Minnema MC, van der Holt B, Kersten MJ, et al: First Interim Analysis of HOVON 76: Lenalidomide Maintenance Following Non Myeloablative Allogeneic Stem Cell Transplantation in Patients with Multiple Myeloma. *Blood (ASH Annual Meeting Abstracts)* 2009; 114: 2285.
57. Kröger N, Zabelina T, Ayuk F, et al: Bortezomib after dose-reduced allogeneic stem cell transplantation for multiple myeloma to enhance or maintain remission status. *Exp Hematol*. 2006; 34:770-775.
58. van de Donk NW, Kroger N, Hegenbart U, et al: Remarkable activity of novel agents bortezomib and thalidomide in patients not responding to donor lymphocyte infusions following nonmyeloablative allogeneic stem cell transplantation in multiple myeloma. *Blood* 2006; 107: 3415-3416.
59. Kröger N, Badbaran A, Lioznov M, et al: Post-transplant immunotherapy with donor-lymphocyte infusion and novel agents to upgrade partial into complete and molecular remission in allografted patients with multiple myeloma. *Exp Hematol*. 2009; 37:791-798.
60. Bruno B, Patriarca F, Sorasio R, et al. Bortezomib with or without dexamethasone in relapsed multiple myeloma following allogeneic hematopoietic cell transplantation. *Haematologica*. 2006;91:837-9.
61. Sun K, Welniak LA, Panoskaltis-Mortari A., et al: Inhibition of acute graft-versus-host disease with retention of graft-versus-tumor effects by the proteasome inhibitor bortezomib. *Proc.Natl.Acad.Sci.U.S.A* 2004; 101:8120-8125.
62. Sun K, Wilkins DE, Anver MR, Sayers TJ, et al: Differential effects of proteasome inhibition by bortezomib on murine acute graft-versus-host disease (GVHD): delayed administration of bortezomib results in increased GVHD-dependent gastrointestinal toxicity. *Blood* 2005; 106:3293-3299.
63. Blanco B, Pérez-Simón JA, Sánchez-Abarca LI, et al: Bortezomib induces selective depletion of alloreactive T lymphocytes and decreases the production of Th1 cytokines. *Blood* 2006; 107: 3575-3583.

64. Blanco B, Pérez-Simón JA, Sánchez-Abarca LI, et al: Treatment with bortezomib of human CD4⁺ T cells preserves natural regulatory T cells and allows the emergence of a distinct suppressor T-cell population. *Haematologica* 2009; 94:975-983.
65. Koreth J, Stevenson KE, Kim HT, et al: Bortezomib, tacrolimus, and methotrexate for prophylaxis of graft-versus-host disease after reduced-intensity conditioning allogeneic stem cell transplantation from HLA mismatched unrelated donors. *Blood* 2009; 114: 3956–3959.
66. Kroger N, Perez-Simon JA, Myint H, et al: Relapse to prior autograft and chronic graft-versus-host disease are the strongest prognostic factors for outcome of melphalan/fludarabine-based dose-reduced allogeneic stem cell transplantation in patients with multiple myeloma. *Biol Blood Marrow Transplant* 2004; 10: 698-708.
67. Lee CK, Badros A, Barlogie B, et al: Prognostic factors in allogeneic transplantation for patients with high-risk multiple myeloma after reduced intensity conditioning. *Exp Hematol* 2003; 31: 73-80.
68. Siegel S, Wagner A, Friedrichs B, et al.: Identification of HLA-A*0201-presented T cell epitopes derived from the oncofetal antigen-immature laminin receptor protein in patients with hematological malignancies. *J Immunol* 2006; 76: 6935-6944.