

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Pleistocene faunas of Sicily: a review.**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/91517> since

*Publisher:*

Archaeopress

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# World Islands in Prehistory

International Insular Investigations

V Deia International Conference of Prehistory

Edited by

W. H. Waldren

and

J. A. Ensenyat

BAR International Series 1095

2002

## PLEISTOCENE FAUNAS OF SICILY: A REVIEW

BONFIGLIO L.(1), MARRA A. C.(1), MASINI F.(2), PAVIA M. (3) PETRUSO D.(2)

(1) *Dipartimento di Scienze della Terra, Università degli Studi di Messina, Via Sperone 31- casella postale 54 - 98166 Messina, Italy*

(2) *Dipartimento di Geologia e Geodesia, Università degli Studi di Palermo, Corso Tukory 131, I-90134 Palermo, Italy*

(3) *Dipartimento di Scienze della Terra, Università degli Studi di Torino, Via Accademia delle Scienze 5, 10123 Torino, Italy*

E-MAIL: laura@labcart.unime.it; fmasini@unipa.it

## INTRODUCTION

Sicily has a rich palaeontological heritage of Pleistocene continental vertebrates, that indicates different faunal complexes inhabited the island through time.

Most of the older studies on Quaternary vertebrates in Sicily have focused on evolutionary and taxonomic aspects (e.g. Pohlig 1893; Pohlig 1909; Vaufrey 1929; Accordi & Colacicchi 1962; Ambrosetti 1968; Aguirre 1968-69; Gliozzi & Malatesta 1984; Brugal 1987) with the aim to identify the continental species from which the Sicilian ones derived. Most of the known vertebrate remains came from cave deposits and little was known about the palaeoenvironmental conditions of the vertebrate-bearing deposits.

Chronological arrangements of the various Pleistocene mammal assemblages of Sicily were based on the assumption of the phyletic derivation of the dwarf elephant *Elephas falconeri* from the middle sized *Elephas mnaidriensis*, which was in turn considered a direct descendent of *Elephas antiquus* (Accordi 1963; Accordi 1965; Accordi & Colacicchi 1962; Ambrosetti et al. 1980). Vaufrey (1929) was inclined to assume a post-Tyrrhenian age for all the vertebrate faunas. Since the studies of Accordi (1957; 1963; 1965) it was thought that most of the size reduction of elephants in Sicily took place during the period preceding the Tyrrhenian. The smallest species (*Elephas falconeri*) was considered to be limited to the early Würm period and to have evolved as a consequence of environmental stress linked to the Würmian climatic cooling (Ambrosetti 1968; Kotsakis 1979). Stratigraphic and taphonomic features have

been generally disregarded although the first valuable scientific paper dealing with the excavations of the Hippopotamus pentlandi bearing deposits of the S. Ciro cave (Palermo), carried out by the abbot Scinà (Scinà 1831), contains many interesting remarks concerning the relationship between the vertebrate-bearing deposits and the underlying marine sands which extend inside and outside the cave.

Vaufrey (1929), Accordi & Colacicchi (1962) and Accordi (1963; 1965) provide also interesting observations concerning the relationship between vertebrate-bearing deposits and littoral marine deposits.

Since 1985, a new synthesis has incorporated new stratigraphic and aminostratigraphic data (Belluomini & Bada 1985; Bada et al. 1991; Burgio & Cani 1988; Bonfiglio 1987; Bonfiglio 1991; Bonfiglio 1992 a; Bonfiglio 1992 b; Bonfiglio & Burgio, 1992; Bonfiglio & Insacco 1992).

These data concern essentially the following aspects:

- (1) Number and composition of the Pleistocene faunal complexes
- (2) Stratigraphic, environmental and taphonomic data
- (3) Chronological data
- (4) Geographic distribution of vertebrate bearing deposits and palaeogeography.

## NUMBER AND COMPOSITION OF PLEISTOCENE FAUNAL COMPLEXES (Table. 1)

On the basis of extensive new evidence, the Pleistocene vertebrate assemblages of Sicily can be arranged into 5 phases or Faunal Complexes (F.C.), spanning from the Early Pleistocene to the Late Glacial (Bonfiglio et al. 2000;

Bonfiglio et al. 2001; Bonfiglio et al. in press).  
MONTE PELLEGRINO FAUNAL COMPLEX.

This is the oldest Quaternary fossil record so far known (late Villafranchian, Burgio & Fiore 1997) and it is documented only in the very restricted geographic area of Monte Pellegrino, close to the town of Palermo. The poorly diversified fauna includes species of the genera Hypolagus, Pannonictis, Leithia, Asoriculus, Pellegrinia, Apodemus.

Hypolagus and Pannonictis are moderately endemic taxa and the potential ancestors appear to be typical European forms, not found so far in northern Africa (Burgio & Fiore

teristics of the African stock of Ctenodactilids, thus indicating a dispersal from that region.

The composition of the Monte Pellegrino fauna - unique for the Mediterranean islands - suggests it may have been derived in part from an older, not locally known, population phase (Messinian age ? Azzaroli 1974; Azzaroli & Guazzone 1979) and partially from younger dispersals from Europe. The different degree of endemism and the different geographical affinity of the taxa, indicate a polyphasic origin (Masini et al. in press). After the first report by Thaler (1972) some of the taxa have been revised more recently

"Monte Pellegrino" F.C.	"Elephas falconeri" F.C.	"Elephas mnaidriensis" F.C.	"Grotta S. Teodoro - Pianetti" F.C.	"Castello" F.C.	Holocene fauna
<i>Testudo graeca</i> <i>Asoriculus</i> sp. <i>Pannonictis arzilla</i> <i>Apodemus maximus</i> <i>Leithia</i> sp. <i>Maltamys</i> cf. <i>gollcheri</i> <i>Pellegrinia</i> <i>panormensis</i> <i>Hypolagus</i> sp.	<i>Discoglossus</i> cf. <i>pictus</i> <i>Bufo</i> cf. <i>viridis</i> <i>Hyla</i> sp. <i>Emys orbicularis</i> <i>Testudo hermanni</i> <i>Geochelone</i> sp. <i>Lacerta</i> <i>siculomelitensis</i> <i>Lacerta viridis</i> <i>Lacerta</i> sp. <i>Coluber</i> cf. <i>viridiflavus</i> <i>Natrix</i> sp. Several species of bats <i>Crociodura esuae</i> <i>Vulpes</i> sp. <i>Nesolutra trinacriae</i> <i>Elephas falconeri</i> <i>Leithia cartei</i> <i>Leithia melitensis</i> <i>Maltamys gollcheri</i>	<i>Discoglossus</i> cf. <i>pictus</i> <i>Emys orbicularis</i> <i>Testudo hermanni</i> <i>Lacerta</i> <i>siculomelitensis</i> Birds <i>Erinaceus europaeus</i> <i>Crociodura</i> aff. <i>esuae</i> <i>Crocota crocuta</i> cf. <i>spelaea</i> <i>Panthera leo</i> <i>Canis lupus</i> <i>Nesolutra trinacriae</i> <i>Ursus</i> cf. <i>arctos</i> <i>Elephas mnaidriensis</i> <i>Sus scrofa</i> <i>Hippopotamus pentlandi</i> <i>Cervus elaphus siciliae</i> <i>Dama</i> <i>carburangelensis</i> <i>Bos primigenius siciliae</i> <i>Bison priscus siciliae</i> <i>Leithia</i> cf. <i>melitensis</i> <i>Maltamys</i> cf. <i>wiendincitensis</i>	Reptiles <i>Testudo</i> sp. Birds Chiroptera <i>Crocota crocuta</i> cf. <i>spelaea</i> <i>Ursus</i> cf. <i>arctos</i> <i>Canis</i> cf. <i>lupus</i> <i>Vulpes vulpes</i> <i>Elephas mnaidriensis</i> <i>Sus scrofa</i> <i>Equus hydruntinus</i> <i>Cervus elaphus siciliae</i> <i>Bos primigenius siciliae</i> <i>Apodemus</i> cf. <i>sylvaticus</i> <i>Microtus (Terricola)</i> ex gr. <i>savii</i> <i>Crociodura</i> cf. <i>sicula</i>	Birds <i>Erinaceus europaeus</i> <i>Crociodura</i> cf. <i>sicula</i> <i>Canis lupus</i> <i>Vulpes vulpes</i> <i>Equus caballus</i> <i>Equus hydruntinus</i> <i>Sus scrofa</i> <i>Cervus elaphus</i> <i>Bos primigenius</i> <i>Microtus (Terricola)</i> ex gr. <i>savii</i> <i>Apodemus</i> sp. <i>Lepus europaeus</i>	<i>Bufo bufo</i> <i>Bufo</i> cf. <i>viridis</i> Turtle indet. <i>Emys orbicularis</i> <i>Lacerta viridis</i> Birds Chiroptera indet. <i>Erinaceus europaeus</i> <i>Crociodura</i> sp. <i>Canis</i> cf. <i>lupus</i> <i>Vulpes vulpes</i> <i>Felis silvestris</i> <i>Martes</i> sp. <i>Mustela</i> cf. <i>nivalis</i> <i>Ursus</i> sp. <i>Sus scrofa</i> <i>Cervus</i> sp. <i>Bos primigenius</i> , <i>Microtus (Terricola)</i> cf. <i>savii</i> <i>Apodemus</i> sp. <i>Arvicola terrestris</i> <i>Mioxus glis</i> <i>Monachus monachus</i>

TABLE 1. FAUNAL COMPLEXES OF SICILY (AFTER BONFIGLIO et al., 2001)

1997); the endemic shrew *Asoriculus burgioi* is of uncertain provenance (Masini & Sarà 1998); *Asoriculus* is well documented in Europe from the Early Pliocene, and it has also been found recently in northern Africa, in Late Pliocene - Early Pleistocene deposits (Geraads 1995; Rzebik-Kowalska 1988).

The endemic dormouse *Leithia* (Thaler 1972), is regarded as one of the direct ancestor of the better known and better described *Leithia* and *Maltamys* from the Middle and Late Pleistocene of Sicily and Malta (Zammit Maempel & De Bruijn 1982). *Leithia* and *Maltamys* are considered by Daams & De Bruijn (1995) as relics of a scarcely known older (Messinian) population phase and these elements have remained isolated in the Sicilian Maltese archipelago for the entire Pliocene and most of the Quaternary.

According to Thaler (1972) *Pellegrinia* has derived charac-

but our knowledge of this very peculiar and intriguing faunal assemblage is still not exhaustive.

ELEPHAS FALCONERI FAUNAL COMPLEX. This Faunal Complex is even poorer than the preceding one in mammalian biodiversity. The poorly diversified fauna includes, besides the pigmy elephant, members of the genera *Crociodura*, *Lutra*, *Leithia*, *Maltamys*, a giant tortoise, and a rich typical endemic avifauna (Pavia 1999; Pavia 2000).

The shrew *Crociodura esuae* is an endemic species, of uncertain bio-geographic affinity (Kotsakis 1986). The genus *Crociodura* occurs widely in the Quaternary of Europe (Reumer 1984; Rzebik-Kowalska 1995) and in northern Africa (Butler 1980). Large mammals include the pigmy elephant *Elephas falconeri*, strongly reduced in size, and a member of the Lutrinae (genus *Lutra*).

Neither the ancestor nor the geographic provenance of *E. falconeri* has been determined unequivocally. Even though its derivation from a Paleoloxodontine elephant is accepted by several authors, its possible origin from European *Elephas antiquus* stock, or from a north African species is still a matter for discussion.

Moreover, also the composition of this faunal complex reveals a polyphasic origin; some taxa are relics from the preceding phase and others are 'newcomers' that probably entered the island through a strongly filtering barrier (Masini et al. in press).

**ELEPHAS MNAIDRIENSIS FAUNAL COMPLEX:** This Faunal Complex is almost completely renewed with respect to the preceding F. C. The pigmy *E. falconeri* is extinct, while the faunal composition is more balanced and includes top predators such as the lion and the spotted 'cave' hyena. The faunal composition is quite similar to that found in the southern Italian peninsula, and the continental forerunner of most of these taxa has been fairly accurately identified. The only significant absence is that of perissodactyls (rhinos and horses). The herbivorous taxa (bison, auroch, fallow deer, red deer, hippo) are moderately modified with respect to the congeneric or conspecific taxa from the Italian mainland and the endemic nature of the fauna is apparent mainly from the modest reduction in size. *E. mnaidriensis* is a smaller, but not extremely modified, descendant of *Elephas antiquus*. The red deer *Cervus elaphus siciliae* is slightly reduced in size (about 20%). *Dama carburangelensis* (previously identified as *Megaceroides carburangelensis*) is probably derived from fallow deer populations (*Dama dama tiberina*) widespread in central and southern Italy during the late Middle Pleistocene (Abbazzi et al. 2001).

A peculiarity of this fauna is the very low diversity of small mammals, which are represented only by survivors from the *E. falconeri* F. C. (*Leithia*, *Maltamys* and *Crocidura esuae*; Petruso 2001; Petruso in progress; Masini et al. in press). The knowledge of the composition of the avifauna is still incomplete (Pavia 2000; Pavia 2001). One endemic species of birds, *Cygnus falconeri* occurs, while the endemic species of the previous F. C. became extinct, except *Grus melitensis* whose persistence is, however, questionable. Typical continental taxa (i. e. Galliforms) are still lacking. On the whole the avifauna assemblages are partially renewed alike the mammal fauna.

#### THE PIANETTI - SAN TEODORO CAVE FAUNAL COMPLEX

The Pianetti - San Teodoro Cave Faunal Complex dates to the last glacial cycle. The faunal history of this period is characterised by extinction events (hippopotamus, endemic dormice and *Crocidura esuae*), and by the dispersal of equids (*Equus hydruntinus*) and of mainland small mammals, which are represented by taxa still occurring in Sicily (*Microtus* (*Terricola*), *Crocidura cf. sicula*, *Apodemus cf. sylvaticus*, *Erinaceus europaeus*) (Bonfiglio et al. 1997; Bonfiglio et al. 2001). Almost all of the large mammals belonging to this F. C. seem to be inherited from *Elephas mnaidriensis* F. C. endemic species of avifauna are lacking

(Pavia 2000).

The faunal composition of this F. C. is probably not completely known, since only a few sites have been discovered so far and one of them (S. Teodoro cave) has not yet been completely investigated.

Castello Faunal Complex. Finally, the Late Glacial associations show a dramatic decrease in diversity, missing all endemic large mammals still occurring in the Pianetti-S. Teodoro F. C. Late glacial faunas, which are similar to continental ones, are associated to lithic artifacts and cultural evidences of late Upper Palaeolithic and, following some authors (Kotsakis 1979), include taxa present in one (*Equus hydruntinus*) or both (*Sus scrofa*) previous faunal complexes.

#### STRATIGRAPHIC, ENVIRONMENTAL AND TAPHONOMIC DATA

Taphonomic data show that Pleistocene vertebrates were distributed in both cave environments and broad, open environment (Bonfiglio 1987; Bonfiglio 1992 b; Bonfiglio 1995; Bonfiglio et al. 1993; Bonfiglio et al. 1996; Bonfiglio et al. 1997; Chilardi & Gilotti 1996) and numerous relationships have been found between the vertebrate bearing deposits and terraced marine deposits, which can be correlated with the  $\delta^{18}O$  isotopic record and the main palaeogeographic events in Sicily (Agnesi et al. 1997; Bonfiglio 1991; Bonfiglio et al. 2000; Bonfiglio et al. in press; Di Maggio et al. 1999).

Actually most of the mammalian fossils of Sicily are found in caves and fissures - rather common in the carbonate mountain ranges of Northern Sicily and in the Hyblean plateau. Another important category of deposits is represented by ancient shorelines, beach deposits and marine terraces, frequently associated with a lagoon or swamp. Limnic deposits related to small freshwater basins also occur, often in relation to coastal and fully marine deposits.

The assemblages of the Monte Pellegrino F.C. and those of S. Teodoro Cave - Pianetti and Castello F.C. 'xes are contained in caves and fissure-filling deposits and they have not relationships with marine deposits. The assemblages of the *Elephas falconeri* and of the *Elephas mnaidriensis* F.C. 'xes are contained in cave deposits as well as in coastal plain and/or in marine littoral deposits.

The biodiversity, the preservation conditions and the concentration of the skeletal remains are very diverse in the different environments (see FIGURE 1).

Littoral marine sands - In the late Middle Pleistocene sandy gravel outside the S. Ciro cave, the rare hippopotamus and elephant remains (molars, limbs) are disarticulated, encrusted by serpulid polychaetes worms and associated with a rich marine fauna containing gastropods, bivalves, ostracods, foraminifera, of littoral a euryhaline environment (Galletti & Scaletta 1991).

An isolated scapula of *Elephas mnaidriensis* comes from Thyrrenian marine deposits at Maddalena peninsula (Siracusa) (Accordi 1963; Di Grande & Scamarda 1973).

Deltaic marine clay sands - At Contrada Cacaladritta (south eastern Sicily) rare skeletal remains of *Hippopotamus pentlandi*, *Elephas cf. mnaidriensis* and *Bos primigenius siciliae*

are concentrated in deltaic Middle Pleistocene clay sands with *Ostrea edulis* (Roda 1965; Ruggieri 1973; Lentini 1984; Bonfiglio et al. 1997).

Deltaic marine gravels and sands - Disarticulated, fragmented, worn and mechanically selected remains of elephant (*Elephas mnaidriensis*), hippopotamus (*Hippopotamus pentlandi*), red deer (*Cervus elaphus siciliae*), bear (*Ursus cf. arctos*), tortoise (*Testudo cf. hermanni*) are contained in the deltaic marine gravel and sandy deposits underlying Tyrrhenian sands with *Strombus bubonius* at Cape Peloro. Serpulids encrust some skeletal elements (Bonfiglio & Berdar 1979; Bonfiglio & Violanti 1986; Marra 2001).

#### FLUVIAL AND LACUSTRINE COASTAL PLAIN DEPOSITS

At Comiso a regressive marine sequence begins with Early Pleistocene marine deposits and ends at the top with sands of brackish environment which, in turn, pass to deposits of limnic environment underlying Middle Pleistocene marine sands (Conti et al. 1980; Carbone et al. 1982). In the different levels of the limnic deposits skeletal elements of *Elephas falconeri*, *Leithia melitensis*, bats, fish, birds, reptiles (*Lacerta sp.*, *Testudo sp.*, *Emys orbicularis*, *Geochelone sp.*) are contained (Bonfiglio & Insacco 1992).

At Contrada Fusco, near Siracusa, gravel, sandy and silty deposits of coastal plain environment contain a rich vertebrate fauna (an elephant of little reduced size, *Elephas mnaidriensis*, *Hippopotamus pentlandi*, *Ursus cf. arctos*, *Crocota crocuta spelaea*, *Lutra trinacriae*, *Leithia melitensis*, *Maltamys wiedincitensis*, *Crocidura esuae*, *Emys orbicularis*, *Testudo cfr. hermanni*, *Lacerta siculomelitensis*, *Natrix sp.*) (Chilardi 1996; Kotsakis 1996 a; Kotsakis 1996 b; Cassoli & Tagliacozzo 1996 a; Cassoli & Tagliacozzo 1996 b). The coastal plain deposits overlie Early Pleistocene bathyal clays and underlie Tyrrhenian calcarenites (Chilardi & Gilotti 1996).

To the coastal plain environment belong the silty, laminated lacustrine deposits at Acquadolci (north eastern Sicily) containing very abundant skeletal elements of the endemic hippo *Hippopotamus pentlandi* associated with scarce remains of *Cervus elaphus siciliae*, *Ursus cfr. arctos*, *Canis lupus*, *Elephas cfr. mnaidriensis*, *Testudo cfr. hermanni*, birds (Bonfiglio 1992a; Bonfiglio 1995).

#### CAVE DEPOSITS CORRELATED WITH MARINE DEPOSITS

The vertebrate assemblage containing abundant remains of *Hippopotamus pentlandi* associated with rare remains of *Elephas mnaidriensis*, *Cervus siciliae*, *Dama carburangelensis*, *Bos primigenius siciliae*, *Canis lupus*, *Ursus cfr. arctos*, *Crocota spelaea* of the S. Ciro cave overlie middle Pleistocene marine sands (Scinà 1831; Fabiani 1928a; Fabiani 1928b).

At Spinagallo cave the lower vertebrate deposits containing abundant remains of *Elephas falconeri* associated with *Leithia melitensis*, *Maltamys gr. gollcheri-wiedincitensis*, *Crocidura esuae*, *Testudo hermanni* overlie Middle Pleistocene littoral calcarenites (Accordi & Colacicchi 1962; Pet-

ronio 1970; Kotsakis 1977; 1986; Kotsakis & Petronio 1981; Di Grande & Raimondo 1984; Bonfiglio 1992 b; Petruso in progress).

Vertebrate bearing deposits overlying marine terraces - At San Vito Lo Capo peninsula mammal remains of the *Elephas falconeri* F.C. and of the *Elephas mnaidriensis* F.C. overlie marine terraces correlated with the IOS 11-15 and with the IOS 5a or 5c respectively. (Di Maggio et al. 1999). In eastern Sicily vertebrate bearing deposits with prevalent hippo overlie late Middle Pleistocene (Coste di Gigia; Bonfiglio 1992 b) or Tyrrhenian (Scodoni, Bonfiglio 1987) abrasion platforms.

#### KARST FISSURE AND CAVE DEPOSITS

Fossil vertebrates contained in fissure-filling deposits were accumulated by different ways. At Contrada Pianetti remains of *Bos primigenius siciliae*, *Cervus elaphus siciliae*, *Canis cf. lupus*, *Ursus cf. arctos*, have been recovered in a small karst cavity (Bonfiglio et al. 1997; Marra & Bonfiglio, 2000). The taphomic features (fragmentary and disarticulated skeletal elements belonging to different species) suggest that detritus and bones scattered on surface filled the cavity by the action of gravity and/or washing. At Contrada Cimillà, abundant remains of *Dama carburangelensis* were associated to scarce remains of *Cervus elaphus siciliae*, *Elephas mnaidriensis*, *Ursus cf. arctos* (Bonfiglio et al., 1993; Abbazzi et al., 2001). The skeletal element representation shows that surface opening of the fissure may act as a trap for living fallow deers, which were abundant at the site by ethological and/or ecological reasons.

In the bone assemblage of the San Teodoro cave large mammals skeletal elements were accumulated by *Crocota crocuta spelaea* which damaged the main part of bones. A so impressing accumulation by hyena in a large cave (floor area: more than 1000 square meters) is an absolutely novelty for insular environments (Bonfiglio et al. 2001; Marra & Bonfiglio 2002).

#### CHRONOLOGICAL DATA

The record of fossil vertebrate faunas from Sicilian Pleistocene deposits allowed the reconstruction of a detailed biochronological scale, based on geochemical and radiometric dating also (Bada et al. 1991; Rhodes 1996).

Where correlated with marine sediments, the assemblages of the *Elephas falconeri* Faunal Complex are associated with deposits dated early Middle Pleistocene; amino-acid racemization dating yielded an age of  $455 \pm 90$  Ky for *Elephas falconeri* from the Spinagallo and Luparello cave deposits (Bada et al. 1991).

Where correlated with sediment of littoral environments, the assemblages of the *E. mnaidriensis* Faunal Complex are associated with deposits dated as late Middle Pleistocene and / or Late Pleistocene (Bonfiglio & Burgio 1992; Bonfiglio et al. 1996).

Bada et al. (1991) assigned an age of  $200 \pm 40$  Ky to this faunal complex. ESR dating for teeth enamel of *Elephas mnaidriensis* and *Hippopotamus pentlandi* from Contrada Fusco (Rhodes 1996) provided an age ranging between  $146.8 \pm 28.7$  and  $88.2 \pm 19.5$  Ky.

## WORLD ISLANDS IN PREHISTORY

The assemblages of the Pianetti- S. Teodoro F. C., that are not correlated to marine deposits, have not still been dated radiometrically.

Geronimo & Costa 1978; Di Geronimo et al. 1980; Grasso & Lentini 1982; Ruggieri et al. 1976; Ruggieri & Unti 1974; Ruggieri & Unti 1977; Sprovieri 1982). The Hyblean Plateau was separated from the northern sector of the is-




FIGURE 1. COMPOSITE MAP OF ITALY, SICILY AND RESEARCH SITES

### GEOGRAPHIC DISTRIBUTION OF VERTEBRATE BEARING DEPOSITS AND PALAEOGEOGRAPHY.

Fossil documentation of Quaternary mammalian populations in Sicily is not evenly distributed in time and space. A trend is evident towards a richer fossil record from the Early Pleistocene to the Late Glacial, the latter fossils being most widespread. The geographic distribution could be related to taphonomical factors and to effective differences in paleogeography in the insular system (Bonfiglio & Burgio 1992), while the reason why the number of sites and stratigraphic levels varies so greatly with respect to time (younger sites are much more numerous) has never been investigated. This pattern, widespread in regions in which tectonic activity is very intense, is probably due to the destruction of older deposits by intense erosion (Masini et al. in press).

In Sicily the establishment of an extensional tectonic regime from the Early Pleistocene onward resulted in the collapse of peripheral zones of the island and led to the creation of a series of deep marine basins which occupied large areas until the end of the Early Pleistocene (Bonfiglio & Piperno 1996; Bonfiglio et al. in press; Di Geronimo 1979; Di

land. The continental limnic succession at Comiso made up by paleosols, lacustrine and aeolian deposits, constitutes the evidence of the first connection of the Hyblean Plateau with the northern sector of Sicily in the Pleistocene (Conti et al. 1980).

South Calabria, which represents the continental connection of Sicily with southern Italian peninsula, in the Plio-Pleistocene was constituted also by three main blocks bounded by deep marine basins (Barrier et al. 1990; Barrier et al. 1993; Fabbri et al. 1980; Bonfiglio et al. in press). Actually the E. falconeri F. C. denotes the occurrence of an insular system made up of strongly geographically isolated small islands, with very difficult and sporadic connections with the mainland.

According to Bonfiglio & Piperno (1996) the distribution of the E. falconeri assemblages in south-eastern and western Sicily and the scattered traces of Lower Palaeolithic artifacts in the Hyblean Plateau and along the area linking it to western Sicily, might suggest that the dispersal events of humans and of some faunal elements have followed the same route. According to the latter authors the hypothesis

of an African provenance for both *E. falconeri* and human populations through the Sicilian-Tunisian isthmus, as a result of tectonic controlled changes of sea bottom depth during the Early Pleistocene (see Alimen 1975; Kelling & Stanley 1972; Vaufrey 1929), should at least merit a fresh and more detailed re-consideration.

From the beginning of the Middle Pleistocene onward, the evolution of Southern Calabria and Sicily was characterised by an uplifting tendency which led, in the late Middle Pleistocene, to the establishment of a palaeogeography very similar to the present one (Barrier et al. 1990; Barrier et al. 1993; Fabbri et al. 1980; Bonfiglio et al. in press). In the late Pleistocene the interaction of neotectonic with eustatic marine cycles probably led to the establishment of an intermittent filter barrier in the area of Messina Straits, which controlled the processes and timing of the Late Pleistocene-Late Glacial vertebrate faunal dispersion in Sicily. In fact the composition of the two younger F. C.'s denotes that temporary connections with southern Italy occurred.

#### DISCUSSION AND CONCLUSION

Changes in faunal associations of islands are controlled by climatic and environmental factors, but mainly by geographical barriers (Azzaroli 1982; Azzaroli 1996; Dermitzakis & Sondaar 1978).

Biochronologic, taphonomic and palaeogeographic evidence regarding Pleistocene vertebrate faunas of Sicily leads to the following considerations:

The two oldest Faunal complexes (Monte Pellegrino and *E. falconeri* F.C.) characterised by strongly endemic faunal assemblages, denote the occurrence of an insular system made up of geographically isolated small islands, with very difficult and sporadic connections with the mainland. Palaeogeographic data do not provide more precise information on the route followed by dispersals for these complexes.

The three dispersals documented in Sicily from the late Middle Pleistocene to the Late Pleistocene prove the possibilities of connection between the island and the continent through the Strait of Messina area. However, these conditions do not fully explain the modifications recorded in the transition between faunal complexes.

As a matter of fact, variations in the three faunal complexes depend on three components:

(1) Dispersal events. Dispersals (or new occurrences) change obviously the faunal composition and are easier during low-stand episodes.

(2) Extinctions. It seems that most of the extinction events recorded in insular complexes corresponds to those occurring in Italian peninsula and in Western Europe (*Elephas antiquus* become globally extinct; lion and hyena disappear from Western Europe; the European bison is now present only in few residual populations living in Middle-Eastern Europe; *Equus hydruntinus* disappear from Italian peninsula at the beginning of the Holocene and became totally extinct; etc.) It is still unclear, however, if extinctions took place at the same time in the Southern Italian peninsula as in the island. Observations from S. Teodoro cave suggest that at least elephant survived in Sicily longer than elsewhere in Europe. A remarkable exception is, however, represented by the extinction of the endemic dormice

(*Leithia* and *Maltamys*) and of the shrew *Crocidura esuae*. These endemics became extinct during the last glacial cycle, prior then the extinction of *Elephas mnaidriensis*, possibly owing to the interaction of several factors, such as dispersal of new small mammals and predators from the mainland, concomitant to a severe climatic deterioration (see Masini et al. this volume, for a discussion).

(3) Environmental changes related to climatic and physical variations. These changes (connection between the island and the mainland during the Glacial maximum, disappearance of coastal plains subsequent to deglaciation; rearrangement of vegetation in response to climatic changes; etc.) represent very probably direct and indirect causes of dispersals and extinctions.

Components 1 and 2 are basic parameters to define trends in faunal changes; component 3 refers to explanatory models of the variations themselves. The entity of impact in extinction directly linked to these changes is still unclear.

Many taxa identified in different sites and attributed to the considered faunal complexes are still little known. Exhaustive morphological and biometrical data are lacking and relationships with continental ancestors have not still been investigated in the details. In particular, data about the degree of endemism of some taxa (bear, spotted cave hyena, wolf, wild boar) have not still been collected and elaborated. Direct comparisons between the best known taxa of *Elephas mnaidriensis* F.C. (ox, deer) and those of Pianetti-S. Teodoro F.C. have not been done; *Equus hydruntinus* from the latter F. C. is still to be described. Detailed descriptions of micromammals from the three faunal complexes, in particular voles and dormice, are also partially lacking. The relationship between micromammals from Sicily and Malta has never been studied in the details (Petrucci in progress), and can bring a new sight on dispersal dynamics internal to the Sicily-Malta archipelago, given that the latter island likely was a promontory of Sicily during low stand phases.

Careful taxonomic investigations on the recorded taxa, whose spreading on the island has been controlled by geographical barriers of different entity, and a correct evaluation of changes in faunal composition can also emerge from the recognition of differences between mainland and island taxa and a comparison with data from the continent (Italy).

These studies, still in progress, could enhance our knowledge on how and to what extent "discontinuous" insularity conditions interact with climatic and environmental changes in determining the faunal changes of Sicily during last phases of Pleistocene.

#### REFERENCES

- ABBAZZI, L., ET AL.,  
2001. A revision of medium and small sized deer from the Middle and Late Pleistocene of Calabria and Sicily. *Bollettino della Società Paleontologica Italiana*, 40(2), 115-126.
- ACCORDI, B.,  
1957. Nuovi resti di ippopotamo nano nel Pleistocene dei dintorni di Siracusa. *Atti Accademia Gioenia di Scienze Naturali*, 11, 99-109.
- ACCORDI, B.,  
1963. Rapporti tra il "Milazziano" della costa iblea (Sicilia sud-



- orientale) e la comparsa di *Elephas mnaidriensis*. *Geologica Romana*, 2, 295-304.
- ACCORDI, B.,  
1965. Some data on the Pleistocene stratigraphy and related pigmy mammalian faunas of eastern Sicily. *Quaternaria*, 6, 415-430.
- ACCORDI, B. AND R. COLACICCHI,  
1962 - Excavations in the pygmy elephants cave of Spinagallo (Siracusa). *Geologica Romana*, 1, 217-230.
- Agnesi, V., T. Macaluso T. and F. Masini,  
1997 . L'ambiente e il clima della Sicilia nell'ultimo milione di anni. In S. Tusa, ed. *Prima Sicilia. Alle origini della società siciliana*, vol. I. Siracusa: Ediprint, 1997, 31-56
- AGUIRRE, E.  
1968-69. Revision sistemática del los Elephantidae por su morfología y morfometría dentaria. *Estudios Geológicos*, 24 (1968), 109-168; 25 (1969), 123-177, 317-367.
- ALIMEN, M. H.,  
1975. Les isthmes hispano-marocain et siculo-tunisien aux temps acheuléens. *L'Anthropologie*, 79, 399-436.
- AMBROSETTI, P.,  
1968. The Pleistocene dwarf elephants of Spinagallo (Siracusa, south eastern Sicily). *Geologica Romana*, 7, 277-398.
- AMBROSETTI, P., A. AZZAROLI & T. KOTSAKIS,  
1980. Mammiferi del Plio-Pleistocene delle isole italiane. In "I Vertebrati fossili italiani", *Catalogo della Mostra*, Verona, 243-248.
- AZZAROLI, A.,  
1974. Late Miocene interchange of terrestrial fauna across the Mediterranean. *Memorie della Società Geologica Italiana*, 13(suppl. 2), 261-265.
- AZZAROLI, A.,  
1982. Insularity and its effects on the terrestrial Vertebrates : evolutionary and biogeographic aspects. In "Palaeontology, essential of Historical Geology", First International Meeting, Venice, June 1981, 193-213.
- AZZAROLI, A.,  
1996. Processes of island colonization by terrestrial organisms. *Biogeographia*, 18, 33-50.
- AZZAROLI, A. & G. GUAZZONE,  
1979. Terrestrial mammals and land connections in the Mediterranean before and during the Messinian. *Palaeogeography Palaeoclimatology Palaeoecology*, 29, 155-167.
- BADA, J. L. ET AL.,  
1991. Isoleucine epimerization ages of Quaternary Mammals of Sicily. *Il Quaternario*, 4(1a), 5-11.
- BARRIER, P. ET AL.,  
1990. Le Déroit de Messine (Italie). Evolution tectono-sédimentaire récente (Pliocene et Quaternarie) et environment actuel. *Documents et Travaux, IGAL*, 11, 1-272.
- BARRIER, P. ET AL.,  
1993. La lagune Pléistocene à *Gobius* sp. du Monte Torre (Calabre Meridionale): signification paléogéographique. *Rivista Italiana di Paleontologia e Stratigrafia*, 99(1), 127-140.
- BELLUOMINI, G. AND J. L. BADA,  
1985. Isoleucine epimerization ages of the dwarf elephants of Sicily. *Geology*, 13, 451-452.
- BONFIGLIO, L.,  
1987. Nuovi elementi faunistici e stratigrafici del Pleistocene superiore dei Nebrodi (Sicilia nord-orientale). *Rivista Italiana di Paleontologia e Stratigrafia*, 93(1), 145-164.
- BONFIGLIO, L.,  
1991. Correlazioni tra depositi a Mammiferi, depositi marini, linee di costa e terrazzi medio e tardopleistocenici nella Sicilia orientale. *Il Quaternario*, 4(1b), 205-214.
- BONFIGLIO, L.,  
1992a. Campagna di scavo 1987 nel deposito pleistocenico a *Hippopotamus pentlandi* di Acquadolci (Sicilia nord-orientale). *Bollettino della Società Paleontologica Italiana*, 30(3), 157-173.
- BONFIGLIO, L.,  
1992b. Middle and Upper Pleistocene Mammal-bearing deposits in south-eastern Sicily: new stratigraphical records from Coste di Gigia (Syracuse). *Geobios*, 14, 189-199.
- BONFIGLIO, L.,  
1995. Taphonomy and depositional setting of Pleistocene mammal-bearing deposits from Acquadolci (North-Eastern Sicily). *Geobios*, 18, 57-68.
- BONFIGLIO, L. AND A. BERDAR,  
1979. Gli elefanti delle ghiaie pleistoceniche di Messina. *Quaternaria*, 21, 139-177.
- BONFIGLIO, L. AND E. BURGIO,  
1992. Significato paleoambientale e cronologico delle mammalofaune pleistoceniche della Sicilia in relazione all'evoluzione paleogeografica. *Il Quaternario*, 5(2), 223-234.
- BONFIGLIO, L. AND G. INSACCO,  
1992. Palaeoenvironmental, paleontologic and stratigraphic significance of Vertebrate remains in Pleistocene limnic and alluvial deposits from South Eastern Sicily. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 95, 195-208.
- BONFIGLIO, L., A. C. MARRA AND F. MASINI,  
2000. The contribution of Quaternary vertebrates to the paleoenvironmental and paleoclimatic reconstructions in Sicily. In: Hart M. B. (ed.) *Climates: Past and Present*. Geological Society, London, Special Publications, 181, 169-182.
- BONFIGLIO, L. AND M. PIPERNO,  
1996. Early Faunal and Human populations. In: Leighton R. (ed.), *Early Societies in Sicily*, University of London, Accordia, 21- 29.
- BONFIGLIO L. & D. VIOLANTI,  
1986 - Prima segnalazione di Tirreniano ed evoluzione pleistocenica del Capo Peloro (Sicilia Nord-Orientale). *Geogr. Fis. Dinam. Quat.*, 6, 3-15.
- BONFIGLIO, L., ET AL.,  
1996. Large mammal remains from late Middle Pleistocene deposits of Sicily: new stratigraphic evidence from the western edge of the Hyblean Plateau (Southeastern Sicily). *Rivista Italiana di Paleontologia e Stratigrafia*, 102(3), 375-384.
- BONFIGLIO, L., ET AL.,  
1993 - New Pleistocene fissure-filling deposits from the Hyblean Plateau (South Eastern Sicily). *Rivista Italiana di Paleontologia e Stratigrafia*, 98(4), 523-540.
- BONFIGLIO, L. ET AL.,  
1997. Large and small mammals, amphibians, reptiles from a new fissure filling deposit of the Hyblean Plateau (South Eastern Sicily). *Bollettino della Società Paleontologica Italiana*, 35(1), 97-122.
- BONFIGLIO, L. ET AL.,  
2001. A new Late Pleistocene vertebrate faunal complex from Sicily (S. Teodoro Cave, North-Eastern Sicily, Italy). *Bollettino della Società Paleontologica Italiana*, 40(2), 149-158.

- BONFIGLIO, L. ET AL.,  
in press: Pleistocene calabrian and sicilian paleobioprovinces. *Geobios Special Memoires*
- BRUGAL, J. P.  
1987. Cas de "nanisme" insulaire chez l'aurochs. 112th Congrès National des Sociétés Savantes, Lyon, 2: 53-66.
- BUGIO, E. & M. CANI,  
1988. Sul ritrovamento di elefanti fossili ad Alcamo (Trapani, Sicilia). *Il Naturalista Siciliano*, 12(3-4), 87-97.
- BUGIO, E. & M. FIORE,  
1997. Pannonictis arzilla (De Gregorio 1886) a "Villafranchian" element in the fauna from Monte Pellegrino (Palermo, Sicily). *Il Quaternario*, 10, 65-74.
- BUTLER, P.M.,  
1980. The giant erinaceid insectivore *Deinogalerix* Freudenthal from the Upper Miocene of the Gargano, Italy. *Scripta Geologica*, 57, 1-72.
- CARBONE, S., ET AL.,  
1982. I terrazzi marini quaternari dell'area iblea (Sicilia sud-orientale). Contributi conclusivi per la realizzazione della Carta Neotettonica d'Italia, Progetto Finalizzato Geodinamica, Napoli, 506, 1-35.
- CASSOLI, P.F. & A. TAGLIACOZZO,  
1996a. L'avifauna. In: Basile, B. & S. Chilardi, eds. *Le ossa dei Giganti. Lo scavo paleontologico di Contrada Fusco*. Siracusa: Arnaldo Lombardi, 1996, 61-67.
- CASSOLI, P.F. & A. TAGLIACOZZO,  
1996. L'ittiofauna. In: Basile, B. & S. Chilardi, eds. *Le ossa dei Giganti. Lo scavo paleontologico di Contrada Fusco*. Siracusa: Arnaldo Lombardi, 1996, 55-54.
- CHILARDI, S.,  
1996. I macromammiferi. In: Basile, B. & S. Chilardi, eds. *Le ossa dei Giganti. Lo scavo paleontologico di Contrada Fusco*. Siracusa: Arnaldo Lombardi, 1996, 73-80.
- CHILARDI, S. AND A. GILOTTI,  
1996. A new important palaeontological site of Mediterranean basin: preliminary results from Contrada Fusco (Siracusa, Southeastern Sicily). Second International Symposium on the Conservation of our Geological Heritage, Rome, abstract book, 24.
- CONTI, M. A., ET AL.,  
1980. Il Pleistocene in facies limnica di Vittoria (Sicilia meridionale). *Geologica Romana*, 18, 93-104.
- DAAMS, R. AND H. DE BRUIJN,  
1995. A classification of the Gliridae (Rodentia) on the basis of dental morphology. *Hystrix*, 6(1-2), 3-50.
- DERMITZAKIS, M. D. AND P. Y. SONDAAR,  
1978. The importance of fossil mammals in reconstructing paleobiogeography with special reference to the Pleistocene Aegean archipelago. *Annales Géologiques des Pays Helléniques*, 29, 808-840.
- DI GERONIMO, I.,  
1979. Il Pleistocene in facies batiale di Valle Palione (Grammichele, Catania). *Bollettino di Malacologia*, 15, 85-156.
- DI GERONIMO, I. AND B. COSTA,  
1978. Il Pleistocene di Monte dell'Apa (Gela). *Rivista Italiana di Paleontologia e Stratigrafia*, 84(4), 1121-1158.
- DI GERONIMO, I. ET AL.,  
1980. Dati preliminari sulla tettonica della Sicilia sud-orientale. Fogli 273 (Caltagirone), 274 (Siracusa), 275 (Scoglitti), 276 (Ragusa) e 277 (Noto). Contributi preliminari alla realizzazione della Carta Neotettonica d'Italia, Progetto Finalizzato Geodinamica, Napoli, 356, 747-773.
- DI GRANDE, A. AND W. RAIMONDO,  
1984. Linee di costa plio-pleistoceniche e schema litostratigrafico del Quaternario siracusano. *Geologica Romana*, 21, 279-309.
- DI GRANDE, A. AND G. SCAMARDA,  
1973. Segnalazione di livelli a Strombus bubonius Lamarck nei dintorni di Augusta (Siracusa). *Bollettino Accademia Gioenia Scienze Naturali in Catania*, 11, s. 4 (9-10), 157-172.
- DI MAGGIO, C., ET AL.,  
1999. Oscillazioni eustatiche, biocronologia dei depositi continentali quaternari e neotettonica nella Sicilia Nord - Occidentale (Penisola di San Vito Lo Capo - Trapani). *Il Quaternario*, 12(1), 25-50.
- FABBRI, A., F. GHISSETTI AND L. VEZZANI  
1980. The Peloritani Calabria range and the Gioia basin in the Calabrian arc (Southern Italy): relationships between land and marine data. *Geologica Romana*, 19, 131-150.
- FABIANI, R.,  
1928a. Cenni sulle raccolte di Mammiferi quaternari del Museo Geologico dell'Università di Palermo e sui risultati di nuovi saggi esplorativi. *Boll. Ass. Miner. Sicil.*, 4, 25-34.
- FABIANI, R.,  
1928 b. Aggiunte alla conoscenza dei mammiferi fossili del Ragusano in Sicilia. Istituto di Geologia della Regia Università di Palermo.
- GALLETTI, L. AND C. SCALETTA,  
1991. Descrizione di una sequenza del Pleistocene superiore con fauna continentale a San Ciro-Maredolce (Palermo). *Il Naturalista siciliano*, 15(1-2), 3-10.
- GERAADS, D.,  
1995. Rongeurs et insectivores (Mammalia) du Pliocene final de Ahl Oughlam (Casablanca, Maroc). *Geobios*, 28(1), 99-115.
- GLIOZZI, E. AND A. MALATESTA,  
1984. A megacerine in the Pleistocene of Sicily. *Geologica Romana*, 21, 311-389.
- GRASSO, M. AND F. LENTINI,  
1982. Sedimentary and tectonic evolution of the eastern Hyblean Plateau (Southeastern Sicily) during late Cretaceous to Quaternary time. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 39, 261-280.
- KELLING, G. AND D. J. STANLEY,  
1972. Sedimentation in the vicinity of the Strait of Gibraltar. In D. J. Stanley, ed., *The Mediterranean Sea. A Natural Sedimentation Laboratory*. Dowden, Hutchinson & Ross, Stroudsburg, 489-519.
- KOTSAKIS, T.,  
1977. I resti di Anfibi e Rettili pleistocenici della grotta di Spinagallo (Siracusa, Italia). *Geologica Romana*, 16, 211-229.
- KOTSAKIS, T.,  
1979. Sulle mammalofaune quaternarie siciliane. *Bollettino del Servizio Geologico Italiano*, 99, 263-276.
- KOTSAKIS, T.,  
1986. *Crociodura* esui n. sp. (Soricidae, Insectivora) du Pléistocène supérieur de Spinagallo (Sicilia orientale, Italie). *Geologica Romana*, 23, 51-64.
- KOTSAKIS, T.,  
1996a. Anfibi e rettili. In: Basile, B. & S. Chilardi, eds. *Le ossa dei Giganti. Lo scavo paleontologico di Contrada Fusco*. Siracusa: Arnaldo Lombardi, 56-60.

- Kotsakis, T., 1996b. I micromammiferi. In: Basile, B. & S. Chilardi, eds. *Le ossa dei Giganti. Lo scavo paleontologico di Contrada Fusco. Siracusa: Arnaldo Lombardi*, 68-72
- KOTSAKIS T. AND C. PETRONIO,  
1981. I Chiroterri del Pleistocene superiore della grotta di Spinagallo (Siracusa, Sicilia). *Bollettino del Servizio Geologico d'Italia*, 101, 49-76.
- LENTINI, F. (Ed.),  
1984. Carta geologica della Sicilia sudorientale. Scala 1:100.000. Ente minerario siciliano. Comunità economica europea. Istituto di Scienze della Terra, Univ. degli Studi di Catania. S.E.L.C.A., Firenze.
- MARRA, A. C.,  
2001. Resti di vertebrati delle ghiaie pleistoceniche della Formazione di Messina (Sicilia Nord-Orientale). *Il Naturalista Siciliano*, s. 4, 25(1-2), 3-13.
- MARRA, A. C. AND L. BONFIGLIO,  
2000. Tafonomia dei depositi di fessura: due esempi dal Plateau Ibleo - Atti del I Seminario Internazionale di Studi su "il carsismo negli Iblei e nell'area sud-mediterranea", *Speleologia Iblea*, 8, 115-119.
- MARRA, A. C. AND L. BONFIGLIO,  
2002. Taphonomy of hyena dens: an "insular" perspective. In De Renzi et al., eds. *Current Topics on Taphonomy and fossilization*, Valencia: Ajuntament de Valencia, *Collecciò Encontres* 5, 2002
- MASINI, F. AND M. SARÀ. M.,  
1998. *Asoriculus burgioi* sp. nov. (Soricidae, Mammalia) from the Monte Pellegrino faunal complex (Sicily). *Acta Zoologica Cracoviensia*, 41(1), 111-124.
- MASINI F., ET AL.,  
in press. The role of coastal areas in the Neogene - Quaternary mammal island populations of the central Mediterranean. *Biogeographia*.
- MASINI F., ET AL.,  
this volume - Vertebrate assemblages of central-western Mediterranean islands during the Pliocene and Quaternary: reflecting on extinction events.
- PAVIA, M.,  
1999. The Middle-Pleistocene Avifauna of Spinagallo Cave (Sicily, Italy): Preliminary Report. *Smithsonian Contributions to Paleobiology*, 89, 125-127
- PAVIA, M.,  
2000. Le avifaune pleistoceniche dell'Italia Meridionale. Unpublished PhD Dissertation, University of Torino, XIII cycle, 150 pags.
- Pavia, M.,  
2001. Middle Pleistocene fossil avifauna from the *Elephas mnai-driensis* Faunal Complex of Sicily (Italy): preliminary results. *The World of Elephants*, 1st International Congress (Rome, October 16th-20th, 2001), Abstract book.
- PETRONIO, C.,  
1970. I roditori pleistocenici della Grotta di Spinagallo (Siracusa). *Geologica Romana*, 9, 149-194.
- PETRUSO, D.,  
2001. I ghiri della Sicilia antica. In Sarà, M. eds. *Ghiri in Sicilia: ecologia e conservazione*. WWF, 27-41.
- PETRUSO, D.,  
in progress. Stratigrafia e Paleogeografia del Quaternario continentale siciliano: il contributo dei vertebrati. PhD Dissertation in Geological Science, Universities of Naples and Palermo.
- POHLIG, H.,  
1893. Eine Elefantenhöhle Siciliens und der erste Nachweis des Cranialdome von *Elephas antiquus*. *Aus den Abhandlungen der k. bayer. Akademie der Wiss*, 18, 73-100.
- POHLIG, H.,  
1909. Ueber zwei neue altpleistocäne formen von *Cervus*. *Zeitschr. deutsch. geol. Ges.*, 61, 250-253.
- Reumer, W. F. J., 1984. Plio-Pleistocene Soricidae from Tegelen and Hungary. *Scripta Geologica*, 73, 1-173.
- RHODES, E. J.,  
1996. ESR dating of tooth enamel. In: Basile, B. & S. Chilardi, eds. *Le ossa dei Giganti. Lo scavo paleontologico di Contrada Fusco. Siracusa: Arnaldo Lombardi*, 39-44.
- RODA, C.,  
1965. La sezione stratigrafica pleistocenica di Niscemi (Caltanissetta). *Atti dell'Accademia Gioenia Scienze Naturali*, s. 6, 17, 37-62.
- RUGGIERI, G.,  
1973. Gli Ostracodi e la stratigrafia del Pleistocene marino mediterraneo. *Bollettino della Società Geologica Italiana*, 92, 213-232.
- RUGGIERI, G. AND M. UNTI,  
1974. Pliocene e Pleistocene nell'entroterra di Marsala. *Bollettino della Società Geologica Italiana*, 93, 723-733.
- RUGGIERI, G. AND M. UNTI,  
1977. Il Quaternario del Pianoro di S. Margherita di Belice (Sicilia). *Bollettino della Società Geologica Italiana*, 96, 803-812.
- RUGGIERI, G., ET AL.,  
1976. La calcarenite di Marsala (Pleistocene inferiore) e i terreni contermini. *Bollettino della Società Geologica Italiana*, 94, 1623-1627.
- RZEBIK-KOWALSKA, B.,  
1988. Soricidae (Mammalia, Insectivora) from the Plio-Pleistocene and Middle Quaternary of Morocco and Algeria. *Folia Quaternaria*, 57, 51-90.
- RZEBIK-KOWALSKA, B.,  
1995. Climate and history of European shrews (family Soricidae). *Acta Zoologica Cracoviensia*, 38, 95-107.
- SCINÀ, D.,  
1831. Rapporto sulle ossa fossili di Mardolce e degli altri contorni di Palermo. Dalla Reale Tipografia di Guerra, Palermo.
- SPROVIERI, R.,  
1982. Considerazioni sul Plio-Pleistocene della Sicilia. In "Guida Geologica della Sicilia Occidentale", *Società Geologica Italiana*, 115-118.
- THALER, L.,  
1972. Les rongeurs (Rodentia et Lagomorpha) du Monte Pellegrino et la question des anciens isthmes de la Sicilie. *Comptes rendus de l'Académie des Sciences*, 274, 188-190.
- VAUFREY, R.,  
1929. Les éléphants nains des îles méditerranéennes et la question des isthmes pléistocènes. *Archives de l'Institut de Paléontologie Humaine*, 6, 1-220.
- ZAMMIT MAEMPEL, G. AND H. DE BRUIJN,  
1982. The Plio-Pleistocene Gliridae from the Mediterranean Islands reconsidered. *Proceedings, Koninklijke Nederlandse Akademie van Wetenschappen*, 85, 113-128.

This title published by

Archaeopress  
Publishers of British Archaeological Reports  
Gordon House  
276 Banbury Road  
Oxford OX2 7ED  
England  
bar@archaeopress.com  
www.archaeopress.com

BAR S1095

*World Islands in Prehistory: International Insular Investigations  
V Deia Conference of Prehistory*

© the individual authors 2002

ISBN 1 84171 473 9

Printed in England by The Basingstoke Press

All BAR titles are available from:

Hadrian Books Ltd  
122 Banbury Road  
Oxford  
OX2 7BP  
England  
bar@hadrianbooks.co.uk

The current BAR catalogue with details of all titles in print, prices and means of payment is available free from Hadrian Books or may be downloaded from [www.archaeopress.com](http://www.archaeopress.com)