

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Effect of storage conditions on chemical and physical characteristics of hazelnut (*Corylus avellana* L.)**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/132246> since 2016-07-05T16:48:38Z

*Published version:*

DOI:10.1016/j.postharvbio.2013.02.014

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in *Effect of storage conditions on chemical and physical characteristics of hazelnut (Corylus avellana L.)*, 81, July 2013, pages 37–43, DOI: 10.1016/j.postharvbio.2013.02.014.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), +DOI: 10.1016/j.postharvbio.2013.02.014

# **Effect of storage conditions on chemical and physical characteristics of hazelnut (*Corylus avellana* L.).**

D. Ghirardello, C. Contessa, N. Valentini, G. Zeppa, L. Rolle, V. Gerbi, R. Botta

Department of Agricultural, Forest and Food Sciences, University of Turin, Via Leonardo da Vinci, 44 – 10095 Grugliasco (TO), Italy.

e-mail address: [daniela.ghirardello@unito.it](mailto:daniela.ghirardello@unito.it), [cecilia.contessa@unito.it](mailto:cecilia.contessa@unito.it), [nadia.valentini@unito.it](mailto:nadia.valentini@unito.it), [giuseppe.zeppa@unito.it](mailto:giuseppe.zeppa@unito.it), [luca.rolle@unito.it](mailto:luca.rolle@unito.it), [vincenzo.gerbi@unito.it](mailto:vincenzo.gerbi@unito.it), [roberto.botta@unito.it](mailto:roberto.botta@unito.it).

Corresponding author:

Daniela Ghirardello, University of Turin, Via Leonardo da Vinci, 44 – 10095 Grugliasco (TO), Italy. Tel.: +39 011 6708551; fax: +39 011 6708549, e-mail: [daniela.ghirardello@unito.it](mailto:daniela.ghirardello@unito.it).

## **Abstract**

This study evaluated the effect of different storage conditions currently used by the industry, on the chemical, physical and sensory characteristics of ‘Tonda Gentile delle Langhe’ hazelnuts, during one year of storage. The traditional method of in-shell preservation in a storage room at ambient temperature was compared with refrigerated storage of shelled nuts at 4 °C and 55% relative humidity, with or without modified atmosphere (1% oxygen, 99% nitrogen). The following parameters were measured: moisture content, lipid content, total phenolic content, and antioxidant capacity of the kernel; acidity and peroxide value of the oil. The kernel resistance to breakage was evaluated by texture analysis using a compression test. The hazelnuts were also evaluated by sensory analysis. The results showed that the acidity and the peroxide value were the most discriminating parameters. After one year of storage, the acidity of hazelnuts stored at ambient temperature (0.47% oleic acid) was higher than the value considered the acceptable limit after storage (0.40% oleic acid), while refrigerated storage maintained a low level of acidity and lipid oxidation, with the best performance in modified atmosphere (0.13% oleic acid; 0.057 O<sub>2</sub> mmol kg<sup>-1</sup>). Sensory analysis after 12 months also showed differences among the three storage

treatments. In-shell storage of hazelnuts at ambient temperature was able to preserve the kernels below threshold limits of acidity and oxidative degradation for up to 8 months, but refrigeration was necessary to maintain high quality for up to one year. The use of modified atmosphere is recommended for long periods of storage.

**Keywords:** *Corylus avellana* L., storage conditions, lipid oxidation, polyphenols, antioxidant, mechanical properties.

## Highlights

The effect of three storage conditions on quality parameters of hazelnuts was studied in commercial scale conditions.

Time and storage conditions affected significantly acidity and peroxide value.

Low temperature and modified atmosphere (1% oxygen, and 99% nitrogen) decreased additively the lipid oxidation.

Low temperature maintained high level of phenolic content and antioxidant capacity.

## 1. Introduction

Italy is the world's second largest producer of hazelnuts (*Corylus avellana* L.) after Turkey. Italian hazelnut cultivars are highly valued by the food industry for the quality and sensory characteristics of their nuts, in particular for use in confectionery.

Storage conditions affect hazelnut quality and are thus a concern for both food industry and direct consumption. The resistance to oxidation of lipids is frequently associated with the shelf-life of foods, but there are many other factors that contribute to defining the quality of hazelnuts, such as appearance, texture, flavor, chemical composition, nutritional value, and of course, food safety.

Hazelnuts are one of the most nutritious nuts that contain valuable amounts of nutrients, among which lipids predominate (Venkatachalam and Sathe, 2006). The particular fatty acid composition

of hazelnuts, rich in monounsaturated fatty acids, primarily oleic acid (Amaral et al., 2006; Parcerisa et al., 1998), has a recognized beneficial effect on human health (King et al., 2008; Sabaté and Ang, 2009; Torabian et al., 2009); although as a fatty food, hazelnuts are easily susceptible to rancidity. During storage, the lipid fraction can be subjected to hydrolysis and oxidation, resulting in undesirable odors and flavors, and in the reduction of the nutritional value of the kernels. Very few research articles discuss the effects of postharvest handling and storage on chemical and physical characteristics of hazelnuts (Mencarelli et al., 2008; San Martín et al., 2001), and focused on rate of lipid oxidation, and on fat content and fatty acid composition changes (Koyuncu, 2004; Koyuncu et al., 2005).

Hazelnuts also contain significant quantities of dietary fiber, mineral elements, and vitamins. Hazelnuts are an excellent natural source of the antioxidant vitamin E due to their  $\alpha$ -tocopherol content (Kornsteiner et al., 2006), and are also rich in other biologically active compounds such as polyphenols (Alasalvar and Shahidi, 2009). Recently, there is much interest in phenolic compounds because of potential health benefits related to their antioxidant and antiradical activities, anti-inflammatory properties, anticarcinogenic and antimutagenic effects, and antiproliferative potential. Although antioxidant capacity and phenolic composition of hazelnut kernels and hazelnut by-products have been extensively investigated (Alasalvar and Shahidi, 2009), there is a lack of data concerning the stability of the phenolic fraction and the antioxidant capacity of hazelnuts during storage.

It is well known that many extrinsic factors, such as humidity and temperature, can affect the quality of hazelnuts. One of the most important factors is moisture, since water activity influences quality parameters, including mold if moisture is too high, shrivel if too low, color changes, and rancidity. Consequently, to ensure a long shelf-life and to extend protection from rancidification processes, the nuts have to be dried, immediately after harvest, to 3.5-5% kernel moisture content (Richardson, 1988). In addition, the relative humidity (RH) during storage must never exceed 70% (Tombesi, 1985). Enzymatic and chemical rancidification processes, and vitamin E degradation, are

considerably retarded at low temperature. Mold and insect activity is virtually eliminated near freezing temperatures. In addition, reduced oxygen levels can positively affect long-term kernel storage or prolong the shelf-life of roasted kernels. As reported by Ebrahim et al. (1994), in-shell and un-roasted kernels may be stored for 24 months with minimal loss in quality at temperatures below 10 °C, while roasted kernels, stored at 0 °C, 5 °C or 10 °C, may only be held for 6 months prior to development of detectable rancidity. Although low temperatures are recognized as effective means to prolong hazelnut storage, usually the nuts are stored at ambient temperature because of the high energy cost for refrigeration. However, Johnson et al. (2009) reported that hazelnuts can also be stored at ambient temperature under 99% nitrogen atmosphere, with effects comparable to storage condition at 3-6 °C and 50-60% RH. Optimal storage conditions may be provided by a combination of low temperature and modified atmosphere (saturated with N<sub>2</sub> and/or CO<sub>2</sub>) or vacuum. Recently, Mencarelli et al. (2008) demonstrated that a high concentration of nitrogen (98-100%) and low temperature (4 °C) are best for maintaining color, firmness, acidity and peroxide values of the kernels.

Storage stability or shelf-life of foods could be defined as maintenance of the sensory and physical characteristics associated with becoming stale (Baixauli et al., 2008). Texture is one of the most important characteristics of edible fruits and vegetables (Kilcast, 2004), that can be affected by postharvest treatments and processing. The mechanical behavior of hazelnuts has been studied to characterize different varieties (Güner et al., 2003; Valentini et al., 2006), or different cracking and roasting systems (Özdemir and Özilgen, 1997; Saklar et al., 1999; Demir and Cronin, 2004). However, very little data are available on the effect of particular storage conditions on hazelnut textural characteristics (Mencarelli et al., 2008; Ghirardello et al., 2009b).

This work focuses on the evaluation of the effects of three different storage conditions on the quality of ‘Tonda Gentile delle Langhe’ hazelnuts harvested in 2009, during one year of storage. Usually, data reported in literature are from pilot storage systems. In this research, nuts were stored in industrial storage rooms, typical of commercial storage.

## **2. Materials and Methods**

### **2.1. Sample preparation**

The experiments were carried out on ‘Tonda Gentile delle Langhe’ hazelnuts harvested in a single orchard, located in Cravanzana (Langhe district, Piedmont, NW Italy). Immediately after harvest, the nuts were dried to about 5% moisture content in the kernels. The nuts, at 10% starting moisture content, were dried in a food dryer with a slow stream of warm air (50 °C) for 8 h. After an additional cooling step of 6 h, the samples were stored in storage rooms provided by Ascopiemonte – Organizzazione Produttori Frutta a Guscio S.c.a.r.l. (Santo Stefano Belbo, Piedmont, Italy). Just before the samples were placed in storage rooms (Beginning), samples of hazelnuts (four replicates of 2 kg each) were analyzed, and data were used as references for all treatments. Hazelnuts were then divided into in-shell and shelled batches. The kernels were obtained using an industrial shelling-machine (P8O Sheller, Chianchia, Cherasco, Italy). The samples were packaged in 25-kg food polypropylene bags.

Three different storage conditions were tested: in-shell hazelnuts were stored at ambient temperatures (ranging between 10 and 26 °C) and 60-80 % RH, while kernels (shelled hazelnuts) were cold stored (4 °C, 55% RH) with or without modified atmosphere (1% oxygen, 99% nitrogen). Analyses were conducted after 8 and 12 months of storage, except for kernels stored under nitrogen. The modified atmosphere chamber was opened only after 1 year, according to the standard methods in the industry. Therefore, in this case the kernels were analyzed at the end of the experiment. At each sampling time, batches of about 2 kg of kernels and 5 kg of in-shell nuts were taken for analysis. Just before analysis, the in-shell hazelnuts were manually cracked and shelled.

### **2.2. Chemical analyses**

Moisture content was determined by vacuum oven at  $70 \pm 1^\circ \text{C}$  (method 934.06; AOAC, 1990). Total fat content was determined by using the Soxhlet petroleum-ether extraction method (method 920.39C; AOAC, 1990).

Titrateable acidity (expressed as the percentage of oleic acid), peroxide value (PV, determined by iodometric titration and expressed as millimoles of active O<sub>2</sub> per kg of oil), and fatty acid composition of the oil were determined on each sample according to the European Official Methods of Analysis (Council Regulation, EEC-N.2568/91). Fatty acids were converted into methyl esters (FAME) and analyzed using a Shimadzu GC-2010 plus gas chromatograph (Shimadzu, Milan, Italy) equipped with a flame ionization detector (FID). Separation was achieved on a Supelco SP<sup>TM</sup> 2560 capillary column (Supelco, Bellefonte, USA), 100 m long, 0.25 mm i.d, 0.2 µm film thickness. The split-splitless injector was used at a split ratio of 1:50. The injector volume of the sample was 1 µL. The injector and detector temperatures were both set at 250 °C. The column temperature was 165 °C for 1 min hold and programmed to increase to 200 °C at a rate of 0.05 °C s<sup>-1</sup> and then held for 45 min. Helium was used as the carrier gas with a flow rate of 16.7 µL s<sup>-1</sup>. Fatty acid peaks were identified by comparing retention times with FAME stock solution. The quantification was performed by internal normalization.

The extraction of antioxidant compounds was conducted by mixing finely ground kernels with 50% ethanol (v/v) in ultrapure water acidified with formic acid (pH 4). Each sample (1 g) was extracted with 10 mL of extraction solvent in a capped glass tube on a VDRL 711 orbital shaker (Asal s.r.l., Milan, Italy) at a constant oscillation (1.67 oscillations s<sup>-1</sup>), at ambient temperature (20-22 °C), for 77 min (Ghirardello et al., 2009a). Afterward, the extracts were centrifuged (15 min at 2700 × g) and the supernatants were filtered through a 0.45-µm pore size syringe filter. The extractions were done in quadruplicate for each sample, and the extracts were stored at -18 °C until analysis.

The total phenolic content (TPC) of kernel extracts was assayed spectrophotometrically by means of the Folin-Ciocalteu method, as modified by Singleton and Rossi (1965). Briefly, 0.5 mL of kernel extract was added into a tube containing 2.5 mL of 10-fold dilute Folin-Ciocalteu reagent. The tube was vortexed and allowed to stand at room temperature for 3 min. Then, 2 mL of Na<sub>2</sub>CO<sub>3</sub> (7.5%, w/v) was added to the mixture. The absorbance was measured at 756 nm with a UV-1700



PharmaSpek UV-Vis spectrophotometer (Shimadzu, Milan, Italy), after 15 min heating at 45 °C (Pinelo et al., 2003). Phenolic content was expressed as grams of gallic acid equivalent (GAE) per kilogram of sample.

To determine the antioxidant capacity of the extracts, the DPPH• (2,2-diphenyl-1-picrylhydrazyl) radical scavenging assay (RSA) was performed by the method described by von Gadov et al. (1997) slightly modified. Seventy-five microliters of sample extract were added to 3 mL of DPPH• methanol solution ( $6.1 \times 10^{-5} \text{ mol L}^{-1}$ ) and incubated for 1 h at room temperature in the dark. The absorbance was measured at 515 nm against methanol solution of DPPH• as a blank. The results were expressed as inhibition percentage (IP) of the DPPH radical calculated according to the following Eq.:

$$\text{IP} = [(A_{0\text{min}} - A_{60\text{min}}) / A_{0\text{min}}] \times 100$$

where  $A_{0\text{min}}$  is the absorbance of the blank at  $t = 0 \text{ min}$ ; and  $A_{60\text{min}}$  is the absorbance of the samples at 60 min. Results were expressed as millimoles of Trolox equivalent (TE) per kilogram of sample, by means of a dose-response curve for Trolox (0-350  $\mu\text{mol}$ ).

### 2.3. Physical analyses

Instrumental texture properties of the kernels were monitored with a Universal Testing Machine (UTM) TA.XT2i® Texture Analyser (Stable Micro System, Godalming, Surrey, UK). The device was equipped with a 50 kg load cell and a HDP/90 platform. The compression test was performed at  $1 \text{ mm s}^{-1}$  constant bar speed with a P/75 circular aluminum flat probe (75mm of diameter) (Valentini et al., 2006). The force-deformation curve was acquired as a graph (Fig. 1) and processed by the Texture Export Exceed software rel. 2.54 (Stable Micro Systems, Godalming, UK). Three replicates of 10 nuts for each storage condition were compressed along the longitudinal axis through the hilum containing the major dimension (length) of the kernel (Güner et al., 2003). The breakage characteristics of hazelnut were expressed according to Saklar et al. (1999) by the following parameters: the rupture force (N), represented by the first fracture point ( $F_1$ ), the slope

( $S_{F1}$ ) of the line between the starting point and the first fracture point ( $N\ m^{-1}$ ), and the rupture energy (mJ), represented by the area under the curve ( $W_1$ ) for the region between the starting point and the first fracture point.

[Fig. 1 about here]

## 2.4. Sensory analyses

The sensory evaluation of the samples was performed with a duo-trio overall difference test (ISO 10399, 2004), by a group of 24 trained panelists. Since ‘Tonda Gentile delle Langhe’ hazelnuts are generally used by the food industry as roasted kernels, the tastings were conducted on roasted samples. To ensure the characteristic texture and aroma, the kernels were roasted (160 °C, 20 min in a ventilated oven) just before the tasting sessions. All samples were furnished in white plastic cups, containing 6-7 whole roasted kernels. Water was provided for palate cleansing.

## 2.5. Statistical analyses

Results were expressed as mean  $\pm$  SD. Analyses of variance were done using SPSS software (version 18.0 for Windows, SPSS Inc., Chicago, Illinois). ANOVA was performed on the chemical and physical data, considering all factors and their interaction (two-way ANOVA). Intra-storage condition and intra-storage time differences were analyzed using one-way ANOVA (factor being storage time and storage condition). Significant differences ( $P < 0.05$ ) among means were determined using the Tukey’s test at a fixed level of  $\alpha = 0.05$ . For the sensory analysis, the total amount of correct responses was compared to the critical number of correct responses in a ‘duo-trio’ difference test for significance with an  $\alpha = 0.05$ .

# 3. Results and Discussion

## 3.1. Chemical analyses

Data on moisture and lipid content of kernels, acidity and peroxide value of oils, are reported in Table 1. The kernel moisture content was nearly stable during storage, except for in-shell hazelnuts stored at room temperature at the twelfth month, probably due to a partial rehydration of the kernels. The moisture content of the kernels never reached 5%, the threshold value for the good preservation of hazelnuts.

The lipid content of the kernels was very stable under all storage conditions. Contrary to data reported by Koyuncu (2004), no significant differences of lipid content were observed between hazelnut stored shelled and in-shell. Table 2 shows the fatty acid compositions of hazelnuts during the storage period. The predominant saturated fatty acid in hazelnuts was palmitic (C16:0), followed by stearic (C18:0). The two most abundant unsaturated fatty acids were oleic (C18:1) and linoleic (C18:2). Their relative amounts were comparable to data reported in literature for the same cultivar (Cristofori et al., 2008). The linoleic acid content varied inversely to the oleic acid content (Koyuncu et al., 2005), and was always less than 9% of total fatty acids, considered a critical threshold value by the food industry (Arcoleo, 1991).

[Table 1 and Table 2 about here]

During storage, the ratio of unsaturated/saturated fatty acids decreased from 0.1203 to 0.1094. Indeed, the total saturated fatty acid content increased from 7.69% to 8.42% at the end of the storage time, while the total unsaturated fatty acid content decreased from 92.30% to 91.69% (Table 2). These changes were significant and related to the decrease of linoleic acid content, probably as the result of its peroxidation and subsequent loss. A similar trend was reported by Koyuncu et al. (2005) for the fatty acid composition of hazelnut kernels stored for one year in vacuum packages.

A significant effect of time and condition of storage on the indices of stability of the lipid fraction was observed. As expected, the acidity and peroxide values increased with storage in all samples (Table 1). Interaction effect of storage condition and storage time was also found significant for the acidity and peroxide value parameters ( $P < 0.001$ ). At the beginning of the experiment, the kernels exhibited very low acidity and peroxide values, close to zero (0.06% oleic

acid;  $O_2$  0.045 mmol kg<sup>-1</sup>), indicating the absence of initial trygliceride hydrolysis and fatty acid oxidation. After 8 months of storage, in-shell hazelnuts stored at ambient temperature showed higher acidity and PV in comparison with cold stored kernels (0.25 and 0.10% of oleic acid,  $O_2$  of 0.082 and 0.050 mmol kg<sup>-1</sup>, respectively). The variation of these two parameters was very small, and the lipid fraction of the hazelnuts maintained its characteristics of freshness and stability. This trend is in agreement with Gattuso et al. (1995), where the PV at eight months of storage of in-shell nuts stored in perforated containers in a dry and airy storage room, was  $O_2$  of 0.10 mmol kg<sup>-1</sup>. After twelve months of storage, the acidity of in-shell hazelnuts stored at ambient temperature was 0.47% oleic acid. This value was higher than the acidity reported for the superior extra-virgin olive oils (0.40% oleic acid) that, in the absence of indications about a critical acidity value for the nut industry, we can choose as a limit of acceptability after storage.

Storage at low temperature permitted to maintain a low level of acidity and lipid oxidation, with the best performance in modified atmosphere (0.13% of oleic acid and  $O_2$  of 0.057 mmol kg<sup>-1</sup>).

The PV is one of the parameters adopted by the nut industry to evaluate the storage aptitude of hazelnut; higher scores are assigned to lots with  $O_2$  values lower than 0.25 mmol kg<sup>-1</sup> (Arcoleo, 1991). In this study the PV was lower than  $O_2$  of 0.25 mmol kg<sup>-1</sup> at all treatments and times, except for in-shell hazelnuts after 12 months of storage. The analysis of variance documented an increasing PV trend for storage time ( $0 < 8 < 12$  months), and for storage conditions (refrigerated under nitrogen < refrigerated < ambient temperature). San Martin et al. (2001) reported, for hazelnuts stored in modified atmosphere conditions, a positive correlation, over time, between oxidative rancidity and oxygen content. Differences in PV with time were noted by the third month of storage, but were almost stable at 6, 9 and 12 months with values of 0.45-0.50. In another study on storage (Mencarelli et al., 2008), prevention of oxidative process in hazelnuts was achieved by modified atmosphere (100% CO<sub>2</sub> or 100% N<sub>2</sub>) at 4 °C. A little increase of PV (from  $O_2$  of 0.075 to 0.15 mmol kg<sup>-1</sup>) with the acidity was reported. Hazelnuts stored for three months at 4 °C under nitrogen (98% N<sub>2</sub>) maintained the initial PV, in accordance with our data.

The total phenolic content decreased significantly at the 8<sup>th</sup> month (GAE of 1.05 and 1.09 g kg<sup>-1</sup> for shelled and in-shell hazelnuts, respectively), then remained almost unchanged, with a slight increase in refrigerated kernels (Table 1). At the twelfth month of storage the higher values of TPC was recorded for refrigerated kernels (GAE of 0.19 g kg<sup>-1</sup>) with significant differences compared to the other storage conditions. Previous studies reported that low temperature and modified atmosphere can effectively prevent the decrease of phenolic content and antioxidant capacity in long-term stored nuts. For example, during 12 months of storage, low temperature (1 °C instead of 20 °C) and packaging atmosphere with 100% N<sub>2</sub>, prevented additively the loss in antioxidants in stored pistachios (Tsantili et al., 2011). Peanuts stored at 20 and 35 °C for up to 4 months had 35% less total phenolics than initially (Talcott et al., 2005), but at 20 °C total phenolics losses were less than those at 35 °C. During one year of storage, Christopoulos and Tsantili (2001) reported a progressive decrease of total phenolics in walnuts. The losses were additively reduced by lower temperature and packaging under elevated N<sub>2</sub> or CO<sub>2</sub>.

The present results showed that the decrease in DPPH• values followed a pattern similar with that of TPC. When data were analyzed by two-way ANOVA (Table 1), only the storage time effect was significant ( $P < 0.001$ ) for both parameters. This result was more evident for antioxidant capacity indices, with an increase of the DDPH• scavenging activity between the 8<sup>th</sup> and the 12<sup>th</sup> month of storage, and the best performance at the 12<sup>th</sup> month for the refrigerated kernels (TE of 6.29 mmol kg<sup>-1</sup>). This was partially in contrast with the hypothesis that storage would decrease the antioxidant potential of hazelnuts due to oxidation of phenolic compounds. A similar behavior was highlighted by Bolling et al. (2010) studying the influence of storage on polyphenol content and antioxidant capacity of California almond skins. They observed an increase of polyphenols and antioxidant capacity in skin extracts of almonds stored for 15 months, without storage temperature effects (4 or 23 °C, 30% RH). They suggested that a dynamic process affected the changes in flavonoid and phenolic acid contents, by an increase of polyphenol extractability, a degradation of polymeric polyphenols and consequently an increase of soluble phenolics, or a polyphenols

synthesis after harvest, just observed in a few foods. Therefore, as reported by Manzocco et al. (2001), the loss of antioxidant capacity of polyphenols is due to their enzymatic or chemical oxidation; however, some authors suggested that partially oxidized polyphenols can exhibit higher antioxidant activity than that of non-oxidized phenols.

### 3.2. Physical and sensory analyses

As reported in a previous study on hazelnuts of the cultivar ‘Tonda Romana’ (Ghirardello et al., 2009b), rupture force (N) and slope at the first fracture point ( $\text{slope}_{F1}$ ,  $\text{N m}^{-1}$ ) are the most discriminating parameters during storage. These indices can be considered important quality markers because they are correlated with sensory characteristics of crispiness and crunchiness (Saklar et al., 2009). The textural parameters measured in this experiment, despite the high variability of the values, showed significant differences from the starting values at both eight and twelve months of storage (Table 3). The two-way ANOVA analysis showed a significant effect ( $P < 0.001$ ) of storage time for the rupture force and  $\text{slope}_{F1}$ , and of storage conditions for  $\text{slope}_{F1}$  ( $P < 0.05$ ); an interaction effect ( $P < 0.01$ ) was observed for the rupture energy. All parameters increased with time in in-shell hazelnuts; for those shelled the highest values of rupture force and rupture energy were registered at 8<sup>th</sup> month. These results are in agreement with those reported by Ghirardello et al. (2009b) showing that the rupture force and the rupture energy of raw hazelnuts refrigerated and cold stored under nitrogen were higher after four and eight months of storage, in comparison to the fresh samples. After 12 months of storage, the in-shell hazelnuts were characterized by the highest values of rupture force and rupture energy, and the lowest value of  $\text{slope}_{F1}$ , therefore they had the highest firmness and resistance to deformation. However, only the rupture energy (mJ) was able to discriminate cold stored kernels, with or without modified atmosphere, from those stored at ambient temperature, characterized by a greater resistance to fracture. Mencarelli et al. (2008) utilized an Instron Universal Testing Machine to study the effect of different temperatures and modified atmospheres on the deformation of hazelnut kernels. They

observed an increase with time of the kernels fracturability for all storage conditions. Though, these data cannot be compared with our results because of the use of different operative conditions during the compression test and of the measurement of different parameters.

[Table 3 about here]

The duo-trio difference test results are reported in Table 4. The obtained results showed that, after 8 months of storage, no sensory differences were found by panelists between hazelnuts stored under refrigerated and ambient conditions. Instead, after 12 months there were sensory differences among all storage conditions, and particularly between the samples refrigerated and refrigerated under nitrogen.

[Table 4 about here]

#### **4. Conclusions**

Acidity and peroxide values in this study indicated the efficacy of low temperature for minimizing lipid oxidation during 8 months of storage. Assays of other important quality parameters did not document significant differences among the samples. Sensory panelists were not able to discriminate the refrigerated hazelnuts from those stored at ambient temperature for 8 months. However, after 12 months, the sensory analysis was able to distinguish the different storage techniques. Again, acidity and peroxide value were the more powerful and discriminating indices. Low temperature and elevated nitrogen atmosphere prevented additively lipid oxidation giving the lowest values of acidity and peroxide value. Furthermore, the use of low temperature was more effective for maintaining high level of phenolic content and antioxidant capacity than low temperature combined with elevated nitrogen atmosphere. The effectiveness of low temperature in delaying the quality decay of hazelnuts is confirmed; refrigeration was effective for maintaining kernel quality for up to one year of storage. On the other hand, when stored as in-shell nuts at ambient temperature, quality was only maintained for a period of about 8 months after harvest.

## Acknowledgements

This work was funded by a grant from Regione Piemonte – Assessorato Agricoltura. Title: “Filiera nocciolo: aspetti agronomici e tecnologici per il miglioramento delle produzioni piemontesi (CORIFIL)”. The authors would like to thank ‘Ascopiemonte’ for supplying the storage rooms, and ‘Chianchia s.r.l.’ that kindly provided the shelling machine.

## References

- Alasalvar, C., Shahidi, F. (Eds.), 2009. Tree nuts: composition, phytochemicals and health effects. CRC Press, Boca Raton, FL.
- Amaral, J.S., Casal, S., Citová, I., Santos, A., Seabra R.M., Oliveira, B.P.P. 2006. Characterization of several hazelnut (*Corylus avellana* L.) cultivars based in chemical, fatty acid and sterol composition. Eur. Food Res. Technol., 222, 274–280.
- AOAC, 1990. Official Methods of Analysis (17th ed.). Association of Official Analytical Chemists, Washington, DC.
- Arcoleo, G., 1991. Caratteristiche e composizione di nocciole di alcune varietà coltivate in Sicilia. La rivista italiana delle sostanze grasse, 68, 257–260.
- Baixaui, R., Salvador, A., Fiszman, S.M., 2008. Textural and colour changes during storage and sensory shelf life of muffins containing resistant starch. Eur. Food Res. Technol. 226, 523–530.
- Bolling, B.W., Blumberg, J.B., Chen, C-Y.O., 2010. The influence of roasting, pasteurization, and storage on polyphenols content and antioxidant capacity of California almond skins. Food Chem. 123, 1040–1047.
- Commission Regulation, 1991. “Commission Regulation (EEC) n 2568/91 of July 11<sup>th</sup> 1991 on the characteristics of olive oil and olive-residue oil and on the relevant methods of analysis”.
- Cristofori, V., Ferramondo, S., Bertazza, G., Bignami, C., 2008. Nut and kernel traits and chemical composition of hazelnut (*Corylus avellana* L.) cultivars. J. Sci. Food Agric. 88, 1091–1098.



- Christopoulos, M.V., Tsantili, E., 2011. Effects of temperature and packaging atmosphere on total antioxidants and colour of walnut (*Juglans regia* L.) kernels during storage. *Sci. Hortic.* 131, 49–57.
- Demir, A.D., Cronin, K., 2004. The thermal kinetics of texture change and the analysis of texture variability for raw and roasted hazelnuts. *Int. J. Food Sci. Tech.* 39, 371–383.
- Ebraheim, K.S., Richardson, D.G., Tetley, R.M., 1994. Effects of storage temperature, kernel intactness and roasting temperature on vitamin E, fatty acids and peroxide value of hazelnuts. *Acta Hort.* 351, 677–684.
- Gattuso, A.M., Indovina, M.C., Arcoleo, G., 1994. Variazioni della composizione chimica del grasso nel periodo di conservazione di semi di nove cvs di nocciole con particolare riferimento all'ossidabilità. *Acta Hort.* 351, 649–656.
- Ghirardello, D., Zeppa, G., Belviso, S., Gerbi, V., 2009a. Determination of total phenolic content in hazelnut kernel extracts and their antioxidant capacity – Evaluation of spectrophotometric methods. *Proceeding 2 EuroFoodChem XV – Food for the Future*, Copenhagen, Denmark, 5-8 July 2009, 142–145.
- Ghirardello, D., Rolle, L., Zeppa, G., 2009b. Effects of storage conditions on hazelnut (*Corylus avellana* L.) textural characteristics. *Acta Hort.* 845 (2), 681–685.
- Güner, M., Dursun, E., Dursun, I.G., 2003. Mechanical behavior of hazelnut under compression loading. *Biosyst. Engng.* 85, 485–491.
- International Organisation for Standardisation. ISO 10399:2004. Sensory analysis – Methodology – Duo-trio test.
- Johnson, J.A., Yahia, E.M., Brandl, D.G., 2009. Dried fruits and tree nuts. In: Yahia, E.M., Raton, B. (Eds.), *Modified and controlled atmospheres for the storage, transportation, and packaging of horticultural commodities*. CRC Press, Taylor & Francis Group, pp. 507–526.
- Kilcast, D. (Ed.), 2004. *Texture in food*. Vol. 2: Solid foods. CRC Press, Boca Raton, FL.

- King, J.C., Blumberg, J., Ingwersen, L., Jenab, M., Tucker, K.L., 2008. Tree nuts and peanuts as components of a healthy diet. *J. Nutr.* 138, 1736S–1740S.
- Kornsteiner, M., Wagner, K-H., Elmadfa, I., 2006. Tocopherols and total phenolics in 10 different nut types. *Food Chem.* 98, 381–387.
- Koyuncu, M.A., 2004. Change of fat content and fatty acid composition of Turkish hazelnuts (*Corylus avellana* L.) during storage. *J. Food Qual.* 27, 304–309.
- Koyuncu, M.A., Islam, A., Küçük, M., 2005. Fat content and fatty acid composition of hazelnut kernels in vacuum packages during storage.
- Manzocco, L., Calligaris, S., Mastrocola, D. Nicoli, M.C., Lerici, C.R., 2001. Review of non-enzymatic browning and antioxidant capacity in processed foods. *Trends Food Sci. Technol.* 131, 49–57.
- Mencarelli, F., Forniti, R., Desantis, D., Bellincontro, A., 2008. Effetti delle atmosfere inerti e della temperatura nella conservazione delle nocciole. *Ingredienti Alimentari*, VII, agosto, 16–21.
- Özdemir, M., Özilgen M., 1997. Comparison of the quality of hazelnuts unshelled with different sizing and cracking systems. *J. Agric. Engng. Res.* 67, 219–227.
- Parcerisa, J., Richardson, D.G, Rafecas, M., Codony, R., Boatella, J., 1998. Fatty acid, tocopherol and sterol content of some hazelnut varieties (*Corylus avellana* L.) harvested in Oregon (USA). *J Chrom. A.* 805, 259–268.
- Pinelo, M., Rubilar, M., Sineiro, J., Núñez, M.J., 2003. Extraction of antioxidant phenolics from almond hulls (*Prunus amygdalus*) and pine sawdust (*Pinus pinaster*). *Food Chem.* 85, 267–273.
- Richardson, D.G., 1988. Hazelnut quality. *Proceedings of the Annual Meeting of the Nut Growers Society of Oregon, Washington and British Columbia.* 73, 83–86.
- Sabaté, J., Ang, Y., 2009. Nuts and health outcomes: new epidemiological evidence. *Am. J. Clin. Nutr.* 89, 1643S–1648S.
- Saklar, S., Ugan, S., Katnas, S., 1999. Instrumental crispness and crunchiness of roasted hazelnuts and correlations with sensory assessment. *J. Food Sci.* 6, 1015–1019.

- San Martín, M.B., Fernández-García, T., Romero, A., López, A., 2001. Effects of modified atmosphere storage on hazelnut quality. *J. Food Process. Pres.* 25, 309–321.
- Singleton, V.L., Rossi, J.A., 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.* 16, 144–158.
- Talcott, S.T., Duncan, C.E., Pozo-Insfran, D.D., Gorbet, D.W., 2005. Polyphenolic and antioxidant changes during storage of normal, mid, and high oleic acid peanuts. *Food Chem.* 89, 77–84.
- Tombesi, A., 1985. *Il nocciolo. Manuale pratico.* Reda, Roma.
- Torabian, S., Haddad, E., Rajaram, S., Banta, J., Sabaté, J., 2009. Acute effect of nut consumption on plasma total polyphenols, antioxidant capacity and lipid peroxidation. *J. Hum. Nutr. Diet.* 22, 64–71.
- Tsantili, E., Kostantinidis, K., Christopoulos, M.V., Roussos, P.A., 2011. Total phenolics and flavonoids and total antioxidant capacity in pistachio (*Pistachia vera* L.) nuts in relation to cultivars and storage conditions. *Sci. Hortic.* 129, 694–701.
- Valentini, N., Rolle, L., Stévigny, C., Zeppa, G., 2006. Mechanical behavior of hazelnuts used for table consumption under compression loading. *J. Sci. Food. Agric.* 86, 1257–1262.
- Venkatachalam, M., Sathe, S.K., 2006. Chemical composition of selected edible nut seeds. *J. Agric. Food Chem.* 54, 4705–4714.
- von Gadow, A., Joubert, E., Hansmann, C.F., 1997. Comparison of antioxidant activity of aspalathin with that of other plant phenols of Rooibos tea (*Aspalathus linearis*),  $\alpha$ -tocopherol, BHT and BHA. *J. Agric. Food Chem.* 45, 632–648.

**Table 1.** Moisture content, lipid content, acidity, peroxide value (PV), total phenolic content (TPC), and antioxidant capacity (DPPH<sup>•</sup>) of the hazelnuts during storage. IS: in-shell hazelnuts; S: shelled hazelnuts (kernel); Refrigerated: stored at 4 °C and 55% RH; Under N<sub>2</sub>: refrigerated under nitrogen (4 °C, 55% RH - 1% O<sub>2</sub>, 99% N<sub>2</sub>); Ambient temp.: stored at ambient temperature (70% RH).

		Beginning	8 <sup>th</sup> month	12 <sup>th</sup> month	$P_{sc}^{\dagger}$	$P_{st}$	$P_{sc} \times P_{st}$
Moisture content (% dry basis)					***	***	***
S	Refrigerated	3.94 ± 0.04 <sup>AB</sup>	3.97 ± 0.05 <sup>B</sup>	3.89 ± 0.01 <sup>aA</sup>			
S	Under N <sub>2</sub>	3.94 ± 0.04		3.97 ± 0.05 <sup>b</sup>			
IS	Ambient temp.	3.94 ± 0.04 <sup>A</sup>	3.97 ± 0.02 <sup>A</sup>	4.95 ± 0.04 <sup>cB</sup>			
Lipid content (% dry basis)					NS	NS	NS
S	Refrigerated	61.28 ± 7.53	62.38 ± 1.27	65.36 ± 3.61			
S	Under N <sub>2</sub>	61.28 ± 7.53		63.74 ± 4.26			
IS	Ambient temp.	61.28 ± 7.53	63.78 ± 2.33	62.69 ± 3.27			
Acidity (% oleic acid)					***	***	***
S	Refrigerated	0.06 ± 0.01 <sup>A</sup>	0.10 ± 0.01 <sup>aB</sup>	0.27 ± 0.01 <sup>bC</sup>			
S	Under N <sub>2</sub>	0.06 ± 0.01 <sup>A</sup>		0.13 ± 0.01 <sup>aB</sup>			
IS	Ambient temp.	0.06 ± 0.01 <sup>A</sup>	0.25 ± 0.02 <sup>bB</sup>	0.47 ± 0.02 <sup>cC</sup>			
PV (O <sub>2</sub> , mmol kg <sup>-1</sup> )					***	***	***
S	Refrigerated	0.045 ± 0.006 <sup>A</sup>	0.050 ± 0.004 <sup>aA</sup>	0.093 ± 0.005 <sup>bB</sup>			
S	Under N <sub>2</sub>	0.045 ± 0.006 <sup>A</sup>		0.057 ± 0.005 <sup>aB</sup>			
IS	Ambient temp.	0.045 ± 0.006 <sup>A</sup>	0.082 ± 0.006 <sup>bB</sup>	0.263 ± 0.012 <sup>cC</sup>			
TPC (GAE, g kg <sup>-1</sup> )					NS	***	NS
S	Refrigerated	1.40 ± 0.29 <sup>B</sup>	1.05 ± 0.05 <sup>A</sup>	1.19 ± 0.04 <sup>bAB</sup>			
S	Under N <sub>2</sub>	1.40 ± 0.29 <sup>B</sup>		1.03 ± 0.04 <sup>aA</sup>			
U	Ambient temp.	1.40 ± 0.29	1.09 ± 0.13	1.07 ± 0.06 <sup>a</sup>			
DPPH <sup>•</sup> (TE, mmol kg <sup>-1</sup> )					NS	***	NS
S	Refrigerated	8.40 ± 1.79 <sup>B</sup>	4.35 ± 0.36 <sup>A</sup>	6.29 ± 0.32 <sup>bAB</sup>			
S	Under N <sub>2</sub>	8.40 ± 1.79 <sup>B</sup>		5.20 ± 0.23 <sup>aA</sup>			
IS	Ambient temp.	8.40 ± 1.79 <sup>B</sup>	3.81 ± 0.94 <sup>A</sup>	5.50 ± 0.28 <sup>aA</sup>			

Data were expressed as mean ± SD (n = 4). Values in the column with different lowercase letters were significantly different at  $P < 0.05$ . Values in the row with different capital letters were significantly different at  $P < 0.05$ .

<sup>†</sup>Probabilities of the effects:  $P$ -level calculated for samples from different storage condition ( $P_{sc}$ ).  $P$ -level calculated for samples from different storage time ( $P_{st}$ ).  $P$ -level calculated from storage condition ( $P_{sc}$ ) × storage time ( $P_{st}$ ).

\*\* Significant at  $P < 0.01$ .

\*\*\* Significant at  $P < 0.001$ .

NS, not significant.

**Table 2.** Percentage of fatty acids, total saturated fatty acids, total unsaturated fatty acids and unsaturated/saturated fatty acids ratio of lipid fractions extracted from hazelnuts during storage. IS: in-shell hazelnuts; S: shelled hazelnuts (kernel); Refrigerated: stored at 4 °C and 55% RH; Under N<sub>2</sub>: refrigerated under nitrogen (4 °C, 55% RH - 1% O<sub>2</sub>, 99% N<sub>2</sub>); Ambient temp.: stored at ambient temperature (70% RH).

		Beginning	8 <sup>th</sup> month	12 <sup>th</sup> month	$P_{sc}^{\dagger}$	$P_{st}$	$P_{sc} \times P_{st}$
C16:0 (palmitic)							
S	Refrigerated	5.74 ± 0.07	5.85 ± 0.17	5.76 ± 0.24			
S	Under N <sub>2</sub>	5.74 ± 0.07 <sup>B</sup>		4.96 ± 0.24 <sup>A</sup>			
IS	Ambient temp.	5.74 ± 0.07	5.78 ± 0.11 <sup>A</sup>	5.35 ± 0.71			
C18:0 (stearic)							
S	Refrigerated	1.85 ± 0.46	2.40 ± 0.96	2.61 ± 0.69			
S	Under N <sub>2</sub>	1.85 ± 0.46 <sup>A</sup>		3.25 ± 0.28 <sup>B</sup>			
IS	Ambient temp.	1.85 ± 0.46 <sup>A</sup>	2.60 ± 0.33 <sup>B</sup>	3.12 ± 0.29 <sup>B</sup>			
C 18:1 (oleic)							
S	Refrigerated	85.70 ± 2.53	85.63 ± 1.34	87.41 ± 0.96			
S	Under N <sub>2</sub>	85.70 ± 2.53		85.82 ± 0.96			
IS	Ambient temp.	85.70 ± 2.53	85.01 ± 0.79	84.60 ± 1.90			
C 18:2 (linoleic)							
S	Refrigerated	6.16 ± 2.03	5.78 ± 0.51	3.76 ± 1.54			
S	Under N <sub>2</sub>	6.16 ± 2.03		4.95 ± 1.15			
IS	Ambient temp.	6.16 ± 2.03	6.26 ± 0.49	6.87 ± 0.91			
C 18:3 (linolenic)							
S	Refrigerated	0.13 ± 0.01	0.11 ± 0.00	0.16 ± 0.05			
S	Under N <sub>2</sub>	0.13 ± 0.01		0.11 ± 0.02			
IS	Ambient temp.	0.13 ± 0.01 <sup>B</sup>	0.11 ± 0.01 <sup>B</sup>	0.00 ± 0.00 <sup>A</sup>			
Total saturated					NS	**	NS
S	Refrigerated	7.69 ± 0.44	8.30 ± 0.78	8.39 ± 0.44			
S	Under N <sub>2</sub>	7.69 ± 0.44 <sup>A</sup>		8.37 ± 0.24 <sup>B</sup>			
IS	Ambient temp.	7.69 ± 0.44	8.42 ± 0.22	8.51 ± 1.00			
Total unsaturated					NS	*	NS
S	Refrigerated	92.30 ± 0.44	91.69 ± 0.78	91.64 ± 0.47			
S	Under N <sub>2</sub>	92.30 ± 0.44 <sup>B</sup>		91.61 ± 0.23 <sup>A</sup>			
IS	Ambient temp.	92.30 ± 0.44	91.59 ± 0.24	91.81 ± 0.95			
Unsat./Sat.					NS	**	NS
S	Refrigerated	12.03 ± 0.73	11.12 ± 1.07	10.95 ± 0.65			
S	Under N <sub>2</sub>	12.03 ± 0.73 <sup>B</sup>		10.95 ± 0.34 <sup>A</sup>			
IS	Ambient temp.	12.03 ± 0.73	10.88 ± 0.31	10.91 ± 1.36			

Data were expressed as mean ± SD (n = 4). Values in the column with different lowercase letters were significantly different at  $P < 0.05$ . Values in the row with different capital letters were significantly different at  $P < 0.05$ .

<sup>†</sup>Probabilities of the effects:  $P$ -level calculated for samples from different storage condition ( $P_{sc}$ ).  $P$ -level calculated for samples from different storage time ( $P_{st}$ ).  $P$ -level calculated from storage condition ( $P_{sc}$ ) × storage time ( $P_{st}$ ).

\*\* Significant at  $P < 0.01$ .

\*\*\* Significant at  $P < 0.001$ .

NS, not significant.

**Table 3.** Rupture force, slope at the first fracture point ( $Slope_{F1}$ ), and rupture energy of the hazelnuts during storage. IS: in-shell hazelnuts; S: shelled hazelnuts (kernel); Refrigerated: stored at 4 °C and 55% RH; Under N<sub>2</sub>: refrigerated under nitrogen (4 °C, 55% RH - 1% O<sub>2</sub>, 99% N<sub>2</sub>); Amb. temp.: stored at ambient temperature (70% RH).

		Beginning	8 <sup>th</sup> month	12 <sup>th</sup> month	$P_{sc}^{\dagger}$	$P_{st}$	$P_{sc} \times P_{st}$
Rupture force (N)					NS	***	NS
S	Refrigerated	91.83 ± 20.91 <sup>A</sup>	105.39 ± 22.19 <sup>B</sup>	99.17 ± 14.83 <sup>AB</sup>			
S	Under N <sub>2</sub>	91.83 ± 20.91		95.88 ± 17.64			
IS	Amb. temp.	91.83 ± 20.91 <sup>A</sup>	97.16 ± 20.29 <sup>AB</sup>	106.39 ± 21.53 <sup>B</sup>			
$Slope_{F1}$ (N m <sup>-1</sup> )					*	***	NS
S	Refrigerated	26063.4 ± 4033.9 <sup>A</sup>	30125.2 ± 6229.1 <sup>B</sup>	33637.3 ± 7082.3 <sup>abC</sup>			
S	Under N <sub>2</sub>	26063.4 ± 4033.9 <sup>A</sup>		36170.7 ± 9248.3 <sup>bB</sup>			
IS	Amb. temp.	26063.4 ± 4033.9 <sup>A</sup>	29420.9 ± 6117.2 <sup>B</sup>	31236.3 ± 7232. <sup>aB</sup>			
Rupture energy (mJ)					NS	NS	**
S	Refrigerated	153.95 ± 68.74 <sup>AB</sup>	181.33 ± 72.06 <sup>B</sup>	137.26 ± 41.70 <sup>aA</sup>			
S	Under N <sub>2</sub>	153.95 ± 68.74 <sup>A</sup>		121.69 ± 51.64 <sup>aB</sup>			
IS	Amb. temp.	153.95 ± 68.74	149.60 ± 59.22	177.14 ± 64.41 <sup>b</sup>			

Data were expressed as mean ± SD (n = 10). Values in the column with different lowercase letters were significantly different at  $P < 0.05$ . Values in the row with different capital letters were significantly different at  $P < 0.05$ .

<sup>†</sup>Probabilities of the effects:  $P$ -level calculated for samples from different storage condition ( $P_{sc}$ ).  $P$ -level calculated for samples from different storage time ( $P_{st}$ ).  $P$ -level calculated from storage condition ( $P_{sc}$ ) × storage time ( $P_{st}$ ).

\* Significant at  $P < 0.05$ .

\*\* Significant at  $P < 0.01$ .


\*\*\* Significant at  $P < 0.001$ .

NS, not significant.

## Figure captions

### Figure 1

Typical force-deformation curve for compressed hazelnut.  $F_1$ : first fracture point (N);  $S_{F1}$ : slope of the line between starting point and the first fracture point (N m<sup>-1</sup>);  $W_1$ : area under the curve between starting point and the first fracture point (mJ).



**Figure 1.**