

## Ovine TRIM5 $\alpha$ Can Restrict Visna/Maedi Virus

P. Jáuregui, H. Crespo, I. Glaria, L. Luján, A. Contreras, S. Rosati, D. de Andrés, B. Amorena, G. J. Towers and R. Reina  
*J. Virol.* 2012, 86(17):9504. DOI: 10.1128/JVI.00440-12.  
Published Ahead of Print 13 June 2012.

---

Updated information and services can be found at:  
<http://jvi.asm.org/content/86/17/9504>

---

	<i>These include:</i>
<b>REFERENCES</b>	This article cites 42 articles, 22 of which can be accessed free at: <a href="http://jvi.asm.org/content/86/17/9504#ref-list-1">http://jvi.asm.org/content/86/17/9504#ref-list-1</a>
<b>CONTENT ALERTS</b>	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), <a href="#">more»</a>

---

---

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>  
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

---

# Ovine TRIM5 $\alpha$ Can Restrict Visna/Maedi Virus

P. Jáuregui,<sup>a</sup> H. Crespo,<sup>a</sup> I. Glaria,<sup>a</sup> L. Luján,<sup>b</sup> A. Contreras,<sup>c</sup> S. Rosati,<sup>d</sup> D. de Andrés,<sup>a\*</sup> B. Amorena,<sup>a</sup> G. J. Towers,<sup>e</sup> and R. Reina<sup>a</sup>

Instituto de Agrobiotecnología, CSIC-UPNA-Gobierno de Navarra, Mutilva Baja, Navarra, Spain<sup>a</sup>; Facultad de Veterinaria, Universidad de Zaragoza, Zaragoza, Spain<sup>b</sup>; Departamento de Epidemiología y Enfermedades Infecciosas, Facultad de Veterinaria, Universidad de Murcia, Murcia, Spain<sup>c</sup>; Dipartimento di Produzione Animali, Epidemiologia ed Ecologia, Università degli Studi di Torino, Turin, Italy<sup>d</sup>; and MRC Centre for Medical Molecular Virology, Infection and Immunity, University College London, London, United Kingdom<sup>e</sup>

**The restrictive properties of tripartite motif-containing 5 alpha (TRIM5 $\alpha$ ) from small ruminant species have not been explored. Here, we identify highly similar TRIM5 $\alpha$  sequences in sheep and goats. Cells transduced with ovine TRIM5 $\alpha$  effectively restricted the lentivirus visna/maedi virus DNA synthesis. Proteasome inhibition in cells transduced with ovine TRIM5 $\alpha$  restored restricted viral DNA synthesis, suggesting a conserved mechanism of restriction. Identification of TRIM5 $\alpha$  active molecular species may open new prophylactic strategies against lentiviral infections.**

Small ruminant lentiviruses (SRLV), including visna/maedi virus (VMV) and caprine encephalitis virus (CAEV), are widespread in sheep and goats, causing a slow progressive disease. Since neither treatment nor efficient vaccines are available, infection is commonly controlled by early diagnosis and culling (23). Recently, the study of host cell restriction factors interfering with the retroviral life cycle, such as the tripartite motif-containing 5 (TRIM5) protein, has gained interest (10, 37). TRIM5 family members bear a RING–B-box–coiled-coil structure consisting of an N-terminal RING domain (with E3 ubiquitin ligase activity), a B-box domain, and a coiled-coil domain (19). The TRIM5 $\alpha$  isoform, which is active against retroviruses, contains a C-terminal PRYSPRY domain that binds retroviral capsid CA (12, 20, 35). This interaction, involving amino acid 332 of TRIM5 $\alpha$  in humans (15) and 334 in monkeys, may explain the high relative rates of nonsynonymous changes of the primate TRIM5 $\alpha$  gene (13). TRIM5 $\alpha$  has been described in primates and several mammals (3, 6, 30, 33, 41) but not in sheep or goats, both of which are infected by SRLV, their own lentivirus. This study aimed to identify and characterize the ovine and caprine TRIM5 $\alpha$  proteins and explore the possible restrictive role of ovine TRIM5 $\alpha$  on VMV infection.

First, we cloned and sequenced ovine and caprine TRIM5 $\alpha$  cDNA sequences. For this, total RNA from ovine skin fibroblasts (SF), bronchoalveolar lavage (BAL) fluid, or lung tissue obtained from domestic sheep of the Assaf ( $n = 3$ ), Churra ( $n = 2$ ), and Rasa Aragonesa ( $n = 4$ ) breeds was purified using TRIzol (Invitrogen) passed through RNeasy minikit columns (Qiagen), before being reverse transcribed with SuperScript II (Invitrogen) using an oligo(dT) primer according to the manufacturer's instructions. To clone the caprine counterpart, cDNA from peripheral blood mononuclear cells (PBMC) from goats of the Roccaverano ( $n = 1$ ) and Murciano-Granadina ( $n = 2$ ) breeds was used. These cDNAs were employed as the PCR template using Phusion high-fidelity DNA polymerase (Finnzymes) with the forward primer TrimEXNFW (5'-TGCA CCTCGAGATGGCTTCAGGAATCCTG-3', XhoI site underlined) and the reverse primer PJ2 (5'-GATCCGGGCCCTCAAC AGCTTGGTGAG-3', ApaI site underlined) following standard thermal profiles. Amplified products were cloned into the TOPO Blunt vector (Invitrogen) as a shuttle/sequencing vector, yielding

a total of 12 ovine and 5 caprine independent sequences. Four ovine sequences were obtained at least twice and were aligned with previously described TRIM5 $\alpha$  sequences (ClustalX and PHYLIP: Phylogeny Inference Package version 3.5c), revealing a conserved structure across species. Analysis of six clones from SF of one Rasa Aragonesa sheep revealed the presence of only two TRIM5 $\alpha$  amino acid sequences (named Ov1 and Ov2), suggesting that these sequences are encoded by a single heterozygous gene. The sequences differed only at a single residue (39) of the PRYSPRY-domain V1 region. Greater levels of amino acid diversity were found in additional sheep and goat sequences (Fig. 1). To examine sequence diversity, phylogenetic trees were produced by the neighbor-joining method with Kimura's correction using 1,000 bootstrap confidence limits. Results with over 950 bootstraps were considered highly likely. As expected, ovine and caprine sequences were closely related, followed by bovine sequences (Table 1), forming a nonprimate TRIM5 $\alpha$  cluster (Fig. 2). Comparison of these sequences revealed greater variation between caprine and ovine TRIM5 $\alpha$  proteins than between ovine sequences (Table 1; Fig. 1A), with the PRYSPRY being the most variable domain. Such variation was higher than expected given that sheep and goats diverged 6 million years ago (16), whereas humans and chimpanzees, which encode more highly related TRIM5 $\alpha$  sequences, diverged 7 million years ago (5). The close relatedness between sheep and goats is consistent with the ability of sheep (VMV) and goat (CAEV) lentiviruses to infect both ruminant species (8, 32). The high variability of both PRYSPRY (6, 34; this work) and CA of SRLV (7, 26) may account for the evolution of both virus and host, involving TRIM5 $\alpha$  and CA interactions, as described for primate lentiviruses (11, 28, 34, 38). Natural selection in ovine and caprine sequences was determined by estimating  $\omega$  (ratio of the rate of nonsynonymous substitutions, dN, to synonymous substitutions,

Received 24 February 2012 Accepted 1 June 2012

Published ahead of print 13 June 2012


Address correspondence to R. Reina, ramses.reina@unavarra.es.

\* Present address: D. de Andrés, Cátedra de Recursos Cinegéticos y Piscícolas, Universidad de Córdoba, Córdoba, Spain.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JVI.00440-12

A



B




FIG 1 TRIM5 $\alpha$  amino acid sequences from ruminants. (A) Alignment of TRIM5 $\alpha$  amino acid sequences from ovine (Ov1 to Ov4), caprine, and bovine species. Ov1 and Ov2 were amplified from skin fibroblasts of one seronegative Rasa Aragonesa breed sheep. Ov3 and Ov4 were amplified from bronchoalveolar lavage fluid cDNA from a seropositive Rasa Aragonesa sheep. RING, B-box-2, coiled-coil, and PRYSPRY domains are shown, and the variable segments (V1, V2, and V3) of this domain are outlined. Amino acid differences between ovine sequences (Ov1 to Ov4), including those corresponding to V1 residue 339 among others, are highlighted in gray. Accession numbers (GenBank) are as follows: Ov1 to Ov4, JN835300 to JN835303; caprine, JQ582845; bovine, DQ380509. (B) Structure of the ovine TRIM5: full-length isoform (top) and an alternatively spliced form (bottom).

TABLE 1 Sequence identity matrix<sup>a</sup>

Sequence type	% Similarity with sequence type									
	Ov2		Ov3		Ov4		Caprine		Bovine	
	T5	PRYSPRY	T5	PRYSPRY	T5	PRYSPRY	T5	PRYSPRY	T5	PRYSPRY
Ov1	99	99	98	98	99	99	91	89	84	76
Ov2			98	97	99	98	91	89	83	75
Ov3					99	99	92	90	84	77
Ov4							92	90	84	77
Caprine									84	75

<sup>a</sup> Sequence identity matrix for the complete coding sequence (CDS) of the TRIM5 $\alpha$  protein (T5) and its PRYSPRY domain, using the ovine (Ov1, Ov2, Ov3, and Ov4) and caprine amino acid sequences obtained and a bovine sequence (DQ380509) available at GenBank.


dS), using three methods: single-likelihood ancestor counting (SLAC), fixed effects likelihood (FEL), and random effects likelihood (REL) implemented in the Datamonkey webserver (21). The existence of strong positive selection observed when comparing other species (6, 17, 29, 34) was not observed in sheep-versus-goat comparisons ( $P \geq 0.064$ ), consistent with viral transmission between these species.

Additional RNA species (6 from a total of 3 sheep) were identified that had a stop codon at residue 347 compared to the full ovine protein TRIM5 $\alpha$  (OvT5 $\alpha$ ), resulting in a reduced number of exons and elongated noncoding sequences at the 3' end, a structure similar to that of human TRIM5 $\gamma$  (2) (Fig. 1B). No splicing donor/acceptor consensus sequences were found in the shortened TRIM5 $\alpha$  sequences under study, suggesting that truncated proteins are indeed splice variants of TRIM5 and that splicing of these forms is conserved between primates and ruminants. Numerous isoforms that are shorter than the antiviral TRIM5 $\alpha$  exist in humans (2) and macaques (4). Some of these lack the PRYSPRY domain and are therefore unrestricted. Like human TRIM5 $\gamma$  (36), the short ovine isoform is likely to act as a dominant negative through lack of a viral binding PRYSPRY domain (2).

To characterize restriction by OvT5, sequences Ov1, Ov2 (both from SF of a seronegative Rasa Aragonesa sheep from a seropositive flock), and Ov4 (obtained from BAL fluid cells of a Rasa Aragonesa seropositive sheep affected with pneumonia) were cloned into the gammaretroviral expression vector pCNCR-HA, using XhoI/ApaI restriction sites. The vector contains the LTR of Moloney murine leukemia virus (MLV), driving expression of an N-terminal hemagglutinin (HA)-tagged OvT5 $\alpha$  protein and the gene for the red fluorescent protein (RFP). The resulting vector (pCTCR-HAOvT5) was packaged into vesicular stomatitis virus G envelope protein (VSV-G)-pseudotyped MLV cores by cotransfection of 293T cells as described previously (3). Culture supernatants containing MLV virions encoding OvT5 were used to transduce the *Mus dunni* tail fibroblast (MDTF) cell line, and HA-tagged OvT5 stably expressing cells were obtained. Cells transduced with empty pCNCR-HA were used as controls. Single cell clones were isolated by limiting dilution, identified by red fluorescence microscopy, expanded, and checked for expression of TRIM5 $\alpha$  proteins by Western blotting (WB) using anti-HA antibodies and quantitative reverse transcriptase PCR (RT-PCR) with forward (qPCR3T5Fw: 5'-TTCCTAGACTATGAGGCTTGCTCTGT-3') and reverse (qPCR3T5Rv: 5'-TTCTGAGGAAAGGAACATGAAGAGA-3') primers, designed within the PRYSPRY region.

$\beta$ -Actin RT-PCR allowed relative quantification using the primers described (24).

Transduced MDTF cells expressed OvT5-HA according to the results of WB (Fig. 3B) and RT-PCR (not shown). These OvT5-expressing clones were subjected to infection by VMV strain Ev1 (27) to study restriction. The dose was determined by titrating Ev1 by inoculation of 10-fold serial dilutions onto ovine SF and visualization of the cytopathic effect by microscopy after 7 days. Titers, calculated by the Reed-Muench method (22), were expressed as 50% tissue culture infectious doses (TCID<sub>50</sub>) per ml. Cells were infected at an apparent multiplicity of infection (MOI) of 0.2, and 16 h later MDTF total DNA was purified using a QIAamp DNA minikit (Qiagen). TaqMan quantitative PCR (qPCR) was used to measure viral DNA synthesis using a plasmid standard curve as described previously (9). Strain Ev1 entered and was reverse transcribed in the MDTF cells. Viral DNA was detected by qPCR (mean copy number/100 ng DNA when infecting at 0.2 TCID<sub>50</sub>/cell,  $9.9 \times 10^2$ ), and viral transcripts were produced according to RT-PCR (not shown). Thus, heterologous MDTF cells were suitable for assessing OvT5-mediated restriction, even though infection was not productive since supernatants had no RT activity up to day 20 postinoculation (not shown). Measurement of viral DNA synthesis indicated that MDTFs transduced with OvT5 $\alpha$  were less permissive to reverse transcription than control MDTFs ( $P < 0.05$ ). We conclude that OvT5 $\alpha$  was able to significantly restrict VMV infectivity. TRIM5 $\alpha$  Ov1 and Ov2 were able to restrict Ev1 whereas TRIM5 $\alpha$  Ov4 was not, despite strong expression detected by immunoblotting (Fig. 3A and B). In the TRIM5 $\alpha$  protein of humans and simians, a single amino acid substitution at position 332 or 334 abrogates TRIM5 $\alpha$ -mediated restriction of particular viruses due to its essential role in viral recognition (13). Specifically, any non-positively charged amino acid at that position, which belongs to a "patch" of positively selected positions, improves CA binding (15). Surprisingly, Ov1 and Ov2 had either a positively charged amino acid (K) or a negative residue (E) at this position (Fig. 1A), but both showed a similar restriction of Ev1 in MDTF cells. Due to the poor alignment of this highly variable region, it is difficult to be sure that this amino acid is analogous to primate TRIM5 position 339, but it is certainly very close and putatively present on the surface of the protein in the highly variable loop that is most important for determining TRIM5 $\alpha$  specificity. Recent studies highlight the importance of domains other than PRYSPRY (18). Significantly, Ov4 had differences in the RING and PRYSPRY domains compared with Ov1 and Ov2




**FIG 2** Phylogenetic tree of TRIM nucleotide sequences of different species. The tree shows that all the sheep and goat TRIM5 sequences described in this work are TRIM5 $\alpha$  orthologues. GenBank accession numbers shown are for sequences from the following sources: JN835300 and JN835301, Rasa Aragonesa skin fibroblasts; JN835302 and JN835303, Rasa Aragonesa bronchoalveolar lavage fluid; JN835304, Rasa Aragonesa lung tissue; JN835305, lung tissue of two Rasa Aragonesa and two Assaf sheep; JN835306, lung tissue of two Assaf and one Churra sheep; JN835307, Assaf lung tissue; JN835308, Rasa Aragonesa lung tissue; JN835309, Rasa Aragonesa lung tissue; JN835310, Churra lung tissue; JN835311, lung tissue of one Assaf and one Churra sheep; JQ582845, peripheral blood mononuclear cells of a Rocaverano goat; JQ582846 to JQ582848, peripheral blood mononuclear cells of a Murciano-Granadina goat; JQ582849, peripheral blood mononuclear cells of a Murciano-Granadina goat.

and may have lost restrictive activity against VMV. Indeed, Ov4 was obtained from an infected sheep, consistent with a permissive TRIM5 genotype.


In addition, we tested the restrictive role of OvT5 against VSV-G-pseudotyped HIV-2 viral vectors encoding green fluorescent protein (GFP), prepared by Fugene-6 transfection of 293T cells as

described previously (10). We infected OvT5-MDTF cells and quantified infection at 48 h by measuring GFP expression using flow cytometry (BD FACScalibur). MDTFs showed decreased levels of HIV-2 infection when expressing Ov2 (Fig. 3A), strongly suggesting a role for OvT5 in protecting sheep from HIV-2 infection.



**FIG 3** Ovine TRIM5 $\alpha$  (OvT5 $\alpha$ )-mediated restriction. (A) MDTF cells expressing OvT5 $\alpha$  were infected with GFP-expressing HIV-2 VSV-G pseudotyped lentiviral vector or VMV strain Ev1. Data represent viral DNA copy numbers (mean values  $\pm$  standard error) per 100 ng of total DNA obtained by qPCR using Ev1-specific primers/probe (9) or infectious units per ml in the case of HIV-2. Cells transduced with pCNCR-HA were used as negative controls. Three independent experiments were performed. (B) Western blot illustrating the expression of different HA-OvT5 sequences in MDTF cells (transduced pCTCR-HA-OvT5 or empty vector CNCR), using anti-HA-tag antibodies.

Since the proteasome has been shown to be involved in TRIM5 $\alpha$  restriction in other species (1, 25, 40), we inhibited the proteasome and examined OvT5-mediated restriction of VMV. MDTF cells expressing TRIM5 $\alpha$  Ov2 or empty vector were treated with prewarmed (37°C) proteasome inhibitor MG132 (Sigma-Aldrich) at a final concentration of 25  $\mu$ M for 1 h before infection with Ev1 at an MOI of 0.2. After 16 h of infection, viral DNA was quantified in duplicate by qPCR as described above, and the experiment was repeated three times. The results indicated that viral DNA was significantly increased ( $P < 0.01$ ) in proteasome-inhib-



**FIG 4** Effect of proteasome inhibition on VMV DNA levels in ovine OvT5 $\alpha$ -expressing cells. MDTF cells expressing restrictive OvT5 $\alpha$  Ov2 or transduced with an empty CNCR-HA vector were treated for 1 h with proteasome inhibitor MG132 (untreated cells were used as a control) before infection with VMV strain Ev1, and viral DNA was measured 16 h after infection by qPCR using Ev1-specific primers/probe. Data represent viral DNA copy numbers (mean  $\pm$  standard error) per 100 ng of total DNA. Three independent experiments were performed.

ited MDTF cells expressing a restrictive OvT5 compared to untreated cells. Either treated or untreated cells transduced with an empty vector showed no viral restriction, having similar levels of viral DNA (Fig. 4). The involvement of the proteasome in OvT5 restriction is in line with findings on other lentiviruses (1, 40).

TRIM5 $\alpha$  has been characterized mostly in the context of restricting heterologous viruses, but as shown here homologous virus restriction may also take place upon overexpression of TRIM5 $\alpha$  (14, 42). This suggests that differences in TRIM5 expression levels as well as intrinsic antiviral specificity may account for differences in permissiveness to infection between individuals (31, 42). Importantly, our study suggests functional differences between polymorphic ovine TRIM5 $\alpha$  variants to restrict both heterologous (HIV-2) and homologous (VMV) viruses. A better understanding of these differences could eventually be used to design SRLV control strategies such as identification and selective breeding of animals that are less permissive to infection, thus avoiding culling and helping to reduce viral load and therefore disease development.

**Nucleotide sequence accession numbers.** The nucleotide sequences obtained in this work were deposited in GenBank under accession numbers JN835300 to JN835311 (ovine) and JQ582845 to JQ582849 (caprine).

**ACKNOWLEDGMENTS**

This work was supported by grants from CICYT (no. AGL2010-22341-C04-01), the Government of Navarra (no. IIQ14064.RI1), the UK Medical Research Centre, and the UK National Institute of Health

Research UCL/UCLH Comprehensive Biomedical Research Centre and fellowships from the Spanish Ministry of Science and Innovation (P.J.) and the Wellcome Trust (G.J.T.).

The CMV-intron MLV packaging plasmid was a gift from F. L Cosset, and HIV-2 vectors were from A. Lever.

## REFERENCES

- Anderson JL, et al. 2006. Proteasome inhibition reveals that a functional preintegration complex intermediate can be generated during restriction by diverse TRIM5 proteins. *J. Virol.* **80**:9754–9760.
- Battivelli E, et al. 2011. Modulation of TRIM5 $\alpha$  activity in human cells by alternatively spliced TRIM5 isoforms. *J. Virol.* **85**:7828–7835.
- Besnier C, Takeuchi Y, Towers G. 2002. Restriction of lentivirus in monkeys. *Proc. Natl. Acad. Sci. U. S. A.* **99**:11920–11925.
- Brennan G, Kozyrev Y, Kodama T, Hu SL. 2007. Novel TRIM5 isoforms expressed by *Macaca nemestrina*. *J. Virol.* **81**:12210–12217.
- Chen FC, Li WH. 2001. Genomic divergences between humans and other hominoids and the effective population size of the common ancestor of humans and chimpanzees. *Am. J. Hum. Genet.* **68**:444–456.
- Fletcher AJ, Hue S, Schaller T, Pillay D, Towers GJ. 2010. Hare TRIM5 $\alpha$  restricts divergent retroviruses and exhibits significant sequence variation from closely related lagomorpha TRIM5 genes. *J. Virol.* **84**:12463–12468.
- Giammarioli M, et al. 2011. Phylogenetic analysis of small ruminant lentivirus (SRLV) in Italian flocks reveals the existence of novel genetic subtypes. *Virus Genes* **43**:380–384.
- Glaria I, et al. 2009. Phylogenetic analysis of SRLV sequences from an arthritic sheep outbreak demonstrates the introduction of CAEV-like viruses among Spanish sheep. *Vet. Microbiol.* **138**:156–162.
- Gonzalez B, et al. 2005. Mucosal immunization of sheep with a maedi-visna virus (MVV) env DNA vaccine protects against early MVV productive infection. *Vaccine* **23**:4342–4352.
- Hatzioannou T, Cowan S, Goff SP, Bieniasz PD, Towers GJ. 2003. Restriction of multiple divergent retroviruses by Lv1 and Ref1. *EMBO J.* **22**:385–394.
- Ikeda Y, Ylinen LM, Kahar-Bador M, Towers GJ. 2004. Influence of gag on human immunodeficiency virus type 1 species-specific tropism. *J. Virol.* **78**:11816–11822.
- James LC, Keeble AH, Khan Z, Rhodes DA, Trowsdale J. 2007. Structural basis for PRYSPRY-mediated tripartite motif (TRIM) protein function. *Proc. Natl. Acad. Sci. U. S. A.* **104**:6200–6205.
- Johnson WE, Sawyer SL. 2009. Molecular evolution of the antiretroviral TRIM5 gene. *Immunogenetics* **61**:163–176.
- Kaummann P, Haggmann I, Dittmar MT. 2006. Human TRIM5 $\alpha$  mediated restriction of different HIV-1 subtypes and Lv2 sensitive and insensitive HIV-2 variants. *Retrovirology* **3**:79.
- Li Y, Li X, Stremlau M, Lee M, Sodroski J. 2006. Removal of arginine 332 allows human TRIM5 $\alpha$  to bind human immunodeficiency virus capsids and to restrict infection. *J. Virol.* **80**:6738–6744.
- Maddox JF. 2005. A presentation of the differences between the sheep and goat genetic maps. *Genet. Sel. Evol.* **37**(Suppl 1):S1–S10.
- Newman RM, et al. 2006. Balancing selection and the evolution of functional polymorphism in Old World monkey TRIM5 $\alpha$ . *Proc. Natl. Acad. Sci. U. S. A.* **103**:19134–19139.
- Ohmine S, et al. 2011. The antiviral spectra of TRIM5 $\alpha$  orthologues and human TRIM family proteins against lentiviral production. *PLoS One* **6**:e16121. doi:10.1371/journal.pone.0016121.
- Ozato K, Shin DM, Chang TH, and Morse HC, 3rd. 2008. TRIM family proteins and their emerging roles in innate immunity. *Nat. Rev. Immunol.* **8**:849–860.
- Perez-Caballero D, Hatzioannou T, Yang A, Cowan S, Bieniasz PD. 2005. Human tripartite motif 5 $\alpha$  domains responsible for retrovirus restriction activity and specificity. *J. Virol.* **79**:8969–8978.
- Pond SK, Muse SV. 2005. Site-to-site variation of synonymous substitution rates. *Mol. Biol. Evol.* **22**:2375–2385.
- Reed LJ, Muench H. 1938. A simple method of estimating fifty per cent endpoints. *Am. J. Hyg.* **27**:493–497.
- Reina R, et al. 2009. Prevention strategies against small ruminant lentiviruses: an update. *Vet. J.* **182**:31–37.
- Reina R, et al. 2007. Association of CD80 and CD86 expression levels with disease status of visna/maedi virus infected sheep. *Viral Immunol.* **20**:609–622.
- Rold CJ, Aiken C. 2008. Proteasomal degradation of TRIM5 $\alpha$  during retrovirus restriction. *PLoS Pathog.* **4**:e1000074. doi:10.1371/journal.ppat.1000074.
- Rosati S, et al. 2004. Antigenic variability of ovine lentivirus isolated in Italy. *Vet. Res. Commun.* **28**(Suppl 1):319–322.
- Sargan DR, et al. 1991. Nucleotide sequence of EV1, a British isolate of maedi-visna virus. *J. Gen. Virol.* **72**(Pt 8):1893–1903.
- Sawyer SL, Emerman M, Malik HS. 2007. Discordant evolution of the adjacent antiretroviral genes TRIM22 and TRIM5 in mammals. *PLoS Pathog.* **3**:e197. doi:10.1371/journal.ppat.0030197.
- Sawyer SL, Wu LI, Emerman M, Malik HS. 2005. Positive selection of primate TRIM5 $\alpha$  identifies a critical species-specific retroviral restriction domain. *Proc. Natl. Acad. Sci. U. S. A.* **102**:2832–2837.
- Schaller T, Hue S, Towers GJ. 2007. An active TRIM5 protein in rabbits indicates a common antiviral ancestor for mammalian TRIM5 proteins. *J. Virol.* **81**:11713–11721.
- Sewram S, et al. 2009. Human TRIM5 $\alpha$  expression levels and reduced susceptibility to HIV-1 infection. *J. Infect. Dis.* **199**:1657–1663.
- Shah C, et al. 2004. Phylogenetic analysis and reclassification of caprine and ovine lentiviruses based on 104 new isolates: evidence for regular sheep-to-goat transmission and worldwide propagation through livestock trade. *Virology* **319**:12–26.
- Si Z, et al. 2006. Evolution of a cytoplasmic tripartite motif (TRIM) protein in cows that restricts retroviral infection. *Proc. Natl. Acad. Sci. U. S. A.* **103**:7454–7459.
- Soares EA, et al. 2010. Evolution of TRIM5 $\alpha$  B30.2 (SPRY) domain in New World primates. *Infect. Genet. Evol.* **10**:246–253.
- Song B, et al. 2005. The B30.2(SPRY) domain of the retroviral restriction factor TRIM5 $\alpha$  exhibits lineage-specific length and sequence variation in primates. *J. Virol.* **79**:6111–6121.
- Stremlau M, et al. 2004. The cytoplasmic body component TRIM5 $\alpha$  restricts HIV-1 infection in Old World monkeys. *Nature* **427**:848–853.
- Stremlau M, Perron M, Welikala S, Sodroski J. 2005. Species-specific variation in the B30.2(SPRY) domain of TRIM5 $\alpha$  determines the potency of human immunodeficiency virus restriction. *J. Virol.* **79**:3139–3145.
- Towers GJ. 2007. The control of viral infection by tripartite motif proteins and cyclophilin A. *Retrovirology* **4**:40.
- Wilson SJ, et al. 2008. Rhesus macaque TRIM5 alleles have divergent antiretroviral specificities. *J. Virol.* **82**:7243–7247.
- Wu X, Anderson JL, Campbell EM, Joseph AM, Hope TJ. 2006. Proteasome inhibitors uncouple rhesus TRIM5 $\alpha$  restriction of HIV-1 reverse transcription and infection. *Proc. Natl. Acad. Sci. U. S. A.* **103**:7465–7470.
- Ylinen LM, et al. 2006. Isolation of an active Lv1 gene from cattle indicates that tripartite motif protein-mediated innate immunity to retroviral infection is widespread among mammals. *J. Virol.* **80**:7332–7338.
- Ylinen LM, Keckesova Z, Wilson SJ, Ranasinghe S, Towers GJ. 2005. Differential restriction of human immunodeficiency virus type 2 and simian immunodeficiency virus SIVmac by TRIM5 $\alpha$  alleles. *J. Virol.* **79**:11580–11587.