

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

CCR5-Δ32 genotype does not improve predictive value of IL28B polymorphisms for treatment response in chronic HCV infection.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/147235> since 2017-09-27T17:00:47Z

Published version:

DOI:10.1038/gene.2013.15

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

***[Genes Immun. 2013 Jul-Aug;14(5):286-90. doi: 10.1038/gene.2013.15. Epub
2013 Apr 18.]***

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<http://www.nature.com/gene/journal/v14/n5/full/gene201315a.html>]

**CCR5-Δ32 genotype does not improve predictive value of IL28B
polymorphisms for treatment response in chronic HCV infection**

Vijayaprakash Suppiah^{1,2}, Nicola J Armstrong^{3,4}, Kate S. O'Connor², Thomas Berg^{5,6}, Martin Weltman⁷, Maria Lorena Abate⁸, Ulrich Spengler⁹, Margaret Bassendine¹⁰, Gregory J. Dore^{11,12}, William L. Irving¹³, Elizabeth Powell^{14,15}, Jacob Nattermann⁹, Stephen Riordan¹⁶, Graeme J. Stewart², Jacob George¹, David R. Booth², Golo Ahlenstiel¹, for the International Hepatitis C Genetics Consortium (IHCGC).

¹ Storr Liver Unit, Westmead Millennium Institute, University of Sydney, Sydney, Australia,

² Institute for Immunology and Allergy Research, Westmead Millennium Institute, University of Sydney, Sydney, Australia, ³Cancer Research Program, Garvan Institute for Medical

Research and ⁴School of Mathematics and Statistics, University of New South Wales, Sydney, Australia, ⁵ Medizinische Klinik m.S. Hepatologie und Gastroenterologie, Charité,

Campus Virchow-Klinikum, Universitätsmedizin Berlin, Germany, ⁶Department of

Hepatology, Clinic for Gastroenterology and Rheumatology, University Clinic Leipzig,

Leipzig, Germany, ⁷Department of Gastroenterology and Hepatology, Nepean Hospital,

Sydney, Australia, ⁸Liver Physiopathology Lab, Department of Internal Medicine, University

of Turin, Turin, Italy, ⁹Department of Internal Medicine I, University of Bonn, Bonn,

Germany, ¹⁰Liver Research Group, Institute of Cellular Medicine, Medical School, Newcastle

University, Newcastle upon Tyne, United Kingdom, ¹¹National Centre in HIV Epidemiology

and Clinical Research, University of New South Wales, Sydney, Australia, ¹²St Vincent's

Hospital, Sydney, Australia, ¹³NIHR Biomedical Research Unit in Gastroenterology and the

Liver, University of Nottingham, Nottingham, United Kingdom, ¹⁴Princess Alexandra

Hospital, Department of Gastroenterology and Hepatology, Woolloongabba, ¹⁵The University

of Queensland, School of Medicine, Princess Alexandra Hospital, Woolloongabba,

Queensland, Australia.

CCR5-Δ32 and IL28B polymorphisms in treatment of HCV infection

Short title: IL28B and CCR5-Δ32 in HCV infection

Character count (title): 131 (no spaces)

Word count (abstract): 200

Word count (total): 3783

Address for Correspondence:

Professor Jacob George

Storr Liver Unit, Westmead Hospital

Hawkesbury Road, Westmead, NSW 2145, Australia

Tel.: +61 2 9845 7705

Fax: +61 2 9635 7582

jacob.george@sydney.edu.au

Abbreviations: ALT, alanine aminotransferase; GGT, gamma glutamyltransferase; HCV, hepatitis C virus; IFN, interferon; SVR, sustained virological response.

Abstract

Background: IL28B polymorphisms strongly predict spontaneous and treatment-induced clearance of HCV infection. A recent study proposed a 32-base pair deletion in the CCR5 gene (CCR5-Δ32) interacting with the IL28B polymorphisms to influence spontaneous HCV clearance. The aim of this study was to clarify the role of CCR5-Δ32 in treatment-induced clearance of chronic hepatitis C (CHC).

Methods: A cross-sectional cohort of 758 Caucasian patients with CHC genotype 1 (341 responders and 417 non-responders) who had received standard of care dual therapy with IFN-α and ribavirin (RBV) were genotyped for the CCR5-Δ32 and IL28B polymorphisms to examine their interaction with respect to treatment response.

Results: CCR5-Δ32 did not influence treatment-induced recovery to IFN-α/RBV in CHC and did not improve prediction of SVR in the context of the IL28B polymorphisms in a multivariate model. CCR5-Δ32 homozygotes were significantly more frequent in those with CHC than healthy controls in the Europeans cohorts (2.9% vs 0.4%, $p < 0.0001$), but not in Australians of European ancestry.

Conclusion: CCR5-Δ32 does not influence treatment response to IFN-α/RBV in the context of IL28B polymorphisms. Although CCR5-Δ32 may affect viral clearance within closely controlled geographic and genetic environments, we found no effect in larger cohorts treated with dual therapy.

Key words: CCR5, IL28B, HCV, SVR

INTRODUCTION

Hepatitis C virus (HCV) is a major cause of blood-borne hepatitis with an estimated worldwide prevalence of 3% ¹. Chronic infection causes hepatic inflammation eventually resulting in fibrosis and cirrhosis, and ultimately liver failure (reviewed in ²). Cytokines and chemokines are key players in this process through a complex network of interactions that regulate innate and adaptive immune responses to viral infection, and recruitment of inflammatory cells to the liver ³. Importantly, lymphocytes infiltrating HCV-infected livers have been shown to express high levels of the CC-chemokines CCL3 (macrophagic inflammatory protein 1α, MIP-1α), CCL4 (macrophagic inflammatory protein 1β, MIP-1β), CCL5 (regulated upon activation, normal T cell expressed and secreted, RANTES) and their receptor, CC-chemokine receptor 5 (CCR5), suggesting a Th1-mediated antiviral response by the host to the virus ⁴.

Several studies have linked HCV proteins to the expression of CCR5 ligands: Soo et al. showed that the expression of full-length HCV led to the induction of both CCR5 mRNA and protein in HeLa, Huh7 and HepG2 cell lines ⁵. An increase in CCL5 levels by CD8+ T cells was reported to be related to the HCV E2 protein ⁶. Finally, chronic HCV infection has been shown to lead to reduced surface expression of CCR5 on peripheral blood T cells ⁷.

CCR5 has been previously identified as the major co-receptor for human immunodeficiency virus type 1 (HIV-1) ⁸. A 32 bp deletion in exon 4 of CCR5 (CCR5-Δ32) leads to truncation and loss of function of the receptor ⁹. Homozygosity for this deletion confers high level (but not complete) resistance to HIV-1 infection, while heterozygotes show delayed progression to AIDS ^{10, 11}. An increased frequency of CCR5-Δ32 homozygosity was observed in HCV infected individuals compared to healthy controls (7.8% vs 1.0%) ¹². In a later Irish study, the authors found that CCR5-Δ32 was significantly associated with increased spontaneous

CCR5-Δ32 and IL28B polymorphisms in treatment of HCV infection

clearance¹³. Recently, Nattermann et al. have shown, in a very well defined patient cohort (n=396) infected with HCV genotype 1 from a single source, that CCR5-Δ32 was associated with a decrease in spontaneous viral clearance¹⁴. Importantly, this effect seemed to be due to an interaction of CCR5-Δ32 with the recently described IL28B *rs12979860* polymorphism, which has also been shown to influence spontaneous recovery from HCV infection^{15, 16}. As IL28B polymorphisms are currently the strongest host genetic markers to predict treatment-induced clearance of HCV infection¹⁷⁻²⁰, we designed this study to examine for a possible interaction between the CCR5-Δ32 mutation and the two most predictive IL28B polymorphisms, *rs8099917* and *rs12979860*, with respect to treatment-induced clearance of HCV infection.

Results

When compared to healthy controls, patients with chronic HCV infection had a significantly increased frequency of CCR5-Δ32 homozygosity (Table 1; $p=0.039$). The association was due to the European cohort, with an increase in CCR5-Δ32 homozygous subjects (2.9% vs 0.4%; $p=0.0014$) and a decrease in heterozygous subjects (11.3% vs 18.0%; $p=0.0020$). Further, the enrichment of CCR5-Δ32 homozygous European Caucasian subjects with CHC was quadruple that of what would have been expected based on the Hardy Weinberg equilibrium ($p < 0.0001$).

For treatment response to standard dual therapy (IFN- α /RBV), there was no significant difference between sustained virological responders (SVR) and non-SVR patients (NSVR) with respect to the frequency of CCR5-Δ32 genotype distribution or allele frequency (Table 2 and Figure 1).

In order to examine if the combination of the CCR5-Δ32 deletion with the IL28B SNPs, *rs12979860* and *rs8099917*, improves prediction of treatment-induced HCV clearance, we used logistic regression modelling as previously described^{14, 21}. We found no evidence for any two-way interactions between either of the two IL28B SNPs individually or in combination with CCR5-Δ32 for NSVR, even though both the IL28B SNPs individually were significantly associated with treatment response (Table 3).

Discussion

The aim of this study was to establish whether there is an interaction between IL28B polymorphisms and CCR5-Δ32 with respect to predicting treatment-induced clearance of chronic HCV infection. Our results clearly show that A) CCR5-Δ32 does not directly influence treatment response of HCV genotype 1 to treatment with IFN-α and RBV and B) there is no interaction between IL28B genotypes and CCR5-Δ32 in this context. This is in line with previous, substantially smaller studies that have not shown a direct effect of CCR5-Δ32 on treatment response to IFN/RBV treatment²¹⁻²⁷. As we have previously shown in a small cohort of patients, the outcome of IFN monotherapy, but not dual therapy, may be affected by CCR5-Δ32²¹. The lack of a CCR5-Δ32 association in that and the current cohort may be due to a RBV effect. The impact of IL28B might not be masked by RBV, suggesting a much stronger role for variants of this gene in treatment outcome. Of note, age, viral load and gender were significantly different between responders and non-responders (data not shown; $P < 0.05$). These clinical parameters are well known to be different between responders and non-responders^{28,29}.

It is well known that a major problem with clinical studies involving CCR5-Δ32 is that patient source does influence the CCR5-Δ32 frequency: CCR5-Δ32 has been mainly described in Caucasian populations and is subject to a strong Northern to Southern European decline in frequency³⁰. Co-exposure to HIV, for which CCR5-Δ32 homozygosity confers a significant degree of resistance, may introduce another possible selection bias as HIV is transmitted via similar routes to HCV. Interestingly enough, CCR5-Δ32 was more common in the chronic HCV patients in Europe, similar to our published data¹². Significant, but opposite, associations of CCR5-Δ32 with spontaneous recovery were identified in two cohorts, one East German the other Irish, both from single source outbreaks and with

geographically and socioeconomically analogous backgrounds^{13, 14}. This suggests that a weak (compared to IL28B) effect of CCR5-Δ32 on spontaneous recovery may be observed in genetically well defined contexts. More importantly, irrespective of these biases, the SVR and NSVR patients were subject to similar pre-selection criteria for therapy and despite this, CCR5-Δ32 did not impact on treatment-induced HCV clearance in our large multicenter cohort, nor was there a significant interaction between IL28B and CCR5-Δ32.

CCR5-Δ32 presents an attractive target for study in terms of HCV pathogenesis: HCV infection induces IFN-α³¹ and one of the interferon induced genes is CCR5³². CCR5 is expressed on Th1 cells and facilitates the migration of T cells primed by antigenic molecules. It has been observed previously that in HCV patients, lymphocytes infiltrating the liver show an increased expression of CCR5⁴. The presence of the CCR5-Δ32 deletion might result in a reduced expression of CCR5 in these patients, impairing their activation and migration to the infected liver. These mechanisms may be the reason why CCR5-Δ32 deletion may be involved in spontaneous clearance and possibly in response to IFN monotherapy²¹. RBV, however, has been postulated to exploit a novel innate mechanism to potentiate the antiviral effects of IFN-α³³ which could potentially also involve a bias favouring the Th1 immune response³⁴. In our multicenter cohort, the combination of RBV with IFN-α clearly overrode any potential effect of the CCR5-Δ32 deletion, whereas IL28B remained a strong predictive marker as previously reported¹⁷⁻²⁰.

In summary, CCR5-Δ32 did not influence treatment response to treatment with IFN-α/RBV and there was no significant interaction with IL28B genotypes. This suggests that CCR5-Δ32 is not relevant for treatment-induced clearance of HCV and that therefore CCR5 expression is likely not essential in this process. New powerful direct acting antivirals for difficult to treat HCV genotype 1 infection are currently being introduced and will likely reduce the impact of

CCR5-Δ32 and IL28B polymorphisms in treatment of HCV infection

host genetics on treatment response³⁵. In view of CCR5 inhibitors currently available for HIV treatment, this is an important observation, as our data suggest that neither genetic nor drug-induced impairment of CCR5 signalling is likely to impair the efficacy of anti-HCV therapy in patients with HIV/HCV co-infection. Indeed, a recent study in HIV/HCV co-infected patients reported no significant changes in viral titres or LFTs during a short term course of a CCR5 inhibitor³⁶.

MATERIALS AND METHODS

SUBJECTS

Ethical approval was obtained from the Human Research Ethics Committees of Sydney West Area Health Service and the University of Sydney. All other sites had ethical approval from their respective ethics committees. Written informed consent was obtained from all participants. Characteristics of the study cohorts have been described elsewhere ¹⁶. Briefly, all treated patients were Caucasian and infected with genotype 1, the majority of these patients received pegylated interferon and ribavirin and had virological response determined 6 months after completion of therapy. The diagnosis of chronic hepatitis C was based on appropriate serology and presence of HCV RNA. Patients received therapy for 48 weeks except if there was less than a 2 log drop in HCV RNA after 12 weeks therapy. Patients were excluded if they were co-infected with HIV or HBV. The control data was obtained from previous studies done on healthy individuals from the same geographical region as the patients in this study ³⁷⁻⁴⁰.

GENOTYPING

The CCR5 Δ32 deletion was genotyped using high resolution melt, as described previously ⁴¹. IL28B data presented in this study have been previously published ¹⁶.

STATISTICAL ANALYSIS

The χ^2 test or Fisher's test where appropriate were used to examine differences in genotype and allele frequencies between SVR versus Non-SVR (NSVR), and chronic hepatitis C (i.e. NSVR plus SVR) versus healthy controls. P values less than 0.05 (two-sided) were regarded as significant. The relationships between CCR5-Δ32 and IL28B SNPs, *rs8099917* and

CCR5-Δ32 and IL28B polymorphisms in treatment of HCV infection

rs12979860, were investigated using logistic regression. Significance of all models was assessed by likelihood ratio tests. Analysis was carried out in R (v2.12).

Acknowledgments: Work on this manuscript was in part supported by research grants from the National Health and Medical Research Council of Australia and by an Australian ARC linkage grant. JG is supported by the Robert W. Storr bequest to the University of Sydney Medical Foundation. We would like to thank all patients for their valuable participation in this study. The IHCGC team includes Monika Michalk, University of Bonn, Germany; Barbara Malik Universitaetsmedizin Berlin, Germany; Tobias Mueller, University Clinic Leipzig, Germany; Patrick McClure and Sherie Smith, University of Nottingham, UK; David Sheridan, Newcastle University, UK; Elizabeth Snape and Vincenzo Fragomeli, Nepean Hospital, Australia; Richard Norris and Dianne How-Chow, St Vincent's Hospital, Australia; Julie R. Jonsson and Helen Barrie, Princess Alexandra Hospital, Australia; Sacha Stelzer-Braid and Shona Fletcher, Prince of Wales Hospital, Australia; Tanya Applegate, Jason Grebely and Gail Matthews, National Centre in HIV Epidemiology and Clinical Research; Mandvi Bharadwaj, Burnet Institute, Australia; Antonina Smedile, University of Turin, Italy. We would also like to thank Reynold Leung for technical assistance.

Conflict of interest: The authors declare that they have no competing financial interests in relation to this work.

References

1. WHO. Hepatitis C. Fact Sheet No. 164. Revised October 2000. <http://www.who.int/mediacentre/factsheets/fs164/en/>. 2000.
2. Hoofnagle JH. Course and outcome of hepatitis C. *Hepatology* 2002; **36**: S21-S29.
3. Lambotin M, Raghuraman S, Stoll-Keller F, Baumert TF, Barth H. A look behind closed doors: interaction of persistent viruses with dendritic cells. *Nat Rev Microbiol* 2010; **8**: 350-360.
4. Shields PL, Morland CM, Salmon M, Qin S, Hubscher SG, Adams DH. Chemokine and chemokine receptor interactions provide a mechanism for selective T cell recruitment to specific liver compartments within hepatitis C-infected liver. *J Immunol* 1999; **163**: 6236-6243.
5. Soo HM, Garzino-Demo A, Hong W, Tan YH, Tan YJ, Goh PY *et al*. Expression of a full-length hepatitis C virus cDNA up-regulates the expression of CC chemokines MCP-1 and RANTES. *Virology* 2002; **303**: 253-277.
6. Nattermann J, Nischalke HD, Feldmann G, Ahlenstiel G, Sauerbruch T, Spengler U. Binding of HCV E2 to CD81 induces RANTES secretion and internalization of CC chemokine receptor 5. *J Viral Hepat* 2004; **11**: 519-526.
7. Lichterfeld M, Leifeld L, Nischalke HD, Rockstroh JK, Hess L, Sauerbruch T *et al*. Reduced CC chemokine receptor (CCR) 1 and CCR5 surface expression on peripheral blood T lymphocytes from patients with chronic hepatitis C infection. *J Infect Dis* 2002; **185**: 1803-1807.
8. Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M *et al*. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 1996; **381**: 661-666.

9. Samson M, Libert F, Doranz BJ, Rucker J, Liesnard C, Farber CM *et al.* Resistance to HIV-1 infection in caucasian individuals bearing mutant alleles of the CCR-5 chemokine receptor gene. *Nature* 1996; **382**: 722-725.
10. Dean M, Carrington M, Winkler C, Huttley GA, Smith MW, Allikmets R *et al.* Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CKR5 structural gene. Hemophilia Growth and Development Study, Multicenter AIDS Cohort Study, Multicenter Hemophilia Cohort Study, San Francisco City Cohort, ALIVE Study. *Science* 1996; **273**: 1856-1862.
11. Stewart G. Chemokine genes--beating the odds. *Nat Med* 1998; 4: 275-257.
12. Woitas RP, Ahlenstiel G, Iwan A, Rockstroh JK, Brackmann HH, Kupfer B *et al.* Frequency of the HIV-protective CC chemokine receptor 5-Delta32/Delta32 genotype is increased in hepatitis C. *Gastroenterology* 2002; **122**: 1721-1728.
13. Goulding C, McManus R, Murphy A, MacDonald G, Barrett S, Crowe J *et al.* The CCR5-delta32 mutation: impact on disease outcome in individuals with hepatitis C infection from a single source. *Gut* 2005; **54**: 1157-1161.
14. Nattermann J, Timm J, Nischalke HD, Olbrich A, Michalk M, Tillmann HL *et al.* The predictive value of IL28B gene polymorphism for spontaneous clearance in a single source outbreak cohort is limited in patients carrying the CCR5Delta32 mutation. *J Hepatol* 2011; **55**: 1201-1206.
15. Thomas DL, Thio CL, Martin MP, Qi Y, Ge D, O'Huigin C *et al.* Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature* 2009; **461**: 798-801.
16. Suppiah V, Gaudieri S, Armstrong NJ, O'Connor KS, Berg T, Weltman M, *et al.* IL28B, HLA-C, and KIR variants additively predict response to therapy in chronic hepatitis C virus infection in a European Cohort: a cross-sectional study. *PLoS Med* 2011; **8**: e1001092.

17. Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ *et al.* Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature* 2009; **461**: 399-401.
18. Suppiah V, Moldovan M, Ahlenstiel G, Berg T, Weltman M, Abate ML *et al.* IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 2009; **41**: 1100-1104.
19. Tanaka Y, Nishida N, Sugiyama M, Kurosaki M, Matsuura K, Sakamoto N *et al.* Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009; **41**: 1105-1109.
20. Rauch A, Kutalik Z, Descombes P, Cai T, Di Iulio J, Mueller T *et al.* Genetic variation in IL28B is associated with chronic hepatitis C and treatment failure: a genome-wide association study. *Gastroenterology* 2010; **138**: 1338-1345.
21. Ahlenstiel G, Berg T, Woitas RP, Grunhage F, Iwan A, Hess L *et al.* Effects of the CCR5-Delta32 mutation on antiviral treatment in chronic hepatitis C. *J Hepatol* 2003; **39**: 245-252.
22. Promrat K, McDermott DH, Gonzalez CM, Kleiner DE, Koziol DE, Lessie M *et al.* Associations of chemokine system polymorphisms with clinical outcomes and treatment responses of chronic hepatitis C. *Gastroenterology* 2003; **124**: 352-360.
23. Hellier S, Frodsham AJ, Hennig BJ, Klenerman P, Knapp S, Ramaley P *et al.* Association of genetic variants of the chemokine receptor CCR5 and its ligands, RANTES and MCP-2, with outcome of HCV infection. *Hepatology* 2003; **38**: 1468-1476.
24. Mascheretti S, Hinrichsen H, Ross S, Buggisch P, Hampe J, Foelsch UR *et al.* Genetic variants in the CCR gene cluster and spontaneous viral elimination in hepatitis C-infected patients. *Clin Exp Immunol* 2004; **136**: 328-333.

25. Wasmuth HE, Werth A, Mueller T, Berg T, Dietrich CG, Geier A *et al.* CC chemokine receptor 5 delta32 polymorphism in two independent cohorts of hepatitis C virus infected patients without hemophilia. *J Mol Med* 2004; **82**: 64-69.
26. Goyal A, Suneetha PV, Kumar GT, Shukla DK, Arora N, Sarin SK. CCR5Delta32 mutation does not influence the susceptibility to HCV infection, severity of liver disease and response to therapy in patients with chronic hepatitis C. *World J Gastroenterol* 2006; **12**: 4721-4726.
27. Glas J, Torok HP, Simperl C, Konig A, Martin K, Schmidt F *et al.* The Delta 32 mutation of the chemokine-receptor 5 gene neither is correlated with chronic hepatitis C nor does it predict response to therapy with interferon-alpha and ribavirin. *Clin Immunol* 2003; **108**: 46-50.
28. Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncales FL, Jr. *et al.* Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002; **347**: 975-982.
29. Hadziyannis SJ, Sette H, Jr., Morgan TR, Balan V, Diago M, Marcellin P *et al.* Peginterferon-alpha2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann Intern Med* 2004; **140**: 346-355.
30. Novembre J, Galvani AP, Slatkin M. The geographic spread of the CCR5 Delta32 HIV-resistance allele. *PLoS Biol* 2005; **3**: e339.
31. Shin EC, Seifert U, Kato T, Rice CM, Feinstone SM, Klotzel PM *et al.* Virus-induced type I IFN stimulates generation of immunoproteasomes at the site of infection. *J Clin Invest* 2006; **116**: 3006-3014.

32. Yang YF, Tomura M, Iwasaki M, Ono S, Zou JP, Uno K *et al.* IFN-alpha acts on T-cell receptor-triggered human peripheral leukocytes to up-regulate CCR5 expression on CD4+ and CD8+ T cells. *J Clin Immunol* 2001; **21**: 402-429.
33. Thomas E, Feld JJ, Li Q, Hu Z, Fried MW, Liang TJ. Ribavirin potentiates interferon action by augmenting interferon-stimulated gene induction in hepatitis C virus cell culture models. *Hepatology* 2001; **53**: 32-41.
34. Tam RC, Pai B, Bard J, Lim C, Averett DR, Phan U *et al.* Ribavirin polarizes human T cell responses towards a Type 1 cytokine profile. *J Hepatol* 1999; **30**: 376-382.
35. Ahlenstiel G, Booth DR, Goerge J. Will IL28B polymorphism remain relevant to DAA treatment paradigms? *Antiviral Therapy* 2012 (in press).
36. Fatkenheuer G, Hoffmann C, Slim J, Rouzier R, Keung A, Li J *et al.* Short-term administration of the CCR5 antagonist vicriviroc to patients with HIV and HCV coinfection is safe and tolerable. *J Acquir Immune Defic Syndr* 2010; **53**: 78-85.
37. Bennetts BH, Teutsch SM, Buhler MM, Heard RN, Stewart GJ. The CCR5 deletion mutation fails to protect against multiple sclerosis. *Hum Immunol* 1997; **58**: 52-59.
38. Libert F, Cochaux P, Beckman G, Samson M, Aksenova M, Cao A *et al.* The *deltaccr5* mutation conferring protection against HIV-1 in Caucasian populations has a single and recent origin in Northeastern Europe. *Hum Mol Genet* 1998; **7**: 399-406.
39. Spagnolo P, Renzoni EA, Wells AU, Copley SJ, Desai SR, Sato H *et al.* C-C chemokine receptor 5 gene variants in relation to lung disease in sarcoidosis. *Am J Respir Crit Care Med* 2005; **172**: 721-728.
40. Oh DY, Jessen H, Kucherer C, Neumann K, Oh N, Poggensee G *et al.* CCR5Delta32 genotypes in a German HIV-1 seroconverter cohort and report of HIV-1 infection in a CCR5Delta32 homozygous individual. *PLoS One* 2008; **3**: e2747.

41. Nischalke HD, Nattermann J, Lichterfeld M, Woitas RP, Rockstroh JK, Sauerbruch T *et al.* Rapid determination of the Delta32 deletion in the human CC-chemokine receptor 5 (CCR5) gene without DNA extraction by lightcycler real-time polymerase chain reaction. *AIDS Res Hum Retroviruses* 2004; **20**: 750-754.

Figure 1.

Response in different CCR5-Δ32 genotypes (WT/WT = homozygous wildtype, Δ32/WT = CCR5-Δ32 heterozygous, Δ32/Δ32 = CCR5-Δ32 homozygous, all Δ32+ = all patients carrying one or two CCR5-Δ32 alleles.

Table 1

CCR5 Genotype	All patients		Australian patients		European patients	
	Healthy Controls (n=836)	Chronic HCV (n=758)	Healthy Controls (n=168)	Chronic HCV (n=245)	Healthy Controls (n=668)	Chronic HCV (n=513)
WT/WT	682 (81.6)	635 (83.8)	137 (82.0)	195 (79.6)	545 (81.6)	440 (85.8)
Δ32/WT	150 (17.9)	108 (14.2)	30 (18.0)	50 (20.4)	120 (18.0)	58 (11.3)
Δ32/ Δ32	4 (0.5)	15 (2.0)	1 (1.0)	0 (0.0)	3 (0.4)	15 (2.9)
Analysis	p = 0.0039		n.s.		p < 0.0001	
Allele						
WT	1514 (90.6)	1378 (90.9)	304 (90.5)	440 (89.9)	1210 (90.6)	938 (91.4)
Δ32	158 (9.4)	138 (9.1)	32 (10.7)	50 (10.2)	126 (9.4)	88 (8.6)

* WT = wildtype, numbers in brackets refers to %

Table 2

CCR5 genotype	All patients		Statistics
	SVR (n=341)	NSVR (n=417)	
WT/WT	287 (84.2)	348 (83.5)	n.s.
Δ32/WT	45 (13.2)	63 (15.1)	
Δ32/ Δ32	9 (2.6)	6 (1.4)	
CCR5 allele	SVR	NSVR	n.s.
WT	619 (90.8)	759 (91.0)	
Δ32	63 (9.2)	75 (9.0)	

* SVR = sustained virological responders, NSVR = non-SVR, WT = wildtype, numbers in brackets refers to %

Table 3

CCR5 + rs8099917	Total patients (n=710)			Australian patients (n=238)			European patients (n=472)		
	Responders (n=318)	Non-responders (n=392)	P value	Responders (n=101)	Non-responders (n=137)	P value	Responders (n=217)	Non-responders (n=255)	P value
WT/WT + G/G	12 (3.8)	25 (6.4)	–	2 (2.0)	7 (5.1)	–	10 (4.6)	18 (7.1)	–
WT/WT + T/G	96 (30.2)	162 (41.3)	0.0022	22 (21.8)	59 (43.1)	0.0008	74 (34.1)	103 (40.4)	–
WT/WT + T/T	161 (50.6)	141 (36.0)	< 0.0001	58 (57.4)	40 (29.2)	< 0.0001	103 (47.5)	101 (39.6)	–
D32/WT + G/G	1 (0.3)	0	–	0	0	–	1 (0.5)	0	–
D32/WT + T/G	13 (4.1)	36 (9.2)	0.011	7 (6.9)	24 (17.5)	0.019	6 (2.8)	12 (4.7)	–
D32/WT + T/T	26 (8.2)	23 (5.9)	–	12 (11.9)	7 (5.1)	–	14 (6.5)	16 (6.3)	–
D32/D32 + G/G	0	0	–	0	0	–	0	0	–
D32/D32 + T/G	1 (0.3)	4 (1.0)	–	0	0	–	1 (0.5)	4 (1.6)	–
D32/D32 + T/T	8 (2.5)	1 (0.3)	–	0	0	–	8 (3.7)	1 (0.4)	–

CCR5 + rs12979860	Total patients (n=791)			Australian patients (n=298)			European patients (n=493)		
	Responders (n=328)	Non-responders (n=401)	P value	Responders (n=101)	Non-responders (n=135)	P value	Responders (n=227)	Non-responders (n=266)	P value
WT/WT + C/C	132 (40.2)	68 (17.0)	< 0.0001	45 (44.6)	21 (15.6)	< 0.0001	87 (38.3)	47 (20.7)	< 0.0001
WT/WT + C/T	118 (36.0)	204 (50.9)	< 0.0001	32 (31.7)	68 (50.4)	0.0051	86 (37.9)	136 (59.9)	0.0032
WT/WT + T/T	25 (7.6)	65 (16.2)	0.0007	5 (5.0)	16 (11.9)	–	20 (8.8)	49 (21.6)	0.0026
D32/WT + C/C	22 (6.7)	13 (3.2)	0.029	8 (7.9)	5 (3.7)	–	14 (6.2)	8 (3.5)	0.0022
D32/WT + C/T	18 (5.5)	40 (10.0)	0.05	10 (9.9)	22 (16.3)	–	8 (3.5)	18 (7.9)	–
D32/WT + T/T	4 (1.2)	6 (1.5)	–	1 (1.0)	3 (2.2)	–	3 (1.3)	3 (1.3)	–
D32/D32 + C/C	3 (0.9)	1 (0.2)	–	0	0	–	3 (1.3)	1 (0.4)	–
D32/D32 + C/T	6 (1.8)	3 (0.7)	–	0	0	–	6 (2.6)	3 (1.3)	–
D32/D32 + T/T	0	1 (0.2)	–	0	0	–	0	1 (0.4)	–

