

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Selective detection of ATP and ADP in aqueous solution by using a fluorescent zinc receptor.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/126354> since 2015-12-29T12:00:42Z

Published version:

DOI:10.1039/c2cc35730e

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Chem. Commun., 48, **2012**, 11419–11421, DOI: 10.1039/c2cc35730e

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://pubs.rsc.org/en/Content/ArticleLanding/2012/CC/c2cc35730e>

Selective Detection of ATP and ADP in Aqueous Solution by using a Fluorescent Zinc Receptor

Maria Strianese,^a Stefano Milione,^{*a} Andrea Maranzana,^b Alfonso Grassi^a and Claudio Pellecchia^a

We report on the successfully use of a new zinc complex for the selective fluorescent detection of ADP and ATP in water. This is achieved by the complementary coordination of the phosphate groups to the metal centre and hydrogen bonding of the adenosine with the coordinated ligand.

Anions play a key role in a variety of environmental and biological processes. In particular among all anions, phosphate and molecules featuring this group are widely studied because of their ubiquitous presence in a range of life processes spanning from energy storage and signal transduction to gene construction.¹

Lately several optical sensors for monitoring anions have been devised.^{2,3} When designing phosphate receptors, possibility of sensing in an aqueous solution should be considered. The solvation of these substrates strongly competes with the process of complex formation, making the recognition of phosphate anions a challenging area of research. Another important point is the selectivity of recognition. Although some examples of fluorescent sensors which are selective over inorganic phosphate anions, such as phosphate and pyrophosphate have been developed³⁻⁸ the recognition and fluorescent sensing of adenosine di- or tri-phosphates over various nucleotide polyphosphates remains as a challenging problem.⁹

Among the various approaches, metal ion complexes are considered to be ideal for phosphate recognition in aqueous solutions.¹⁰ Many reported phosphate sensors involve fluorescent complexes with lanthanoid metals such as Eu(III), Tb(III) and Yb(III), having long lifetime of fluorescence and large Stokes' shifts.¹¹⁻¹³ Alternatively, the use of a transition metal offers the advantage of inducing fluorescence enhancement which is more desirable in sensing than fluorescence quenching. Recently Schiff base zinc complexes have been reported as efficient fluorescent sensors for adenosine nucleotides.¹⁴ In this contest, we thought of synthesizing a Zn complex featuring a Schiff base ligand with alcoholic functions on the ligand skeleton that could be able to establish intramolecular hydrogen bonds with the guest of interest. It is expected that additional functionalities for multiple non-covalent interactions with the substrates can contribute to the stability of the host-guest adduct leading to an enhancement of specificity and selectivity of detection. Our Schiff base ligand of choice is a derivative of pyridoxal with ethylenediamine (*pyr₂enH₂*, = N,N'-ethylenebis(pyridoxylideneiminato), Fig. 1).¹⁵




Fig. 1. Synthesis of *pyr₂enZn*

The target zinc complex (*pyr₂enZn*) was obtained in good yield by mixing equimolar amount of deprotonated ligand and $\text{Zn}(\text{CH}_3\text{COO})_2$ in MeOH at room temperature. The stoichiometry was confirmed by ESI-MS where the major peak at 421.17 *m/z* units corresponds to the mononuclear $[(\text{pyr}_2\text{enZn})\text{H}]^+$ species. No evidence for binuclear species in solution was found. The metal complex is soluble in dimethyl sulfoxide and acidic water. The ¹H NMR spectrum of *pyr₂enZn* in DMSO-*d*₆ indicated the presence of sharp signals consistent with the expected symmetric coordination of the ligand. Diagnostic resonances at 7.39 and 8.86 ppm for the iminic and aromatic protons, respectively, together with a singlet at 3.80 ppm for the protons on the diimine bridge were observed. These assignments were based on literature data¹⁵ and corroborated by means of mono- and bidimensional NMR experiments.

In DMSO-*d*₆ solution, the methylenic protons of the CH₂OH group are equivalent and show scalar coupling with the hydroxyl proton, the CH₂OH group gives rise to a doublet and a triplet at 4.55 and 5.20 ppm with a vicinal coupling constant ³*J*_{HH} of 4.7 Hz (Fig. S1, ESI†). When switching to D₂O, the same methylenic protons produce an AB pattern consisting of two doublets at 5.33 and 5.19 ppm with a geminal coupling constant ²*J*_{HH} of 13.8 Hz (Fig. S2, ESI†). Most likely, this change indicates a dependence of the speed of rotation of the CH₂OH group upon the intermolecular interactions that occur in solution. *Pyr₂enZn* was further characterized via Uv-vis and fluorescence spectroscopy. The electronic absorption spectra of the free ligand and of *pyr₂enZn* were measured both in DMSO and in MilliQ water (Fig. S5-S6, ESI†). In DMSO an intense band was observed at 378 nm. In H₂O *pyr₂enZn* displayed a more structured spectrum with a major peak at 223 nm and less intense absorption bands at 315 and 363 nm due to $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ intraligand transitions.^{16,17}

The capability of *pyr₂enZn* to act as recognition element for phosphate anions in aqueous solutions was first studied by ³¹P NMR spectroscopy. Addition of the nucleoside ADP, ATP, GTP, CTP, UTP or TTP to a saturated D₂O solution of *pyr₂enZn*

resulted into a clear and significant down field shift in the ^{31}P NMR spectrum with respect to the signals of the free anions. In the case of ADP the β -P-atom was shifted of 3.66 ppm whereas the α -P-atom underwent smaller shift ($\Delta\delta = 0.99$ ppm, Fig. 2). For ATP, the β -P-atom and γ -P-atom underwent consistent shifts (respectively $\Delta\delta = 1.13$ and 1.50 ppm) whereas the shift for α -P-atom was relatively small ($\Delta\delta = 0.19$ ppm) strongly suggesting that the α -P center was not interacting with the zinc center. Similar shifts were observed for GTP, CTP, UTP and TTP (Fig. S7-S10, Table S1, ESI †). These behaviors are fully consistent with the coordination of the tested nucleosides to the metal centre.^{14;18;19}




Fig 2. ^{31}P NMR spectra of Na_2AMP , Na_2ADP and Na_2ATP and after addition of pyr_2enZn (rt, D_2O). [pyr_2enZn] = 25 mM.

Significantly, when adding AMP or an inorganic phosphate (PO_4^{3-} , $\text{P}_2\text{O}_7^{4-}$, HPO_4^{2-} , H_2PO_4^-) to a saturated D_2O solution of pyr_2enZn no difference could be detected in the ^{31}P NMR spectra of the phosphate salt (Fig. 2 and Fig. S11-S13, ESI †). Further insights for the interaction of selected nucleosides with pyr_2enZn were gained from ^1H NMR studies. While the addition of AMP or inorganic phosphates to a saturated D_2O solution of pyr_2enZn did not produce any relevant change in the ^1H NMR spectrum of pyr_2enZn , the addition of ADP, ATP, GTP, CTP, UTP or TTP gave rise to an upfield shift in the ^1H NMR pattern of the complex. The largest shifts were observed for the aromatic $\text{C}(\text{H})\text{N}$ and imine $\text{HC}=\text{N}$ resonances (Fig. 14-17, ESI †). For example, in the case of ADP, the aromatic $\text{C}(\text{H})\text{N}$ proton, resonated at 8.19 ppm in the case of pyr_2enZn whereas it shifted at 7.82 ppm in the case of $\text{pyr}_2\text{enZn}/\text{ADP}$.

Intriguingly when adding ADP or ATP to a saturated D_2O solution of pyr_2enZn we observed a clear change in the multiplicity of the signal due the methylenic protons (at circa 5.3 ppm) of the CH_2OH group (Fig. 3). In addition to the geminal coupling, a scalar correlation with the imine ($J = 5.1$ Hz) and with the aromatic $\text{C}(\text{H})\text{N}$ protons ($J = 8.5$ Hz) was observed. The ^1H - ^1H COSY spectrum of the $\text{pyr}_2\text{enZn}/\text{ADP}$ adduct establishing the coupling partners is also displayed in Figure 3. Similar results were obtained with ATP (Fig. S18, ESI †). This finding suggests that the OH groups on the ligand framework participate within the interaction process between pyr_2enZn and ADP or ATP. The hydrogen-bonding between the OH and the nucleosides made the rotation of the CH_2OH group particularly hindered. On successive dilution the initial signal multiplicity is gradually restored, suggesting the reversibility of the recognition/interaction process. Significantly, when adding GTP, CTP, UTP or TTP to a saturated D_2O solution of pyr_2enZn , the multiplicity of the CH_2OH remained unchanged (Fig. S19-S22, ESI †).




Fig. 3. Change in multiplicity for the CH_2OH group and ex-panded region of 2D COSY spectrum of $\text{pyr}_2\text{enZn}/\text{ADP}$. ([pyr_2enZn] = 25 mM [Na_2ADP] = 70 mM, rt, D_2O).

To gain insights into the mode of binding of phosphorylated anions to the Zn complex and to assess the contributions of the alcoholic functions to the binding, we studied the interaction of pyr_2enZn with ADP, ATP and, for comparison, $\text{P}_2\text{O}_7^{4-}$ anions by Density Functional Theory, making use of the M06 functional. In the obtained adducts, the metal ion binds two hard phosphate oxygens of the phosphorylated species saturating its coordination sphere, the calculated O-Zn bond distances are 2.10 Å (Fig. S23-25, ESI †). ADP and ATP form structurally similar complexes: the organic moieties of the nucleotides are located above the coordinated ligand. In particular in the $\text{pyr}_2\text{enZn}/\text{ADP}$ adduct (Fig. 4) the amino group of the adenosine fragment (the hydrogen bond donor site) is close to the hydroxyl group of the ligand (the hydrogen bond acceptor site) interacting with it.




Fig. 4. Optimized structure of *pyr₂enZn*/ADP adduct. The water molecules were omitted for clarity of representation.

The distance between -NH₂ and OH, peculiar to hydrogen bonds, is about 2.1 Å. The bonding between the phosphates and hydroxyls in these adducts is not feasible because of the large distance between the two groups. The interaction between these groups is mediated by an hydrogen bonded water molecule.




Fig. 5. Emission spectra of *pyr₂enZn* (λ_{exc} 315 nm) after progressive additions of Na₂ADP (rt, MOPS). [Zn] = 5.5 mM.

Interactions between *pyr₂enZn* with ADP, ATP, AMP, UTP, CTP, TTP, GTP, PO₄³⁻, P₂O₇⁴⁻, HPO₄²⁻, H₂PO₄⁻, Cl⁻, F⁻ in MilliQ water solutions were also studied by means of fluorescence spectroscopy. When exciting an aqueous solution of *pyr₂enZn* at the maximum of its absorption (λ = 315 nm) and adding an excess of ADP a fast and consistent enhancement (~80%) of the initial fluorescence intensity was observed. When repeating the same experiment in the same conditions as before and adding an excess of ATP instead of ADP an enhancement of ~75% of the initial fluorescence intensity value was registered. On the other hand, adding an excess of UTP, CTP, TTP, GTP, resulted in a 75%, 35%, 75%, 82% fluorescence quenching respectively (Fig. S26, ESI†). Addition of an excess of an inorganic phosphate, either HPO₄²⁻ or H₂PO₄⁻ or P₂O₇⁴⁻ or PO₄³⁻, resulted in a 15%, 20%, 40% or 51% fluorescence quenching respectively (Fig. S27, ESI†). Addition of an excess of Cl⁻ or of F⁻ or of AMP did not result in any significant change of the fluorescence intensity.‡

The selective fluorescent detection of adenosine di- or tri-phosphate over other nucleotides and inorganic phosphates should include the contribution of the hydrogen-bonding interactions between OH groups on the ligand framework and ATP or ADP. A possible explanation for this finding is the interruption of photoinduced electron transfer (PET) of electronic density from the lone pairs on the oxygen atom to the π -system of the fluorophore resulting in luminescence enhancement.²⁰ This explanation is also supported by the ¹H NMR data by which we found that the multiplicity of the CH₂OH signal changes only in the case of addition of ATP and ADP.

To determine the binding affinity of *pyr₂enZn* for adenosine di- or tri-phosphates, ADP and ATP titrations were performed while monitoring the fluorescence intensity of the system. Figure 5 shows the ADP fluorescence titration of *pyr₂enZn* in the range of 0 - 3.5 eq. A similar fluorescence trend was obtained when titrating *pyr₂enZn* with ATP (Fig. S28, ESI†). The apparent equilibrium constants of association (K_a) were $158.7 \pm 1.4 \text{ M}^{-1}$ for the *pyr₂enZn* /ADP system and of $434.8 \pm 0.5 \text{ M}^{-1}$ for the *pyr₂enZn* /ATP construct. The greater interaction of ATP is probably due to the greater electronic charge density on the phosphate groups than that for ADP.

The detection limit of the proposed system for ADP and ATP was also determined and found to be in the micromolar range (Fig. S29-30 ESI†).

In summary a Zn²⁺-pyridoxal derivative has been synthesized and properly characterized via Uv-vis, fluorescence and NMR spectroscopy. Its potential in molecular recognition of adenosine phosphates over various nucleoside polyphosphates (UTP, CTP, TTP, GTP) and inorganic phosphates (P₂O₇⁴⁻, PO₄³⁻, H₂PO₄⁻ and HPO₄²⁻) has been investigated by means of ³¹P NMR, ¹H NMR and fluorescence spectroscopy. Clear evidences that *pyr₂enZn* can be successfully used for the selective fluorescent detection of ADP and ATP have been provided. It was shown that the interaction of the nucleoside polyphosphate with *pyr₂enZn* occurs through the complementary chelation of the phosphate groups to the metal centre and the hydrogen bonding between the adenosine and the alcoholic groups on the ligand framework. In order to gain independent evidence for the binding of ADP/ATP to *pyr₂enZn* and to model the adducts formed, DFT calculations were undertaken.

The OFF-ON switching of fluorescence together with the possibility of performing the measurements in aqueous solution add advantages to our system.

The authors thank Prof. Francesco De Riccardis and Prof. Moshe Kol for useful discussions. Dr. Patrizia Oliva and Dr. Patrizia Iannece of the Department of Chemistry (University of Salerno) for their technical assistance

Notes and references

^a Dipartimento di Chimica e Biologia, Università di Salerno, via Ponte don Melillo I-84084 Fisciano (SA), Italy.

E-mail: smilione@unisa.it

^b Dipartimento di Chimica, Università di Torino, via Pietro Giuria 7. I-10125 Torino (TO), Italy

† Electronic Supplementary Information (ESI) available: [Synthetic procedures, NMR and UV spectra, cartesian coordinates]. See DOI: 10.1039/b000000x/

‡ No significant changes in the fluorescence intensities were observed in the absence of phosphates or when aqueous solutions of free phosphates were excited at 315 nm.

- 1 A. E. Hargrove, S. Nieto, T. Zhang, J. L. Sessler, E. V. Anslyn, *Chem.Rev.* 2011, **111**, 6603.
- 2 M. Wenzel, J. R. Hiscock, P. A. Gale, *Chem.Soc.Rev.* 2012, **41**, 480.
- 3 G. Ambrosi, M. Formica, V. Fusi, L. Giorgi, A. Guerri, E. Macedi, M. Micheloni, P. Paoli, R. Pontellini, P. Rossi, *Inorg.Chem.* 2009, **48**, 5901.
- 4 H. K. Cho, D. H. Lee, J. I. Hong, *Chem Commun.* 2005, 1690.
- 5 S.K. Kim, D.H. Lee, J.I. Hong, J. Yoon, *Acc.Chem Res.* 2009, **42**, 23.
- 6 G. Ambrosi, M. Formica, V. Fusi, L. Giorgi, E. Macedi, M. Micheloni, P. Paoli, R. Pontellini, P. Rossi, *Chem.Eur.J.* 2011, **17**, 1670.
- 7 Z. Xu, N. J. Singh, J. Lim, J. Pan, H. N. Kim, S. Park, K. S. Kim, J. Yoon, *J.Am.Chem.Soc.* 2009, **131**, 15528.
- 8 S. Khatua, S. H. Choi, J. Lee, K. Kim, Y. Do, D. G. Churchill, *Inorg.Chem.* 2009, **48**, 2993.
- 9 M. J. Kim, K. M. Swamy, K. M. Lee, A. R. Jagdale, Y. Kim, S. J. Kim, K. H. Yoo, J. Yoon, *Chem.Commun.* 2009, 7215.
- 10 M. Schñferling, O. Wolfbeis, *Chem.Eur.J.* 2007, **13**, 4342.
- 11 Y. Miao, J. Liu, F. Hou, C. J. Jiang, *Lumin.* 2006, **116**, 67.
- 12 C. Yin, F. Gao, F. Huo, P. Yang, *Chem.Commun.* 2004, 934.
- 13 M. Cano, L. Rodriguez, J. C. Lima, F. Pina, C. A. Dalla, C. Pasquini, L. Schiaffino, *Inorg.Chem.* 2009, **48**, 6229.
- 14 M.F. Hawthorne, X. Yang, Z. Zheng, *Pure Appl.Chem.* 1994, 245.
- 15 I. Correia, P. J. Costa, M. T. Duarte, R. T. Henriques, M. F. Piedade, L. F. Veiros, T. Jakusch, T. Kiss, A. Dornyei, M. M. Castro, C. F. Geraldès, F. Avecilla, *Chem. Eur. J.* 2004, **10**, 2301.
- 16 B. Bosnich, *J.Am.Chem.Soc.* 1968, **90**, 627.
- 17 H. E. Smith, J. R. Neergaard, E. P. Burrows, F. M. Chen, *J.Am.Chem.Soc.* 1974, **96**, 2908.

Reference List

- (1) Ambrosi, G.; Formica, M.; Fusi, V.; Giorgi, L.; Guerri, A.; Macedi, E.; Micheloni, M.; Paoli, P.; Pontellini, R.; Rossi, P., *Inorg.Chem* 2009, **48**, 5901-5912.
- (2) Hargrove, A. E.; Nieto, S.; Zhang, T.; Sessler, J. L.; Anslyn, E. V., *Chem.Rev.* 2011, **111**, 6603-6782.
- (3) Wenzel, M.; Hiscock, J. R.; Gale, P. A., *Chem.Soc.Rev.* 2012, **41**, 480-520.
- (4) Cho, H. K.; Lee, D. H.; Hong, J. I., *Chem Commun.(Camb.)* 2005, 1690-1692.
- (5) Kim, S. K.; Lee, D. H.; Hong, J. I.; Yoon, J., *Acc.Chem Res.* 2009, **42**, 23-31.
- (6) Ambrosi, G.; Formica, M.; Fusi, V.; Giorgi, L.; Macedi, E.; Micheloni, M.; Paoli, P.; Pontellini, R.; Rossi, P., *Chemistry* 2011, **17**, 1670-1682.
- (7) Xu, Z.; Singh, N. J.; Lim, J.; Pan, J.; Kim, H. N.; Park, S.; Kim, K. S.; Yoon, J., *J Am.Chem Soc.* 2009, **131**, 15528-15533.
- (8) Khatua, S.; Choi, S. H.; Lee, J.; Kim, K.; Do, Y.; Churchill, D. G., *Inorg.Chem* 2009, **48**, 2993-2999.
- (9) Xu, Z.; Spring, D. R.; Yoon, J., *Chem.Asian J.* 2011, **6**, 2114-2122.
- (10) Kim, M. J.; Swamy, K. M.; Lee, K. M.; Jagdale, A. R.; Kim, Y.; Kim, S. J.; Yoo, K. H.; Yoon, J., *Chem Commun.(Camb.)* 2009, 7215-7217.
- (11) Schñferling, M.; Wolfbeis, O., *Chem.Eur.J.* 2007, **13**, 4342-4349.
- (12) Miao, Y.; Liu, J.; Hou, F.; Jiang, C., *Journal of Luminescence* 2006, **116**, 67-72.
- (13) Yin, C.; Gao, F.; Huo, F.; Yang, P., *Chem.Commun.* 2004, 934-935.
- (14) Cano, M.; Rodriguez, L.; Lima, J. C.; Pina, F.; Dalla, C. A.; Pasquini, C.; Schiaffino, L., *Inorg.Chem* 2009, **48**, 6229-6235.
- (15) Correia, I.; Costa, P. J.; Duarte, M. T.; Henriques, R. T.; Piedade, M. F.; Veiros, L. F.; Jakusch, T.; Kiss, T.; Dornyei, A.; Castro, M. M.; Geraldès, C. F.; Avecilla, F., *Chemistry* 2004, **10**, 2301-2317.
- (16) Bosnich, B., *J.Am.Chem.Soc.* 1968, **90**, 627-632.

- (17) Smith, H. E.; Neergaard, J. R.; Burrows, E. P.; Chen, F. M., *J.Am.Chem.Soc.* 1974, **96**, 2908-2916.
- (18) Kim, K.; Ha, Y.; Kaufman, L. +.; Churchill, D. G., *Inorg.Chem.* 2011, **51**, 928-938.
- (19) Rodríguez, L.; Lima, J. â. C.; Parola, A. J.; Pina, F.; Meitz, R.; Aucejo, R.; Garcia-Espanlâa, E.; Llinares, J. ü. M.; Soriano, C.; Alarcólin, J., *Inorg.Chem.* 2008, **47**, 6173-6183.
- (20) de Silva, A. P.; Gunaratne, H. Q. N.; Gunnlaugsson, T.; Huxley, A. J. M.; McCoy, C. P.; Rademacher, J. T.; Rice, T. E., *Chem.Rev.* 1997, **97**, 1515-1566.