

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**A new record of *Pycnodactylopsis tomentosa* (Fåhraeus 1842) (Coleoptera: Curculionidae) from Sindh, Pakistan.**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/144651> since

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

## Note

*Pakistan J. Zool.*, vol. 46(3), pp. 888, 2014.

### **A New Record of the Weevil *Pycnodactylopsis tomentosa* (Fåhraeus, 1842) (Coleoptera: Curculionidae) from Sindh, Pakistan**

**Zubair Ahmed,<sup>1,\*</sup> Massimo Meregalli<sup>2</sup> and  
Hanya Raza<sup>1</sup>**

<sup>1</sup>*Department of Zoology, Federal Urdu University  
of Arts, Science & Technology, Karachi, Pakistan*

<sup>2</sup>*Department of Life Sciences and Systems Biology,  
University of Torino, Italy*

The superfamily of weevils, Curculionoidea, contains about 62,000 described species and subspecies worldwide (Oberprieler *et al.*, 2007). The tribe Lixinae: Cleonini includes almost 600 species and subspecies, mainly distributed in the Palaearctic and Afrotropical regions, with some expansion to the Indo-Malayan region and Northamerica. The genus *Pycnodactylopsis* Voss, 1963 includes eight species. Six of the eight species are exclusive to the Afrotropical and Indo-Malayan fauna (Meregalli, unpublished data). The type species of *Pycnodactylopsis*, *P. schauffelei* Voss, 1963 is a synonym of *P. albogilva* (Gyllenhal, 1834), a species present in subsaharan Africa (Meregalli, 2013). One of the two Palaearctic species is *P. tomentosa* (Fåhraeus, 1842). This is the type species of *Louwia* Alonso-Zarazaga and Lyal, 1999, proposed as a subgenus of *Pycnodactylopsis* and reduced to synonym of the latter by Meregalli (2013). *P. tomentosa* was reported from Iran (Legalov *et al.*, 2010) and has a broad distribution in the Canary Islands, Northern Africa and India. Hashmi and Tashfeen (1992) and Chaudhry *et al.* (1966) listed species of weevils from Pakistan but never recorded this species. In August 2013 the first author collected one female specimen of *P. tomentosa* from Thar Desert Sindh, Pakistan, which represents a new country record. The specimen, determined by the second author, is deposited in the Federal Urdu University of Arts, Science and Technology, Karachi, Pakistan. This research work is part of B.S. thesis of the third author, who is

studying in the department of Zoology, the Federal Urdu University of Arts, Science & Technology, Karachi, Pakistan.

#### References

- Chaudhry, G-U., Chaudhry, M.I. and Khan, S.M., 1966. *Survey of insect fauna of forests of Pakistan. Final technical report*. Biological Sciences Research Division. Pakistan Forest Institute, Peshwar, pp. 167.
- Hashmi, A.A. and Tashfeen, A., 1992. *Proc. Pakistan Congr. Zool.*, **12**: 133-170.
- Legalov, A.A., Ghahari, H. and Arzanov, Yu. G., 2010. *Am. Zool. J.*, **2**: 191-244.
- Meregalli, M., 2013. In: *Catalogue of Palaearctic Coleoptera. 8. Curculionidae, II* (eds. I. Loebl and A. Smetana). Brill Academic Publishers, Leiden, The Netherlands, pp. 97-102.
- Oberprieler, R.G., Marvaldi, A.E. and Anderson, R.S., 2007. *Zootaxa*, **1668**: 491-520.

(Received 19 February 2014)

\* Corresponding author: zbrahmed36@gmail.com