

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

T-type channel-mediated neurotransmitter release

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/146621> since

Published version:

DOI:10.1007/s00424-014-1489-z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Emilio Carbone, Chiara Calorio, David H. F. Vandael
T-type channel-mediated neurotransmitter release
PFLUGERS ARCHIV (2014) 466
DOI: 10.1007/s00424-014-1489-z

The definitive version is available at:

<http://link.springer.com/content/pdf/10.1007/s00424-014-1489-z>

T-type channel-mediated neurotransmitter release

Emilio Carbone[§], Chiara Calorio, David H.F. Vandael

*Department of Drug Science
Lab of Cellular Physiology and Molecular Neuroscience
NIS Center, CNISM Unit, University of Torino, 10125 Torino, Italy*

[§] *Correspondence to:* Emilio Carbone

Key words: LVA calcium channels, T-type channel secretion coupling, vesicle exocytosis, neurons, synaptic transmission, neuroendocrine chromaffin cells

Acknowledgments: The authors thank the Italian M.I.U.R., Regione Piemonte, Università di Torino, Compagnia di San Paolo di Torino for funding.

Abstract

Besides controlling a wide variety of cell functions, T-type channels have been shown to regulate neurotransmitter release in peripheral and central synapses and neuroendocrine cells. Growing evidence over the last ten years suggests a key role of Cav3.2 and Cav3.1 channels in controlling basal neurosecretion near resting conditions and sustained release during mild stimulations. In some case, the contribution of low-voltage activated (LVA) channels is not directly evident but requires either the activation of coupled presynaptic receptors, block of ion channels or chelation of metal ions. Concerning the coupling to the secretory machinery, T-type channels appear loosely coupled to neurotransmitter and hormone release. In neurons, Cav3.2 and Cav3.1 channels mainly control the asynchronous appearance of “minis” (mIPSCs and mEPSCs). The same loose coupling is evident from membrane capacity and amperometric recordings in chromaffin cells and melanotropes where the low-threshold driven exocytosis possesses the same linear Ca^{2+} -dependence of the other voltage-gated Ca^{2+} channels (Cav1 and Cav2) that is strongly attenuated by slow calcium buffers. The intriguing issue is that, despite not expressing a consensus “synprint” site, Cav3.2 channels do interact with syntaxin-1A and SNAP-25 and, thus, may form nanodomains with secretory vesicles that can be regulated at low voltages. In this review, we discuss all the past and recent issues related to T-type channel-secretion coupling in neurons and neuroendocrine cells.

T-types and other non-canonical presynaptic Ca^{2+} channels regulate neurotransmitter release

T-type Ca^{2+} channels are transient, low-voltage activated Ca^{2+} channels that control Ca^{2+} entry during depolarizations near resting potential. Due to their biophysical properties and widespread expression, T-type channels control key functions like: low-threshold spikes, oscillatory cell activity, muscle contraction, hormone and neurotransmitter release, cell growth, differentiation and proliferation (see [21, 28, 61, 75, 78, 94] for a review). Given this, they are now proposed as therapeutic targets for a variety of diseases like: hypertension, angina pectoris, heart failure, atrial fibrillation, neuropathic pain, epilepsy, sleep disorders, obesity and cancer [39].

T-type channels possess unique gating properties that allow their electrophysiological identification: *i*) activation is voltage-dependent and channels open at very low voltages [19], *ii*) inactivation is voltage-dependent and complete within tens of milliseconds above 0 mV, *iii*) deactivation is slow at potentials near rest [18, 20], *iv*) Ca^{2+} and Ba^{2+} are carried equally well and single channel conductance is 3 to 4-fold lower than Cav1 and Cav2 channels [74], and *v*) activation and steady-state inactivation overlap at potentials near resting, generating a sizeable ‘window Ca^{2+} current’ [61].

Molecular cloning of T-type channels has provided evidence for the existence of three different pore forming $\alpha 1$ subunits (Cav3.1, Cav3.2, Cav3.3) with only 25% amino acid homology but similar pore structure organization to Cav1 (L-type) and Cav2 (N, P/Q, R-type) channels [78]. Cav3 channels differ from Cav1 and Cav2 channels for two other important singularities: *i*) they are not surrounded by β , $\alpha_2\delta$ and γ auxiliary subunits [62] and, *ii*) they lack the consensus “synprint site” (synaptic protein interaction site) [78] which allows Cav2.1 and Cav2.2 channels to bind SNARE proteins (syntaxin-1A/B, SNAP-25) and non-SNARE proteins (synaptotagmin) (see [22] for a review). This binding is crucial to give rise to Ca^{2+} nanodomains with docked presynaptic vesicles, ensuring high rates of vesicle release during rapid synaptic activity [73, 99].

Despite Cav2.1 (P/Q-type) and Cav2.2 (N-type) are the dominant Ca^{2+} channel species controlling presynaptic vesicle release in most neurons, there is increasing evidence that also other Ca^{2+} channel isoforms lacking the consensus synprint site (Cav2.3, Cav1 and Cav3) contribute to

presynaptic Ca^{2+} elevations and vesicle release. There are now abundant proofs that Cav3.2 (R-type) channels are localized at the presynaptic terminals [29, 51] and trigger neurotransmitter release by enhancing presynaptic Ca^{2+} levels [10, 31, 37, 55, 97, 104]. The same is true for Cav1 (L-type) channels, which regulate glutamate release in rod photoreceptor terminals [103] and co-localize at the hot spots of bipolar cell terminals [6]. L-type channels also regulate transmitter release in isolated hippocampal single buttons [71], cultured GABAergic hippocampal neurons [53] and cerebral cortical neurons [54] during high-frequency presynaptic activity. To date there is also overwhelming evidence that Cav3 channels are also directly involved in neurotransmitter release in an increasing number of neuronal preparations and neuroendocrine cells, making undisputable their key role in neurotransmitter release [17, 21, 102]. Here we summarize earlier and recent findings that support a direct role of T-type channels in the control of Ca^{2+} -dependent transmitter release and discuss recently suggested molecular mechanisms of T-type channel coupling to synaptic vesicles that do not occur through the consensus synprint site [100].

T-type channels and neurotransmitter release: earlier evidence (1997-2005)

The first report describing a role of T-type channels on neurotransmitter release was on the activity of the oscillatory heart GABAergic interneurons of the leech, where fast inactivating presynaptic T-type currents were shown to regulate the graded response of heart interneurons [3, 50]. At nearly the same time, Bao et al., 1998 [5] showed that, T-type channels highly expressed at the soma of afferent nociceptive neurons [94], are also involved in the control of synaptic activity between primary afferents and second-order nociceptive neurons in the dorsal horn of the spinal cord. Presynaptic T-type channels control the frequency of spontaneous glutamatergic EPSCs mediated by AMPA/kainate receptors while the latency and amplitude of AP-evoked EPSCs is controlled by HVA channels. Interestingly, the frequency of mEPSCs increases greatly with mild KCl depolarizations and is prevented by mibefradil, suggesting a critical role of T-type channels in basal nociception activity near rest. A role for T-type channels in the control of mEPSCs is relevant since the frequency of events can markedly increase during hyperalgesia to reach the threshold of nociception responses (see below).

Clear evidence of the involvement of T-type channels in fast neurotransmitter release came also from early studies at the reciprocal synapses that retinal bipolar cells form with A17 and AII amacrine cells [77, 88]. Bipolar cells express both L-type (Cav1.4) and T-type channels, which give rise to Ca^{2+} currents of comparable amplitude: at -40 mV for T-type and -10 mV for L-type channels. Correcting for the driving force, single channel conductance and open probability this implies comparable expression densities of the two channels. Evidence for the involvement of T-type channel-mediated exocytosis comes from membrane capacitance measurements (ΔC) [see [72] for details] in which glutamate release is directly measured as an increase of membrane surface area during presynaptic vesicle fusion. Pan et al. [77] estimated a ΔC of 7 fF attributed to T-type channels, corresponding to the fusion of about 90 vesicles of 50 nm diameter during pulses of 500 ms at -40 mV. T-type channels could be recruited only if the cells were previously hyperpolarized to -80 mV and the ΔC was fully blocked by mibefradil and spared by nimodipine.

T-type channels play a unique functional role also at the large spines of axonless granule cells (GCs) that form dendrodendritic reciprocal synapses with the mitral and tufted cells (M/TCs) of olfactory bulbs. These terminals express sufficient high densities of T-type channels to regulate fast GABAergic responses [32]. Small inhibitory post-synaptic potentials (IPSPs) with short latency and slow decay time constant can be recorded in whole-cell clamped mitral cells. The IPSCs are fully blocked by bicuculline and strongly attenuated by mibefradil, suggesting direct involvement of T-type channels on the inhibitory GABA_A-mediated control of mitral cells by GCs stimulation. GABA release from GC dendrites can be induced also by Ca^{2+} entry through NMDA receptors activation and Ca^{2+} release from intracellular stores [33]. The co-participation of T-type channels to GABA release from GC spines thus strengthens the GC-mediated lateral inhibition of M/TCs. It is

also worth noticing that, high-densities of T-type channels at distal and proximal dendrites ensure the possibility of evoking sub-threshold Ca^{2+} transients during back-propagation of low-threshold spikes from the soma to the spines. Under physiological conditions this warrants a better graded control of olfactory bulb function.

Early observations on the involvement of T-type channels in neurosecretion include also the original report by Harkins et al. [44] in a mouse pheochromocytoma (MPC) cell line. In this work newly expressed Cav3.2 channels produce Ca^{2+} current densities that generate membrane capacitance changes associated to vesicle secretion with comparable Ca^{2+} -efficiency to Cav2.2 (N-type) and Cav1.2 (L-type) channels. Since MPC cells express most proteins involved in Ca^{2+} -dependent neurotransmitter release (SNAP-25 and syntaxin), the rationale of these findings is that in MPC cell lines all voltage-gated Ca^{2+} channels can functionally couple to catecholamines release sites and control fast exocytosis, regardless of the availability of the “synprint site” that is present only in N and P/Q-type channels. Similar conclusions are drawn in rat melanotropes where T-type currents can drive membrane capacitance changes associated with vesicle secretion, with the same Ca^{2+} -dependence of HVA channels (L-, N-, P/Q- and R-type). This, together with the findings on chromaffin cells discussed in the following sections, allows to conclude that in neurosecretory cells there is no preferential Ca^{2+} channel coupling to secretory vesicles. The presence of T-type channels in neuroendocrine cells is thus mainly to ensure functional secretion over a broader range of membrane voltages: from near resting conditions where a T-type “window current” warrants sufficient basal hormone release to very positive potentials where HVA channels are maximally activated.

T-type channels-mediated transmitter release in neurons: recent findings (2006-2014)

Following the above early reports, there has been an increasing interest on the role of T-type channels in neurotransmitter release in the last eight years. Here we illustrate the main findings on presynaptic T-type channels controlling GABAergic and glutamatergic synapses of central neurons and T-type channels-mediated hormone secretion in neuroendocrine cells.

Cav3.2-mediated control of excitatory synapses in nociceptive dorsal-horn neurons

T-type channels are widely expressed both at the soma of small diameter nociceptive afferent neurons and at the presynaptic terminals of nociceptive dorsal horn neurons in lamina I-II, where they control the activity of mEPSCs [5] (see [94] for a review). T-type channels are also expressed at the soma of the lamina I neurons that are responsible for forwarding nociceptive signals to different regions of the brain as part of the ascending pain pathway [49]. In this way, Cav3 channels have a wide control of pain-processing pathways both at the pre- and postsynaptic level of nociceptive projecting neurons. Until recently the identity of the presynaptic Cav3 channel type involved in such pain responses was unknown. This has been solved recently, by showing that: *i*) Cav3.2 channels are expressed at the presynaptic terminals of nociceptive projecting neurons [52], *ii*) the frequency of mEPSCs is strongly attenuated by TTA-P2 and Ni^{2+} and *iii*) TTA-P2 have no action on the mEPSCs of Cav3.2^{-/-} dorsal horn neurons. Interestingly, T-type channels control specifically the nociceptive neurons of superficial laminae I-II but not the excitatory transmission of non-nociceptive neurons of deeper laminae (III-V), which are regulated by Cav2.1 and Cav2.2 channels. The Cav3.2-mediated regulation of mEPSCs is also remarkably more robust in superficial dorsal horn neurons of animal models of painful diabetic neuropathy as compared to healthy age-matched animals. The frequency of the mEPSCs increases drastically in diabetic animals and can thus critically trigger the transmission of hyperalgesic signals through the ascending pain pathway.

Cav3.2-mediated ATP release in dorsal root ganglion sandwich synapses

Subpopulations of dorsal root ganglion (DRG) neurons can communicate with their immediate neighbors via neuron–glial cell–neuron "sandwich synapses" [81]. Stimulation of one neuron leads to transglial activation of its partner by a purinergic synaptic contact between the stimulated neuron and the glial cell and a glutamatergic synaptic contact between the glial cell and the receiving neuron [83]. Release of ATP from the stimulated neuron is attributed to a classical mechanism involving Ca^{2+} entry via Cav3.2 channels, while the release of glutamate from the glia is mediated by a Gq/PLC pathway causing intracellular Ca^{2+} increase [82]. In the case of ATP, the Cav3.2-mediated release is blocked by moderate depolarization (-50 mV) or low-concentration of Ni^{2+} and is insensitive to specific blockers and toxins of Cav1 and Cav2 channels [84]. Synaptic transmission is particularly sensitive to the holding potential as expected from the presence of T-type channels at the somatic presynaptic terminal. The presence of Cav3.2 to gate the transmission at the DRG "sandwich synapses" thus broadens the possibility of modulating nociceptive sensory signals via sub-threshold voltage oscillations in addition to AP evoked responses. Taken together, the Cav3.2-driven release pathway at the DRG neuron–glial cell–neuron may play a critical role in the transmission of abnormal pain sensation.

Cav3.1-mediated GABA release at hippocampal perisomatic interneurons

T-type channels have also a key role in the release of GABA from hippocampal perisomatic-targeting interneurons [93]. More specifically, immune-histochemically identified Cav3.1 channels are co-localized with $\alpha_3\beta_4$ nicotinic receptors (nAChR) at the synaptic regions of parvalbumin expressing GABAergic interneurons. Local applications of ACh activate Cav3.1 currents, which in turn trigger bursts of mIPSCs that can transiently reduce pyramidal cell excitability. The inhibitory response is only partially attenuated by mibefradil, low doses of Ni^{2+} and TTA-P2, suggesting that T-type channels contribute only to a fraction of the overall mIPSCs bursting response. The remaining part is associated to a Ca^{2+} -induced release from presynaptic Ca^{2+} stores. In addition, bath application of EGTA-AM strongly reduces the nAChR-induced inhibitory response, suggesting loose coupling between T-type channels and GABA release. This is different from the HVA-driven pathway in which few Cav2.1 (P/Q-type) channels are sufficient to elicit synaptic activity from parvalbumin basket cells [11]. The Cav3.1-driven transmission occurs regardless of HVA channel Ca^{2+} influx and it may thus operate in parallel with the normal AP-driven HVA channel-dependent mechanism. This ensures a broader control of inhibitory interneurons on pyramidal cell excitability in hippocampus.

Cav3.2-driven glutamate release in cortical neurons

Presynaptic Cav3.2 channels are also expressed in glutamatergic cortical synaptic terminals contacting entorhinal cortical layer III pyramidal neurons [48]. These terminals, express high densities of the hyperpolarization-activated cyclic-nucleotide gated channel type 1 (HCN1) that partially depolarize the terminal by passing a net inward current. A high resting potential (-50 mV) inactivates functionally available T-type channels, reduces basal presynaptic Ca^{2+} levels and attenuates the frequency of spontaneous mEPSCs. Block of HCN1 channels by ZD-7288 or genetic deletion of the channel causes a net increase in the frequency of spontaneous mEPSCs that is effectively blocked by mibefradil (10 μM), TTA-A2 (1 μM) and TTA-P2 (1 μM) [87, 96]. Applications of ω -agatoxin IVA, ω -conotoxin GVIA and SNX-482 have no effect on the ZD-7288-induced increase of mEPSCs frequency, suggesting a dominant presynaptic control of Cav3 channels on the increased mEPSCs activity following HCN1 block. Given that changes in synaptic strength are pivotal to the induction and maintenance of synaptic plasticity and that entorhinal cortical layer III neurons are involved in memory formation and spatial navigation [80], these findings indicate that a HCN1/Cav3.2-driven excitatory transmission may critically contribute to such physiological processes.

T-type channel-mediated GABA release of fast spiking interneurons in dentate granule cells

In line with the above findings T-type channels are also known to play a role at the mono- and di-synaptic GABAergic inputs of dentate granule cells [41]. Zinc conditions the functioning of these synapses, but its action is widely unexplored. Zinc inhibits ionotropic receptors commonly found at central synapses, as well as a number of voltage-gated ion channels including Cav3 channels [91, 95]. In fast spiking GABAergic interneurons Zn blocks somatic T-type Ca^{2+} currents. Thus, zinc chelation enhances these currents, decreases spike threshold and broadens AP half-width. Zinc also reversibly depresses GABAergic transmission while zinc chelation enhances the inhibitory response. Since zinc chelation is prevented only by T-type and not by HVA channel blockers, a reasonable assumption is that T-type channels controls GABA release of fast spiking interneurons. Although lacking a convincing immunoreactivity proof of presynaptically expressed Cav3 channels, the reported electrophysiological findings together with the evidence of a dense localization of Cav3 channel gene transcripts in the cell granule layer [92] and parvalbumin-immunoreactive interneurons [7, 86], strongly support a role of T-type channels in GABA transmission.

T-type channel-driven dopamine release from the somata of dopaminergic neurons

T-type channels control also dopamine release from the soma of dopaminergic neurons in the substantia nigra *pars compacta*, as shown by amperometric recordings with carbon fiber microelectrodes [57]. Prolonged depolarizations induced by high KCl solutions cause massive release of dopamine in the form of bursts of amperometric events that are associated to the quantal dopamine release. Mibefradil (10 μM), reduces the frequency of amperometric spikes thus suggesting a direct role of T-type channels in dopamine release in substantia nigra *pars compacta* neurons.

T-type channel-driven synaptic exocytosis in immature auditory hair cells

A role of T-type channels on vesicle exocytosis has been reported also in chick auditory hair cells during development [63]. Immature chick hair cells (embryonic *days 10–12*) possess a low-voltage-activating, rapidly inactivating T-type current and a high-voltage-activating, non-inactivating L-type current to control vesicle exocytosis. Synaptic activity evoked by T-type channels and measured through membrane capacitance changes (ΔC) displays a fast release component but lacks the slow sustained release component. This suggests an efficient recruitment of nearby synaptic vesicles and a less efficient action on more distant vesicles by T-type channel Ca^{2+} currents in immature hair cells. During hair cell development, the participation of T-type currents progressively decreases, whereas that of L-type increases. A property observed also in other cell types, including chromaffin cells (see below [64]). Of relevance is the observation that with maturation (from E12 to P2) the Ca^{2+} -efficiency of the T-type channel increases markedly, although less steeply than L-type channels. These latter dominate during development.

T-type channels-driven hormone release in neuroendocrine cells

In line with the early reports in MPC cells [44] and melanotropes [67] described above and those on chromaffin cells described below [15, 16, 38], two recent reports confirm that T-type channels can effectively drive hormone secretion in neuroendocrine cells. Using markers of prostate secretion and FM1-43 fluorescence imaging of membrane trafficking, Gackiere et al [35] have shown unequivocally that prostate cell differentiation is associated with an increase in Ca^{2+} -dependent secretion that critically relies on Cav3.2 channel activity. Cav3.2 channels up-regulation may account for the alteration of secretion during prostate cancer development. T-type channels promote the secretion of potential mitogenic factors and participate to the progression of the disease toward an androgen-independent stage. In line with this, Zanatta et al. [106] show that exocytosis in immature rat Sertoli cells is modulated by the reverse T3 thyroid hormone (rT3) through the activation of T-type channels. Cav3 channels are expressed in Sertoli cells [60] and are up-regulated by rT3 through an integrin-mediated action and consequent PKC and MEK activation.

T-type channels secretion coupling in adrenal chromaffin cells

Despite widely distributed in most nervous, muscular and epithelial tissues, T-type channels are weakly expressed or absent in adult chromaffin cells. In bovine chromaffin cells of adult animals, the mRNA encoding for Cav3.1 and Cav3.2 is clearly expressed [36], but functional T-type channels are mainly absent [1, 4, 13, 23, 24]. T-type channels are usually expressed and carry sufficient Ca^{2+} currents in embryonic and neonatal RCCs [9, 64, 90]. They are also available in a small percentage of adult RCCs [47]. T-type channels have been reported to be absent in adult mouse chromaffin cells (MCCs) [45, 69, 70, 98]. Preliminary data from our lab, however, clearly show that the I/V characteristics of adult MCCs display the typical “low-threshold shoulder” associated with T-type channels during ramp commands (see Fig. 4 in [40]). It is thus very likely that availability of T-type channels may critically depend on chromaffin cell isolation, culture cell conditions or exogenous stimulations, as in the case of applied chronic stressors which up-regulate Cav3.2 channels in RCCs [16, 38, 76, 90] and MCCs [46, 89].

When expressed, T-type channels appear effectively coupled to chromaffin secretory granules and control a sizeable amount of “low-threshold” catecholamine release [15, 16, 38, 59, 68]. The Ca^{2+} -dependence of T-type channel driven secretion is linear and strikingly similar to that of Cav1 channels [15], which is comparable to that of Cav2.1 and Cav2.2 channels [14]. Independently of the low voltage range of activation, T-type channels are thus closely distributed near the docked secretory vesicles, just like Cav1 and Cav2 channels and control catecholamine secretion (see Fig. 1 for a T-type channel vs. vesicles arrangement derived from [15, 16, 38]). Thus, by lowering the threshold of AP generation T-type channels broaden the interval of voltage control of catecholamine release. This extended Ca^{2+} channel secretion coupling may be relevant when chromaffin cells depolarize steadily or fire at very high frequencies during stressful conditions.

T-type channels and low-threshold exocytosis: an adaptive response to stress

As for the nicotinic receptors and gap-junction channels [42, 43] also T-type channels undergo functional remodeling during stress [17, 21, 66]. This is particularly evident in chromaffin cells of adult animals as an adaptive response to stress conditions like chronic and intermittent hypoxia [16, 90], cAMP/PKA, β -adrenoreceptor and high-frequency sympathetic stimulation [46, 59, 76, 89]. T-type channel remodeling occurs also during the transition from intra- to extra-uterine life [64], when newborns experience episodic hypoxia conditions and catecholamine secretion switches from a non-neurogenic to a neurogenic control mode [85].

A common molecular mechanism in response to stressors is a marked recruitment of functioning Cav3.2 channels. Up-regulation of Cav3.2 channels lowers the threshold of AP firing [76], increases the amount of catecholamines secreted at low-voltages [16, 38] and mobilizes a RRP equivalent to that of L-type channels with the same Ca^{2+} -dependence (see Fig. 1). An increased density of T-type channels is also expected to decrease the resting potential and increase the firing frequency, as it occurs in spontaneously firing rat chromaffin cells in the adrenal-gland slices from stressed rats [25]. Indeed, an increased density of functioning T-type channels is a remodeling mechanism that many cells and tissues develop in response to stressors. Beside hypoxia, β -AR and high-frequency sympathetic stimulation other stressors (glucocorticoids, aldosterone, ACTH, 17- β estradiol) and stress-induced conditions (hypertension, cardiomyopathies, diabetic and chronic visceral pain) lead to the up-regulation of Cav3.1 and Cav3.2 channels in neuronal, muscular and endocrine cells (see Table 1 in [65]). This reinforces the view that T-type are “stress-activated channels” [17, 21, 27, 89].

As listed above, there are multiple ways by which Cav3.2 channels are either activated or up-regulated by stressors in chromaffin cells. They act either acutely within minutes (sustained sympathetic stimulation, PACAP release) or slowly, requiring hours or days (chronic and

intermittent hypoxia, β_1 -AR stimulation, VIP, aldosterone, ACTH). The rapidly developing mechanism (minutes) is most evident in mouse innervated adrenal gland slices [46]. This pathway is activated by PACAP, released during high-frequency splanchnic stimulation, and mediated by cAMP-activated exchange proteins (Epac) that stimulate PLC and PKC through a non-canonical cAMP-dependent pathway. The PKC target is the phosphorylation of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger [59] and a fast recruitment of Cav3.2 channels [46] (Fig. 2a). The former induces a small depolarization near resting potential (ΔV_m) sufficient to activate a robust “low-threshold” Cav3.2 current that triggers exocytosis.

The second pathway, slowly developing, is a long lasting mechanism that requires hours or days to progress. This is the case for chronic/intermittent hypoxia and cell exposure to β_1 -AR agonists (adrenaline, isoprenaline), which both increase the levels of CACNA1H mRNA (Cav3.2 gene). Their action is prevented by protein synthesis inhibitors [15, 16, 38, 76, 90], suggesting net recruitment of newly synthesized Cav3.2 channels (Fig. 2b). For the case of β_1 -AR stimulation, Cav3.2 recruitment is mediated by a cAMP-dependent (PKA-independent) Epac pathway which leads to the activation of transcription factors [76], most likely through a Rap-mediated pathway and the phosphorylation of ERK [8]. This cAMP-dependent mechanism is completely different from the fast cAMP-dependent up-regulation of Cav1.2 and Cav1.3 channels observed in MCCs [66].

Concerning chronic and intermittent hypoxia, recent findings indicate the involvement of hypoxia-inducible factors. HIF-2 α is involved in the up-regulation of Cav3.2 in PC12 cells exposed to chronic hypoxia [30]. In line with this, overnight incubations with the unspecific HIF activator desferrioxamine mimics the effects of hypoxia in PC12 cells [30] and RCCs [15]. In addition to HIFs, also NADHP-oxidases (NOX2, NOX4) and reactive-oxygen species [90] appear to play an important role in the recruitment of Cav3.2 in neonatal RCCs subjected to intermittent hypoxia [61]. A hypothetical scheme in which NOX, ROS and HIF are linked together is shown in Fig. 2c. The scheme is derived from [105] in which ROS is shown to activate PLC, DAG and PKC, favoring the stabilization and translocation of HIF-1 α to the nucleus to start transcription and gene expression.

T-type channel-driven catecholamine release and O₂ sensing during development

Chromaffin cells possess an inborn chemosensitivity to O₂ levels during fetal life, when the control of hormone release from the adrenal medulla is mostly non-neurogenic [85] and effectively regulated by the gap-junction coupling existing between neighboring cells [25]. During this developmental phase, chromaffin cells respond to acute hypoxia with a robust catecholamine release, mostly controlled by Cav3.2 channels [64]. High densities of T-type channels and enhanced electrical coupling facilitate action potential generation and propagation, as well as the synchronous release of catecholamines from cell populations of the adrenal medulla. This increases the amount of released adrenaline that helps sustaining the cardiovascular response to hypoxia [42, 43]. Of great significance, however, is the observation that cholinergic innervation of the adrenal gland, following postnatal development [85], causes a loss of functional Cav3.2 channels [64] paralleled by an increased density of α_7 -built nAChRs [12]. But even more interesting is the observation that O₂-sensitivity and Cav3.2 channel expression is reestablished in chromaffin cells after adrenal gland denervation, suggesting an opposing correlation between nicotinic AChRs expression and Cav3.2 channels down-regulation or vice versa [64].

This apparent coupling between AChRs and Cav3.2 is worth being investigated since T-type channels co-localize with a number of ion channels. In central neurons, Cav3.2 channels are co-localized to voltage-gated (Kv4) [2], Ca²⁺-gated K⁺ channels (KCa1.1, KCa3.1) [34, 79] and HCN1 channels [48] to widen the voltage range where K⁺, Ca²⁺ and HCN1 channels can control the shape and frequency of AP trains. Cav3.2 channels are also effectively inhibited by closely associated TRPV1 channels in small diameter DRG neurons, which would predict a TRPV1-mediated

analgesic action via an increased threshold of nociceptive signals [26]. Indeed, a TRPV1-mediated inhibition of Cav3.2 channels has been recently proposed to regulate the analgesic effect of paracetamol in supraspinal DRG neurons [56]. All these findings widen the range of operation of T-type channels in the regulation of cell excitability and underline once more the amazing functional importance of these voltage-gated Ca^{2+} channels in the control of vital cell functions.

Cav3.2 channel binding to syntaxin-1A and SNAP-25: a likely mechanism for fast neurosecretion?

It is surprising that despite the overwhelming evidence in favor of a role of T-type channels in neurotransmitter release there are still few data available on the molecular mechanism that triggers these events. Most data in presynaptic terminals, neuronal somata and neuroendocrine cells suggest loose coupling between T-type channels and secretory vesicles. This contradicts the idea of the formation of “nanodomains” between T-type channels, SNARE proteins and vesicles. However, recent reports clearly show that Cav3.2 channels biochemically interact with syntaxin-1A and SNAP-25 [100] at the carboxy-terminal domain. Electrophysiological recordings show that syntaxin-1A binding to Cav3.2 channels causes marked changes to channel gating, resembling those regarding N- and P/Q-type channels (see [22, 101] for a review). Most relevant is the observation that the Cav3.2-syntaxin-SNAP-25 interaction somehow affects a reconstituted T-type-driven exocytosis in MPC cells. Expression of Cav3.2 in MPC9/3L cells, which possess native SNARE proteins [44], produces a marked “low-threshold” membrane capacitance (ΔC) increase that is fully prevented by uncoupling T-type channels from SNARE proteins. Taken together these findings suggest that Cav3.2 channels co-localize with SNARE proteins and may thus potentially form membrane nanodomains with secretory vesicles able to trigger vesicle fusion and neurotransmitter release.

Concluding remarks

There is clearly no doubt that T-type channels contribute to neurotransmitter release in neurons and neuroendocrine cells. The growing evidence over the last ten years suggests a key role of these channels in controlling basal neurosecretion near resting conditions and sustained release during mild stimulations. In some cases, the contribution of T-type channels is not direct but requires either the activation of coupled presynaptic receptors [93], the block of ion channels [48] or the chelation of metal ions [41]. The effect is nevertheless remarkable and in all cases of physio-pathological significance. Concerning the coupling to the secretory apparatus, T-type channels appear loosely coupled to neurotransmitter and hormone release. In central neurons, T-type channels mainly control the asynchronous appearance of mIPSCs and mEPSCs while AP-evoked IPSCs and EPSCs are dominated by P/Q and N-type channels. The same loose coupling is evident from the membrane capacity changes in chromaffin cells and melanotropes where the low-threshold-driven exocytosis possesses the same Ca^{2+} dependence of the other HVA calcium channels that is strongly attenuated by slow calcium buffers [58]. The intriguing issue is that, despite not expressing a consensus synprint site, T-type channels do interact with syntaxin-1A and SNAP-25 and, thus, may form nanodomains with secretory vesicles that can be regulated at low voltages. Future studies will clarify this issue, may be by showing that a fraction of T-type channels couple directly to the vesicle via SNARE-proteins and the remaining does not and that the picture may be different in presynaptic terminal and neurosecretory cells.

Figure Legends

Figure 1 – Loose coupling of T- and L-type channels to exocytosis in chromaffin cells.

Left) Schematic drawing illustrating the loose coupling of T- and L-type channels to secretory granules in adrenal chromaffin cells. The two voltage-gated channels are coupled to secretion with the same efficacy for Ca^{2+} . This implies that on average they are at an equal distance from fused vesicles ready for release. For more details see ref [15]. *Top-right*) Membrane capacity changes (ΔC) versus the corresponding quantity of calcium charge (pC) for cAMP-treated RCCs expressing T- and R-type channels and incubated in a medium containing ω -toxins and nifedipine (black circles, solid line) compared to the ΔC increases induced by L- and R-type channels expressed in control RCCs treated with ω -toxins (red squares, dashed line). The linear regressions have nearly equal slope (adapted from [15] and [38]). *Bottom-right*) ΔC recordings driven by T-type currents passing during step depolarization of 150 ms to -20 mV (black traces) and L-type currents during pulse of 200 ms to +10 mV (red traces) (adapted from [15] and [38]).

Figure 2 - Cell signaling pathways leading to fast or slow recruitment of Cav3.2 channel-mediated secretion in chromaffin cells.

a) PACAP-stimulated secretion in mouse chromaffin cells through the acute phosphoregulation of Cav3.2 channels [46, 59, 89]. PACAP activation of the PAC1 receptor elevates cAMP that activates Epac within minutes (not shown). The subsequent activation of PLC and PKC leads to the phosphorylation of a sodium-calcium exchanger (NCX) that causes a small cell depolarization (ΔV_m), the opening of T-type channels and release of catecholamines. T-type channels may also be directly up-regulated by PKC activation. *b*) cAMP and β_1 -AR mediated up-regulation of Cav3.2 channels through an Epac-mediated pathway that activates transcription factors and CACNA1H gene expression in rat chromaffin cells [76]. Cav3.2 channel recruitment leads to a well resolved “low-threshold” exocytosis [38]. The involvement of Rap and ERK in the modulatory pathway is only hypothetical (?). *c*) Cell signaling pathway leading to CACNA1H gene expression and Cav3.2 channels recruitment during hypoxia in rat chromaffin cells [16]. The final result of chronic/intermittent hypoxia is the activation of transcription factors (HIF, CREB, etc...) and CACNA1H gene expression that leads to Cav3.2 channels recruitment and associated exocytosis. The schematic pathway induced by hypoxia (NOX, ROS, PLC, PKG, HIF) is adapted from [65, 105].

References

1. Albillos A, Garcia AG, Gandia L (1993) omega-Agatoxin-IVA-sensitive calcium channels in bovine chromaffin cells. *FEBS Lett* 336:259-62 DOI
2. Anderson D, Mehaffey WH, Iftinca M, Rehak R, Engbers JDT, Hameed S, Zamponi GW, Turner RW (2010) Regulation of neuronal activity by Cav3-Kv4 channel signaling complexes. *Nature Neuroscience* 13:333-U14 DOI 10.1038/nn.2493
3. Angstadt JD, Calabrese RL (1991) Calcium currents and graded synaptic transmission between heart interneurons of the leech. *Journal of Neuroscience* 11:746-759 DOI
4. Artalejo CR, Dahmer MK, Perlman RL, Fox AP (1991) Two types of Ca²⁺ currents are found in bovine chromaffin cells: facilitation is due to the recruitment of one type. *J Physiol* 432:681-707 DOI
5. Bao JP, Li JJ, Perl ER (1998) Differences in Ca²⁺ channels governing generation of miniature and evoked excitatory synaptic currents in spinal laminae I and II. *Journal of Neuroscience* 18:8740-8750 DOI
6. Berntson A, Taylor WR, Morgans CW (2002) Molecular identity, synaptic localization, and physiology of calcium channels in retinal bipolar cells. *J Neurosci Res* 71:146-51 DOI 10.1002/jnr.10459
7. Blasco-Ibanez JM, Martinez-Guijarro FJ, Freund TF (2000) Recurrent mossy fibers preferentially innervate parvalbumin-immunoreactive interneurons in the granule cell layer of the rat dentate gyrus. *Neuroreport* 11:3219-25 DOI
8. Borland G, Smith BO, Yarwood SJ (2009) EPAC proteins transduce diverse cellular actions of cAMP. *Br J Pharmacol* 158:70-86 DOI 10.1111/j.1476-5381.2008.00087.x
10.1111/j.1476-5381.2008.00087.x. Epub 2009 Feb 6.
9. Bornaud R, Hidalgo J, Yu H, Jaimovich E, Shimahara T (2001) Low threshold T-type calcium current in rat embryonic chromaffin cells. *J Physiol* 537:35-44 DOI
10. Breustedt J, Vogt KE, Miller RJ, Nicoll RA, Schmitz D (2003) Alpha1E-containing Ca²⁺ channels are involved in synaptic plasticity. *Proc Natl Acad Sci U S A* 100:12450-5 DOI 10.1073/pnas.2035117100
11. Bucurenciu I, Kulik A, Schwaller B, Frotscher M, Jonas P (2008) Nanodomain coupling between Ca²⁺ channels and Ca²⁺ sensors promotes fast and efficient transmitter release at a cortical GABAergic synapse. *Neuron* 57:536-545 DOI 10.1016/j.neuron.2007.12.026
12. Buttigieg J, Brown S, Holloway AC, Nurse CA (2009) Chronic nicotine blunts hypoxic sensitivity in perinatal rat adrenal chromaffin cells via upregulation of KATP channels: role of alpha7 nicotinic acetylcholine receptor and hypoxia-inducible factor-2alpha. *J Neurosci* 29:7137-47 DOI 10.1523/jneurosci.0544-09.2009
13. Carabelli V, Carra I, Carbone E (1998) Localized secretion of ATP and opioids revealed through single Ca²⁺ channel modulation in bovine chromaffin cells. *Neuron* 20:1255-68 DOI

14. Carabelli V, Giaccipoli A, Baldelli P, Carbone E, Artalejo AR (2003) Distinct potentiation of L-type currents and secretion by cAMP in rat chromaffin cells. *Biophys J* 85:1326-37 DOI 10.1016/s0006-3495(03)74567-6
15. Carabelli V, Marcantoni A, Comunanza V, Carbone E (2007) Fast exocytosis mediated by T- and L-type channels in chromaffin cells: distinct voltage-dependence but similar Ca²⁺ - dependence. *Eur Biophys J* 36:753-62 DOI 10.1007/s00249-007-0138-2
16. Carabelli V, Marcantoni A, Comunanza V, de Luca A, Diaz J, Borges R, Carbone E (2007) Chronic hypoxia up-regulates alpha1H T-type channels and low-threshold catecholamine secretion in rat chromaffin cells. *J Physiol* 584:149-65 DOI 10.1113/jphysiol.2007.132274
17. Carbone E, Giaccipoli A, Marcantoni A, Guido D, Carabelli V (2006) A new role for T-type channels in fast "low-threshold" exocytosis. *Cell Calcium* 40:147-54 DOI 10.1016/j.ceca.2006.04.019
18. Carbone E, Lux HD (1984) A low voltage-activated calcium conductance in embryonic chick sensory neurons. *Biophys J* 46:413-8 DOI 10.1016/s0006-3495(84)84037-0
19. Carbone E, Lux HD (1984) A low voltage-activated, fully inactivating Ca channel in vertebrate sensory neurones. *Nature* 310:501-2 DOI
20. Carbone E, Lux HD (1987) Kinetics and selectivity of a low-voltage-activated calcium current in chick and rat sensory neurones. *J Physiol* 386:547-70 DOI
21. Carbone E, Marcantoni A, Giaccipoli A, Guido D, Carabelli V (2006) T-type channels-secretion coupling: evidence for a fast low-threshold exocytosis. *Pflugers Arch* 453:373-83 DOI 10.1007/s00424-006-0100-7
22. Catterall WA (1999) Interactions of presynaptic Ca²⁺ channels and snare proteins in neurotransmitter release. *Ann N Y Acad Sci* 868:144-59 DOI
23. Cena V, Nicolas GP, Sanchez-Garcia P, Kirpekar SM, Garcia AG (1983) Pharmacological dissection of receptor-associated and voltage-sensitive ionic channels involved in catecholamine release. *Neuroscience* 10:1455-62 DOI
24. Cesetti T, Hernandez-Guijo JM, Baldelli P, Carabelli V, Carbone E (2003) Opposite action of beta1- and beta2-adrenergic receptors on Ca(V)1 L-channel current in rat adrenal chromaffin cells. *J Neurosci* 23:73-83 DOI
25. Colomer C, Lafont C, Guerineau NC (2008) Stress-induced Intercellular Communication Remodeling in the Rat Adrenal Medulla. In: Kvetnansky R, Aguilera G, Goldstein D, Jezova D, Krizanova O, Sabban EL, Pacak K (eds) *Stress, Neurotransmitters, and Hormones: Neuroendocrine and Genetic Mechanisms*. pp. 106-111
26. Comunanza V, Carbone E, Marcantoni A, Sher E, Ursu D (2011) Calcium-dependent inhibition of T-type calcium channels by TRPV1 activation in rat sensory neurons. *Pflugers Arch* 462:709-22 DOI 10.1007/s00424-011-1023-5
27. Comunanza V, Marcantoni A, Vandael DH, Mahapatra S, Gavello D, Carabelli V, Carbone E (2010) CaV1.3 as pacemaker channels in adrenal chromaffin cells: specific role on exo- and endocytosis? *Channels (Austin)* 4:440-6 DOI

28. Cueni L, Canepari M, Adelman JP, Luthi A (2008) Ca(2+) signaling by T-type Ca(2+) channels in neurons. *Pflugers Arch* 457:1161-72 DOI 10.1007/s00424-008-0582-6
29. Day NC, Shaw PJ, McCormack AL, Craig PJ, Smith W, Beattie R, Williams TL, Ellis SB, Ince PG, Harpold MM, Lodge D, Volsen SG (1996) Distribution of alpha 1A, alpha 1B and alpha 1E voltage-dependent calcium channel subunits in the human hippocampus and parahippocampal gyrus. *Neuroscience* 71:1013-24 DOI
30. Del Toro R, Levitsky KL, Lopez-Barneo J, Chiara MD (2003) Induction of T-type calcium channel gene expression by chronic hypoxia. *J Biol Chem* 278:22316-24 DOI 10.1074/jbc.M212576200
31. Dietrich D, Kirschstein T, Kukley M, Pereverzev A, von der Brölie C, Schneider T, Beck H (2003) Functional specialization of presynaptic Cav2.3 Ca²⁺ channels. *Neuron* 39:483-96 DOI
32. Egger V, Svoboda K, Mainen ZF (2003) Mechanisms of lateral inhibition in the olfactory bulb: Efficiency and modulation of spike-evoked calcium influx into granule cells. *Journal of Neuroscience* 23:7551-7558 DOI
33. Egger V, Svoboda K, Mainen ZF (2005) Dendrodendritic synaptic signals in olfactory bulb granule cells: Local spine boost and global low-threshold spike. *Journal of Neuroscience* 25:3521-3530 DOI 10.1523/jneurosci.4746-04.2005
34. Engbers JDT, Anderson D, Asmara H, Rehak R, Mehaffey WH, Hameed S, McKay BE, Kruskic M, Zamponi GW, Turner RW (2012) Intermediate conductance calcium-activated potassium channels modulate summation of parallel fiber input in cerebellar Purkinje cells. *Proceedings of the National Academy of Sciences of the United States of America* 109:2601-2606 DOI 10.1073/pnas.1115024109
35. Gackiere F, Bidaux G, Delcourt P, Van Coppenolle F, Katsogiannou M, Dewailly E, Bavencoffe A, Van Chuoï-Mariot MT, Mauroy B, Prevarskaya N, Mariot P (2008) CaV3.2 T-type calcium channels are involved in calcium-dependent secretion of neuroendocrine prostate cancer cells. *J Biol Chem* 283:10162-73 DOI 10.1074/jbc.M707159200
36. Garcia-Palomero E, Renart J, Andres-Mateos E, Solis-Garrido LM, Matute C, Herrero CJ, Garcia AG, Montiel C (2001) Differential expression of calcium channel subtypes in the bovine adrenal medulla. *Neuroendocrinology* 74:251-61 DOI 54692
37. Gasparini S, Kasyanov AM, Pietrobon D, Voronin LL, Cherubini E (2001) Presynaptic R-type calcium channels contribute to fast excitatory synaptic transmission in the rat hippocampus. *J Neurosci* 21:8715-21 DOI
38. Giacippoli A, Novara M, de Luca A, Baldelli P, Marcantoni A, Carbone E, Carabelli V (2006) Low-threshold exocytosis induced by cAMP-recruited CaV3.2 (alpha1H) channels in rat chromaffin cells. *Biophys J* 90:1830-41 DOI 10.1529/biophysj.105.071647
39. Giordanetto F, Knerr L, Wallberg A (2011) T-type calcium channels inhibitors: a patent review. *Expert Opinion on Therapeutic Patents* 21:85-101 DOI 10.1517/13543776.2011.536532
40. Gosso S, Gavello D, Giachello CN, Franchino C, Carbone E, Carabelli V (2011) The effect of CdSe-ZnS quantum dots on calcium currents and catecholamine secretion in mouse chromaffin cells. *Biomaterials* 32:9040-50 DOI 10.1016/j.biomaterials.2011.08.031

41. Grauert A, Engel D, Ruiz AJ (2014) Endogenous zinc depresses GABAergic transmission via T-type Ca²⁺ channels and broadens the time window for integration of glutamatergic inputs in dentate granule cells. *Journal of Physiology-London* 592:67-86 DOI 10.1113/jphysiol.2013.261420
42. Guerineau NC, Desarmenien MG (2010) Developmental and Stress-Induced Remodeling of Cell-Cell Communication in the Adrenal Medullary Tissue. *Cellular and Molecular Neurobiology* 30:1425-1431 DOI 10.1007/s10571-010-9583-z
43. Guerineau NC, Desarmenien MG, Carabelli V, Carbone E (2012) Functional chromaffin cell plasticity in response to stress: focus on nicotinic, gap junction, and voltage-gated Ca²⁺ channels. *J Mol Neurosci* 48:368-86 DOI 10.1007/s12031-012-9707-7
44. Harkins AB, Cahill AL, Powers JF, Tischler AS, Fox AP (2003) Expression of recombinant calcium channels support secretion in a mouse pheochromocytoma cell line. *Journal of Neurophysiology* 90:2325-2333 DOI 10.1152/jn.00425.2003
45. Hernandez-Guijo JM, de Pascual R, Garcia AG, Gandia L (1998) Separation of calcium channel current components in mouse chromaffin cells superfused with low- and high-barium solutions. *Pflugers Arch* 436:75-82 DOI
46. Hill J, Chan SA, Kuri B, Smith C (2011) Pituitary adenylate cyclase-activating peptide (PACAP) recruits low voltage-activated T-type calcium influx under acute sympathetic stimulation in mouse adrenal chromaffin cells. *J Biol Chem* 286:42459-69 DOI 10.1074/jbc.M111.289389
47. Hollins B, Ikeda SR (1996) Inward currents underlying action potentials in rat adrenal chromaffin cells. *J Neurophysiol* 76:1195-211 DOI
48. Huang Z, Lujan R, Kadurin I, Uebele VN, Renger JJ, Dolphin AC, Shah MM (2011) Presynaptic HCN1 channels regulate Ca(V)_{3.2} activity and neurotransmission at select cortical synapses. *Nature Neuroscience* 14:478-U114 DOI 10.1038/nn.2757
49. Ikeda H, Heinke B, Ruscheweyh R, Sandkuhler J (2003) Synaptic plasticity in spinal lamina I projection neurons that mediate hyperalgesia. *Science* 299:1237-40 DOI 10.1126/science.1080659
50. Ivanov AI, Calabrese RL (2000) Intracellular Ca²⁺ dynamics during spontaneous and evoked activity of leech heart interneurons: Low-threshold Ca currents and graded synaptic transmission. *Journal of Neuroscience* 20:4930-4943 DOI
51. Iwasaki S, Takahashi T (1998) Developmental changes in calcium channel types mediating synaptic transmission in rat auditory brainstem. *J Physiol* 509 (Pt 2):419-23 DOI
52. Jacus MO, Uebele VN, Renger JJ, Todorovic SM (2012) Presynaptic Ca(V)_{3.2} Channels Regulate Excitatory Neurotransmission in Nociceptive Dorsal Horn Neurons. *Journal of Neuroscience* 32:9374-9382 DOI 10.1523/jneurosci.0068-12.2012
53. Jensen K, Jensen MS, Lambert JD (1999) Role of presynaptic L-type Ca²⁺ channels in GABAergic synaptic transmission in cultured hippocampal neurons. *J Neurophysiol* 81:1225-30 DOI
54. Jensen K, Mody I (2001) L-type Ca²⁺ channel-mediated short-term plasticity of GABAergic synapses. *Nat Neurosci* 4:975-6 DOI 10.1038/nn722

55. Kamp MA, Krieger A, Henry M, Hescheler J, Weiergraber M, Schneider T (2005) Presynaptic 'Ca_v2.3-containing' E-type Ca channels share dual roles during neurotransmitter release. *Eur J Neurosci* 21:1617-25 DOI 10.1111/j.1460-9568.2005.03984.x
56. Kerckhove N, Mallet C, Francois A, Boudes M, Chemin J, Voets T, Bourinet E, Alloui A, Eschalier A (2014) Ca_v3.2 calcium channels: The key protagonist in the supraspinal effect of paracetamol. *Pain* 10.1016/j.pain.2014.01.015
57. Kim Y, Park MK, Uhm D-Y, Chung S (2007) Modulation of T-type Ca²⁺ channels by corticotropin-releasing factor through protein kinase C pathway in MN9D dopaminergic cells. *Biochemical and Biophysical Research Communications* 358:796-801 DOI 10.1016/j.bbrc.2007.04.198
58. Klingauf J, Neher E (1997) Modeling buffered Ca²⁺ diffusion near the membrane: Implications for secretion in neuroendocrine cells. *Biophysical Journal* 72:674-690 DOI
59. Kuri BA, Chan SA, Smith CB (2009) PACAP regulates immediate catecholamine release from adrenal chromaffin cells in an activity-dependent manner through a protein kinase C-dependent pathway. *J Neurochem* 110:1214-25 DOI 10.1111/j.1471-4159.2009.06206.x
60. Lalevee N, Pluciennik F, Joffre M (1997) Voltage-dependent calcium current with properties of T-type current in Sertoli cells from immature rat testis in primary cultures. *Biol Reprod* 56:680-7 DOI
61. Lambert RC, Bessaih T, Crunelli V, Leresche N (2013) The many faces of T-type calcium channels. *Pflugers Arch* 10.1007/s00424-013-1353-6
62. Leuranguer V, Bourinet E, Lory P, Nargeot J (1998) Antisense depletion of beta-subunits fails to affect T-type calcium channels properties in a neuroblastoma cell line. *Neuropharmacology* 37:701-8 DOI
63. Levic S, Dulon D (2012) The temporal characteristics of Ca²⁺ entry through L-type and T-type Ca²⁺ channels shape exocytosis efficiency in chick auditory hair cells during development. *J Neurophysiol* 108:3116-23 DOI 10.1152/jn.00555.2012
64. Levitsky KL, Lopez-Barneo J (2009) Developmental change of T-type Ca²⁺ channel expression and its role in rat chromaffin cell responsiveness to acute hypoxia. *J Physiol* 587:1917-29 DOI 10.1113/jphysiol.2009.168989
65. Mahapatra S, Calorio C, Vandael DH, Marcantoni A, Carabelli V, Carbone E (2012) Calcium channel types contributing to chromaffin cell excitability, exocytosis and endocytosis. *Cell Calcium* 51:321-30 DOI 10.1016/j.ceca.2012.01.005
66. Mahapatra S, Marcantoni A, Zuccotti A, Carabelli V, Carbone E (2012) Equal sensitivity of Cav1.2 and Cav1.3 channels to the opposing modulations of PKA and PKG in mouse chromaffin cells. *J Physiol* 590:5053-73 DOI 10.1113/jphysiol.2012.236729
67. Mansvelder HD, Kits KS (2000) All classes of calcium channel couple with equal efficiency to exocytosis in rat melanotropes, inducing linear stimulus-secretion coupling. *J Physiol* 526 Pt 2:327-39 DOI

68. Marcantoni A, Carabelli V, Comunanza V, Hoddah H, Carbone E (2008) Calcium channels in chromaffin cells: focus on L and T types. *Acta Physiol (Oxf)* 192:233-46 DOI 10.1111/j.1748-1716.2007.01815.x
69. Marcantoni A, Carabelli V, Vandael DH, Comunanza V, Carbone E (2009) PDE type-4 inhibition increases L-type Ca(2+) currents, action potential firing, and quantal size of exocytosis in mouse chromaffin cells. *Pflugers Arch* 457:1093-110 DOI 10.1007/s00424-008-0584-4
70. Marcantoni A, Vandael DH, Mahapatra S, Carabelli V, Sinnegger-Brauns MJ, Striessnig J, Carbone E (2010) Loss of Cav1.3 channels reveals the critical role of L-type and BK channel coupling in pacemaking mouse adrenal chromaffin cells. *J Neurosci* 30:491-504 DOI 10.1523/jneurosci.4961-09.2010
71. Murakami N, Ishibashi H, Katsurabayashi S, Akaike N (2002) Calcium channel subtypes on single GABAergic presynaptic terminal projecting to rat hippocampal neurons. *Brain Res* 951:121-9 DOI
72. Neher E, Marty A (1982) Discrete changes of cell membrane capacitance observed under conditions of enhanced secretion in bovine adrenal chromaffin cells. *Proc Natl Acad Sci U S A* 79:6712-6 DOI
73. Neher E, Sakaba T (2008) Multiple roles of calcium ions in the regulation of neurotransmitter release. *Neuron* 59:861-72 DOI 10.1016/j.neuron.2008.08.019
74. Nilius B, Hess P, Lansman JB, Tsien RW (1985) A novel type of cardiac calcium-channel in ventricular cells. *Nature* 316:443-446 DOI 10.1038/316443a0
75. Nilius B, Talavera K, Verkhratsky A (2006) T-type calcium channels: The never ending story. *Cell Calcium* 40:81-88 DOI 10.1016/j.ceca.2006.04.011
76. Novara M, Baldelli P, Cavallari D, Carabelli V, Giaccipoli A, Carbone E (2004) Exposure to cAMP and beta-adrenergic stimulation recruits Ca(V)₃ T-type channels in rat chromaffin cells through Epac cAMP-receptor proteins. *J Physiol* 558:433-49 DOI 10.1113/jphysiol.2004.061184
77. Pan ZH, Hu HJ, Perring P, Andrade R (2001) T-type Ca²⁺ channels mediate neurotransmitter release in retinal bipolar cells. *Neuron* 32:89-98 DOI 10.1016/s0896-6273(01)00454-8
78. Perez-Reyes E (2003) Molecular physiology of low-voltage-activated T-type calcium channels. *Physiological Reviews* 83:117-161 DOI 10.1152/physrev.00018.2002
79. Rehak R, Bartoletti TM, Engbers JDT, Berecki G, Turner RW, Zamponi GW (2013) Low Voltage Activation of KCa_{1.1} Current by Cav₃-KCa_{1.1} Complexes. *Plos One* 8:10.1371/journal.pone.0061844
80. Remondes M, Schuman EM (2004) Role for a cortical input to hippocampal area CA1 in the consolidation of a long-term memory. *Nature* 431:699-703 DOI 10.1038/nature02965
81. Rozanski GM, Kim H, Li Q, Wong FK, Stanley EF (2012) Slow chemical transmission between dorsal root ganglion neuron somata. *European Journal of Neuroscience* 36:3314-3321 DOI 10.1111/j.1460-9568.2012.08233.x

82. Rozanski GM, Li Q, Kim H, Stanley EF (2013) Purinergic transmission and transglial signaling between neuron somata in the dorsal root ganglion. *European Journal of Neuroscience* 37:359-365 DOI 10.1111/ejn.12082
83. Rozanski GM, Li Q, Stanley EF (2013) Transglial transmission at the dorsal root ganglion sandwich synapse: glial cell to postsynaptic neuron communication. *European Journal of Neuroscience* 37:1221-1228 DOI 10.1111/ejn.12132
84. Rozanski GM, Nath AR, Adams ME, Stanley EF (2013) Low voltage-activated calcium channels gate transmitter release at the dorsal root ganglion sandwich synapse. *Journal of Physiology-London* 591:5575-5583 DOI 10.1113/jphysiol.2013.260281
85. Seidler FJ, Slotkin TA (1985) Adrenomedullary function in the neonatal rat - responses to acute-hypoxia. *Journal of Physiology-London* 358:1-16 DOI
86. Seress L, Gallyas F (2000) The use of a sodium tungstate developer markedly improves the electron microscopic localization of zinc by the Timm method. *J Neurosci Methods* 100:33-9 DOI
87. Shipe WD, Barrow JC, Yang Z-Q, Lindsley CW, Yang FV, Schlegel K-AS, Shu Y, Rittle KE, Bock MG, Hartman GD, Tang C, Ballard JE, Kuo Y, Adarayan ED, Prueksaritanont T, Zrada MM, Uebele VN, Nuss CE, Connolly TM, Doran SM, Fox SV, Kraus RL, Marino MJ, Gratifieds VK, Vargas HM, Bunting PB, Hasbun-Manning M, Evans RM, Koblan KS, Renger JJ (2008) Design, synthesis, and evaluation of a novel 4-aminomethyl-4-fluoropiperidine as a T-type Ca(2+) channel antagonist. *Journal of Medicinal Chemistry* 51:3692-3695 DOI 10.1021/jm800419w
88. Singer JH, Diamond JS (2003) Sustained Ca²⁺ entry elicits transient postsynaptic currents at a retinal ribbon synapse. *Journal of Neuroscience* 23:10923-10933 DOI
89. Smith CB, Eiden LE (2012) Is PACAP the major neurotransmitter for stress transduction at the adrenomedullary synapse? *J Mol Neurosci* 48:403-12 DOI 10.1007/s12031-012-9749-x
90. Souvannakitti D, Nanduri J, Yuan G, Kumar GK, Fox AP, Prabhakar NR (2010) NADPH oxidase-dependent regulation of T-type Ca²⁺ channels and ryanodine receptors mediate the augmented exocytosis of catecholamines from intermittent hypoxia-treated neonatal rat chromaffin cells. *J Neurosci* 30:10763-72 DOI 10.1523/jneurosci.2307-10.2010
91. Sun HS, Hui K, Lee DW, Feng ZP (2007) Zn²⁺ sensitivity of high- and low-voltage activated calcium channels. *Biophys J* 93:1175-83 DOI 10.1529/biophysj.106.103333
92. Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, Bayliss DA (1999) Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J Neurosci* 19:1895-911 DOI
93. Tang A-H, Karson MA, Nagode DA, McIntosh JM, Uebele VN, Renger JJ, Klugmann M, Milner TA, Alger BE (2011) Nerve Terminal Nicotinic Acetylcholine Receptors Initiate Quantal GABA Release from Perisomatic Interneurons by Activating Axonal T-Type (Ca_v3) Ca²⁺ Channels and Ca²⁺ Release from Stores. *Journal of Neuroscience* 31:13546-13561 DOI 10.1523/jneurosci.2781-11.2011
94. Todorovic SM, Jevtovic-Todorovic V (2013) Neuropathic pain: role for presynaptic T-type channels in nociceptive signaling. *Pflugers Archiv-European Journal of Physiology* 465:921-927 DOI 10.1007/s00424-012-1211-y

95. Traboulsie A, Chemin J, Chevalier M, Quignard JF, Nargeot J, Lory P (2006) Subunit-specific modulation of T-type calcium channels by zinc. *J Physiol* 578:159-71 DOI 10.1113/jphysiol.2006.114496
96. Uebele VN, Nuss CE, Santarelli VP, Garson SL, Kraus RL, Barrow JC, Stauffer SR, Koblan KS, Renger JJ, Aton S, Seibt J, Dumoulin M, Jha SK, Coleman T, Frank MG (2009) T-type calcium channels regulate cortical plasticity in-vivo (vol 20, pg 257, 2009). *Neuroreport* 20:973-973 DOI 10.1097/WNR.0b013e32832f573b
97. Urbano FJ, Piedras-Renteria ES, Jun K, Shin HS, Uchitel OD, Tsien RW (2003) Altered properties of quantal neurotransmitter release at endplates of mice lacking P/Q-type Ca²⁺ channels. *Proc Natl Acad Sci U S A* 100:3491-6 DOI 10.1073/pnas.0437991100
98. Vandael DH, Zuccotti A, Striessnig J, Carbone E (2012) Ca(V)1.3-driven SK channel activation regulates pacemaking and spike frequency adaptation in mouse chromaffin cells. *J Neurosci* 32:16345-59 DOI 10.1523/jneurosci.3715-12.2012
99. Weber AM, Wong FK, Tufford AR, Schlichter LC, Matveev V, Stanley EF (2010) N-type Ca²⁺ channels carry the largest current: implications for nanodomains and transmitter release. *Nat Neurosci* 13:1348-50 DOI 10.1038/nn.2657
100. Weiss N, Hameed S, Fernandez-Fernandez JM, Fablet K, Karmazinova M, Poillot C, Proft J, Chen L, Bidaud I, Monteil A, Huc-Brandt S, Lacinova L, Lory P, Zamponi GW, De Waard M (2012) A Ca(v)3.2/Syntaxin-1A Signaling Complex Controls T-type Channel Activity and Low-threshold Exocytosis. *Journal of Biological Chemistry* 287:2810-2818 DOI 10.1074/jbc.M111.290882
101. Weiss N, Zamponi GW (2012) Regulation of voltage-gated calcium channels by synaptic proteins. *Adv Exp Med Biol* 740:759-75 DOI 10.1007/978-94-007-2888-2_33
102. Weiss N, Zamponi GW (2013) Control of low-threshold exocytosis by T-type calcium channels. *Biochimica Et Biophysica Acta-Biomembranes* 1828:1579-1586 DOI 10.1016/j.bbamem.2012.07.031
103. Wilkinson MF, Barnes S (1996) The dihydropyridine-sensitive calcium channel subtype in cone photoreceptors. *J Gen Physiol* 107:621-30 DOI
104. Wu LG, Borst JG, Sakmann B (1998) R-type Ca²⁺ currents evoke transmitter release at a rat central synapse. *Proc Natl Acad Sci U S A* 95:4720-5 DOI
105. Yuan G, Nanduri J, Khan S, Semenza GL, Prabhakar NR (2008) Induction of HIF-1 alpha expression by intermittent hypoxia: involvement of NADPH oxidase, Ca²⁺ signaling, prolyl hydroxylases, and mTOR. *J Cell Physiol* 217:674-85 DOI 10.1002/jcp.21537
106. Zanatta AP, Zanatta L, Goncalves R, Zamoner A, Silva FR (2013) Rapid responses to reverse T(3) hormone in immature rat sertoli cells: calcium uptake and exocytosis mediated by integrin. *PLoS One* 8:e77176 DOI 10.1371/journal.pone.0077176






Fig. 1






Fig. 2