

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effect of microwaves on the in situ hydrodistillation of four different Lamiaceae

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/143075> since

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in *COMPTES RENDUS CHIMIE*, 17, 2014, 10.1016/j.crci.2013.11.007.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.crci.2013.11.007

The definitive version is available at:

<http://www.journals.elsevier.com/comptes-rendus-chimie/>

Effect of microwaves on the *in situ* hydrodistillation of four different Lamiaceae.

Arianna Binello^a, Laura Orio^a, Giuseppe Pignata^b, Silvana Nicola^b, Farid Chemat^c, Giancarlo Cravotto^{*a}.

^a Dipartimento di Scienza e Tecnologia del Farmaco, Università di Torino, Via P. Giuria 9, 10235 Torino, Italy

^b Dipartimento AGRIFORFOOD, VEGMAP, Università di Torino, Via Leonardo da Vinci, 44, 10095 Grugliasco, TO, Italy

^c Université d'Avignon et Pays de Vaucluse, INRA, UMR 408, GREEN Extraction Team, F-84000 Avignon, France.

*Corresponding author Tel.: +39 011 670 7684; fax: +39 011 670 7687.

E-mail address: giancarlo.cravotto@unito.it

Abstract

The development of alternative techniques to classic hydrodistillation (HD) has been prompted because of the drawbacks of the more traditional technique. These drawbacks include: partial thermal degradation, high energy consumption and the fact that it is a time-consuming process. *In situ* microwave-generated hydrodistillation (MGH) and microwave hydrodiffusion and gravity (MHG) are suitable methods which may improve the preparation of essential oils. In this work, we report a comparison study of HD, MGH and MHG used in the extraction of four plant species cultivated in Piedmont (Italy): lavender, oregano, basil and sage. Both microwave-assisted procedures gave excellent results; in particular the essential oils obtained under MHG were very similar to those obtained with HD. In MGH, the polarity and the physico-chemical properties of the extracted compounds caused bigger differences in essential oil composition.

Keywords: *in situ* microwave-generated hydrodistillation, hydrodistillation, microwave hydrodiffusion and gravity, *Lamiaceae*, essential oils.

1. Introduction

The use of essential oils (EO) in the food and pharmaceutical industries has a long history. EO are a complex mixture of secondary metabolites, including terpenes, phenolics and alcohols. Although alternative processes are known, the vast majority of EO are produced by hydrodistillation under vapor steam. Hydrodistillation (HD) is a simple form of steam distillation which is often used to isolate non-water soluble, high boiling point and natural products. In HD, botanicals are fully submerged in water, producing a "soup", whose steam contains the aromatic compounds. The EO of the distillate are then extracted and analyzed. This is the most ancient method of distillation and the most versatile. The long treatment time and local overheating may cause partial degradation or hydrolysis resulting in EO with a burnt note and the loss of more volatile compounds [1]. HD seems to work best for powders (ie, spice powders, ground wood, etc.) and very tough materials like roots, wood or nuts.

Over the last few years, more laboratories have successfully applied microwave-assisted extraction (MAE) [2,3] and *in situ* microwave-generated hydrodistillation (MGH) [4], which are both now well recognized as innovative green and efficient techniques [5]. Dielectric heating is a non-classical energy source that is capable of reducing extraction times because of its selective volumetric heating that also gives excellent product quality [6,7]. In MGH, molecular motion within polar or ionic components causes heating. The rapid delivery of energy to the sample allows the temperature to rise rapidly in the plant cell and, after cell membrane collapse, the essential oil is released and carried away by the water vapor; the condensation and collection of the product, followed by EO isolation conclude the extractive operation [8]. Another green technique for EO extraction, microwave hydrodiffusion and gravity (MHG) developed by Chemat *et al.* [9,10], combines microwave heating and earth's gravity at atmospheric pressure. Thanks to hydrodiffusion and earth's gravity, the extract can drip out from the microwave reactor, falling through a perforated Pyrex disc and is cooled by a cooling system; water and essential oils are collected and easily separated into a suitable vessel, traditionally called the "Florentine flask".

The present work deals with Lamiaceae, cosmopolitan plants containing about 236 genera and 6900 to 7200 species. These plants are frequently aromatic in all parts and also are widely cultivated for decorative foliage or food use. Lavender, Greek oregano, sweet basil and sage contain fine essential oils used in cosmetics, food and pharmaceuticals, and were thus chosen for this study. In a previous work [11], we reported the results obtained with HD and MGH in the extraction of fresh and dry mint leaves. We herein report the results of a new comparison study on the effectiveness of MGH and MHG on four different genotypes (dry material): *Lavandula angustifolia* Mill. (lavender), *Origanum vulgare* L. ssp. *hirtum* (Link) Ietswaart (Greek oregano), *Ocimum basilicum* L. (sweet basil), *Salvia officinalis* L. ssp. *Lavandulifolia* (sage). Yields and the chemical profiles of the main components are described and commented on.

2. Results and discussion

2.1. Extraction yield and time

The percentage of dry plant material (% DM) is reported in Table 1. Oregano and basil leaves had lower dry matter levels than the leaves of the other genotypes tested, probably because *Origanum vulgare* and *Ocimum basilicum* have more tender leaves than the other species; in particular, oregano DM was twice as high as in previous experiments when leaves were harvested from much older plants [12]. *Salvia officinalis* L. ssp. *Lavandulifolia* plants were harvested at pre-blossoming stage and their DM were in line with previous experiments [13].

Table 1
Dry matter percentage (DM).

| | DM |
|---|-------|
| <i>Lavandula angustifolia</i> Mill. (leaves + flowers) | 24.85 |
| <i>Origanum vulgare</i> L. ssp. <i>hirtum</i> (Link) Ietswaart (leaves) | 12.65 |
| <i>Ocimum basilicum</i> L. (leaves) | 14.43 |
| <i>Salvia officinalis</i> L. ssp. <i>lavandulifolia</i> (leaves) | 22.50 |

HD, MGH and MHG EO yields are listed in Table 2. No striking differences in the three methods were found in oil yield for *Ocimum basilicum*. HD and MHG of *Origanum vulgare* and *Salvia officinalis* gave the best results, while MGH gave a better yield for *Lavandula angustifolia* distillation, probably due to the presence of the flowers. Considering the extraction times, MHG seems to be more efficient than HD, which is the reference method in EO extraction.

Table 2
HD vs MGH vs MHG EO yield (^w/_w% dry weight).

| | HD ^a | MGH ^b | MHG ^b |
|---|-----------------|------------------|------------------|
| <i>Lavandula angustifolia</i> Mill. (leaves + flowers) | 0.05 | 0.13 | 0.05 |
| <i>Origanum vulgare</i> L. ssp. <i>hirtum</i> (Link) Ietswaart (leaves) | 0.75 | 0.08 | 0.73 |
| <i>Ocimum basilicum</i> L. (leaves) | 0.06 | 0.01 | 0.06 |
| <i>Salvia officinalis</i> L. ssp. <i>lavandulifolia</i> (leaves) | 1.36 | 0.02 | 1.32 |

^a extraction time: 45 min, ^b extraction time: 20 min;

2.2. Composition of essential oils

The EO extracted by MGH, MHG and HD are all rather similar in their composition. In general, a similar number of volatile secondary metabolites was found in the essential oils techniques, but there are some differences in their relative amounts.

In the case of *Lavandula angustifolia* Mill. (Table 3), the compounds extracted via the three methods are roughly the same. A higher oxygenated monoterpene content was found under MHG. These include borneol (13.72%), linalool (8.14%), carvone (3.91%), criptone (3.56%), camphor (3.55%) and 4-terpineol (1.57%), whereas HD mainly leads to higher quantities of α -cadinol (3.95%), α -terpineol (2.14%), carvacrol (1.54) and phellandral (0.69%). MGH extracts highlight a high yield in coumarin (16.59%), 7-methoxycoumarin (11.81%) and caryophyllene oxide (4.26%). Borneol is the most abundant compound found in the MHG profile, while coumarin is the most abundant in HD and MGH.

Table 3Main components of *Lavandula angustifolia* Mill. L. dried material EO.

| Compound | HD (%) | MGH (%) | MHG (%) |
|-----------------------------------|--------|---------|---------|
| 1,8-Cineole | t | 1.22 | 1.35 |
| Linalool | 5.11 | 3.59 | 8.14 |
| Camphor | 1.42 | 1.69 | 3.55 |
| Borneol | 9.68 | 7.14 | 13.72 |
| 4-Terpineol | 0.99 | 0.49 | 1.57 |
| Cryptone | 2.71 | 1.39 | 3.56 |
| α -Terpineol | 2.14 | 0.74 | 1.86 |
| <i>trans</i> -Carveol | 0.56 | t | 0.66 |
| Cuminic aldehyde | 0.78 | 0.61 | 0.96 |
| Carvone | 2.03 | 2.11 | 3.91 |
| Phellandral | 0.69 | 0.39 | 0.57 |
| Carvacrol | 1.54 | 0.76 | 1.11 |
| Piperitenone | 0.47 | 0.28 | 0.43 |
| β -Caryophyllene | 0.43 | 1.29 | 0.33 |
| Coumarin | 10.52 | 16.59 | 9.41 |
| <i>trans</i> - β -Farnesene | 0.21 | 0.25 | 0.15 |
| β -Ionone | 0.59 | 0.45 | 0.52 |
| Dihydroactinidiolide | 1.10 | 0.94 | 0.44 |
| Caryophyllene oxide | 2.48 | 4.26 | 2.90 |
| α -Cadinol | 3.95 | 3.68 | 3.01 |
| 7-Methoxycoumarin | 2.93 | 11.81 | 1.66 |

Linalool, carvacrol and 4-terpineol are the major components in *Origanum vulgare* L. ssp. *hirtum* (Link) Ietswaart dried leaves (Table 4), but their abundance is different in HD, MGH and MHG essential oils. While linalool is the main compound in MHG (26.66%), carvacrol appears to be the most abundant in MGH and HD (35.01% and 26.70% respectively), and the quantity of 4-terpineol is significantly more abundant in MGH (18.05%) than in HD (7.44%) and MHG (4.99).

trans-Sabinene hydrate, *p*-cymene and linalyl acetate amounts are higher in HD and MHG but, overall, MGH extraction seemed to be more selective, although in a lower oil yield.

Table 4Main components of *Origanum vulgare* L. ssp. *hirtum* (Link) Ietswaart dried leaf EO.

| Compound | HD (%) | MGH (%) | MHG (%) |
|-----------------------------------|--------|---------|---------|
| α -Thujene | 0.30 | t | 0.25 |
| β -Pinene | 1.78 | t | 0.15 |
| Sabinene | 1.68 | t | 1.63 |
| β -Myrcene | 0.81 | 0.11 | 0.87 |
| α -Phellandrene | 0.14 | t | 0.12 |
| α -Terpinene | 2.63 | 0.95 | 1.58 |
| <i>p</i> -Cymene | 2.48 | 0.64 | 2.32 |
| β -Phellandrene | 1.14 | 0.28 | 1.11 |
| γ -Terpinene | 6.02 | 4.15 | 4.51 |
| <i>trans</i> -Sabinene hydrate | 3.57 | 1.17 | 4.57 |
| α -Terpinolene | 1.31 | 1.08 | 0.63 |
| Linalol | 21.92 | 6.80 | 26.66 |
| 1-Octen-3-yl acetate | 0.79 | 0.73 | 0.93 |
| Borneol | 0.10 | t | 0.12 |
| 4-Terpineol | 7.44 | 18.05 | 4.99 |
| α -Terpineol | 2.13 | 3.67 | 2.27 |
| Carvacrol methyl ether | 0.11 | 0.01 | 0.12 |
| Linalyl acetate | 3.93 | 0.88 | 4.50 |
| Thymol | 0.20 | 0.88 | 0.27 |
| Carvacrol | 26.70 | 35.01 | 22.80 |
| β -Caryophyllene | 2.08 | 2.33 | 2.28 |
| α -Humulene | 0.30 | 0.32 | 0.30 |
| Germacrene D | 1.46 | 1.66 | 1.77 |
| Bicyclogermacrene | 1.39 | 1.70 | 1.66 |
| β -Bisabolene | 2.78 | 3.12 | 2.94 |
| γ -Cadinene | 0.45 | 0.41 | 0.38 |
| δ -Cadinene | 0.64 | 0.83 | 0.68 |
| Dihydroactinidiolide | 0.13 | 0.12 | 0.24 |
| <i>cis</i> - α -Bisabolene | 0.10 | t | 0.13 |
| Viridiflorol | 0.13 | 0.24 | 0.14 |

Linalool, eugenol and α -cadinol are the main components in the EO of *Ocimum basilicum* L.

leaves. The relative amounts of the main compounds detected are reported in Table 5.

Linalool is the most abundant component in all three extracts, with preponderance in MHG (22.96%) and in MGH (24.68%). A similar, but not identical, trend can be seen for eugenol for which HD, MGH and MHG extracts show 13.57%, 15.19% and 15.93%, respectively; in this case the difference between the two non conventional techniques is very low.

MW irradiation of oregano that typically has a low moisture content, may generate localized hot-spots. These high-energy microenvironments may promote the conversion of linalool to 4-

terpineol, in particular in the case of MGH. In table 4, it is possible to observe a higher content of 4-terpineol, the typical degradation product of linalool *via* the pathway described in figure 1.




Figure.1. Conversion of linalool to 4-terpineol

As far as α -cadinol is concerned, extractive efficacy is quite similar for each method (HD 9.60%, MGH 9.88%, MHG 9.93%). The greatest yield in 1,8-cineole was detected in MHG EO (6.77 %), while HD seems to be slightly better for some alcohols, such as borneol, α -terpineol and geraniol. The gas chromatograms obtained from the *Ocimum basilicum* extracts, therefore, overlap quite extensively.

Table 5
Main components of *Ocimum basilicum* dried leaf EO.

| Compound | HD (%) | MGH (%) | MHG (%) |
|-----------------------------|--------|---------|---------|
| 1,8-Cineole | 2.49 | 2.38 | 6.77 |
| <i>cis</i> -Linalol oxide | 1.32 | 1.12 | t |
| <i>trans</i> -Linalol oxide | 1.13 | 0.93 | t |
| Linalool | 18.72 | 24.68 | 22.96 |
| Camphor | 0.62 | 0.56 | 0.47 |
| Borneol | 2.83 | 2.48 | 1.69 |
| α -Terpineol | 2.03 | 1.81 | 1.10 |
| Geraniol | 1.16 | 0.52 | t |
| Bornyl acetate | 1.16 | 1.57 | 1.86 |
| Eugenol | 13.57 | 15.19 | 15.93 |
| β -Elemene | 1.66 | 1.11 | 1.31 |
| α -Bergamotene | 0.94 | 1.06 | 1.13 |
| Germacrene-D | 1.91 | 1.56 | 1.75 |
| γ -Cadinene | 2.60 | 2.39 | 2.38 |
| α -Cadinol | 9.60 | 9.88 | 9.93 |

For *Salvia officinalis* L. ssp. *Lavandulifolia* dried leaf EO (Table 6), HD and MHG have comparable profiles and the same compounds were found in similar relative amounts; camphor (17.60% and 15.31% HD and MHG, respectively), α -thujone (12.61% and 12.12%) and 1,8-cineole

(11.61% and 10.25%) are the most abundant compounds; a higher percentage of α -pinene was found in HD EO (5.60% vs 2.96% MHG), while manool is more abundant in MHG (6.31% vs 2.53% HD).

The low MGH extract yield is also reflected in the limited number of compounds present in the oil; the relative abundance of the extracted compounds is rather peculiar: in fact, viridiflorol (16.39%) and manool (28.14%) are by far the most concentrated.

Table 6

Main components of *Salvia officinalis* L. ssp. *Lavandulifolia* dried leaf EO.

| Compound | HD (%) | MGH (%) | MHG (%) |
|-------------------------|--------|---------|---------|
| α -Pinene | 5.60 | t | 2.96 |
| Camphene | 3.72 | t | 2.26 |
| β -Pinene | 0.59 | t | 0.39 |
| β -Myrcene | 0.73 | t | 0.53 |
| 1,8-Cineole | 11.61 | t | 10.25 |
| γ -Terpinene | 0.13 | t | 0.14 |
| α -Terpinolene | 0.32 | t | 0.28 |
| α -Thujone | 12.61 | t | 12.12 |
| β -Thujone | 4.60 | t | 4.42 |
| Camphor | 17.60 | 0.77 | 15.31 |
| Borneol | 3.62 | t | 3.68 |
| 4-Terpineol | 0.73 | t | 0.40 |
| α -Terpineol | 0.86 | t | 1.14 |
| Myrtenol | 0.72 | t | 0.65 |
| Carvone | 0.18 | t | 0.28 |
| Bornyl acetate | 1.49 | t | 1.50 |
| Carvacrol | 1.04 | 2.64 | 0.20 |
| γ -Caryophyllene | 0.08 | t | 0.08 |
| β -Caryophyllene | 6.07 | 5.39 | 5.39 |
| α -Humulene | 8.33 | 7.17 | 5.77 |
| Viridiflorene | 0.36 | t | 0.28 |
| Caryophyllene oxide | 0.61 | 4.55 | 1.00 |
| Viridiflorol | 5.42 | 16.39 | 5.89 |
| Manool | 2.53 | 28.14 | 6.31 |

The differences in the three extraction methods confirm that microwaves are able accelerate the extraction processes and afford high quality oils.

2.2.1. Evaluation of MGH and MHG



Essential oils can be classified into two main groups: terpene containing hydrocarbons, such as monoterpenes, sesquiterpenes and diterpenes; and oxygenated compounds, such as esters,

aldehydes, ketones, alcohols, phenols, oxides, acids and lactones. During distillation, aromatic plants, treated with boiling water or steam, release their essential oils via the vapor distillation of immiscible liquids, namely water and EO. Thus the essential oil components, whose boiling points normally range from 200 to 300°C, are evaporated at temperatures close to that of boiling water. The amount of essential oil produced depends on four main factors: the distillation time, the temperature, the operating pressure and, most of all, the type and obviously the quality of the plant material. EO composition is strongly affected by water solubility and dielectric loss properties of the different compounds rather than their boiling point [14]. Microwaves have the peculiarity of deeply penetrating the vegetal matrix, reducing heating times and thus it is a fast and green technology for the solvent-free extraction of naturally occurring aromatic compounds.

By analyzing the results of HD and MGH of *Origanum vulgare* L., we can compare the relative abundance of linalool (a polar compound) and γ -terpinene (a non-polar compound) (Table 7).

Table 7

MHG “selective” extraction of linalool vs γ -terpinene in *Origanum vulgare* L. ssp. *hirtum* (Link) Ietswaart dried leaves.

| | Linalool | γ -Terpinene |
|---------------------------|---|---|
| Structure |  |  |
| % MHG | 26.66 | 4.51 |
| % HD | 21.92 | 6.02 |
| Boiling point (°C) | 220 | 183 |
| Solubility in water (g/L) | 0.57 | 0.001 |
| Dipolar moment (D) | 22 | 13 |
| Dielectric constant | 1.98 | 0.72 |

Most likely, the presence of a continuous vapor stream during the MHG process confirms the main role of the “hydrodiffusion” phenomenon in the extract composition that may intensify the effect of microwave irradiation. With MHG it is possible to obtain EO in a more concentrated volume and shorter heating time.

2.3. Cost, energy and environmental impact

The results obtained from this study have confirmed that MGH and MGH extractions are more advantageous than HD in terms of overall costs and energy. As previously reported, with our instrumentation the energies required for HD and MGH are 1.5 kWh and 0.25 kWh respectively, these figures correspond to a carbon footprint of 1200 and 200 g CO₂ for HD and MGH, respectively [11,15]. As reported by Chemat *et al.* [9] the energy required to perform an extraction with MHG is 0.2 kWh and the calculated quantity of carbon dioxide emitted is 160 g. In a recent paper we successfully experimented a 75 L-pilot MW reactor demonstrating the scaling up feasibility of MGH and MGH extractions [16]. Improvements in product efficiency, energy saving, process enhancement and maintenance costs, place this technology in a first rank for process intensification of plant eco-extraction.

3. Conclusion

The constant evolution of microwave-assisted extraction techniques strongly enhance the quality of EO and of the process itself giving it a lower environmental impact.

EO composition profiles from MHG and HD are normally super-imposable, while different selectivity occurs in MGH because it uses a lower amount of steam.

4. Experimental

4.1. Plant materials

All the plants came from the experimental centre at the Agriforfood Department of the University of Turin (Grugliasco, TO, Italy). The studied genotypes were transplanted in the field from rooted

cuttings grown in greenhouses at the nursery stage in spring 2009 (*Salvia officinalis* L. ssp. *lavandulifolia*) or spring 2011 (*Lavandula angustifolia* Mill., *Origanum vulgare* L. ssp. *hirtum* (Link) Ietswaart) or from seedlings grown in greenhouses at the nursery stage in spring 2011 (*Ocimum basilicum* L.). All cuttings were obtained from same mother plant genotype, while for sweet basil, the seeds were obtained from the same commercial seed lot. The plants were grown in the same environment, in the experimental centre (44°53'11.67''N; 7°41'7.00''E–231 m a.s.l., Carmagnola, TO, Italy), and soil (sandy-loam, according to the USDA criteria). Plant density for each species was ca. 4 plant/m² and plants were grown according to the standard cultural practices used in the local area. Harvesting took place and was scalar according to the balsamic time from 25 June to 8 September 2011. Dried samples were obtained through an oven-drying process as set out in previous experiments and described in Tibaldi *et al.* [13,17]. EO content was calculated as the ratio in weight (w/w) between the cold oil collected from the burette and the original weight of the material.

4.2. Hydrodistillation (HD)

Essential oil extraction via HD was carried out according to Tibaldi *et al.*[12,13,17] with some slight modifications. The equipment used was assembled by Exacta + Optech (San Prospero, MO, Italy), namely an electrical heating mantle (500 W; Electro Thermal, Thermo Scientific, Waltham, MA, USA), a 2-L Pyrex[®] round flask (with 1.5 L deionized water), a 4-L modified Pyrex[®] glass round flask filled with the plant material, a condenser and a graduated burette to collect the oil.

An aliquot of 100-300 g dried material was placed in the steam distillation system. Each distillation lasted 105 min, including 45 min in which the leaves or flowers were steamed by the boiling water in the glass balloon, previous tests with a longer heating time did not increase EO yield. The 45 min distillation time started when the first drop of liquid, condensed in the cooling column, dropped into the graduated burette.

4.3. In situ Microwave-generated hydrodistillation (MGH)

MGH was performed in a NEOS-GRMW oven (Milestone srl, Sorisole, BG, Italy).

Specific equipment and software, a videocamera and an infrared pyrometer were used.

Extractive conditions were chosen on the base of a series of preliminary tests aimed to optimize yields and to limit the formation of degradation products.

The plant material was placed in a 4 L beaker, covered with a polypropylene lid with a central hole connected via a glass tube to the external condenser. The latter was connected to a chiller, with a circulating fluid maintained at around 5-6°C. An aliquot of 100-300 g dried material was placed in a 4 L Pyrex[®] glass beaker and put in the microwave oven. Dried leaves or flowers were previously rewetted under slight steam flow by means of a commercial domestic device ('Vaporetto Polti – Italy') equipped with a flow regulator (calculated swelling: 2 mL g⁻¹).

Each distillation lasted 45 min, including 20 min in which the material was steamed for plant rehydration. The 20 min distillation time started when the first drop of liquid, condensed in the cooling column, dropped into the graduated burette.

4.4. Microwave Hydrodistillation and gravity (MHG)

MHG was performed in a NEOS-GRMW oven (Milestone srl, Sorisole, BG, Italy), under conditions defined in a series of preliminary tests.

100-300 g of dried leaves or flowers are placed on a grid and put in a modified 4 L Pyrex[®] glass beaker with a hole at the bottom, put in the microwave oven and connected to an external round bottom flask with a condenser. The latter was connected to a chiller, with circulating fluid, maintained at around 5–6°C. Dried materials were previously rewetted under slight steam flow by means of a commercial domestic device ('Vaporetto Polti – Italy') equipped with a flow regulator (calculated swelling: 2 mL g⁻¹) and kept wet with this steam flow during microwave irradiation. Each distillation lasted 45 min, including 20 min in which the leaves were steamed for the plant

rehydration. The 20 min distillation time started when the first drop of liquid, condensed in the cooling column, dropped into the round bottom flask below the oven.

4.5. Analytical apparatus

Equipment and analysis conditions are as previously described [11]. All gas chromatographic mass spectra were recorded on an Agilent 6850 gas chromatograph (Agilent Technologies, Palo Alto, CA, USA), fitted with an MS Agilent Network 5973, which uses a 30 m long capillary column HP 5-MS (5% phenyl methyl siloxane, i.d. 0.25 mm, film thickness 0.25 μm), a 1:20 split ratio, a 250°C injector temperature and helium as the carrier gas (1.2 mL min⁻¹ flow).

The gas chromatographic parameters were set up as follows; initial temperature 50°C, rate 3°C min⁻¹, final temperature 80°C, rate 10°C min⁻¹, final temperature 300°C, held for 10 min. The mass spectrometric parameters were as follows; low mass 40, high mass 800, MS quad 150°C, MS source 230°C.

4.6. Sample preparation

The samples were prepared by dissolving 0.04 mL essential oil in 1 mL chloroform.

The samples (1 μL) were injected by an automatic injector (Agilent 7683B series).

The identification of products was achieved using the NIST 05 library (National Institute of Standards and Technology), Wiley Library and confirmed by comparing the retention times with those of the relative standard.

4.7. Chemicals

Gas chromatographic grade chloroform was purchased from Sigma Aldrich (Milan, Italy).

Acknowledgements

The present work was supported by the ALCOTRA project “Eco-extraction transfrontaliere”.

References

- [1] M.E. Lucchesi, F. Chemat, J. Smadja, *J. Chromatog. A* 1043 (2004) 323.
- [2] S. Périno-Issartier, C. Ginies, G. Cravotto, F. Chemat, *J. Chromatogr. A*, 1305 (2013) 41.
- [3] Microwave-assisted extraction for bioactive compounds: Theory and practice. Ed.: F. Chemat G. Cravotto (2013), XII, 238 pp. Springer Science, New York, U.S.A..
- [4] S. Rezvanpanah, K. Rezaei, S.H. Razavi, S. Moini, *Food Sci. Technol. Res.* 14(3) (2008) 311.
- [5] F. Chemat, M. Abert-Vian, G. Cravotto, *Int. J. Mol. Sci.*, 13 (2012) 8615.
- [6] A. Farhat, C. Ginies, M. Romdhane, F. Chemat, *J. Chromatog. A* 1216 (2009) 5077.
- [7] B. Uysal, F. Sozmen, B.S. Buyuktas, *Nat. Prod. Commun.* 5 (2010) 111.
- [8] S. Chemat, H. Ait-Amar, A. Lagha, D.C. Esveld, *Chem. Eng. Process.* 44 (2005) 1320.
- [9] M.A. Vian, X. Fernandez, F. Visinoni, F. Chemat, *J. Chromatog. A* 1190 (2008) 14.
- [10] N. Bousbia, M.A. Vian, M.A. Ferhat, E. Petitcolas, B.Y. Meklati, F. Chemat, *Food Chem.* 114 (2009) 355.
- [11] L. Orio, G. Cravotto, A. Binello, G. Pignata, S. Nicola, F. Chemat, *J. Sci. Food Agric.* 92 (2012) 3085.
- [12] G. Tibaldi, E. Fontana, S. Nicola, *Ind. Crops and Prod.* 34 (2011) 1516.
- [13] G. Tibaldi, E. Fontana, S. Nicola, *J. Food Agric. Envir.* 8(3&4) (2010) 790.
- [14] P. Sandra, C. Bicchi, A. Koedam, *Capillary gas chromatography in essential oil analysis*, Hueting Verlag, New York, 1987.
- [15] M.A. Ferhat, B.Y. Meklati, F. Chemat, *Flavour Fragr. J.* 22 (2007) 494.
- [16] A. Filly, X. Fernandez, M. Minuti, F. Visinoni, G. Cravotto, F. Chemat, *Food Chem.* (2013), <http://dx.doi.org/10.1016/j.foodchem.2013.10.139>
- [17] G. Tibaldi, E. Fontana, S. Nicola, *J. Sci. Food Agric* 92 (2013) 580.