

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Metalloproteinases and Metalloproteinase Inhibitors in Age-Related Diseases.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/149038> since 2016-10-06T12:27:25Z

Published version:

DOI:10.2174/13816128113196660701

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Current Pharmaceutical Design 20(18):2993-3018; 2014. DOI: 10.2174/13816128113196660701

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://www.eurkaselect.com/115822/article>

Metalloproteinases and Metalloproteinase Inhibitors in Age-Related Diseases

Simona Gargiulo, Paola Gamba, Giuseppe Poli and Gabriella Leonarduzzi

Department of Clinical and Biological Sciences, University of Turin, Orbassano (Turin), Italy

*Address correspondence to this author at the Department of Clinical and Biological Sciences, Faculty of Medicine San Luigi Gonzaga, University of Turin, Regione Gonzole 10, 10043, Orbassano (Turin), Italy. Tel. +39 011 6705422; Fax: +39 011 6705424;

E-mail: giuseppe.poli@unito.it (G. Poli).

Running title: **Metalloproteinases in Age-Related Diseases**

Abstract: Degradation of the extracellular matrix is an important feature of embryonic development, morphogenesis, angiogenesis, tissue repair and remodeling. It is precisely regulated under physiological conditions, but when dysregulated it becomes a cause of many diseases, including atherosclerosis, osteoarthritis, diabetic vascular complications, and neurodegeneration. Various types of proteinases are implicated in extracellular matrix degradation, but the major enzymes are considered to be metalloproteinases such as matrix metalloproteinases (MMPs) and disintegrin and metalloproteinase domain (ADAMs) that include ADAMs with a thrombospondin domain (ADAMTS). This review discusses involvement of the major metalloproteinases in some age-related chronic diseases, and examines what is currently known about the beneficial effects of their inhibitors, used as new therapeutic strategies for treating or preventing the development and progression of these diseases.

Keywords: metalloproteinases, metalloproteinase inhibitors, atherosclerosis, diabetes mellitus, neurodegenerative diseases, osteoarthritis

1. INTRODUCTION

Metalloproteinases are a large family of important endopeptidases, which include matrixins (matrix metalloproteinases, or MMPs) and adamalysins (a disintegrin and metalloproteinase domain, or ADAMs) [1, 2].

The subunits of ADAMs, which are transmembrane proteases, comprise a catalytic domain at the end of the extracellular extension, which comprises three domains: a disintegrin, a cysteine-rich domain, and a number of epidermal growth factor repeats. The cytoplasmic tail attached to the epidermal growth factor domain protrudes through the membrane, and signals cell-surface events to the cytoplasm [3]. The catalytic domain contains a typical zinc-binding consensus motif; for example, in the case of ADAM-10, HEXGHXXGXXHD.

As the name “ADAM” indicates, the disintegrin or integrin-binding domain binds these molecules to the membrane integrins, and the metalloproteinase domain provides the protease function [4]. Thus ADAMs are important in events at the cell surface, because they attach to integrins on the cell surface and carry out metalloproteinase functions. Their primary functions are cleavage of extracellular matrix (ECM) molecules and of the extracellular domains of many cell-surface membrane proteins, a process known as “ectodomain shedding” [4-6]. Shedding of the tumor necrosis factor- α (TNF- α) receptor, and of interleukin-6 (IL-6), L-selectin, and syndecans, has also been shown to be a function of ADAMs [7]. They are also important in intracellular signaling and cell adhesion [8]. ADAMs are thus implicated in cell proliferation, migration, differentiation, and survival [9]. Of the more than 30 members of this family, only ADAM-10, -15, and -17 have been characterized in vascular cells [10]. ADAM-10 has also been identified in distinct areas of the human brain [11, 12] and peripheral structures [13, 14] (Table 1). ADAMs with a thrombospondin domain (ADAMTS) form another group of metalloproteinases. ADAMTS-4 and ADAMTS-5 have been shown to degrade aggrecan, a proteoglycan of joint cartilage ECM, and to be involved in spinal cord injury [15, 16]. These enzymes are regulated at multiple levels, through control of gene expression, mRNA splicing, and protein processing, as well as through regulation of the expression of various naturally-occurring inhibitors.

With regard to MMPs, there are at least 25 mammalian MMPs, 14 of which have been characterized in vascular cells. Although mammalian MMPs have overlapping specificities for structural ECM components, they are classified into five groups, by differences in their primary structure and substrate specificity. These groups are: interstitial collagenases, gelatinases, stromelysins/matrilysins, membrane-type MMPs (MT-MMPs), and others MMPs. Further, they are assigned MMP numbers, and some members also have trivial names (Table 2). MMPs are extracellular proteins, but recent studies have reported that MMP-1 [17], MMP-2 [18] and MMP-11 [19] are intracellular, and may act on intracellular proteins. Moreover, most MMPs are expressed as inactive, latent proforms, although MMP-11, -21, -23, and -27, and the MT-MMPs, have a furin recognition sequence before the catalytic domain, and are therefore likely to be activated intracellularly and secreted as active enzymes [20, 21].

The protein structure of MMPs follows a basic pattern. A typical MMP consists of a propeptide of about 80 amino acids at the N-terminal, which is attached to a signal peptide, a catalytic metalloproteinase domain of about 170 amino acids, a linker peptide of variable length (also called the “hinge region”) and a hemopexin domain of about 200 amino acids, which may be attached to a transmembrane domain. Exceptions to this are MMP-7, MMP-26 and MMP-23; these MMPs lack the linker peptide and the hemopexin domain, and MMP-23 has a unique cysteine-rich domain and an immunoglobulin-like domain after the metalloproteinase domain. The two gelatinases, MMP-2 and MMP-9, have three repeats of a fibronectin type II motif in the metalloproteinase domain. Further, MMPs have a zinc binding motif - HEXXHXXGXXH - in the catalytic domain, and a “cysteine switch” motif - PRCGXPD - in the propeptide: the three histidines in the zinc binding motif, and the cysteine in the propeptide, coordinate with the catalytic zinc ion. This Cys-Zn²⁺ coordination keeps proMMPs inactive, preventing catalysis of the zinc atom by water-molecule binding. The

catalytic domain also contains a conserved methionine, forming a “Met-turn” situated eight residues after the zinc binding motif, whose role is to support the structure around the catalytic zinc. The zinc binding motif and the “Met-turn” are also conserved in ADAM family members [20, 21].

Traditionally, MMPs are thought to play a central role in degrading and remodeling ECM components, including fibronectin, laminin, proteoglycans and collagen [21-23]. In recent years, however, findings from several research groups have established that MMPs cleave a wide range of extracellular and bioactive ECM and non-ECM substrates, and that they regulate the activity of these proteins [24, 25]. The established functions of MMPs include releasing cytokines and growth factors from the cell membrane or ECM, cleaving growth factor receptors from the cell surface, activating proforms of cytokines (*e.g.* TNF- α and IL-1 β), activating death receptors at cell membranes, shedding cell adhesion molecules, and activating other MMPs and MMP inhibitors, as well as other signaling molecules [24-28]. They are also involved in regeneration, myelin formation, angiogenesis, and axonal growth [29]. The ability of MMPs to modify the structural integrity of tissues is thus essential for certain physiological and pathological processes: MMPs are potent controllers of physiological processes (cell migration, proliferation, differentiation, growth and development) and of pathological processes (tissue remodeling in response to injury, inflammatory processes, neovascularization, cancer progression, apoptosis, etc.) [21, 30]. In this context, and in brief, MMPs are important in normal development, remodeling, wound healing, and also in a wide variety of pathological processes, including the spread of metastatic cancer cells, arthritic destruction of joints, cardiovascular and neurodegenerative diseases, diabetes mellitus, and lung diseases. In these latter pathological processes, which are characterized by an inflammatory response, the abnormal expression and activation of these proteases lead to ECM breakdown.

As cited above, MMPs are secreted as latent enzymes and require activation, which is tightly regulated so as to prevent tissue damage. The activities of most MMPs are very weak or negligible in normal steady-state tissues, but rapidly increase in response to inflammatory and oxidative stimuli. Their activity can be regulated at four levels: induction of MMP genes, vesicle trafficking and secretion, activation of latent proforms, and complexing with specific endogenous tissue inhibitors of metalloproteinases (TIMPs). Activation of MMPs is, thus, an important regulatory step in MMP activity.

MMPs are synthesized as pre-proenzymes, but during translation the signal peptide is removed, generating proMMPs; thus most MMPs are secreted from the cell in the form of proMMPs. MMPs may be stored intracellularly in vesicles, or secreted directly into the intracellular space. The presence of a proteinase-susceptible “bait” region in the propeptide allows tissue and plasma proteinases or opportunistic bacterial proteinases to activate proMMPs. A small number of MMPs, including the membrane-bound MT-MMPs, are proteolytically activated inside the cell by furin proteases, but most MMPs are activated in the extracellular space. Besides proteolytic cleavage, a change in configuration of the propeptide region can activate the enzymes. The proMMPs thus possess a furin-like proprotein convertase recognition sequence, RX[K/R]R, at the end of the propeptide, and are likely to be activated intracellularly and then secreted or cell-surface-bound as active enzymes. ProMMPs can also be activated by oxidants, such as reactive oxygen species (ROS) and nitric oxide (NO) [31] by reacting with the cysteine of the “cysteine switch” in the propeptide, and this activation process takes place under inflammatory and oxidative conditions [32]. Moreover, they can be activated by lysosomal proteases [20], by urokinase type plasminogen activators [33], by angiotensin [34] or by hyperglycemia [35]. In addition, control of the inducible MMP genes can occur at the promoter region, which contains binding sites for transcription factors such as activator protein-1 (AP-1), nuclear factor- κ B (NF- κ B), and polyoma enhancer A binding protein-3 (PEA-3); in turn, these are responsive to free radicals, protein kinases, and cytokines,

suggesting that these genes may be induced during inflammation [36, 37]. Activated MMPs can also activate other MMPs, in a stepwise activation cascade [20, 30, 38, 39].

Thus MMP activities are regulated by two major types of endogenous inhibitors: α_2 -macroglobulin and TIMPs.

Human α_2 -macroglobulin is a plasma glycoprotein of 725 kDa that inhibits most proteinases by entrapping the proteinase within the macroglobulin, after proteolysis of the bait region of the inhibitor; the complex is rapidly cleared by low density lipoprotein (LDL) receptor-related protein-1-mediated endocytosis. MMP activities in the fluid phase are primarily regulated by α_2 -macroglobulin. MMP-1 reacts with α_2 -macroglobulin more readily than with TIMPs [40].

TIMPs are potent and selective tissue inhibitors of MMPs, consisting of 184-194 amino acids [21, 41]. They are subdivided into an N-terminal and a C-terminal subdomain. Each domain contains three conserved disulfide bonds, and the N-terminal domain folds as an independent unit with MMP inhibitory activity. The N-terminal cysteine is particularly important for inhibition, since its free α -amino group and carbonyl function displace the catalytic water molecule from the essential Zn^{2+} ion at the MMP active site. There are at least four members of the TIMP family (TIMP-1, -2, -3, and -4) that are often secreted by the same cells that secrete MMPs; their expression is closely regulated during embryonic development and tissue remodeling [30, 42]. The four members of the TIMP family have many similarities and overlapping specificities, but their biochemical properties and local expression patterns are distinctive [41]. TIMP-1, -2, and -4 are secreted in soluble form, while TIMP-3 is associated with the ECM. Moreover, their activity is stimulated by platelet-derived growth factor (PDGF) and tumor growth factor β (TGF β) and is regulated by several cytokines. TIMPs form tight inhibitory 1:1 complexes with MMPs [42]. These interactions are generally rather non-selective, meaning that TIMPs inhibit all MMPs, at least to some extent; however, certain TIMPs have weaker or stronger inhibitory effects on specific proteinases [42]. TIMP-1 mainly inhibits MMP-9 as well as MMP-3, while it only weakly inhibits MMP-14 (MT1-MMP), MMP-16 (MT3-MMP), MMP-18 (MT5-MMP) and MMP-19; TIMP-2 inhibits MMP-2 and, paradoxically, at low concentrations contributes to activating proMMP-2. TIMP-3 is the only TIMP bound to the ECM and cell surface. TIMP-3 inhibits several membrane-bound molecules with sheddase functions, such as MMP-3, MMP-7, and MMP-14. ADAMs also contain an MMP-like catalytic domain, which in some cases remains catalytically active. In general, TIMPs do not bind or inhibit the catalytic site of ADAMs, although TIMP-3 inhibits ADAM-10 and ADAM-17, and TIMP-1 inhibits ADAM-10 [43]. TIMP-4 is chiefly localized in vascular tissue [21].

Several other proteins have been reported to inhibit selected members of the MMP family, although the inhibition mechanism of these protein is still unclear: the secreted form of amyloid β precursor protein (APP) inhibits MMP-2 [44]; a C-terminal fragment of procollagen C-proteinase enhancer protein inhibits MMP-2 [45], and RECK (reversion-inducing cysteine-rich protein with kazal motifs), a GPI-anchored glycoprotein that suppresses angiogenesis, inhibits MMP-2, MMP-9 and MMP-14 [41, 46]. Moreover, MMP expression is determined at a transcriptional level by various cytokines and growth factors [47]. In a number of tissue types, some cytokines and growth factors, including IL-1, TNF- α and PDGF, stimulate MMP expression, while others are inhibitory, *e.g.* TGF β .

The balance between production, activation, and inhibition of metalloproteinases is critical in maintaining ECM integrity. When proteolytic activity is greater than inhibition caused by TIMPs or other inhibitors, ECM breakdown occurs. Conversely, if inhibitors are too strongly expressed and proteolysis is restricted, there is a build up of ECM proteins, with fibrosis.

2. METALLOPROTEINASES IN AGE-RELATED DISEASES

2. 1. Metalloproteinases in Atherosclerosis

Uncontrolled ECM remodeling of the myocardium and vasculature, by MMPs and other proteolytic enzymes, are features of cardiovascular disorders such as atherosclerosis, stroke, stenosis, left ventricular hypertrophy, heart failure, and aneurysm [20, 48-51].

A number of MMP gene polymorphisms have also been shown to contribute to inter-individual susceptibility and outcome of these cardiovascular disorders. Genetic polymorphisms may, for example, affect MMP expression levels by conferring protection or propensity to vulnerable plaques [52-54]. Moreover, studies using MMP gene knockout mice have indicated that MMP-2 and MMP-9 play key roles in cardiac rupture after myocardial infarction [55, 56]. A critical role of MMP-2 and MMP-9 has also been demonstrated in the development of abdominal aortic aneurysm, using MMP gene deletion mice [57]. MMPs, in particular MMP-1 and MMP-2, might also be involved in triggering acute coronary syndrome, via their ability to promote platelet activation and aggregation [58, 59]. Finally, *Timp-3* deficiency in mice disrupts matrix homeostasis, and causes spontaneous left ventricular dilation, cardiomyocyte hypertrophy, and contractile dysfunction [60].

With regard to the multifactorial disease atherosclerosis, it is characterized by the development of atherosclerotic plaques in susceptible sites of the arterial wall. It is initiated by cholesterol-rich lipid retention and accumulation, oxidation, and modification, which combine to provoke chronic inflammation. Atherosclerotic plaque growth, due to lipid accumulation, smooth muscle cells (SMCs) proliferation, and matrix synthesis, may in turn narrow the arterial lumen and ultimately causing stenosis or thrombosis [61]. The final clinical outcome depends on whether a plaque becomes unstable, leading to acute disruption of the surface and exposure of the thrombogenic core to luminal blood flow [62]. In this context, our understanding of the pathogenesis of atherosclerotic lesions has improved dramatically, helping to clarify the mechanisms of plaque formation and the differences between stable and unstable plaque morphology [63-66]. In particular, it has now been established that inflammation is a key feature in all stages of the disease, especially in plaque destabilization, which leads to plaque rupture [67]. Alongside the inflammatory response, which triggers activation of macrophage-derived MMPs, also apoptosis of vascular cells, especially that of macrophages and SMCs, contributes to plaque destabilization [68, 69].

A variety of intrinsic and extrinsic factors predisposes an atherosclerotic plaque to instability and acute disruption. Intrinsic factors characterizing a plaque as vulnerable are: a large lipid core, increased inflammatory-cell infiltration (particularly by monocytes/macrophages), content of foam cells and of T lymphocytes, and reduced collagen and vascular SMCs content associated with increased matrix proteolysis. Extrinsic features include increased blood pressure, hemodynamic shear stress, and vasospasm [62, 70-74]. Plaque rupture tends to occur at the shoulder region, which is associated with cap thinning and macrophage infiltration [75]; the shoulder region is also the area of the plaque exposed to the greatest shear stress [74]. Macrophages control many of the inflammatory processes within the plaque, and are the principal cells responsible for the production of MMPs [49]. MMPs are the predominant proteolytic enzymes, thought to participate in weakening the connective tissue matrix in the intima, leading to plaque rupture and acute thrombosis [76, 77]. Of note, human peripheral blood monocytes express MMP-8, -9, -14 and -19, TIMP-1 and TIMP-2 constitutively, but activated macrophages and foam cells express high levels of several MMPs, in response to adhesion and pro-inflammatory mediators, either directly, through mitogen-activated protein kinase (MAPK) activation and the NF- κ B pathway, or indirectly, with prostaglandin E₂ (PGE₂) involvement [78, 79]. The upregulation of MMP secretion by differentiated macrophages can be stimulated by contact with endothelial cells (ECs) [80] and by adhesion to matrix components, such as collagen [81]. Contact-mediated upregulation of MMPs seems therefore to be the first stage of MMP induction in activated macrophages. Intracellular accumulation of lipids, typical of foam cells, thus increases MMP expression in macrophages [82, 83] and, in turn, foam cells produce ROS that trigger the transformation

of pro-MMP into active MMP. Several pro-atherogenic cytokines and growth factors, including IL-1 β , TNF- α , macrophage colony-stimulating factor (M-CSF) and PDGF respectively, as well as other inflammatory mediators increase macrophage MMP expression [78, 84]. Moreover, immune cells might upregulate MMPs in activated macrophages, including MMP-1, -3, -8 and -11, and MMPs co-localize with CD40 expression in atherosclerotic plaques [85-88]. Moreover, it has been demonstrated that MMP secretion can be induced by both innate immune system, through action on Toll-like receptors, and the acquired immune system, through actions of interferon γ and a variety of interleukins (mainly IL-1, -4, -10, and -13) [78]. IL-8 can for example release a local imbalance between MMPs and TIMPs by inhibition of TIMP-1 expression in macrophages [89].

Besides macrophages, which release mainly MMP-1, -3, -8, -9, -11, -12, and -14, as well as TIMPs, other vascular cells produce and secrete various MMPs [20, 49]. The principal MMPs produced by vascular SMCs are MMP-1, -2, -3, -8, -9, and -14. MMP-2, rather than MMP-9, is constitutively expressed in vascular SMCs, but both MMP-2 activation and MMP-9 induction are rapidly triggered by vascular injury [90, 91]. Both inflammatory and immune activation of MMPs are also reported for SMCs. MMP-1, -3, and -9 are inducible by inflammatory cytokines and growth factors [92-94] as well as by immune cells [88, 95]. Atherosclerotic plaque resident T lymphocytes may also be a direct source of MMP-1, -2, -3, -9 [96, 97]. Mast cells, such as SMCs and ECs, constitutively secrete MMP-2 and upregulate MMP-9 secretion, in response to inflammatory stimuli. MMP-9 induction occurs both through cell-contact-dependent mechanisms with activated T cells, and through autocrine TNF- α secretion [98]. Moreover, inflammatory cytokines and growth factors produced by activated macrophages induce expression of MMPs in ECs, including MMP-1, -3, -2, -3, -7, -8, -9, -10, -11, -13, -14, -15, and -16 [99-101]. Ligation of CD40, expressed by ECs, thus upregulates MMP-1, -3, and -9 and increases activation of MMP-2 [102]. Oxidized LDLs also upregulate MMP-1 in human vascular ECs and human coronary artery ECs, whereas they downregulate TIMP-1 [103]. In addition, ECs express co-activators of MMPs, including urokinase plasminogen activator receptor, tissue plasminogen activator, CD44, and RECK, as well as TIMPs [101]. Furthermore, adventitial fibroblasts produce MMP-1, -3, and -9, while several MMPs have been identified in platelets, including MMP-1, -2, -3, -9, and -14, as well as TIMP-1, -2 and -4, which can modulate platelet activation and aggregation [104]. It has been reported that thrombin, which is produced in large quantities during plaque rupture, leads to MMP-2 activation and that, in turn, MMP-2 may increase platelet activation, leading to further activation of thrombin and to secondary MMP-2 activation, in a feedback mechanism [104, 105]. It has also been shown that oxidized LDLs stimulate MMP-9 and MMP-14 release by macrophages and SMCs [106, 107] and increase MMP-1 expression in ECs [108]. In this context, the mechanism responsible for expression and activation of vascular-cell-derived MMPs seems to involve both inflammatory and immune upregulation, as well as oxidative stress.

The implications of MMP overactivity on atherosclerotic plaque destabilization have been confirmed in *apolipoprotein E (ApoE)* knockout mice, by overexpressing individual MMP genes or by deleting TIMPs [77, 109, 110]. In the same type of mice, macrophage overexpression of active MMP-9 was found to induce acute plaque disruption, without significantly affecting lesion size or macrophage content [111]. Another study showed that overexpression of MMP-9 had no effect on the size of early carotid lesions, but disrupted advanced lesions, especially those caused by hypercholesterolemia [112]. It thus appears that MMP-1 overexpression in macrophages reduces atherosclerotic progression in *ApoE* deficient mice, by reducing the amount of collagenous matrix accumulation [113]. Studies in *ApoE* knockout mice with deleted MMP genes have illustrated that MMPs play a dual role in fibrous cup formation and plaque destabilization. Deletion of MMP-13 collagen deposition and greater plaque stability [114, 115]. Conversely, MMP-2 deletion lowered plaque stability in the aortic root, probably due to MMP-2's ability to aid the

migration of vascular SMCs and to build the fibrous cup [116]. *Mmp-3* deficiency produces more stable atherosclerotic plaques and increases plaque size [117]; in the aorta of *Mmp-9* deficient mice, though, plaque size was reduced, as were macrophage content and elastin degradation [118]. Moreover, increased plaque size, macrophage content and buried fibrous layers were observed in the brachiocephalic artery of *ApoE/Mmp-3*, *ApoE/Mmp-7*, and *ApoE/Mmp-9* double knockout mice [119]. Recruitment of vascular SMCs was also reduced in plaques of both *Mmp-3* and *Mmp-9* deficient mice, which is consistent with impaired intimal thickening in these mice [120]. Conversely, MMP-7 deletion increased SMC proliferation in the plaques of *ApoE* knockout mice [119], consistent with the involvement of MMP-7 in vascular SMC apoptosis [121]. With regard to MMP-12, its deletion caused more stable lesions in the brachiocephalic artery [119] and reduced elastin degradation in the aortic arch [118]. MMP-12 deletion also decreased foam cells apoptosis and reduced calcification into plaques [122]. Conversely, MMP-12 overexpression has also been shown to increase plaque size and inflammation in rabbits [123]. In addition, deletion of MMP-8 reduced inflammation and aortic atherosclerosis [124] while deletion of MMP-13 or MMP-14 had little effect on vascular SMC or macrophage content but, contributed to plaque stability by increasing collagen content [115, 125].

Degradation of ECM by MMPs exerts an influence at various stages of atherosclerotic plaque development [126]. In the initial stages of atherosclerosis after vascular injury, matrix degradation, presumably induced by MMP dysregulation (mainly MMP-2 and MMP-9), is thought to be combined with changes in permeability, macrophages activation and increased SMC migration [20, 49, 127]. For example, monocytes and T-lymphocytes adhering to ECs respectively enhance production of MMP-9 and that of MMP-2 [128, 129]. This enhancement is associated with EC basement membrane degradation, and penetration of the intima by monocytes and T-lymphocytes, leading to enhanced EC permeability [80, 100]. Moreover, MMP-2 and MMP-9 initially, and MMP-14 subsequently, promote the migration and proliferation of vascular SMCs, which could increase fibrous cup thickness and promote stability [127, 130, 131]. MMP-3 has also recently been implicated in this process, via its ability to stimulate MMP-9 activation [120]. Induction of MMPs in vascular SMCs, by a combination of inflammatory cytokines and growth factors, may be the key to matrix remodeling [94]. The regulatory mechanisms by which matrix remodeling can control the SMC cycle, involving growth factors and kinases, and by which it can also regulate plaque stabilization, has been studied in detail [132]. Conversely, MMP-7 mediated cadherin cleavage leads apoptosis of vascular SMCs, with possible decreased intima formation [121]. Large amounts of ECM are deposited in the fibrous cup during atherosclerotic plaque development, which provides its structural integrity and stability, while MMP secretion by vascular cells promotes macrophage invasion, thereby increasing plaque inflammation [133, 134]. During atherosclerotic plaque progression, ECs, vascular SMCs, and foam-cell macrophages secrete a variety of cytokines that perpetuate the recruitment and activation of inflammatory cells, which in turn attract modified vascular SMCs to the neointima [135]. In later atherosclerosis stages, an excess of MMPs versus inhibitors significantly contributes to ECM degradation, rendering the plaque more prone to rupture. Numbers of MMP-1, -2, -3, -7, -9, -12 and -14 positive macrophages, and of SMCs in the plaque increase in parallel with plaque progression [82, 126, 136]. The various different proteolytic enzymes have mainly been observed at the shoulder region of the plaque, where types I and III collagen are degraded; the cells playing the central role in this process are macrophages. Many MMPs (MMP-1, -2, -3, -7, -8, -9, -11, -12, -13, -14, and -16) are thus detectable in the macrophage-rich shoulder region of highly inflamed atheromatous plaques. MMP-7 and MMP-12, however, are detected in a greater concentration in macrophages close to the necrotic core [20, 96, 137, 138]. Co-localization of cleaved collagen with collagenases MMP-1, -8, and -13 suggests these enzymes are active [88, 139]. Interestingly, it has been observed that cells isolated from human plaques overexpress MMP-1 and MMP-3, caused by activation of Toll-like receptor 2 [140]. Moreover, isolated foam cells from cholesterol-fed rabbits induce expression of MMP-1, -3, -

12 and -14 [83, 141, 142]. However, a study carried out in endarterectomy biopsies showed that increased MMP-2 was mainly associated with SMCs and plaque stability, while MMP-8 and MMP-9 were associated with the presence of macrophages, and unstable plaques prone to rupture [138]. Increased MMP expression has also been observed after coronary angioplasty, suggesting that MMP expression may be involved in the formation of restenotic lesions [143]. It has also been reported that increased TIMP-1 levels are consistently present in human atherosclerotic plaques, where they are mainly found in areas of calcification [144] and that increased circulating levels of TIMP-1 are related to stable coronary, carotid and peripheral artery atherosclerosis [145, 146]. Of note, it has been shown that only pro-MMP-2, MMP-14, TIMP-1 and TIMP-2 are constitutively found in normal arteries; no activity was detected by *in situ* zymography [20].

Angiogenesis in the adventitia underlying plaques, and within the plaque itself, is another feature of the pathogenesis of atherosclerosis; it is associated with plaque progression and destabilization. The angiogenic fibroblast growth factors (FGFs) promote EC migration and upregulation of MMP-1, -2, -3, -7, -9, -10, -11, and -13 [147, 148] whereas vascular endothelial growth factor (VEGF) induce MMP-2 expression [149]. Thrombin also displays pro-angiogenic effects, by up-regulating MMP-1 and MMP-3 [150] and by enhancing MMP-2 activation via MMP-14 [151].

Endothelial erosion is another process associated with plaque rupture that occurs in highly stenotic and fibrotic plaques. It has been suggested that overproduction of MMPs (principally of MMP-2 and MMP-9) from inflamed or otherwise dysfunctional ECs weakens their interaction with their underlying basement membrane, thereby causing erosion. Moreover, low average shear stress and disturbed flow are also associated with endothelial dysfunction and erosion [101].

Of the various MMPs, MMP-9 has recently been the focus of growing interest in connection with human illnesses, including cardiovascular disorders associated to atherosclerosis [152]. It has then been found that polymorphisms of the MMP-9 gene are linked to atherosclerosis and to complicated coronary lesions [153-155]. Moreover, deficiency of *Mmp-9* reduces the formation of atherosclerotic lesions in *ApoE*-deficient mice [118]: MMP-9 is the enzyme most specifically associated with atherosclerotic plaque instability and rupture. However, MMP-9 is not only important for ECM degradation; it also plays a role in ECM organization [129, 156].

Upregulation of intraplaque MMP-9 leads to increased plaque hemorrhage and rupture in mouse models [111, 112]. MMP-9 has also been found to be highly expressed in unstable human plaques [157, 158], whereas TIMP-1 and TIMP-2 levels remained unchanged, and there was a significant increase in the MMP-9/TIMP-1 ratio [157]. In humans, analysis of coronary atherectomy specimens found higher levels of active intraplaque MMP-9 in patients with unstable angina (i.e. unstable coronary plaques) compared to patients with stable angina (i.e. stable coronary plaques) [159]: MMP-9 has been detected in the vulnerable shoulder region, and in areas of foam-cell formation in atherosclerotic plaques [78]. Moreover, raised levels of MMP-9 and MMP-2 in the coronary arteries carrying the culprit lesion, add further evidence to show the role of these enzymes in plaque rupture, and in the precipitation of an acute vascular event [160]. A significant increase in MMP-9 immunopositivity has also been demonstrated in atherosclerotic lesions of the aorta and carotid artery of rabbits fed a high-cholesterol diet [161]. Increased plasma levels of circulating MMP-9, as well as MMP-2, also correlate with the clinical symptoms of plaque instability and rupture in the coronary and cerebral circulation [162, 163] as well as being found in patients affected by myocardial infarction [164, 165], suggesting that MMP-9 may be useful as a biomarker for acute coronary syndrome [166, 167].

As several studies have also reported, oxidized LDL have been demonstrated to induce expression and activity of MMP-9, and to decrease its endogenous inhibitor, TIMP-1, in human macrophages and ECs [106, 168-170]. A causal

association between oxidized LDL autoantibodies and serum MMP-9 levels *in vivo* has also been demonstrated [171]. Oxysterols, which are cholesterol oxidation products, are present in considerable amounts in oxidized LDL, and appear to be implicated in the pathogenesis of atherosclerosis [172-175]. In this context, several studies have shown that, in cells of the macrophage lineage, oxysterols can initiate specific signal transduction pathways, which are relevant to the development of atherosclerosis [176-179]. A recent study has demonstrated that an oxysterol mixture, of composition similar to that found in advanced human carotid plaques [173], can significantly contribute to destabilizing the fibrotic plaque, by increasing expression and activity of MMP-9, without interfering with expression and synthesis of TIMP-1 or TIMP-2. The consequent net imbalance of the MMP-9/TIMP-1,-2 ratio would in this case then trigger an excessive proteolytic ECM degradation within the advanced lesion, and contribute to plaque instability and likely rupture [180].

2.1.1. Inhibition of Metalloproteinases as Therapeutic Strategies for Atherosclerosis

Although MMP involvement in atherosclerotic pathology and in other vascular diseases goes beyond simple excessive matrix degradation, MMP inhibition may be of therapeutic benefit [101, 181]. Several physiological mediators are present in the vasculature, where they suppress MMP secretion in normal tissues, and in conditions of injury and inflammation. The athero-protective agent NO, produced by ECs, might inhibit MMP-9 production from vascular SMCs [182]. The anti-inflammatory cytokine TGF β inhibits induction of MMP-1, -3, -7, although paradoxically it upregulates MMP-13 [183], and it may upregulate TIMP-3 [93]. Moreover, interferon- α and/or - γ inhibit the induction of MMP-1, -3, -9, and -13, thereby presumably reducing the contribution of immune mechanisms to MMP induction in SMCs and in macrophages [85, 95].

Furthermore, recent data have shown that treatment with a low-molecular-weight heparin decreases the levels of MMP-9 in patients with abdominal aortic aneurysm [184]. Again, angiotensin converting enzyme inhibitors (ACEIs) and angiotensin II receptor blockers (AIIIRBs) reduce the raised circulating MMP-9 levels in patients with stable coronary artery disease [185], and the levels of this enzyme are also reduced by ACEI captopril in patients with acute myocardial infarction [186]. Further, since MMP activity may be induced by ROS, antioxidant therapy can be useful to modulate MMPs, including MMP-9, as has been demonstrated in a hypercholesterolemic rabbit model [187].

Targeting specific cytokines or signaling pathways that are involved in mediating MMP upregulation could thus reduce MMP activity in atherosclerotic plaques. For example, inhibition of the CD40 ligand stabilizes plaques in mice, possibly due in part to effects on MMPs [188]. The MMP pathway can also be interrupted by targeting the inflammatory response. One specific target is PGE₂: reducing PGE₂ synthesis, for example with indomethacin or other anti-inflammatory drugs, can contribute to reducing MMP synthesis [189, 190]. Because PGE₂ production is involved in MMP transcription, antagonists at the PGE₂ receptor could also be useful in stabilizing plaques [191, 192]. Aspirin and other cyclooxygenase inhibitors also inhibit MMP production by monocytes [78], although cyclooxygenase independent pathways of MMP production may reduce the beneficial impact of these compounds in man [79]. In addition, MMP-1, -3 and -14 are expressed via NF- κ B, therefore inhibition of this transcription factor might reduce expression of these MMPs [83].

Taken together, these findings show that there is a clear potential for the application of TIMPs as endogenous inhibitors [193] through gene therapy [41]. It has been demonstrated that adenovirus-mediated gene transfer aimed at overexpressing TIMPs can reduce MMP activity, intimal thickening, and plaque destabilization in various models. For example, overexpression of TIMP-1 in a mouse model of atherosclerosis caused a reduction in the lesion and macrophage content at the aortic root [194]. Another study examined whether short-term overexpression of TIMP-1 or TIMP-2 would attenuate atherosclerotic plaque development and instability, in *ApoE* knockout mice fed a high-fat diet.

Analysis of brachiocephalic artery plaques showed that overexpression of TIMP-2, but not TIMP-1 infection, resulted in marked reduction in lesion area compared with control animals. TIMP-2 significantly reduced migration and apoptosis of macrophages and foam cells, inhibiting atherosclerotic plaque development and destabilization, whereas TIMP-1 failed to exert similar effects [133]. Moreover, a recombinant TIMP-1 has been shown to reduce the activity of MMP-1, -2, -9, and -3 in the shoulder and core regions of the plaque [101]. Finally, overexpression of TIMP-3 markedly reduced neointima formation both *in vitro* and *in vivo*, by promoting cell apoptosis [195, 196].

Furthermore, several observations suggest that statins, which are potent lipid-lowering drugs, may exert their beneficial effects on the arterial wall in part by means of their effects on the inflammatory response and on MMP and TIMP production. In particular, statins contribute to increasing plaque stability by inhibiting MMP secretion [197]. Statins have been reported to prevent atherosclerosis progression and coronary events, by inhibiting expression and secretion of MMP-1, -2, -3, and -9, in macrophages and SMCs *in vitro*, and in rabbit and human atherosclerotic lesions [198-201]. Cerivastatin also suppresses macrophage growth and reduces MMP-1, -3, and -9 expression in rabbits, while conserving the collagen [202]. Treatment with pravastatin before carotid endarterectomy reduced plaque lipid content, inflammation, MMP activity and cell death, also increasing the collagen content [200]. Statins are also known to exert anti-inflammatory and cardioprotective effects in *ApoE* knockout mice [203]. In addition, statin treatment has been shown to cause a significant reduction of C reactive protein (CRP) circulating levels, which are correlated to the severity of atherosclerosis [204, 205].

Another strategy considers antibody-based inhibitors of MMPs. MMP-2 neutralizing antibodies have shown protective activity in hearts exposed to pro-inflammatory cytokines or ischemia/reperfusion injury. MMP-14 blocking antibodies have also been proposed as targets [206]. Alternatively, synthetic MMP inhibitors have been developed and tested. Synthetic MMP inhibitors are potent Zn^{2+} -chelating mimickers of collagen, and the majority of them are broad-spectrum inhibitors that suppress the activity of a large number of different MMPs [207]. More than 50 synthetic MMP inhibitors have to date been considered for possible clinical development [101]. Among synthetic MMP inhibitors are the hydroxamic acid derived inhibitors, such as BB-94 (batimastat), BB1101, BB-2293, BB-2516 (marimastat), and CPD-845 (CT-1746) [208]. Batimastat and marimastat are competitive MMP-inhibitors but, despite the promising results obtained with both inhibitors, their lack of oral bioavailability has precluded their long-term use; marimastat had better oral bioavailability [209] than batimastat [210]. Moreover, non-selective MMP inhibitors [211] or more selective MMP inhibitors, including pyrimidine-2,4,6-trione derivative Ro-28-2653, inhibit MMP-2, -9, and -14, reducing BBB breakdown in rat models of stroke [212]. Likewise, Ro-28-3555 (trocade) is a selective inhibitor of MMP-1 [213] and IW449 is a selective inhibitor of MMP-2 [208]. Other synthetic inhibitors include PD166793, OPB-3206 [214], BAY12-9566, AG-3340, KBR-7785, KBR-8301, GM-6001 (ilomastat or gelardin) metastat and AE-941 (neovastat) [215]. Moreover, 1,10-phenanthroline is a small organic compound inhibiting a broad range of MMPs. The selective MMP-12 inhibitor (RXP470.1) was used to treat established plaques in *ApoE* knockout mice; it blocked plaque enlargement, decreased lipid core formation, improved the ratio of SMCs to macrophages, and also reduced macrophage apoptosis, calcification and medial elastin breaks [122]. Further, a highly-selective inhibitor of MMP-13 also inhibited collagenolysis, thus preserving the collagen content in plaques [216] to a very similar degree as has been observed to occur in both mice transgenic for a collagenase-resistant mutant of mouse collagen I [114] and *Mmp-13* knockout mice [115].

The tetracycline antibiotic doxycycline is also a broad-spectrum MMP inhibitor, reducing both synthesis and activity of MMPs [217, 218]. Its main mode of action is by binding to the active zinc site, resulting in conformational change in the enzyme structure with loss of activity. Clinical trials using doxycycline showed a significant reduction in

MMP-1 levels in carotid endarterectomy samples [219] and of plasma MMP-9 levels in patients with coronary artery disease [220]. Again doxycycline, appeared to be well-tolerated in another clinical trial for the treatment of abdominal aortic aneurysms [221, 222]. However, in all these clinical trials, the clinical benefit was low or remains to be demonstrated. Doxycycline increased SMC adhesion and reduced SMC migration; moreover, it limited the reorganization of fibrillar collagen matrices [217]. Doxycycline and the MMP inhibitors Ro-31-4724 and Ro-31-7467 have also been shown to reduce SMC proliferation *in vivo* [223] and to promote SMC apoptosis [224].

Because angiogenesis within plaques is associated with plaque progression and vulnerability [225] and because MMPs participate in this feature of atherosclerosis, they might be another target for therapy. It has been reported that marimastat inhibits angiogenesis in both collagen and fibrin matrices, by inhibiting MMP expression [149].

2.2. Metalloproteinases and Diabetic Vascular Complications

Type 2 diabetes mellitus is a frequent vascular risk in cardiovascular events, including atherosclerosis, and also in microvascular complications [226]. Considerable evidence points to a role for MMPs and TIMPs in the atherosclerotic process; however, the relationship between MMPs/TIMPs and diabetic angiopathy is less well defined. It is probable that atherosclerosis and diabetes mellitus share common pathways of MMP synthesis and plaque destabilization, including inflammatory pathways and molecules. However, recent *in vitro* and *in vivo* studies have demonstrated that hyperglycemia, either directly, or indirectly via oxidative stress or advanced glycation end-products (AGEs) regulates MMP's expression and activity. Disruption of the ECM may enhance monocyte and vascular SMC migration, which may aggravate atherosclerosis in diabetes mellitus.

The exposure of ECs, macrophages, and SMCs to high concentrations of glucose induces dysregulation of the MMP/TIMP balance. High glucose concentrations have been demonstrated to induce expression of MMP-2 in arterial vasculature *in vivo* [227] and in cultured vascular SMCs [228]. In addition, hyperglycemia induces expression of MMP-1 and MMP-2 in ECs, and expression of MMP-9 in macrophages, decreasing expression of MMP-3 but having no effect on TIMP-1 expression [35]. Another study has found MMP-2 and MMP-9 expression not to be affected, but in a high-glucose environment their activity increased in ECs from umbilical cords [229]. Interestingly, it has also been shown that the effects of hyperglycemia on MMP-2 activity were further enhanced in vascular SMCs that were exposed to intermittent, rather than constant, high-glucose concentrations, more closely resembling a pathophysiological condition [230]. Moreover, increased synthesis of active and latent forms of MMP-2 and MMP-9 was observed in aortic specimens and blood samples from diabetic rats, as well as in cultured ECs, but not in vascular SMCs or macrophages [231]. It was subsequently shown that incubation with plasma LDL from patients with type 2 diabetes significantly increases expression of MMP-9 in monocytes [232]. It has also been suggested that high glucose levels might upregulate transcriptional factors, such as AP-1, or growth factors, such as TGF β , which in turn enhance MMP gene transcription [233,234].

MMP gene transcription may also be regulated by oxidative stress, through several mechanisms such as alteration of NO synthase activity, and AGE formation [235], as well as by genetic polymorphism [236]. There is increasing evidence to suggest that AGEs, and the interrelationship with their receptors (RAGEs), influence several signaling pathways, which are involved in vascular dysfunction [237]. AGEs are usually localized on the surface of macrophages and bind to the ECM. It has been demonstrated that matrix-glycation products increase monocyte infiltration, especially in the shoulder regions of plaques, and that they stimulate macrophages to release cytokines, thus accelerating the inflammatory response [238, 239]. In the shoulder region, there is thus an accumulation of macrophages, which contain the majority of RAGEs; the overexpression of RAGEs, which sustains inflammation,

triggers vascular SMCs and macrophages to produce MMP-2 and MMP-9 [238]. The preponderance of macrophages coincides with reduced collagen content and with MMP-2 and MMP-9 overexpression in human diabetic plaques [238].

The MMP/TIMP system may interfere with the subcellular insulin signaling pathways. Imbalance of the insulin signaling cascades may be involved in the atherosclerosis-promoting effect of insulin resistance, possibly by promoting MMP-2 and MMP-9 overexpression and compromising the expression of TIMP-3 [240]. Conversely, it has been shown that, in healthy subjects, insulin infusion acutely suppresses plasma levels of MMP-9 and VEGF levels. VEGF is known to contribute to proliferative retinopathy, as well as to plaque evolution and rupture [241]. Thus, insulin might act to prevent atherosclerosis and diabetic complications through its anti-proteolytic and anti-angiogenic properties. High glucose levels, acting via the AGE/RAGE interaction and altered insulin signaling, thus provide a stimulus to inflammation and ECM degradation within atherosclerotic plaques.

High circulating levels of MMP-2, -8 and -9 have been found in patients with diabetes [242-244] and peripheral arterial disease [245] or acute coronary syndrome [246], while TIMP-1 levels are reported to be lower in diabetic patients than in non-diabetic subjects [247], although other studies have found normal levels of MMP-1, -3, -9 and TIMP-1 in diabetic patients [248, 249].

Apart from the role of MMPs in diabetic macrovascular complications, such as atherosclerosis, MMPs seem to play a key role in the development of the diabetic microvascular complications known as microangiopathies, which include diabetic retinopathy, diabetic nephropathy, and diabetic peripheral neuropathy.

Although normal retinas express MMP-1 and TIMP-2, retinas from diabetic patients show above-normal concentrations of activated MMP-1, -2, -3 and -9, as well as of TIMP-1, -2, and -3 [250, 251]. These processes may contribute to retinal neovascularization. High concentrations of MMP-8, -9, and -14 have also been found in the urine of patients with diabetic nephropathy [252], leading to investigation of the expression of MMP in diabetic kidneys and mesangial cells. Accumulation of ECM within the glomerulus contributes to diabetic renal dysfunction, and the amount and composition of mesangial matrix in diabetic nephropathy reflects the imbalance between synthesis and degradation of the ECM [253]. Additionally, hyperglycemia reduces MMP expression whereas it accelerates TIMP expression, consequently suppressing ECM degradation, leading to the accumulation of matrix components in the glomerular mesangium [254]. Moreover, it has been observed that the increased AGE formation within glomeruli renders the ECM less susceptible to degradation, by reducing MMP activity, and thus playing a role in the pathogenesis of diabetic nephropathy [255]. Interestingly, alongside their direct role in ECM turnover, MMPs have been demonstrated to release or activate various growth factors that have been associated with renal hypertrophy, tubular cell proliferation, and renal fibrosis, and which contribute to development of the renal abnormalities characteristic of diabetic nephropathy [256].

2.2.1. Inhibition of Metalloproteinases as Therapeutic Strategies for Diabetic Vascular Complications

There is increasing interest in the influence of anti-diabetic drugs on the MMPs/TIMPs balance. Rosiglitazone, a thiazolidinedione, reduces circulating levels of MMP-9, IL-6, white blood cells and other inflammatory markers in type 2 diabetes [257, 258], subsequently reducing neointimal hyperplasia [259]. Moreover, gliclazide reduces oxidized LDL-mediated MMP-9 expression in human aortic ECs *in vitro* [168] and pioglitazone, compared with placebo, significantly decreases plasma MMP-9 levels in diabetic patients with coronary artery disease [260]. Thiazolidinediones may also influence MMP-1 expression in vascular ECs [261].

It has also been reported that both ACEIs and AIIRBs exert beneficial effects by preventing or slowing the progression of diabetic nephropathy, favoring, for example, MMP-2 activity [262]. In addition, it has been shown that diabetic patients benefit from statin treatment [263-265]. In this context, it is speculated that statins may exert their beneficial

effects in diabetic patients through a similar mechanism to that occurring in atherosclerosis: they appear to protect from diabetic complications by decreasing lipid levels, lipid oxidation, inflammation, MMP expression, and cell death, and by increasing the TIMP and collagen content of human atherosclerotic plaques, thus contributing to their stability. Statins may also act by reducing vascular SMC migration and proliferation [266], and exerting an inhibitory action on MMP-1, -3, and -9 secretion from the same cells [267]. Statin treatment also causes significant suppression of MMP activity and significant activation of TIMPs (MMP-2 and TIMP-2, respectively) preventing, for example, glomerular ECM accumulation [268, 269]. Statins are also reported to possess anti-inflammatory action: in particular they reduce the serum concentrations of CRP, which is considered to be a marker of vascular risk [270-272]. These drugs can also improve renal function and, of note, they lower serum urate concentrations, which can influence MMP activity [273-275].

Peroxisome proliferator activator receptors (PPARs) are nuclear receptors that regulate fatty acid oxidation, adipocyte differentiation, and insulin sensitivity, as well as atherosclerosis pathogenesis; PPAR ligands are thus another compound that is used in the treatment of diabetes mellitus. The thiazolidinediones, which are insulin-sensitizing drugs, are PPAR γ activators, while fibric acid derivatives (fibrates) have lipid-lowering properties, and activate PPAR α [276]. There is evidence suggesting that both a PPAR α activator (fenofibrate) and a PPAR γ activator (rosiglitazone) inhibit MMP-9 expression in vascular SMCs and in monocytes/macrophages [277, 278]. Fibrates may also decrease plasma levels of inflammatory markers (*e.g.* CRP) that, in turn, influence MMP expression [279, 280]. Based on these findings, it has been suggested that the fibrate-plus-statin combination could be a promising therapeutic strategy for diabetic atherosclerosis [281].

2.3. Metalloproteinases and Neurodegenerative Diseases

An increasing body of evidence points to the critical role of inflammation in the neurodegenerative process. During neuroinflammation, molecular cascades occur whose goal is to remove damaged cells and prepare the brain for repair, but the overactivated and/or chronically activated state of the microglia contributes to neuronal death and dysfunction. Microglia can be activated by MMPs as well as by amyloid β (A β) and α -synuclein [282, 283].

MMPs and ADAMs are important in acute and chronic neuroinflammation, and recent studies have linked their actions to neurodegenerative disorders that are often associated with vascular cognitive impairment, such as Alzheimer's and Parkinson's diseases [23, 284-286].

Several key MMPs and ADAMs have been implicated in neuroinflammation: MMP-2, MMP-3, MMP-9, MMP-14, ADAM-9, ADAM-10, and ADAM-17. However, with regard to ADAM functions in the brain, they are also implicated in cell survival, proliferation, differentiation, and migration, as well as in axonal growth and myelination.

Brain cells express both constitutive and inducible MMPs in response to cellular stress [7]. For example, MMP-2 is a constitutively expressed molecule that is normally present in brain tissue and in the cerebrospinal fluid (CSF). MMP-9 is normally expressed in brain tissue at low levels, but is markedly upregulated by various inflammatory stimuli (*e.g.* cytokines and growth factors) in many brain disorders; conversely, MMP-9 is not present in the CSF [7,23].

The initial phases of neuroinflammation are characterized by activation of constitutive proteases that begin the process of disassembling the ECM, opening the blood-brain barrier (BBB), preventing normal cell signaling, and initiating cell death by apoptosis [287].

In the active form, MMPs play a number of important roles in normal development, but they are highly destructive in case of inflammation of the central nervous system (CNS). These enzymes, indeed, increase the

permeability of the BBB by attacking the ECM, basal lamina, and tight junctions proteins in ECs; this increased permeability is a feature of the acute neuroinflammatory response, and allows cells to enter the CNS, contributing to white-matter damage. When matrix proteins around the neurons are degraded, there is loss of contact and cell death by anoikis [32]. Proteolysis of the matrix protein of blood vessels and brain cells by MMPs, in particular by the constitutive MMP-2, and subsequently by the inducible MMP-1, -3, and -9, increases the risk of cerebral edema, hemorrhage, and cell death [288]: it has been shown that MMP-2, -3, and -9 increase permeability of the BBB [287, 289-292] leading to the infiltration of inflammatory cells, such as neutrophils, which play an important role in neuroinflammation [293]. Loss of oxygen and energy substrates releases glutamate into the extracellular space, initiating molecular events in the injured cells that might result in loss of membrane integrity and necrosis. Inhibitors of MMPs can reduce damage to the BBB, and thus reduce cerebral edema and hemorrhage [290]. *Mmp-9* or *Mmp-3* knockout mice were also found to have reduced infarct size and significantly less BBB damage and neutrophil infiltration [289, 293, 294]. Direct injection of TNF- α into the rat brain results in a dramatic increase in the expression and activation of MMP-9 and MMP-3, which is associated with a significant opening of the BBB. Microglia and neurons surrounding the injection site are the major cellular sources of MMP-3 and MMP-9, through cyclooxygenase-derived product involvement, following intracerebral TNF- α administration [295, 296]. Moreover, in the brain, microglia and astrocytes are major sources of many pro-inflammatory cytokines, and of other mediators that stimulate increased MMP production [297, 298]. Secondly, MMPs are involved in tissue repair, driving angiogenic and neurogenic processes [299, 300] and in the end ECM remodeling occurs, with impenetrable scar tissue formation that blocks re-growth and re-projection of axons.

However, much has been learned about the function of the MMPs in the brain, by using cell cultures such as neurons, ECs, astrocytes and microglia. The two major inducible MMPs that have been identified in the neuroinflammatory response are MMP-3 and MMP-9. Rat brain ECs stimulated with lipopolysaccharide (LPS) showed induction of MMP-9 [301]; cultured rat astrocytes stimulated with LPS, IL-1 β , TNF- α , or bradykinin secreted normal MMP-2, while expression and activity of MMP-9 was upregulated [302-305]. Other studies have further demonstrated that several external stimuli can upregulate MMP-9 expression, via the MAPK/AP-1 pathway, in different cell types [306-308]. Expression of MMP-9 can also be induced by oxidized LDL in rat brain astrocytes, through the MAPK-extracellular signal-regulated kinase kinase (MEK)1/2, and the phosphatidylinositol 3 (PI3)/Akt-c-Jun N-terminal kinase (JNK)1/2 signaling pathways, leading to AP-1 activation [309]. This data implies that oxidized LDL might play a crucial role in the development of brain injuries and CNS diseases. Oxidized LDL, indeed, has been reported to exhibit a wide range of biological activities, including alteration of neuronal apoptosis, capillary homeostasis, and modulation of inflammatory protein activity in various brain cells [310-312]. Moreover, astrocytes also release MMP-1 when stimulated with IL-1 β , and MMP-1 has been shown to be toxic to human neurons in culture [313]. In addition, incubation of this cell type with the 1-40 fragment of A β induced MMP-9 and MMP-3 production, suggesting that MMPs may be involved in amyloid processing in AD [314]. Stimulated astrocytes might also be induced to produce the inactive form of MMP-9, thus failing to produce its active form; however, cultures of microglia and astrocytes stimulated with LPS have been shown to produce the active form of MMP-9 and MMP-3 [303, 315, 316]. This evidence stresses that the microglia are necessary for activation of the proMMP-9, possibly through MMP-3 or free radical production [315]. It has also been suggested that the effects of LPS on the MMPs expression are due in part to the formation of pro-inflammatory cytokines. TNF- α and IL-1 β produce a significant increase in the production of MMP-3 and MMP-9 in cultured astrocytes and microglia [302, 317]. In LPS-stimulated astrocytes, smaller increases of MMP-10, -12, and -13 were also seen [318].

Of note, a number of immunohistochemistry studies, conducted in human autopsy brain tissue from patients with various pathologic conditions, or in animal models, have identified the cell types expressing the MMPs in the brain. In patients with multiple sclerosis and cerebral infarction, antibodies to MMP-9 were localized in blood vessels and neutrophils, but in more chronic lesions, MMP-2 and MMP-7 were the prominent enzymes found in the inflammatory cells. Astrocytes around the infarctions immunostained positive for MMP-2 [319]. Furthermore, in the progressive form of dementia, due to atherosclerosis of the blood vessels with demyelination of the white matter, tissue macrophages stained for MMP-3 in regions of damaged white matter [320]. There is some evidence to indicate a role for MMP-3 in neurodegeneration, because it plays a critical role as an intercellular signaling molecule that modulates neuroinflammatory responses [321-323].

In neuronal cells, MMP-3 expression is increased in response to cell stress, and the active MMP-3 participates in apoptotic signaling [324] and triggers microglia activation and production of pro-inflammatory cytokines [321] triggering the formation of ROS [322, 323]. Active MMP-3 may also induce microglial activation near the site where apoptosis occurs, to promote clearance of apoptotic cells [321]. Despite these data, the exact molecular mechanisms through which active MMP-3 activates microglia cells are still not clear. In this connection, it has been shown that MMP-3 deficient mice display a significant reduction in microglia activation following the *in vivo* administration of the neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) [322]. This would appear to indicate that, in the extracellular space, MMP-3 triggers the microglia to produce pro-inflammatory and cytotoxic molecules, which in turn contribute to neuronal damage and to the removal of damaged neurons by phagocytosis. In addition, MMP-3 produced by the activated microglia can be released into the extracellular matrix, further exacerbating the neuroinflammatory process [324]. In contrast, overexpression of TIMP-1 results in the attenuation of apoptosis in neuronal cells, along with the suppression of MMP-3 activity [325]. TIMP-1 is also neuroprotective, both against excitotoxic neuronal death [326] and against traumatic and ischemic brain injury in mice [327]. In an animal model of temporary ischemia-reperfusion, MMP-3 was mainly seen in the neurons and microglia, while MMP-9 was present in the neurons. After 3 weeks of reperfusion, the principal MMP present was MMP-2, in astrocytes [315].

In this connection, evidence is emerging that MMPs and ADAMs play roles in neurodegenerative diseases, including in the damage to white matter in patients with vascular cognitive impairment, in degradation of amyloid peptides in AD, and in apoptosis of dopaminergic neurons in PD.

a) Vascular cognitive impairment

In vascular cognitive impairment, MMPs are induced by hypoxic hypoperfusion in the white matter [328]. The brain tissues of patients with vascular cognitive impairment express hypoxia-inducible factor 1 α (HIF1 α). HIF1 α increases during hypoxia or ischemia and, because hypoxia seems to play a crucial part in vascular cognitive impairment, an understanding of the role of HIF1 α is important.

The increase in HIF1 α leads to the expression of many genes implicated in injury and repair [329]. HIF1 α can activate furin, an intracellular convertase that activates several enzymes involved in injury, including the MMPs; these increase permeability of the BBB and demyelination in the white matter, as well as elevating vasoconstriction, with subsequent marked hypoxia [330, 331]. Conversely, HIF1 α can stimulate the expression of genes involved in repair, such as VEGF and TGF β [332, 333]. It has also been observed that patients with vascular cognitive impairment have increased expression of MMPs in the white matter, particularly around blood vessels in regions with loss of myelin. Gliotic regions have reactive astrocytes that overexpress MMP-2; macrophages around damaged blood vessels are immunopositive for MMP-3 [320]. The MMPs might damage blood vessels, disrupting the BBB, and thus activating

microglia and recruiting macrophages that contribute to white-matter injury. Demyelination of white matter might also occur, through MMP-mediated mechanisms, mainly by MMP-2, MMP-3 and MMP-9 [320, 330, 334]. Moreover, patients with vascular cognitive impairment have high concentrations of MMP-9 in the CSF [335]. Patients with vascular cognitive impairment also have increased levels of endothelin-1 in the white matter, where it is highly expressed by MMP-2 [336, 337]; endothelin-1 is a strong vasoconstrictor that may compromise blood flow to the deep white matter [338].

b) Parkinson's Disease

Parkinson's disease (PD), a progressive neurodegenerative disorder, is characterized by the selective loss of dopaminergic neurons in the substantia nigra, and degeneration of projecting nerve fibers in the striatum, leading to extrapyramidal motor dysfunction, which is associated with microglia activation [339]. MMPs have been implicated in the death of dopaminergic neurons; in particular, various experimental models have linked increased MMP-3 levels with PD. An animal model of PD showed an increase in MMP-3 immunoreactivity in the substantia nigra region [340, 341]. It has also been shown that, *in vitro*, apoptotic dopaminergic neurons release MMP-3, which acts as a microglia-activating molecule. This evidence suggest that, in addition to ECM degradation, MMP-3 is a signaling molecule in the neuronal apoptotic process, mediating the interaction between apoptotic neurons and microglia, and consequently causing neuroinflammation [321]. The activated microglia release pro-inflammatory cytokines, including TNF- α , which directly induce neuronal death, contributing to neuronal degeneration. It has also been suggested that, in addition to the extracellularly-triggered apoptotic mechanisms, MMP-3 might also act intracellularly in apoptotic signaling in the dopaminergic neurons [342]. Furthermore, MMP-3 may play an important role in the pathophysiology of PD, by contributing to the generation of the toxic α -synuclein aggregates [340, 343].

Finally, MMP-3 mediates BBB disruption, which would allow infiltration of blood immune cells such as neutrophils to the damaged region [293]. Indeed, BBB leakage has been observed in PD patients [344] as well as in animal models of PD [345, 346]. Interestingly, the substantia nigra region is reportedly more prone to BBB disruption and neutrophil infiltration than is the cortical region [347].

c) Alzheimer's Disease

Alzheimer's disease (AD), the most common form of dementia, is characterized by the progressive loss of neurons and synapses, and by extracellular deposits of A β in the form of senile plaques, A β deposits in the cerebral blood vessels, and intracellular inclusions of hyperphosphorylated tau in the form of neurofibrillary tangles (NFT) [348, 349]. Several mechanisms contribute to AD development and progression, and deposition of improperly processed amyloid is thought to be a major factor in its pathophysiology. Amyloid precursor protein (APP) comprises a transmembrane and an extracellular component, and is degraded into various fragments by the secretases. The physiological pathway results in the cleavage of APP by α -secretase, which produce a soluble fragment (sAPP α) that can be broken down for clearance, and a membrane-bound COOH terminal fragment (CTF). Both β -secretase and γ -secretase act together to produce an ectodomain derivative (sAPP β) and A β peptides (A β ₁₋₄₀ and A β ₁₋₄₂) that can aggregate to form insoluble dimmers, oligomers, and subsequently fibrils, which are deposited extracellularly and intracellularly, forming senile plaques [350, 351]. Both MMPs and ADAMs have been implicated in APP shedding [352].

Three members of the ADAM family have been shown to act as an α -secretase: ADAM-9, ADAM-10, and ADAM-17; these enzymes cleave APP into the soluble sAPP α for clearance [353, 354]. ADAM-17 also activates TNF- α , which induces cell death, contributing to neurodegeneration [355].

Overexpression of ADAM-9 has been reported to increase sAPP α release, although mice lacking ADAM-9 revealed no difference in the production of the α -secretase cleavage product of APP [356]. The impact of ADAM-9 on sporadic AD [357] might therefore rely on a more indirect mechanism: ADAM-9 has been shown to proteolytically process ADAM-10 [358-360]. Unlike ADAM-9, ADAM-10 was found to have constitutive and regulated α -secretase activity [354, 361].

With regard to ADAM-10, it has also been observed, in an AD mouse model in which the animals were crossbred with ADAM-10 transgenic mice, that plaque pathology was greatly attenuated, production of α -secretase was enhanced, and subsequently also that of sAPP α [361]. Furthermore, these mice had increased learning and memory potential [361, 362], which might correlate with the observed enhanced cholinergic and glutamatergic synaptogenesis [363]. Conversely, mice with a dominant negative mutant of ADAM-10 had lowered sAPP α levels, accompanied by an enhanced number of plaques [361] and learning deficiencies [364]. In addition, axonal guidance is conveyed by ADAM-10, as has been shown for retinal and peripheral axons [365, 366], and this enzyme regulates axon withdrawal by ephrin cleavage [367, 368]. However, it remains a matter of controversy whether there is a substantial decline of neuronal ADAM-10 in ageing or in the pathological context.

Experiments performed with ADAM-17-deficient cells indicated a participation of ADAM-17 in the regulated [369, 370] and the constitutive α -secretase pathways [371, 372]. Although there is no evidence that ADAM-17 acts as an *in vivo* APP-sheddase in transgenic mice, ADAM-17-positive neurons are found to co-localize with amyloid plaques in AD brains, supporting its role as an α -secretase [373].

In the CNS, MMPs are produced by astrocytes, microglia, neurons, ECs, oligodendrocytes and leukocytes [301, 374, 375]. Elevated levels of MMPs have been reported in the cortex and hippocampus of AD patients compared with controls [376]. Several studies have suggested that MMPs, like ADAMs, participate in the formation and clearance of A β [377, 378]. It has been shown that MMPs, such as MMP-2, -3, and -9, are induced endogenously by the toxic amyloid fragments (A β ₁₋₄₀ and A β ₁₋₄₂) in astrocytes, microglia, mixed hippocampal neurons, and blood vessels [379-381]. Moreover, it has been demonstrated that the upregulation of MMP-9, expressed by SK-N-SH cells in the presence of A β ₁₋₄₀, is mediated by $\alpha_3\beta_1$ and $\alpha_2\beta_1$ integrin receptors, and that MMP-9 can directly process APP, by interacting on the cell surface with α -secretase-like activity, increasing sAPP α release and substantially reducing levels of secreted A β peptides [382]. These findings indicate that MMP-9 might act as a neuroprotective mechanism whereby the shedding of APP to soluble fragments precludes the formation of aggregating A β peptides. In addition, metal-mediated activation of MAPKs resulted in upregulated MMP-2 and MMP-3 activity, in turn leading to enhanced cleavage of extracellular A β , and preventing its accumulation [383-385].

In the area around the amyloid plaques, the microglia are activated and contribute to neuronal death by releasing inflammatory molecules [386]; elevated production of MMPs in the brain tissue of AD patients is part of the inflammatory response. It has been shown that expression of MMP-2, -3 and -9 is increased in astrocytes around amyloid plaques compared with areas without A β deposition [377]. MMP-3, for example, has been detected around senile plaques, in the grey matter and in the interstitium between myelinated axons and astrocytes in the white matter of AD patients [387]. MMP-3 is also significantly elevated in the plasma and CSF of AD patients [388]. More recently, the CSF of cognitively-healthy individuals with risk markers for future AD has been found to have higher MMP-3

levels and a higher MMP-3/TIMP-1 ratio than healthy individuals without risk markers [389]. Furthermore, in *postmortem* AD brain tissue, MMP-9 has been found expressed in hippocampal neurons around the amyloid plaques, and in the NFT and vascular wall [289, 390]; it has been suggested that MMPs are synthesized in response to A β , and that their activation degrades the peptide *in vivo*, reducing its aggregation [390]. Moreover, plasma concentrations of MMP-9 are increased in AD patients, and it has been suggested that this may be due to ECs releasing the enzyme in response to circulating A β [391]. Conversely, MMP-9 concentrations are not increased in the CSF [335]. Active MMP-9 has been reported to degrade synthetic A β ₁₋₄₀ *in vitro*, acting directly at the α -secretase cleaving site [390], and also to cleave extracellular and fibrillar A β through BBB [377, 380]. In this connection, *Mmp-2* and *Mmp-9* knockout mice had higher levels of A β in the brain tissue than wild-type mice, and treatment with the MMP inhibitor GM-6001 increased A β in the transgenic mice, overexpressing the Swedish variant of APP (mutations at positions 670 and 671) [377]. Besides MMP-9, increased expression of MMP-3 has also been detected in hippocampal neurons, around amyloid plaques in the cortex, and in the interstitium of white matter [387]. It has also been suggested that MMP-14, -16, and -18 might play important roles in regulating APP function, inducing cleavage and shedding of the APP ectodomain when co-expressed with APP adaptor protein Fe65 [352].

2.3.1. Therapeutic Strategies for Inhibiting Metalloproteinases in Neurodegenerative Diseases

Currently, no therapy is available clinically that delays the neurodegenerative process itself, and therefore the development of selective inhibitors and/or other therapeutic strategies against the proteolytic enzyme actions, together with a knowledge of their side-effects, will be of great interest to contrast the progression of neurodegenerative diseases.

In vascular cognitive impairment, treatment with a drug that blocks MMPs, such as minocycline, might be useful to decrease white-matter damage. The possible use of MMP inhibitors in treating neurodegenerative diseases such as AD and PD is even more speculative. Regarding AD, the role of MMPs and ADAMs is complex because of their dual function, in breaking-down amyloid to form A β , and in clearance of the peptide from the brain. MMP inhibitors could thus interfere with A β clearance. With regard to PD, the use of MMP inhibitors might be promising, since death of dopaminergic neurons appears to be stimulated when activated microglia release MMP.

The major synthetic MMP inhibitors are based on a hydroxamate structure [392]. These compounds interfere with the action of the zinc catalytic domain in the MMP molecule; however, the use of these inhibitors in clinical trials gave controversial results. More encouraging results have been obtained with other inhibitors, in terms of reducing BBB damage, infarct size, and cell death, in animal models of CNS diseases [23]. For example, TIMP-2 blocked the MMP-2-induced BBB opening in MMP-induced brain injury [393]. Damage to the BBB was blocked with hydroxamate treatment [394-396]. BB-94 blocked BBB opening after intracerebral injection of TNF- α , and BB-1101 was effective in LPS-induced BBB injury [295, 397] as well as in BBB opening after stroke, in the rat brain [398]. In addition, GM-6001 reduced BBB injury and improved the outcome [394, 399]. Moreover, a highly-specific thiirane gelatinase inhibitor of MMP-9, namely SB-3CT, rescues laminin from proteolysis, and protects neurons from apoptosis, in cerebral ischemia [400]. MMP inhibitors can also protect the brain from hemorrhagic complications of alteplase, a recombinant tissue plasminogen activator, by reducing BBB permeability and preventing alteplase from entering the brain and activating MMPs [401, 402]. Although there are MMP inhibitors that are selective for MMP-2 and MMP-9 [403], most available MMP inhibitors are broad-spectrum drugs [207]. Indeed, although MMP-3 is thought to contribute to neurodegeneration via multiple mechanisms, development of selective inhibitors for this proteolytic enzyme has been difficult, because the MMP family members share many structural characteristics. Pyrone-based

inhibitors with potency against MMP-3 and selectivity for MMP-1, -2 and -3 have been studied [404]; because MMP-3 prefers a more acidic environment than other MMPs, more acidic inhibitors, such as the carboxylates, have also been shown specific for MMP-3 [405].

However, the problem with these inhibitors, especially if used for long-term therapy, is that as side-effects they might block important functions, for example remodeling the ECM, leading to excessive fibrosis, and in particular causing joint stiffness [406]. Short-term use of MMP inhibitors may be less controversial in neurological disorders, particularly for treating cerebrovascular diseases. In addition, most non-peptidic MMP inhibitors reported thus far contain hydroxamic acid, carboxylic acids, phosphonates or thiols, and may not easily cross the BBB. Moreover, the most commonly used MMP inhibitors have poor solubility, and further studies are needed to improve delivery systems; the development of selective MMP inhibitors, with high BBB penetration rates and good solubility, would be of great benefit.

Alternatively, because MMP-3 gene expression is induced to a dramatic extent in response to cellular stress, while it is very low in basal conditions, it should be possible to suppress its upregulation without altering basal levels. In this connection, modification of MMP-3 gene expression could be a promising novel approach to providing neuroprotection. Tetracycline derivatives that penetrate the BBB, such as doxycycline and minocycline, downregulate the MMP-3 expression induced by cell stress, cytokines and other stimuli, with subsequent attenuation of neuroinflammation and apoptosis of dopaminergic neurons, both *in vitro* and *in vivo* [407-411]. Other compounds, such as ghrelin (an endogenous ligand for growth hormone secretagogue receptor 1a), exendin-4 (analog of glucagon-like peptide-1) or glycitein (a bacterial metabolite of the isoflavone glycitin) have also been found to attenuate MMP-3 expression, as well as loss of nigrostriatal dopaminergic neurons and hippocampal neurons [412-415].

Furthermore, because of its involvement in the non-amyloidogenic processing of APP, ADAM-10 overexpression or activation in the brain might be beneficial for the treatment of neurodegenerative diseases, in particular AD. In ADAM-10 overexpressing mice it has been shown that cortical synaptogenesis is enhanced [363]; in AD model mice, long-term potentiation deficiency is rescued [361], and learning, as well as memory, is positively influenced by ADAM-10 [362]. ADAM-10 activity can be enhanced, for example, by cholesterol depletion [416, 417] or by statin application [416]. Various results obtained from *Adam-10* transgenic mice have suggested that increasing ADAM-10 activity might be a valuable alternative to other strategies for treating AD, such as inhibiting β - or γ -secretase, or immunization. However, α -secretase activation must be moderate in extent, and needs to be closely monitored, since overexpression of ADAM-10 in the brain might affect homeostasis of the entire organism [354]. Again, also in this case, the crucial question is whether there are any side-effects connected with enhanced ADAM-10 activity in the brain or in peripheral tissues.

2.4. Metalloproteinases and Osteoarthritis

Osteoarthritis (OA) is a chronic degenerative joint disease that causes disability in the elderly, in the form of pain, stiffness and loss of function in articulating joints. OA is characterized by changes in the anatomy of load-bearing joints, which lead to degradation of the articular cartilage by proteolytic enzymes, inflammation of the synovium (synovitis), changes to subchondral bone, and growth of new bone and cartilage (osteophytes) at the joint edge [418-420]. The primary causes of OA are mechanical factors such as joint injury and obesity, with other risk factors, including age (associated intraarticular crystal deposition, muscle weakness and peripheral neuropathy), gender, and genetics, contributing to disease development and progression [418, 421, 422].

In OA, degradation of ECM molecules, which are synthesized and catabolized by chondrocytes, exceeds their synthesis, resulting in a net decrease in the amount of cartilage matrix, and eventually leading to total or partial erosion of the cartilage. Among the articular cartilage components, aggrecan and collagen are the main molecules that are slowly degraded. Aggrecan is a large proteoglycan, containing numerous chondroitin sulphate and keratan sulphate glycosaminoglycan moieties, which are important for the molecular function since they draw water into the cartilage matrix, giving it the ability to withstand compressive deformation during joint articulation. Degradation of aggrecan is an important manifestation of OA [15]. The depletion of aggrecan from articular cartilage is confirmed by the release of aggrecan catabolites into the synovial fluid [423]. Along with aggrecan breakdown, degradation of collagen, in particular of type II collagen, which provides strength to the tissue, is also a central feature of OA [424, 425]. Aggrecan is lost in the initial phases of the disease, while collagen is lost at the later stages.

Molecules mediating matrix degradation, including MMPs and ADAMTSs, are upregulated in OA chondrocytes [426, 427]. Studies on transgenic mice have confirmed the central role of ADAMTS-4 and ADAMTS-5 in aggrecan degradation, and that of the collagenolytic MMP-13 in collagen degradation [16].

Although ADAMTS-1, -8, -9, -15, -16 and -18 can degrade aggrecan *in vitro* [428-431] ADAMTS-5 is the most active “aggrecanase” *in vitro*, followed by ADAMTS-4 [432]. ADAMTS-4 and ADAMTS-5 are thus considered to be the major enzymes responsible for pathological cleavage of aggrecan at the Glu³⁷³-Ala³⁷⁴ bound in the interglobular domain [433-435]. Of note, studies in synovial cells suggest that ADAMTS-5 is constitutively expressed, whereas ADAMTS-4 is only induced by pro-inflammatory cytokines [436, 437].

The pathological importance of ADAMTS-5 in the development of OA was demonstrated by the finding that *Adamts-5* knockout mice develop less severe cartilage damage, both in a murine surgical model of OA, and in an antigen-induced arthritis model [438, 439]. In comparison, *Adamts-4* knockout mice did not show any significant suppression of aggrecanase activity in an arthritic model [440], indicating that ADAMTS-5 is the primary aggrecanase, at least in mice. Conversely, the role of ADAMTS-4 as aggrecanase appears more evident in humans [441, 442]. It has then been observed that ADAMTS-4 mRNA is induced in chondrocytes by IL-1 [443], while *Adamts-4/Adamts-5* double knockout mice are protected from cartilage degradation by IL-1, but not by retinoic acid, suggesting that other aggrecans apart from ADAMTS-4 and -5 are capable of retinoic acid-induced cartilage breakdown, at least in animal models [444]. In addition, suppressing ADAMTS-4 and/or ADAMTS-5 in human cartilage explants, via transfection of these ADAMTSs by siRNAs, significantly decreased aggrecan release and catabolism induced by a combination of IL-1 β , TNF- α and oncostatin M [441].

MMPs normally present in articular cartilage and bone, such as MMP-1, -2, -3, -7, -8, -9 and -13, are also thought to be capable of degrading cartilage components, primarily type II collagen. For example, fibrillar collagens, which are highly stable molecules, can be degraded by the collagenolytic MMP-1, -8, -13 and -14 and by cathepsin K. However, MMP-13 is thought to be the primary collagenase in OA, with its expression increased in OA cartilage [424, 445, 446], whereas *Mmp-13* knockout mice were protected by MMP-13 collagen degradation in a surgical OA model [447]. MMP-13 is also up-regulated during chondrocyte terminal differentiation, and deficiency of *Mmp-13* results in impaired endochondral ossification [448, 449].

However, in addition to ADAMTS-4 and -5 and MMP-13, expression of various other proteolytic enzymes (*i.e.* MMPs, ADAMs and ADAMTSs) has been reported to increase in OA. ADAM-8, for example, has been suggested to contribute to OA pathogenesis, by cleaving fibronectin, generating fragments that stimulate further cartilage catabolism [450]. MMP-3 is, then, the most abundantly secreted enzyme in OA cartilage, and it is known to induce activation of other MMPs, such as MMP-1 and MMP-13, raising the possibility that it may contribute to OA by

activating latent collagenases [436]. It has also been suggested that MMP-3 promotes collagenase activation, and either direct or indirect MMP-mediated aggrecan cleavage [451]. Higher levels of MMP-3 have, indeed, been found in the synovial fluid and serum of OA patients, compared with normal individuals [452, 453]. Expression of MMP-1 and MMP-8 as well as MMP-13 was also detected by immunostaining in the superficial zone of OA cartilage [454].

Fragmentation of aggrecan, type II collagen, fibronectin and hyaluronan reveals cryptic epitopes, which also stimulate proteolytic enzymes including MMPs and ADAMTSs. Proteolytic fragments also stimulate the release of NO, chemokines and cytokines [455]. Of note, it has been observed that human chondrocytes cultured with intact monomeric type II collagen induced expression of MMP-1, -3, -13, -14, as well as that of IL-1 β , IL-6, and IL-8 [456]. Similarly, collagen fragments generated during arthritis may influence matrix turnover. A fragment of type II collagen has been reported to upregulate levels of MMP-2, -3, -9 and -13 in bovine chondrocyte explants [457] and in human chondrocytes [458]. These results may suggest that type II collagen causes sequential activation of MMPs and cytokines during cartilage damage.

It is increasingly accepted that inflammation plays a role in OA pathogenesis [459-463]. Inflammatory cells, cytokines and growth factors that are not normally present in the cartilage matrix are found in OA patients [437, 464]. Inflammatory cytokines, for example, can increase chondrocyte expression of ADAMTS-4 and ADAMTS-5, leading to cartilage damage [465]. The cytokine IL-1 is suggested to be a principal mediator of joint damage in OA. Chondrocytes from OA cartilage display high levels of IL-1 α and IL-1 β , but are also more sensitive to IL-1. Moreover, IL-1 has the ability to stimulate chondrocytes to degrade both aggrecan and collagen. In cultured chondrocytes or cartilage explants, ADAMTS-4 has been shown to be induced following stimulation with IL-1, TNF- α or TGF β [441, 443, 466, 467] whereas ADAMTS-5 was upregulated by IL-1 α in an immortalized human chondrocyte cell line [468]. Using a model of cultured human synovial cells from digested OA synovium, it was observed that ADAMTS-4 gene expression is dependent on TNF- α , and that IL-1 is produced by the synovial macrophages, whereas the level of ADAMTS-5 is not significantly change by either of the two pro-inflammatory cytokines [437]. In contrast, monocytes from wild-type mice, but not those from *Il-1* deficient mice, upregulated ADAMTS-5 mRNA in chondrocytes, without affecting ADAMTS-4. This data suggest that murine ADAMTS-4 is unresponsive to IL-1 [469]. The role of NF- κ B in regulating ADAMTS gene expression has recently been clarified: ADAMTS-4, but not ADAMTS-5, has several NF- κ B binding sites. Of note, it has also been reported that IL-1 and TNF- α increase ADAMTS-4 expression in a NF- κ B dependent manner [470].

Moreover, IL-1 β is known to stimulate expression of MMPs, for example MMP-1 and -13, in OA cartilage [436,471]. The role of IL-1, together with TNF- α , in matrix degradation has been clarified [471]. Of note, when IL-1 is combined with TNF- α and the two are injected simultaneously, there is enhanced cartilage destruction, which exceeds the effects observed with either cytokine alone [472]. The combination of IL-1 and oncostatin M also upregulates matrix-degrading proteinases in cartilage [473]. In addition to inducing the synthesis of MMPs and other proteinases by chondrocytes, IL-1 and TNF- α increase the synthesis of PGE₂, by stimulating expression or activity of COX-2, microsomal PGE synthase-1, and soluble phospholipase A2 (PLA2); they also upregulate NO production via inducible NO synthetase (iNOS), contributing to the inflammatory response [474]. *Il-1 β* knockout mice showed protection against OA induced by destabilization of the medial meniscus [475]. Further, IL-17 and -18 are also thought to play a role in cartilage catabolism [476]. Osteoarthritic chondrocytes also overexpress TGF β , which has dual effects: it can be protective as well as deleterious for articular cartilage. This dual effect can be explained by the fact that TGF β can signal via different receptors and related the Smad signaling routes [477]. Importantly, signaling via anaplastic

lymphoma kinase (ALK)1, but not that via ALK5, stimulates MMP-13 expression by chondrocytes. In cartilage of aging mice, and in experimental OA models, it has been found that the ALK1/ALK5 ratio is significantly increased, favoring TGF β signaling via the Smad1/5/8 (ALK1) route, and augmenting changes in chondrocyte differentiation and MMP-13 expression [478]. Moreover, in human OA cartilage there is a significant correlation between ALK1 and MMP-13 expression [477]. Chemokines have also been reported to play a role in OA [479, 480]; expression of chemokine receptors has been demonstrated in OA chondrocytes [481] and synovial cells [482].

MMPs are strongly inhibited by all four of the mammalian TIMPs [483] but, conversely, ADAMTS-4 and ADAMTS-5 are effectively inhibited only by TIMP-3 [484-486]. As TIMP-3 can inhibit MMPs and ADAMTSs, it is considered to be a central inhibitor of cartilage degradation. The addition of TIMP-3 blocks cartilage degradation in explant cultures [487]. The key role played by TIMP-3 in cartilage protection has been confirmed by the finding that *Timp-3* knockout mice develop increased cartilage degradation upon aging [488] and display increased cartilage damage in an antigen-induced arthritis model [489]. In addition, TIMP-3 protein levels are reduced in human OA cartilage [490]. With regard to the other TIMPs, unlike their action on MMPs, TIMP-1 has been shown to partially inhibit glycosaminoglycan release from human cartilage, whereas TIMP-2 has no effect on cartilage components [465, 484]; the level of TIMP-4 is decreased in OA cartilage [446].

2.4.1. Therapeutic Strategies for Inhibiting Metalloproteinases in Osteoarthritis

OA remains a disease with insufficient disease-modifying treatments. With an increasing number of people suffering from the disease, the identification of novel therapeutic targets is a priority.

The routine therapies to moderate OA clinically are intraarticular injection of hyaluronic acid, which has the ability to aggregate aggrecan, or of steroids, or oral administration of anti-inflammatory molecules. These conventional OA therapeutics, however, do not inhibit the underlying tissue catabolism, and thus they allow the disease to progress into irreversible ECM loss and chronic disability.

Since the central role of aggrecanases (*e.g.* ADAMTS-4 and -5) and collagenases (*e.g.* MMP-13) in cartilage degradation has now been verified by studies on transgenic mice, identification of the key role played by these proteinases is now of importance for the successful development of specific inhibitors to be used as therapeutic agents in OA. It has been reported that blocking aggrecanase cleavage in the interglobular domain of aggrecan diminishes aggrecan loss and cartilage erosion in knockin mice, in surgically-induced OA and also in a model of inflammatory arthritis, and the procedure also appeared to stimulate cartilage repair following acute inflammation [435].

The design of small molecule inhibitors or protein antagonist inhibitors, to block the increase catabolism of matrix components in OA, is thus an area of considerable interest for the pharmaceutical industry. New inhibition strategies using small molecules inhibitors and TIMPs, engineered to increase their inhibitory specificity, or using new reagents such as ribozymes and siRNAs, which repress expression of specific enzymes, are thus now under consideration [491]. However, disappointing results from clinical trials with small molecule inhibitors have highlighted the critical importance of inhibitor specificity, and the need to better identify the individual enzymes responsible for joint destruction. The potential side-effects of inhibiting these enzymes, which are expressed in a number of tissue, are also still unclear. However, among agents targeting aggrecanase inhibitors, α -amino hydroxamate has been discovered to be a potent and selective aggrecanase inhibitor [492]; in subsequent attempts to optimize the potency and pharmacokinetic profile of this inhibitor, anti-succinate hydroxamates containing cyclic P1 substituents were identified [493]. Other potential selective synthetic inhibitors of ADAMTS-5, as well as of ADAMTS-4, have been investigated, *e.g.* a series of 5-((1H-pyrazol-4-yl)methylene)-2-thioxothiazolidin-4-one [494, 495].

Studies on mice with specific gene ablations have also identified a network of factors and cellular signals that regulate MMP-13 and ADAMTS expression in chondrocytes [16]. Given that many pathways can stimulate an increase in proteinase expression, the development of inhibitors targeting the effector proteinases, and their use in combination, may block cartilage damage more effectively than therapies targeting a single activating factor. For example, it has been reported that the OA process is driven by loss of the Smad2/3 block on differentiation in articular chondrocytes, leading to progression of chondrocyte differentiation and an autolytic phenotype. In the early stages of OA, some chondrocytes will have progressed in their differentiation to an OA chondrocytes phenotype, triggered by a loss of the Smad2/3 block; others will still be in a quiescent, healthy state of differentiation [477]. The latter cells could be targets for therapy, to block further progression of the OA process: blocking the progression of chondrocyte differentiation will block further expansion of the OA process in the remaining healthy cartilage; compounds specifically stimulating the Smad2/3 route could be developed as a therapeutic strategy [496, 497]. An alternative therapy could be stimulation of one of the other Smad2/3 routes in chondrocytes. In this connection, signaling via the activin ALK4 and ALK7 receptors leads to activation of the Smad2/3 pathway [498]. These receptors could be potential targets to enhance Smad2/3 signaling in chondrocytes in OA. Alternatively, ALK1, which is involved in vessel formation, or the Smad1/5/8 pathway, which is associated with the activity of ALK1, ALK2, ALK3 and ALK6, could be blocked in chondrocytes (for example with kinase inhibitors) with subsequent blocking of aberrant chondrocyte differentiation [477]. These possible therapeutic strategies, however, might trigger side-effects in other tissues: stimulation of Smad2/3 pathway using TGF β mimetics, or of the ALK4/7 pathway, might result in excessive induction of fibrosis; blocking ALK1 might reduce blood vessel formation; general inhibition of the Smad1/5/8 pathway might interfere with bone metabolism [477].

Furthermore, ADAMTS-4 is induced by IL-1 and TNF- α , and depends on the NF- κ B [470, 499, 500], and it has been reported that treatment of bovine cartilage explants with small-molecule I κ B kinase inhibitor leads to the prevention of IL-1-induced aggrecan degradation. These data suggest that aggrecan degradation occurs in a NF- κ B dependent manner [501]. Conversely, ADAMTS-5 has been found to be NF- κ B independent and to lack κ B elements on its promoter [500]. The differential regulation of ADAMTS-4 and ADAMTS-5 could thus have important implications for the development of disease-modifying OA drugs [502].

In addition, a more complete knowledge of the pathways and receptors used by endogenous matrix molecules in concert with cytokines and chemokines [455] may improve future possibilities for developing new therapeutic strategies for OA.

3. CONCLUSIONS

Metalloproteinases are a large family of enzymes that have been implicated in the pathophysiology of several chronic diseases related to aging, including atherosclerosis, type 2 diabetes mellitus, neurodegenerative diseases, and osteoarthritis (Figure 1). They thus represent potential therapeutic targets. In order to develop therapeutic strategies against the action of these proteolytic enzymes, it is fundamental to understand the regulation of and roles played by MMPs, ADAMs and ADAMTSs, and their endogenous inhibitors, especially of TIMPs, in all phases of the pathogenesis of these age-related diseases. More research is thus needed to clarify the effects of MMPs, not only on ECM degradation, but also on cell types involved in the pathogenesis of these diseases.

The design of agents that may treat or prevent the excessive degradation of ECM components, induced by MMP dysregulation, or the effects of MMPs on cells is now an area of considerable interest for the pharmaceutical industry. However, the use of broad spectrum synthetic inhibitors of MMPs, or of other exogenous therapeutic agents,

has not always replicated the effects of TIMPs in preclinical models and in clinical trials, possibly due to their side-effects, but also because these proteolytic enzymes possess reparative as well as pathogenetic properties. In this connection, the emphasis is therefore shifting to the development of MMP-modulating agents of restricted specificity. Despite this new impetus for developing more beneficial therapeutic strategies, more scientific studies are essential to improve our knowledge of protease function, thus helping to identify targets for targeted pharmacology therapy.

ACKNOWLEDGMENTS

The authors thank the Italian Ministry of University (Prin 2009), the CRT Foundation, Turin, and the University of Turin, Italy, for supporting this work.

REFERENCES

- [1] Swarnakar S, Mishra A, Chaudhuri SR. The gelatinases and their inhibitors: the structure-activity relationships. *EXS* 2012; 103: 57-82.
- [2] Weber S, Saftig P. Ectodomain shedding and ADAMs in development. *Development* 2012; 139: 3693-709.
- [3] Yong VW. Metalloproteinases: mediators of pathology and regeneration in the CNS. *Nat Rev Neurosci* 2005; 6: 931-44.
- [4] Kheradmand F, Werb Z. Shedding light on sheddases: role in growth and development. *Bioessays* 2002; 24: 8-12.
- [5] Schlöndorff J, Blobel CP. Metalloprotease-disintegrins: modular proteins capable of promoting cell-cell interactions and triggering signals by protein-ectodomain shedding. *J Cell Sci* 1999; 112: 3603-17.
- [6] Seals DF, Courtneidge SA. The ADAMs family of metalloproteases: multidomain proteins with multiple functions. *Genes Dev* 2003; 17: 7-30.
- [7] Yong VW, Power C, Forsyth P, Edwards DR. Metalloproteinases in biology and pathology of the nervous system. *Nat Rev Neurosci* 2001; 2: 502-11.
- [8] Blobel CP. Remarkable roles of proteolysis on and beyond the cell surface. *Curr Opin Cell Biol* 2000; 12: 606-12.
- [9] Alfandari D, McCusker C, Cousin H. ADAM function in embryogenesis. *Semin Cell Dev Biol* 2009; 20: 153-63.
- [10] Drey Mueller D, Pruessmeyer J, Groth E, Ludwig A. The role of ADAM-mediated shedding in vascular biology. *Eur J Cell Biol* 2012; 91: 472-85.
- [11] Marcinkiewicz M, Seidah NG. Coordinated expression of beta-amyloid precursor protein and the putative beta-secretase BACE and alpha-secretase ADAM10 in mouse and human brain. *J Neurochem* 2000; 75: 2133-43.
- [12] Kärkkäinen I, Rybnikova E, Pelto-Huikko M, Huovila AP. Metalloprotease-disintegrin (ADAM) genes are widely and differentially expressed in the adult CNS. *Mol Cell Neurosci* 2000; 15: 547-60.
- [13] Dallas DJ, Genever PG, Patton AJ, Millichip MI, McKie N, Skerry TM. Localization of ADAM10 and Notch receptors in bone. *Bone* 1999; 25: 9-15.
- [14] McKie N, Edwards T, Dallas DJ. Expression of members of a novel membrane linked metalloproteinase family (ADAM) in human articular chondrocytes. *Biochem Biophys Res Commun* 1997; 230: 335-9.
- [15] Huang K, Wu LD. Aggrecanase and aggrecan degradation in osteoarthritis: a review. *J Int Med Res* 2008; 36: 1149-60.
- [16] Troeberg L, Nagase H. Proteases involved in cartilage matrix degradation in osteoarthritis. *Biochim Biophys Acta* 2012; 1824: 133-45.

- [17] Limb GA, Matter K, Murphy G *et al.* Matrix metalloproteinase-1 associates with intracellular organelles and confers resistance to lamin A/C degradation during apoptosis. *Am J Pathol* 2005; 166: 1555-63.
- [18] Kwan JA, Schulze CJ, Wang W *et al.* Matrix metalloproteinase-2 (MMP-2) is present in the nucleus of cardiac myocytes and is capable of cleaving poly (ADP-ribose) polymerase (PARP) in vitro. *FASEB J* 2004; 18: 690-2.
- [19] Luo D, Mari B, Stoll I, Anglard P. Alternative splicing and promoter usage generates an intracellular stromelysin 3 isoform directly translated as an active matrix metalloproteinase. *J Biol Chem* 2002; 277: 25527-36.
- [20] Newby AC. Dual role of matrix metalloproteinases (matrixins) in intimal thickening and atherosclerotic plaque rupture. *Physiol Rev* 2005; 85: 1-31.
- [21] Nagase H, Visse R, Murphy G. Structure and function of matrix metalloproteinases and TIMPs. *Cardiovasc Res* 2006; 69: 562-73.
- [22] Sternlicht MD, Werb Z. How matrix metalloproteinases regulate cell behavior. *Annu Rev Cell Dev Biol* 2001; 17: 463-516.
- [23] Rosenberg GA. Matrix metalloproteinases in neuroinflammation. *Glia* 2002; 39: 279-91.
- [24] Mott JD, Werb Z. Regulation of matrix biology by matrix metalloproteinases. *Curr Opin Cell Biol* 2004; 16: 558-64.
- [25] Parks WC, Wilson CL, López-Boado YS. Matrix metalloproteinases as modulators of inflammation and innate immunity. *Nat Rev Immunol* 2004; 4: 617-29.
- [26] Egeblad M, Werb Z. New functions for the matrix metalloproteinases in cancer progression. *Nat Rev Cancer* 2002; 2: 161-74.
- [27] Cauwe B, Van den Steen PE, Opdenakker G. The biochemical, biological, and pathological kaleidoscope of cell surface substrates processed by matrix metalloproteinases. *Crit Rev Biochem Mol Biol* 2007; 42: 113-85.
- [28] Vargová V, Pytliak M, Mechírová V. Matrix metalloproteinases. *EXS* 2012; 103: 1-33.
- [29] Agrawal SM, Lau L, Yong VW. MMPs in the central nervous system: where the good guys go bad. *Semin Cell Dev Biol* 2008; 19: 42-51.
- [30] Visse R, Nagase H. Matrix metalloproteinases and tissue inhibitors of metalloproteinases: structure, function, and biochemistry. *Circ Res* 2003; 92: 827-39.
- [31] Rajagopalan S, Meng XP, Ramasamy S, Harrison DG, Galis ZS. Reactive oxygen species produced by macrophage-derived foam cells regulate the activity of vascular matrix metalloproteinases in vitro. Implications for atherosclerotic plaque stability. *J Clin Invest* 1996; 98: 2572-9.
- [32] Gu Z, Kaul M, Yan B *et al.* S-nitrosylation of matrix metalloproteinases: signaling pathway to neuronal cell death. *Science* 2002; 297: 1186-90.
- [33] Carmeliet P, Moons L, Herbert JM *et al.* Urokinase but not tissue plasminogen activator mediates arterial neointima formation in mice. *Circ Res* 1997; 81: 829-39.
- [34] Browatzki M, Larsen D, Pfeiffer CA *et al.* Angiotensin II stimulates matrix metalloproteinase secretion in human vascular smooth muscle cells via nuclear factor-kappaB and activator protein 1 in a redox-sensitive manner. *J Vasc Res* 2005; 42: 415-23.
- [35] Death AK, Fisher EJ, McGrath KC, Yue DK. High glucose alters matrix metalloproteinase expression in two key vascular cells: potential impact on atherosclerosis in diabetes. *Atherosclerosis* 2003; 168: 263-9.
- [36] Nelson AR, Fingleton B, Rothenberg ML, Matrisian LM. Matrix metalloproteinases: biologic activity and clinical implications. *J Clin Oncol* 2000; 18: 1135-49.

- [37] Chakraborti S, Mandal M, Das S, Mandal A, Chakraborti T. Regulation of matrix metalloproteinases: an overview. *Mol Cell Biochem* 2003; 253: 269-85.
- [38] Schmidt R, Bültmann A, Ungerer M *et al.* Extracellular matrix metalloproteinase inducer regulates matrix metalloproteinase activity in cardiovascular cells: implications in acute myocardial infarction. *Circulation* 2006; 113: 834-41.
- [39] Yoon YW, Kwon HM, Hwang KC *et al.* Upstream regulation of matrix metalloproteinase by EMMPRIN; extracellular matrix metalloproteinase inducer in advanced atherosclerotic plaque. *Atherosclerosis* 2005; 180: 37-44.
- [40] Cawston TE, Mercer E. Preferential binding of collagenase to alpha 2-macroglobulin in the presence of the tissue inhibitor of metalloproteinases. *FEBS Lett* 1986; 209: 9-12.
- [41] Baker AH, Edwards DR, Murphy G. Metalloproteinase inhibitors: biological actions and therapeutic opportunities. *J Cell Sci* 2002; 115: 3719-27.
- [42] Brew K, Dinakarpanian D, Nagase H. Tissue inhibitors of metalloproteinases: evolution, structure and function. *Biochim Biophys Acta* 2000; 1477: 267-83.
- [43] Amour A, Slocombe PM, Webster A *et al.* TNF-alpha converting enzyme (TACE) is inhibited by TIMP-3. *FEBS Lett* 1998; 435: 39-44.
- [44] Higashi S, Miyazaki K. Identification of a region of beta-amyloid precursor protein essential for its gelatinase A inhibitory activity. *J Biol Chem* 2003; 278: 14020-8.
- [45] Mott JD, Thomas CL, Rosenbach MT, Takahara K, Greenspan DS, Banda MJ. Post-translational proteolytic processing of procollagen C-terminal proteinase enhancer releases a metalloproteinase inhibitor. *J Biol Chem* 2000; 275: 1384-90.
- [46] Oh J, Takahashi R, Kondo S *et al.* The membrane-anchored MMP inhibitor RECK is a key regulator of extracellular matrix integrity and angiogenesis. *Cell* 2001; 107: 789-800.
- [47] Mauviel A. Cytokine regulation of metalloproteinase gene expression. *J Cell Biochem* 1993; 53: 288-95.
- [48] Spinale FG. Matrix metalloproteinases: regulation and dysregulation in the failing heart. *Circ Res* 2002; 90: 520-30.
- [49] Newby AC. Metalloproteinases and vulnerable atherosclerotic plaques. *Trends Cardiovasc Med* 2007; 17: 253-8.
- [50] Raffetto JD, Khalil RA. Matrix metalloproteinases and their inhibitors in vascular remodeling and vascular disease. *Biochem Pharmacol* 2008; 75: 346-59.
- [51] Busti C, Falcinelli E, Momi S, Gresele P. Matrix metalloproteinases and peripheral arterial disease. *Intern Emerg Med* 2010; 5: 13-25.
- [52] Ye S. Influence of matrix metalloproteinase genotype on cardiovascular disease susceptibility and outcome. *Cardiovasc Res* 2006; 69: 636-45.
- [53] Ghaderian SM, Akbarzadeh Najar R, Tabatabaei Panah AS. Genetic polymorphisms and plasma levels of matrix metalloproteinases and their relationships with developing acute myocardial infarction. *Coron Artery Dis* 2010; 21: 330-5.
- [54] Kunz J. Matrix metalloproteinases and atherogenesis in dependence of age. *Gerontology* 2007; 53: 63-73.
- [55] Romanic AM, Harrison SM, Bao W *et al.* Myocardial protection from ischemia/reperfusion injury by targeted deletion of matrix metalloproteinase-9. *Cardiovasc Res* 2002; 54: 549-58.
- [56] Matsumura S, Iwanaga S, Mochizuki S, Okamoto H, Ogawa S, Okada Y. Targeted deletion or pharmacological inhibition of MMP-2 prevents cardiac rupture after myocardial infarction in mice. *J Clin Invest* 2005; 115: 599-609.

- [57] Longo GM, Xiong W, Greiner TC, Zhao Y, Fiotti N, Baxter BT. Matrix metalloproteinases 2 and 9 work in concert to produce aortic aneurysms. *J Clin Invest* 2002; 110: 625-32.
- [58] Galt SW, Lindemann S, Allen L *et al.* Outside-in signals delivered by matrix metalloproteinase-1 regulate platelet function. *Circ Res* 2002; 90: 1093-9.
- [59] Sawicki G, Sanders EJ, Salas E, Wozniak M, Rodrigo J, Radomski MW. Localization and translocation of MMP-2 during aggregation of human platelets. *Thromb Haemost* 1998; 80: 836-9.
- [60] Fedak PW, Smookler DS, Kassiri Z *et al.* TIMP-3 deficiency leads to dilated cardiomyopathy. *Circulation* 2004; 110: 2401-9.
- [61] Insull W Jr. The pathology of atherosclerosis: plaque development and plaque responses to medical treatment. *Am J Med* 2009; 122: S3-14.
- [62] Loftus I, Thompson M. Plaque biology: interesting science or pharmacological treasure trove? *Eur J Vasc Endovasc Surg* 2008; 36: 507-16.
- [63] Aikawa M, Libby P. The vulnerable atherosclerotic plaque: pathogenesis and therapeutic approach. *Cardiovasc Pathol* 2004; 13: 125-38.
- [64] Virmani R, Kolodgie FD, Burke AP, Farb A, Schwartz SM. Lessons from sudden coronary death: a comprehensive morphological classification scheme for atherosclerotic lesions. *Arterioscler Thromb Vasc Biol* 2000; 20: 1262-75.
- [65] Virmani R, Burke AP, Farb A, Kolodgie FD. Pathology of the vulnerable plaque. *J Am Coll Cardiol* 2006; 47: C13-8.
- [66] Halvorsen B, Otterdal K, Dahl TB *et al.* Atherosclerotic plaque stability--what determines the fate of a plaque? *Prog Cardiovasc Dis* 2008; 51: 183-94.
- [67] Buffon A, Biasucci LM, Liuzzo G, D'Onofrio G, Crea F, Maseri A. Widespread coronary inflammation in unstable angina. *N Engl J Med* 2002; 347: 5-12.
- [68] Clarke MC, Figg N, Maguire JJ *et al.* Apoptosis of vascular smooth muscle cells induces features of plaque vulnerability in atherosclerosis. *Nat Med* 2006; 12: 1075-80.
- [69] Tabas I. Consequences and therapeutic implications of macrophage apoptosis in atherosclerosis: the importance of lesion stage and phagocytic efficiency. *Arterioscler Thromb Vasc Biol* 2005; 25: 2255-64.
- [70] Dickson BC, Gotlieb AI. Towards understanding acute destabilization of vulnerable atherosclerotic plaques. *Cardiovasc Pathol* 2003; 12: 237-48.
- [71] Lutgens E, van Suylen RJ, Faber BC *et al.* Atherosclerotic plaque rupture: local or systemic process? *Arterioscler Thromb Vasc Biol* 2003; 23: 2123-30.
- [72] Nakamura M, Lee DP, Yeung AC. Identification and treatment of vulnerable plaque. *Rev Cardiovasc Med* 2004; 2: S22-33.
- [73] Wang LX, Lü SZ, Zhang WJ, Song XT, Chen H, Zhang LJ. Coronary spasm, a pathogenic trigger of vulnerable plaque rupture. *Chin Med J (Engl)* 2011; 124: 4071-8.
- [74] Gijssen F, van der Giessen A, van der Steen A, Wentzel J. Shear stress and advanced atherosclerosis in human coronary arteries. *J Biomech* 2013; 46: 240-7.
- [75] Toutouzas K, Synetos A, Nikolaou C, Tsiamis E, Tousoulis D, Stefanadis C. Matrix metalloproteinases and vulnerable atheromatous plaque. *Curr Top Med Chem* 2012; 12: 1166-80.
- [76] Loftus IM, Naylor AR, Bell PR, Thompson MM. Matrix metalloproteinases and atherosclerotic plaque instability. *Br J Surg* 2002; 89: 680-94.

- [77] Bäck M, Ketelhuth DF, Agewall S. Matrix metalloproteinases in atherothrombosis. *Prog Cardiovasc Dis* 2010; 52: 410-28.
- [78] Newby AC. Metalloproteinase expression in monocytes and macrophages and its relationship to atherosclerotic plaque instability. *Arterioscler Thromb Vasc Biol* 2008; 28: 2108-14.
- [79] Reel B, Sala-Newby GB, Huang WC, Newby AC. Diverse patterns of cyclooxygenase-independent metalloproteinase gene regulation in human monocytes. *Br J Pharmacol* 2011; 163: 1679-90.
- [80] Amorino GP, Hoover RL. Interactions of monocytic cells with human endothelial cells stimulate monocytic metalloproteinase production. *Am J Pathol* 1998; 152: 199-207.
- [81] Galt SW, Lindemann S, Medd D *et al.* Differential regulation of matrix metalloproteinase-9 by monocytes adherent to collagen and platelets. *Circ Res* 2001; 89: 509-16.
- [82] Galis ZS, Sukhova GK, Kranzhöfer R, Clark S, Libby P. Macrophage foam cells from experimental atheroma constitutively produce matrix-degrading proteinases. *Proc Natl Acad Sci USA* 1995; 92: 402-6.
- [83] Chase AJ, Bond M, Crook MF, Newby AC. Role of nuclear factor-kappa B activation in metalloproteinase-1, -3, and -9 secretion by human macrophages in vitro and rabbit foam cells produced in vivo. *Arterioscler Thromb Vasc Biol* 2002; 22: 765-71.
- [84] Feinberg MW, Jain MK, Werner F *et al.* Transforming growth factor-beta 1 inhibits cytokine-mediated induction of human metalloelastase in macrophages. *J Biol Chem* 2000; 275: 25766-73.
- [85] Mach F, Schönbeck U, Bonnefoy JY, Pober JS, Libby P. Activation of monocyte/macrophage functions related to acute atheroma complication by ligation of CD40: induction of collagenase, stromelysin, and tissue factor. *Circulation* 1997; 96: 396-9.
- [86] Schönbeck U, Mach F, Sukhova GK *et al.* Expression of stromelysin-3 in atherosclerotic lesions: regulation via CD40-CD40 ligand signaling in vitro and in vivo. *J Exp Med* 1999; 189: 843-53.
- [87] Wu L, Fan J, Matsumoto Si, Watanabe T. Induction and regulation of matrix metalloproteinase-12 by cytokines and CD40 signaling in monocyte/macrophages. *Biochem Biophys Res Commun* 2000; 269: 808-15.
- [88] Herman MP, Sukhova GK, Libby P *et al.* Expression of neutrophil collagenase (matrix metalloproteinase-8) in human atheroma: a novel collagenolytic pathway suggested by transcriptional profiling. *Circulation* 2001; 104: 1899-904.
- [89] Moreau M, Brocheriou I, Petit L, Ninio E, Chapman MJ, Rouis M. Interleukin-8 mediates downregulation of tissue inhibitor of metalloproteinase-1 expression in cholesterol-loaded human macrophages: relevance to stability of atherosclerotic plaque. *Circulation* 1999; 99: 420-6.
- [90] George SJ, Zaltsman AB, Newby AC. Surgical preparative injury and neointima formation increase MMP-9 expression and MMP-2 activation in human saphenous vein. *Cardiovasc Res* 1997; 33: 447-59.
- [91] Johnson JL, van Eys GJ, Angelini GD, George SJ. Injury induces dedifferentiation of smooth muscle cells and increased matrix-degrading metalloproteinase activity in human saphenous vein. *Arterioscler Thromb Vasc Biol* 2001; 21: 1146-51.
- [92] Galis ZS, Muszynski M, Sukhova GK *et al.* Cytokine-stimulated human vascular smooth muscle cells synthesize a complement of enzymes required for extracellular matrix digestion. *Circ Res* 1994; 75: 181-9.
- [93] Fabunmi RP, Baker AH, Murray EJ, Booth RF, Newby AC. Divergent regulation by growth factors and cytokines of 95 kDa and 72 kDa gelatinases and tissue inhibitors or metalloproteinases-1, -2, and -3 in rabbit aortic smooth muscle cells. *Biochem J* 1996; 315: 335-42.

- [94] Bond M, Chase AJ, Baker AH, Newby AC. Inhibition of transcription factor NF-kappaB reduces matrix metalloproteinase-1, -3 and -9 production by vascular smooth muscle cells. *Cardiovasc Res* 2001; 50: 556-65.
- [95] Schönbeck U, Mach F, Sukhova GK *et al.* Regulation of matrix metalloproteinase expression in human vascular smooth muscle cells by T lymphocytes: a role for CD40 signaling in plaque rupture? *Circ Res* 1997; 81: 448-54.
- [96] Galis ZS, Sukhova GK, Lark MW, Libby P. Increased expression of matrix metalloproteinases and matrix degrading activity in vulnerable regions of human atherosclerotic plaques. *J Clin Invest* 1994; 94: 2493-503.
- [97] Yakubenko VP, Lobb RR, Plow EF, Ugarova TP. Differential induction of gelatinase B (MMP-9) and gelatinase A (MMP-2) in T lymphocytes upon alpha(4)beta(1)-mediated adhesion to VCAM-1 and the CS-1 peptide of fibronectin. *Exp Cell Res* 2000; 260: 73-84.
- [98] Baram D, Vaday GG, Salamon P, Drucker I, Hershkoviz R, Mekori YA. Human mast cells release metalloproteinase-9 on contact with activated T cells: juxtacrine regulation by TNF-alpha. *J Immunol* 2001; 167: 4008-16.
- [99] Hanemaaijer R, Koolwijk P, le Clercq L, de Vree WJ, van Hinsbergh VW. Regulation of matrix metalloproteinase expression in human vein and microvascular endothelial cells. Effects of tumour necrosis factor alpha, interleukin 1 and phorbol ester. *Biochem J* 1993; 296: 803-9.
- [100] Hojo Y, Ikeda U, Takahashi M *et al.* Matrix metalloproteinase-1 expression by interaction between monocytes and vascular endothelial cells. *J Mol Cell Cardiol* 2000; 32: 1459-68.
- [101] Newby AC. Matrix metalloproteinase inhibition therapy for vascular diseases. *Vascul Pharmacol* 2012; 56: 232-44.
- [102] Mach F, Schönbeck U, Fabunmi RP *et al.* T lymphocytes induce endothelial cell matrix metalloproteinase expression by a CD40L-dependent mechanism: implications for tubule formation. *Am J Pathol* 1999; 154: 229-38.
- [103] Huang Y, Song L, Wu S, Fan F, Lopes-Virella MF. Oxidized LDL differentially regulates MMP-1 and TIMP-1 expression in vascular endothelial cells. *Atherosclerosis* 2001; 156: 119-25.
- [104] Santos-Martínez MJ, Medina C, Jurasz P, Radomski MW. Role of metalloproteinases in platelet function. *Thromb Res* 2008; 121: 535-42.
- [105] Gresele P, Falcinelli E, Momi S. Potentiation and priming of platelet activation: a potential target for antiplatelet therapy. *Trends Pharmacol Sci* 2008; 29: 352-60.
- [106] Xu XP, Meisel SR, Ong JM *et al.* Oxidized low-density lipoprotein regulates matrix metalloproteinase-9 and its tissue inhibitor in human monocyte-derived macrophages. *Circulation* 1999; 99: 993-8.
- [107] Rajavashisth TB, Xu XP, Jovinge S *et al.* Membrane type 1 matrix metalloproteinase expression in human atherosclerotic plaques: evidence for activation by proinflammatory mediators. *Circulation* 1999; 99: 3103-9.
- [108] Huang Y, Mironova M, Lopes-Virella MF. Oxidized LDL stimulates matrix metalloproteinase-1 expression in human vascular endothelial cells. *Arterioscler Thromb Vasc Biol* 1999; 19: 2640-7.
- [109] Silence J, Collen D, Lijnen HR. Reduced atherosclerotic plaque but enhanced aneurysm formation in mice with inactivation of the tissue inhibitor of metalloproteinase-1 (TIMP-1) gene. *Circ Res* 2002; 90: 897-903.
- [110] Lemaître V, Soloway PD, D'Armiento J. Increased medial degradation with pseudo-aneurysm formation in apolipoprotein E-knockout mice deficient in tissue inhibitor of metalloproteinases-1. *Circulation* 2003; 107: 333-8.
- [111] Gough PJ, Gomez IG, Wille PT, Raines EW. Macrophage expression of active MMP-9 induces acute plaque disruption in apoE-deficient mice. *J Clin Invest* 2006; 116: 59-69.

- [112] de Nooijer R, Verkleij CJ, von der Thüsen JH *et al.* Lesional overexpression of matrix metalloproteinase-9 promotes intraplaque hemorrhage in advanced lesions but not at earlier stages of atherogenesis. *Arterioscler Thromb Vasc Biol* 2006; 26: 340-6.
- [113] Lemaître V, O'Byrne TK, Borczuk AC, Okada Y, Tall AR, D'Armiento J. ApoE knockout mice expressing human matrix metalloproteinase-1 in macrophages have less advanced atherosclerosis. *J Clin Invest* 2001; 107: 1227-34.
- [114] Fukumoto Y, Deguchi JO, Libby P *et al.* Genetically determined resistance to collagenase action augments interstitial collagen accumulation in atherosclerotic plaques. *Circulation* 2004; 110: 1953-9.
- [115] Deguchi JO, Aikawa E, Libby P *et al.* Matrix metalloproteinase-13/collagenase-3 deletion promotes collagen accumulation and organization in mouse atherosclerotic plaques. *Circulation* 2005; 112: 2708-15.
- [116] Kuzuya M, Nakamura K, Sasaki T, Cheng XW, Itohara S, Iguchi A. Effect of MMP-2 deficiency on atherosclerotic lesion formation in apoE-deficient mice. *Arterioscler Thromb Vasc Biol* 2006; 26: 1120-5.
- [117] Silence J, Lupu F, Collen D, Lijnen HR. Persistence of atherosclerotic plaque but reduced aneurysm formation in mice with stromelysin-1 (MMP-3) gene inactivation. *Arterioscler Thromb Vasc Biol* 2001; 21: 1440-5.
- [118] Lutun A, Lutgens E, Manderveld A *et al.* Loss of matrix metalloproteinase-9 or matrix metalloproteinase-12 protects apolipoprotein E-deficient mice against atherosclerotic media destruction but differentially affects plaque growth. *Circulation* 2004; 109: 1408-14.
- [119] Johnson JL, George SJ, Newby AC, Jackson CL. Divergent effects of matrix metalloproteinases 3, 7, 9, and 12 on atherosclerotic plaque stability in mouse brachiocephalic arteries. *Proc Natl Acad Sci USA* 2005; 102: 15575-80.
- [120] Johnson JL, Dwivedi A, Somerville M, George SJ, Newby AC. Matrix metalloproteinase (MMP)-3 activates MMP-9 mediated vascular smooth muscle cell migration and neointima formation in mice. *Arterioscler Thromb Vasc Biol* 2011; 31: e35-44.
- [121] Williams H, Johnson JL, Jackson CL, White SJ, George SJ. MMP-7 mediates cleavage of N-cadherin and promotes smooth muscle cell apoptosis. *Cardiovasc Res* 2010; 87: 137-46.
- [122] Johnson JL, Devel L, Czarny B *et al.* A selective matrix metalloproteinase-12 inhibitor retards atherosclerotic plaque development in apolipoprotein E-knockout mice. *Arterioscler Thromb Vasc Biol* 2011; 31: 528-35.
- [123] Liang J, Liu E, Yu Y *et al.* Macrophage metalloelastase accelerates the progression of atherosclerosis in transgenic rabbits. *Circulation* 2006; 113: 1993-2001.
- [124] Laxton RC, Hu Y, Duchene J *et al.* A role of matrix metalloproteinase-8 in atherosclerosis. *Circ Res* 2009; 105: 921-9.
- [125] Schneider F, Sukhova GK, Aikawa M *et al.* Matrix-metalloproteinase-14 deficiency in bone-marrow-derived cells promotes collagen accumulation in mouse atherosclerotic plaques. *Circulation* 2008; 117: 931-9.
- [126] Galis ZS, Khatri JJ. Matrix metalloproteinases in vascular remodeling and atherogenesis: the good, the bad, and the ugly. *Circ Res* 2002; 90: 251-62.
- [127] Newby AC. Matrix metalloproteinases regulate migration, proliferation, and death of vascular smooth muscle cells by degrading matrix and non-matrix substrates. *Cardiovasc Res* 2006; 69: 614-24.
- [128] Cho A, Reidy MA. Matrix metalloproteinase-9 is necessary for the regulation of smooth muscle cell replication and migration after arterial injury. *Circ Res* 2002; 91: 845-51.
- [129] Galis ZS, Johnson C, Godin D *et al.* Targeted disruption of the matrix metalloproteinase-9 gene impairs smooth muscle cell migration and geometrical arterial remodeling. *Circ Res* 2002; 91: 852-9.

- [130] Choi ET, Collins ET, Marine LA *et al.* Matrix metalloproteinase-9 modulation by resident arterial cells is responsible for injury-induced accelerated atherosclerotic plaque development in apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol* 2005; 25: 1020-5.
- [131] Filippov S, Koenig GC, Chun TH *et al.* MT1-matrix metalloproteinase directs arterial wall invasion and neointima formation by vascular smooth muscle cells. *J Exp Med* 2005; 202: 663-71.
- [132] Wu YJ, Bond M, Sala-Newby GB, Newby AC. Altered S-phase kinase-associated protein-2 levels are a major mediator of cyclic nucleotide-induced inhibition of vascular smooth muscle cell proliferation. *Circ Res* 2006; 98: 1141-50.
- [133] Johnson JL, Baker AH, Oka K *et al.* Suppression of atherosclerotic plaque progression and instability by tissue inhibitor of metalloproteinase-2: involvement of macrophage migration and apoptosis. *Circulation* 2006; 113: 2435-44.
- [134] Johnson JL, Fritsche-Danielson R, Behrendt M *et al.* Effect of broad-spectrum matrix metalloproteinase inhibition on atherosclerotic plaque stability. *Cardiovasc Res* 2006; 71: 586-95.
- [135] Tedgui A, Mallat Z. Cytokines in atherosclerosis: pathogenic and regulatory pathways. *Physiol Rev* 2006; 86: 515-81.
- [136] Katsuda S, Kaji T. Atherosclerosis and extracellular matrix. *J Atheroscler Thromb* 2003; 10: 267-74.
- [137] Choudhary S, Higgins CL, Chen IY *et al.* Quantitation and localization of matrix metalloproteinases and their inhibitors in human carotid endarterectomy tissues. *Arterioscler Thromb Vasc Biol* 2006; 26: 2351-8.
- [138] Sluijter JP, Pulskens WP, Schoneveld AH *et al.* Matrix metalloproteinase 2 is associated with stable and matrix metalloproteinases 8 and 9 with vulnerable carotid atherosclerotic lesions: a study in human endarterectomy specimen pointing to a role for different extracellular matrix metalloproteinase inducer glycosylation forms. *Stroke* 2006; 37: 235-9.
- [139] Sukhova GK, Schönbeck U, Rabkin E *et al.* Evidence for increased collagenolysis by interstitial collagenases-1 and -3 in vulnerable human atheromatous plaques. *Circulation* 1999; 99: 2503-9.
- [140] Monaco C, Gregan SM, Navin TJ, Foxwell BM, Davies AH, Feldmann M. Toll-like receptor-2 mediates inflammation and matrix degradation in human atherosclerosis. *Circulation* 2009; 120: 2462-9.
- [141] Thomas AC, Sala-Newby GB, Ismail Y, Johnson JL, Pasterkamp G, Newby AC. Genomics of foam cells and nonfoamy macrophages from rabbits identifies arginase-I as a differential regulator of nitric oxide production. *Arterioscler Thromb Vasc Biol* 2007; 27: 571-7.
- [142] Johnson JL, Sala-Newby GB, Ismail Y, Aguilera CM, Newby AC. Low tissue inhibitor of metalloproteinases 3 and high matrix metalloproteinase 14 levels defines a subpopulation of highly invasive foam-cell macrophages. *Arterioscler Thromb Vasc Biol* 2008; 28: 1647-53.
- [143] Ikeda U, Shimada K. Matrix metalloproteinases and coronary artery diseases. *Clin Cardiol* 2003; 26: 55-9.
- [144] Orbe J, Fernandez L, Rodríguez JA *et al.* Different expression of MMPs/TIMP-1 in human atherosclerotic lesions. Relation to plaque features and vascular bed. *Atherosclerosis* 2003; 170: 269-76.
- [145] Noji Y, Kajinami K, Kawashiri MA *et al.* Circulating matrix metalloproteinases and their inhibitors in premature coronary atherosclerosis. *Clin Chem Lab Med* 2001; 39: 380-4.
- [146] Beaudoux JL, Giral P, Bruckert E, Bernard M, Foglietti MJ, Chapman MJ. Serum matrix metalloproteinase-3 and tissue inhibitor of metalloproteinases-1 as potential markers of carotid atherosclerosis in infraclinical hyperlipidemia. *Atherosclerosis* 2003; 169: 139-46.
- [147] Partridge CR, Hawker JR Jr, Forough R. Overexpression of a secretory form of FGF-1 promotes MMP-1-mediated endothelial cell migration. *J Cell Biochem* 2000; 78: 487-99.

- [148] Holnthoner W, Kerenyi M, Gröger M, Kratochvill F, Petzelbauer P. Regulation of matrilysin expression in endothelium by fibroblast growth factor-2. *Biochem Biophys Res Commun* 2006; 342: 725-33.
- [149] Burbridge MF, Cogé F, Galizzi JP, Boutin JA, West DC, Tucker GC. The role of the matrix metalloproteinases during in vitro vessel formation. *Angiogenesis* 2002; 5: 215-26.
- [150] Duhamel-Clérin E, Orvain C, Lanza F, Cazenave JP, Klein-Soyer C. Thrombin receptor-mediated increase of two matrix metalloproteinases, MMP-1 and MMP-3, in human endothelial cells. *Arterioscler Thromb Vasc Biol* 1997; 17: 1931-8.
- [151] Lafleur MA, Hollenberg MD, Atkinson SJ, Knäuper V, Murphy G, Edwards DR. Activation of pro-(matrix metalloproteinase-2) (pro-MMP-2) by thrombin is membrane-type-MMP-dependent in human umbilical vein endothelial cells and generates a distinct 63 kDa active species. *Biochem J* 2001; 357: 107-15.
- [152] Rybakowski JK. Matrix metalloproteinase-9 (MMP9)-A mediating enzyme in cardiovascular disease, cancer, and neuropsychiatric disorders. *Cardiovasc Psychiatry Neurol* 2009; 2009: 904836.
- [153] Pöllänen PJ, Karhunen PJ, Mikkelsen J *et al.* Coronary artery complicated lesion area is related to functional polymorphism of matrix metalloproteinase 9 gene: an autopsy study. *Arterioscler Thromb Vasc Biol* 2001; 21: 1446-50.
- [154] Morgan AR, Zhang B, Tapper W, Collins A, Ye S. Haplotypic analysis of the MMP-9 gene in relation to coronary artery disease. *J Mol Med (Berl)* 2003; 81: 321-6.
- [155] Blankenberg S, Rupprecht HJ, Poirier O *et al.* Plasma concentrations and genetic variation of matrix metalloproteinase 9 and prognosis of patients with cardiovascular disease. *Circulation* 2003; 107: 1579-85.
- [156] Johnson C, Galis ZS. Matrix metalloproteinase-2 and -9 differentially regulate smooth muscle cell migration and cell-mediated collagen organization. *Arterioscler Thromb Vasc Biol* 2004; 24: 54-60.
- [157] Loftus IM, Naylor AR, Goodall S *et al.* Increased matrix metalloproteinase-9 activity in unstable carotid plaques. A potential role in acute plaque disruption. *Stroke* 2000; 31: 40-7.
- [158] Chen F, Eriksson P, Hansson GK *et al.* Expression of matrix metalloproteinase 9 and its regulators in the unstable coronary atherosclerotic plaque. *Int J Mol Med* 2005; 15: 57-65.
- [159] Brown DL, Hibbs MS, Kearney M, Loushin C, Isner JM. Identification of 92-kD gelatinase in human coronary atherosclerotic lesions. Association of active enzyme synthesis with unstable angina. *Circulation* 1995; 91: 2125-31.
- [160] Hojo Y, Ikeda U, Katsuki Ta, Mizuno O, Fujikawa H, Shimada K. Matrix metalloproteinase expression in the coronary circulation induced by coronary angioplasty. *Atherosclerosis* 2002; 161: 185-92.
- [161] Mathapati S, Arumugam SB, Verma RS. High cholesterol diet increases MMP9 and CD40 immunopositivity in early atherosclerotic plaque in rabbits. *Acta Histochem* 2010; 112: 618-23.
- [162] Fukuda D, Shimada K, Tanaka A *et al.* Comparison of levels of serum matrix metalloproteinase-9 in patients with acute myocardial infarction versus unstable angina pectoris versus stable angina pectoris. *Am J Cardiol* 2006; 97: 175-80.
- [163] Alvarez B, Ruiz C, Chacón P, Alvarez-Sabin J, Matas M. Serum values of metalloproteinase-2 and metalloproteinase-9 as related to unstable plaque and inflammatory cells in patients with greater than 70% carotid artery stenosis. *J Vasc Surg* 2004; 40: 469-75.
- [164] Eldrup N, Grønholdt ML, Sillesen H, Nordestgaard BG. Elevated matrix metalloproteinase-9 associated with stroke or cardiovascular death in patients with carotid stenosis. *Circulation* 2006; 114: 1847-54.
- [165] Jefferis BJ, Whincup P, Welsh P *et al.* Prospective study of matrix metalloproteinase-9 and risk of myocardial infarction and stroke in older men and women. *Atherosclerosis* 2010; 208: 557-63.

- [166] Ferroni P, Basili S, Martini F *et al.* Serum metalloproteinase 9 levels in patients with coronary artery disease: a novel marker of inflammation. *J Investig Med* 2003; 51: 295-300.
- [167] Konstantino Y, Nguyen TT, Wolk R, Aiello RJ, Terra SG, Fryburg DA. Potential implications of matrix metalloproteinase-9 in assessment and treatment of coronary artery disease. *Biomarkers* 2009; 14: 118-29.
- [168] Li L, Renier G. The oral anti-diabetic agent, gliclazide, inhibits oxidized LDL-mediated LOX-1 expression, metalloproteinase-9 secretion and apoptosis in human aortic endothelial cells. *Atherosclerosis* 2009; 204: 40-6.
- [169] Hua Y, Xue J, Sun F, Zhu L, Xie M. Aspirin inhibits MMP-2 and MMP-9 expressions and activities through upregulation of PPARalpha/gamma and TIMP gene expressions in ox-LDL-stimulated macrophages derived from human monocytes. *Pharmacology* 2009; 83: 18-25.
- [170] Kang JH, Kim JK, Park WH *et al.* Ascochlorin suppresses oxLDL-induced MMP-9 expression by inhibiting the MEK/ERK signaling pathway in human THP-1 macrophages. *J Cell Biochem* 2007; 102: 506-14.
- [171] Kalela A, Koivu TA, Höyhty M *et al.* Association of serum MMP-9 with autoantibodies against oxidized LDL. *Atherosclerosis* 2002; 160: 161-5.
- [172] Garenc C, Julien P, Levy E. Oxysterols in biological systems: the gastrointestinal tract, liver, vascular wall and central nervous system. *Free Radic Res* 2010; 44: 47-73.
- [173] Leonarduzzi G, Poli G, Sottero B, Biasi F. Activation of the mitochondrial pathway of apoptosis by oxysterols. *Front Biosci* 2007; 12: 791-9.
- [174] Sottero B, Gamba P, Gargiulo S, Leonarduzzi G, Poli G. Cholesterol oxidation products and disease: an emerging topic of interest in medicinal chemistry. *Curr Med Chem* 2009; 16: 685-705.
- [175] Poli G, Sottero B, Gargiulo S, Leonarduzzi G. Cholesterol oxidation products in the vascular remodeling due to atherosclerosis. *Mol Aspects Med* 2009; 30: 180-9.
- [176] Leonarduzzi G, Gamba P, Sottero B *et al.* Oxysterol-induced up-regulation of MCP-1 expression and synthesis in macrophage cells. *Free Radic Biol Med* 2005; 39: 1152-61.
- [177] Leonarduzzi G, Vizio B, Sottero B *et al.* Early involvement of ROS overproduction in apoptosis induced by 7-ketocholesterol. *Antioxid Redox Signal* 2006; 8: 375-80.
- [178] Leonarduzzi G, Gamba P, Gargiulo S *et al.* Oxidation as a crucial reaction for cholesterol to induce tissue degeneration: CD36 overexpression in human promonocytic cells treated with a biologically relevant oxysterol mixture. *Aging Cell* 2008; 7: 375-82.
- [179] Leonarduzzi G, Gargiulo S, Gamba P *et al.* Molecular signaling operated by a diet-compatible mixture of oxysterols in up-regulating CD36 receptor in CD68 positive cells. *Mol Nutr Food Res* 2010; 54: S31-41.
- [180] Gargiulo S, Sottero B, Gamba P, Chiarotto E, Poli G, Leonarduzzi G. Plaque oxysterols induce unbalanced up-regulation of matrix metalloproteinase-9 in macrophagic cells through redox-sensitive signaling pathways: Implications regarding the vulnerability of atherosclerotic lesions. *Free Radic Biol Med* 2011; 51: 844-55.
- [181] Lim CS, Shalhoub J, Gohel MS, Shepherd AC, Davies AH. Matrix metalloproteinases in vascular disease--a potential therapeutic target? *Curr Vasc Pharmacol* 2010; 8: 75-85.
- [182] Upchurch GR Jr, Ford JW, Weiss SJ *et al.* Nitric oxide inhibition increases matrix metalloproteinase-9 expression by rat aortic smooth muscle cells in vitro. *J Vasc Surg* 2001; 34: 76-83.
- [183] Uría JA, Jiménez MG, Balbín M, Freije JM, López-Otín C. Differential effects of transforming growth factor-beta on the expression of collagenase-1 and collagenase-3 in human fibroblasts. *J Biol Chem* 1998; 273: 9769-77.

- [184] Grzela T, Brawura-Biskupski-Samaha R, Jelenska MM, Szmidt J. Low molecular weight heparin treatment decreases MMP-9 plasma activity in patients with abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg* 2008; 35: 159-61.
- [185] Schieffer B, Bunte C, Witte J *et al.* Comparative effects of AT1-antagonism and angiotensin-converting enzyme inhibition on markers of inflammation and platelet aggregation in patients with coronary artery disease. *J Am Coll Cardiol* 2004; 44: 362-8.
- [186] Yamamoto D, Takai S, Miyazaki M. Inhibitory profiles of captopril on matrix metalloproteinase-9 activity. *Eur J Pharmacol* 2008; 588: 277-9.
- [187] Galis ZS, Asanuma K, Godin D, Meng X. N-acetyl-cysteine decreases the matrix-degrading capacity of macrophage-derived foam cells: new target for antioxidant therapy? *Circulation* 1998; 97: 2445-53.
- [188] Lutgens E, Gorelik L, Daemen MJ *et al.* Requirement for CD154 in the progression of atherosclerosis. *Nat Med* 1999; 5: 1313-6.
- [189] Miralles M, Wester W, Sicard GA, Thompson R, Reilly JM. Indomethacin inhibits expansion of experimental aortic aneurysms via inhibition of the cox2 isoform of cyclooxygenase. *J Vasc Surg* 1999; 29: 884-92.
- [190] Walton LJ, Franklin IJ, Bayston T *et al.* Inhibition of prostaglandin E2 synthesis in abdominal aortic aneurysms: implications for smooth muscle cell viability, inflammatory processes, and the expansion of abdominal aortic aneurysms. *Circulation* 1999; 100: 48-54.
- [191] Cipollone F, Fazio ML, Iezzi A *et al.* Association between prostaglandin E receptor subtype EP4 overexpression and unstable phenotype in atherosclerotic plaques in human. *Arterioscler Thromb Vasc Biol* 2005; 25: 1925-31.
- [192] Pavlovic S, Du B, Sakamoto K *et al.* Targeting prostaglandin E2 receptors as an alternative strategy to block cyclooxygenase-2-dependent extracellular matrix-induced matrix metalloproteinase-9 expression by macrophages. *J Biol Chem* 2006; 281: 3321-8.
- [193] Zucker S, Cao J, Chen WT. Critical appraisal of the use of matrix metalloproteinase inhibitors in cancer treatment. *Oncogene* 2000; 19: 6642-50.
- [194] Rouis M, Adamy C, Duverger N *et al.* Adenovirus-mediated overexpression of tissue inhibitor of metalloproteinase-1 reduces atherosclerotic lesions in apolipoprotein E-deficient mice. *Circulation* 1999; 100: 533-40.
- [195] George SJ, Lloyd CT, Angelini GD, Newby AC, Baker AH. Inhibition of late vein graft neointima formation in human and porcine models by adenovirus-mediated overexpression of tissue inhibitor of metalloproteinase-3. *Circulation* 2000; 101: 296-304.
- [196] George SJ, Wan S, Hu J, MacDonald R, Johnson JL, Baker AH. Sustained reduction of vein graft neointima formation by ex vivo TIMP-3 gene therapy. *Circulation* 2011; 124: S135-42.
- [197] Forrester JS, Libby P. The inflammation hypothesis and its potential relevance to statin therapy. *Am J Cardiol* 2007; 99: 732-8.
- [198] Bellosta S, Via D, Canavesi M *et al.* HMG-CoA reductase inhibitors reduce MMP-9 secretion by macrophages. *Arterioscler Thromb Vasc Biol* 1998; 18: 1671-8.
- [199] Aikawa M, Rabkin E, Okada Y *et al.* Lipid lowering by diet reduces matrix metalloproteinase activity and increases collagen content of rabbit atheroma: a potential mechanism of lesion stabilization. *Circulation* 1998; 97: 2433-44.
- [200] Crisby M, Nordin-Fredriksson G, Shah PK, Yano J, Zhu J, Nilsson J. Pravastatin treatment increases collagen content and decreases lipid content, inflammation, metalloproteinases, and cell death in human carotid plaques: implications for plaque stabilization. *Circulation* 2001; 103: 926-33.

- [201] Luan Z, Chase AJ, Newby AC. Statins inhibit secretion of metalloproteinases-1, -2, -3, and -9 from vascular smooth muscle cells and macrophages. *Arterioscler Thromb Vasc Biol* 2003; 23: 769-75.
- [202] Fukumoto Y, Libby P, Rabkin E *et al.* Statins alter smooth muscle cell accumulation and collagen content in established atheroma of watanabe heritable hyperlipidemic rabbits. *Circulation* 2001; 103: 993-9.
- [203] Scalia R, Gooszen ME, Jones SP *et al.* Simvastatin exerts both anti-inflammatory and cardioprotective effects in apolipoprotein E-deficient mice. *Circulation* 2001; 103: 2598-603.
- [204] Jialal I, Stein D, Balis D, Grundy SM, Adams-Huet B, Devaraj S. Effect of hydroxymethyl glutaryl coenzyme a reductase inhibitor therapy on high sensitive C-reactive protein levels. *Circulation* 2001; 103: 1933-5.
- [205] Ridker PM, Rifai N, Lowenthal SP. Rapid reduction in C-reactive protein with cerivastatin among 785 patients with primary hypercholesterolemia. *Circulation* 2001; 103: 1191-3.
- [206] Devy L, Dransfield DT. New strategies for the next generation of matrix-metalloproteinase inhibitors: selectively targeting membrane-anchored MMPs with therapeutic antibodies. *Biochem Res Int* 2011; 2011: 191670.
- [207] Hu J, Van den Steen PE, Sang QX, Opdenakker G. Matrix metalloproteinase inhibitors as therapy for inflammatory and vascular diseases. *Nat Rev Drug Discov* 2007; 6: 480-98.
- [208] Rosenberg GA, Estrada EY, Mobashery S. Effect of synthetic matrix metalloproteinase inhibitors on lipopolysaccharide-induced blood-brain barrier opening in rodents: Differences in response based on strains and solvents. *Brain Res* 2007; 1133: 186-92.
- [209] Denis LJ, Verweij J. Matrix metalloproteinase inhibitors: present achievements and future prospects. *Invest New Drugs* 1997; 15: 175-85.
- [210] Parsons SL, Watson SA, Brown PD, Collins HM, Steele RJ. Matrix metalloproteinases. *Br J Surg* 1997; 84: 160-6.
- [211] Sood RR, Taheri S, Candelario-Jalil E, Estrada EY, Rosenberg GA. Early beneficial effect of matrix metalloproteinase inhibition on blood-brain barrier permeability as measured by magnetic resonance imaging countered by impaired long-term recovery after stroke in rat brain. *J Cereb Blood Flow Metab* 2008; 28: 431-8.
- [212] Nagel S, Heinemann PV, Heiland S, Koziol J, Gardner H, Wagner S. Selective MMP-inhibition with Ro 28-2653 in acute experimental stroke--a magnetic resonance imaging efficacy study. *Brain Res* 2011; 1368: 264-70.
- [213] Maquoi E, Sounni NE, Devy L *et al.* Anti-invasive, antitumoral, and antiangiogenic efficacy of a pyrimidine-2,4,6-trione derivative, an orally active and selective matrix metalloproteinases inhibitor. *Clin Cancer Res* 2004; 10: 4038-47.
- [214] Shono T, Motoyama M, Tatsumi K *et al.* A new synthetic matrix metalloproteinase inhibitor modulates both angiogenesis and urokinase type plasminogen activator activity. *Angiogenesis* 1998; 2: 319-29.
- [215] Wojtowicz-Praga S. Clinical potential of matrix metalloprotease inhibitors. *Drugs R D* 1999; 1: 117-29.
- [216] Quillard T, Tesmenitsky Y, Croce K *et al.* Selective inhibition of matrix metalloproteinase-13 increases collagen content of established mouse atherosclerosis. *Arterioscler Thromb Vasc Biol* 2011; 31: 2464-72.
- [217] Franco C, Ho B, Mulholland D *et al.* Doxycycline alters vascular smooth muscle cell adhesion, migration, and reorganization of fibrillar collagen matrices. *Am J Pathol* 2006; 168: 1697-709.
- [218] Sapadin AN, Fleischmajer R. Tetracyclines: nonantibiotic properties and their clinical implications. *J Am Acad Dermatol* 2006; 54: 258-65.
- [219] Axisa B, Loftus IM, Naylor AR *et al.* Prospective, randomized, double-blind trial investigating the effect of doxycycline on matrix metalloproteinase expression within atherosclerotic carotid plaques. *Stroke* 2002; 33: 2858-64.

- [220] Brown DL, Desai KK, Vakili BA, Nouneh C, Lee HM, Golub LM. Clinical and biochemical results of the metalloproteinase inhibition with subantimicrobial doses of doxycycline to prevent acute coronary syndromes (MIDAS) pilot trial. *Arterioscler Thromb Vasc Biol* 2004; 24: 733-8.
- [221] Dodd BR, Spence RA. Doxycycline inhibition of abdominal aortic aneurysm growth: a systematic review of the literature. *Curr Vasc Pharmacol* 2011; 9: 471-8.
- [222] Golledge J, Norman PE. Current status of medical management for abdominal aortic aneurysm. 2011; 217: 57-63.
- [223] Bendeck MP, Conte M, Zhang M, Nili N, Strauss BH, Farwell SM. Doxycycline modulates smooth muscle cell growth, migration, and matrix remodeling after arterial injury. *Am J Pathol* 2002; 160: 1089-95.
- [224] Cowan KN, Jones PL, Rabinovitch M. Elastase and matrix metalloproteinase inhibitors induce regression, and tenascin-C antisense prevents progression, of vascular disease. *J Clin Invest* 2000; 105: 21-34.
- [225] Eriksson EE. Intravital microscopy on atherosclerosis in apolipoprotein e-deficient mice establishes microvessels as major entry pathways for leukocytes to advanced lesions. *Circulation* 2011; 124: 2129-38.
- [226] Forbes JM, Cooper ME. Mechanisms of diabetic complications. *Physiol Rev* 2013; 93: 137-88.
- [227] Chung AW, Hsiang YN, Matzke LA, McManus BM, van Breemen C, Okon EB. Reduced expression of vascular endothelial growth factor paralleled with the increased angiostatin expression resulting from the upregulated activities of matrix metalloproteinase-2 and -9 in human type 2 diabetic arterial vasculature. *Circ Res* 2006; 99: 140-8.
- [228] Hao F, Yu JD. High glucose enhance expression of matrix metalloproteinase-2 in smooth muscle cells. *Acta Pharmacol Sin* 2003; 24: 534-8.
- [229] Tarallo S, Beltramo E, Berrone E, Dentelli P, Porta M. Effects of high glucose and thiamine on the balance between matrix metalloproteinases and their tissue inhibitors in vascular cells. *Acta Diabetol* 2010; 47: 105-11.
- [230] Sun J, Xu Y, Dai Z, Sun Y. Intermittent high glucose enhances proliferation of vascular smooth muscle cells by upregulating osteopontin. *Mol Cell Endocrinol* 2009; 313: 64-9.
- [231] Uemura S, Matsushita H, Li W *et al.* Diabetes mellitus enhances vascular matrix metalloproteinase activity: role of oxidative stress. *Circ Res* 2001; 88: 1291-8.
- [232] Worley JR, Hughes DA, Dozio N, Gavrilovic J, Sampson MJ. Low density lipoprotein from patients with Type 2 diabetes increases expression of monocyte matrix metalloproteinase and ADAM metalloproteinase genes. *Cardiovasc Diabetol* 2007; 6: 21.
- [233] Wilmer WA, Cosio FG. DNA binding of activator protein-1 is increased in human mesangial cells cultured in high glucose concentrations. *Kidney Int* 1998; 53: 1172-81.
- [234] Pascal MM, Knott RM, Forrester JV. Glucose mediated regulation of transforming growth factor beta in human retinal endothelial cells. *Biochem Soc Trans* 1996; 24: 228S.
- [235] Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature* 2001; 414: 813-20.
- [236] Jormsjö S, Ye S, Moritz J *et al.* Allele-specific regulation of matrix metalloproteinase-12 gene activity is associated with coronary artery luminal dimensions in diabetic patients with manifest coronary artery disease. *Circ Res* 2000; 86: 998-1003.
- [237] Yan SF, Ramasamy R, Naka Y, Schmidt AM. Glycation, inflammation, and RAGE: a scaffold for the macrovascular complications of diabetes and beyond. *Circ Res* 2003; 93: 1159-69.
- [238] Cipollone F, Iezzi A, Fazia M *et al.* The receptor RAGE as a progression factor amplifying arachidonate-dependent inflammatory and proteolytic response in human atherosclerotic plaques: role of glycemic control. *Circulation* 2003; 108: 1070-7.

- [239] Jacob SS, Shastry P, Sudhakaran PR. Influence of non-enzymatically glycated collagen on monocyte-macrophage differentiation. *Atherosclerosis* 2001; 159: 333-41.
- [240] Federici M, Menghini R, Mauriello A *et al.* Insulin-dependent activation of endothelial nitric oxide synthase is impaired by O-linked glycosylation modification of signaling proteins in human coronary endothelial cells. *Circulation* 2002; 106: 466-72.
- [241] Dandona P, Aljada A, Mohanty P, Ghanim H, Bandyopadhyay A, Chaudhuri A. Insulin suppresses plasma concentration of vascular endothelial growth factor and matrix metalloproteinase-9. *Diabetes Care* 2003; 26: 3310-4.
- [242] Sundström J, Evans JC, Benjamin EJ *et al.* Relations of plasma matrix metalloproteinase-9 to clinical cardiovascular risk factors and echocardiographic left ventricular measures: the Framingham Heart Study. *Circulation* 2004; 109: 2850-6.
- [243] Marx N, Froehlich J, Siam L *et al.* Antidiabetic PPAR gamma-activator rosiglitazone reduces MMP-9 serum levels in type 2 diabetic patients with coronary artery disease. *Arterioscler Thromb Vasc Biol* 2003; 23: 283-8.
- [244] Gonçalves FM, Jacob-Ferreira AL, Gomes VA *et al.* Increased circulating levels of matrix metalloproteinase (MMP)-8, MMP-9, and pro-inflammatory markers in patients with metabolic syndrome. *Clin Chim Acta* 2009; 403: 173-7.
- [245] Signorelli SS, Malaponte G, Libra M. Plasma levels and zymographic activities of matrix metalloproteinases 2 and 9 in type II diabetics with peripheral arterial disease. *Vasc Med* 2005; 10: 1-6.
- [246] Derosa G, D'Angelo A, Scalise F. Comparison between metalloproteinases-2 and -9 in healthy subjects, diabetics, and subjects with acute coronary syndrome. *Heart Vessels* 2007; 22: 361-70.
- [247] Zayani Y, Allal-Elasmi M, Jacob MP *et al.* Abnormal circulating levels of matrix metalloproteinases and their inhibitors in diabetes mellitus. *Clin Lab* 2012; 58: 779-85.
- [248] Baugh MD, Gavrilovic J, Davies IR, Hughes DA, Sampson MJ. Monocyte matrix metalloproteinase production in Type 2 diabetes and controls--a cross sectional study. *Cardiovasc Diabetol* 2003; 2: 3.
- [249] Portik-Dobos V, Anstadt MP, Hutchinson J, Bannan M, Ergul A. Evidence for a matrix metalloproteinase induction/activation system in arterial vasculature and decreased synthesis and activity in diabetes. *Diabetes* 2002; 51: 3063-8.
- [250] Das A, McGuire PG, Eriqat C *et al.* Human diabetic neovascular membranes contain high levels of urokinase and metalloproteinase enzymes. *Invest Ophthalmol Vis Sci* 1999; 40: 809-13.
- [251] Salzmann J, Limb GA, Khaw PT *et al.* Matrix metalloproteinases and their natural inhibitors in fibrovascular membranes of proliferative diabetic retinopathy. *Br J Ophthalmol* 2000; 84: 1091-6.
- [252] Lauhio A, Sorsa T, Srinivas R *et al.* Urinary matrix metalloproteinase -8, -9, -14 and their regulators (TRY-1, TRY-2, TATI) in patients with diabetic nephropathy. *Ann Med* 2008; 40: 312-20.
- [253] Thrailkill KM, Clay Bunn R, Fowlkes JL. Matrix metalloproteinases: their potential role in the pathogenesis of diabetic nephropathy. *Endocrine* 2009; 35: 1-10.
- [254] Phillips AO, Steadman R, Morrissey K, Martin J, Eynstone L, Williams JD. Exposure of human renal proximal tubular cells to glucose leads to accumulation of type IV collagen and fibronectin by decreased degradation. *Kidney Int* 1997; 52: 973-84.
- [255] McLennan SV, Martell SK, Yue DK. Effects of mesangium glycation on matrix metalloproteinase activities: possible role in diabetic nephropathy. *Diabetes* 2002; 51: 2612-8.
- [256] Zhuang S, Kinsey GR, Rasbach K, Schnellmann RG. Heparin-binding epidermal growth factor and Src family kinases in proliferation of renal epithelial cells. *Am J Physiol Renal Physiol* 2008; 294: F459-68.

- [257] Haffner SM, Greenberg AS, Weston WM, Chen H, Williams K, Freed MI. Effect of rosiglitazone treatment on nontraditional markers of cardiovascular disease in patients with type 2 diabetes mellitus. *Circulation* 2002; 106: 679-84.
- [258] Goldstein BJ, Weissman PN, Wooddell MJ, Waterhouse BR, Cobitz AR. Reductions in biomarkers of cardiovascular risk in type 2 diabetes with rosiglitazone added to metformin compared with dose escalation of metformin: an EMPIRE trial sub-study. *Curr Med Res Opin* 2006; 22: 1715-23.
- [259] Lee CS, Kwon YW, Yang HM *et al.* New mechanism of rosiglitazone to reduce neointimal hyperplasia: activation of glycogen synthase kinase-3beta followed by inhibition of MMP-9. *Arterioscler Thromb Vasc Biol* 2009; 29: 472-9.
- [260] Forst T, Karagiannis E, Lübben G *et al.* Pleiotropic and anti-inflammatory effects of pioglitazone precede the metabolic activity in type 2 diabetic patients with coronary artery disease. *Atherosclerosis* 2008; 197: 311-7.
- [261] Game BA, Xu M, Lopes-Virella MF, Huang Y. Regulation of MMP-1 expression in vascular endothelial cells by insulin sensitizing thiazolidinediones. *Atherosclerosis* 2003; 169: 235-43.
- [262] Ruilope LM, Segura J. Losartan and other angiotensin II antagonists for nephropathy in type 2 diabetes mellitus: a review of the clinical trial evidence. *Clin Ther* 2003; 25: 3044-64.
- [263] Papadakis JA, Millionis HJ, Press M, Mikhailidis DP. Treating dyslipidaemia in non-insulin-dependent diabetes mellitus -- a special reference to statins. *J Diabetes Complications* 2001; 15: 211-26.
- [264] Athyros VG, Papageorgiou AA, Symeonidis AN. Early benefit from structured care with atorvastatin in patients with coronary heart disease and diabetes mellitus. *Angiology* 2003; 54: 679-90.
- [265] Collins R, Armitage J, Parish S, Sleight P, Peto R. MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Lancet* 2003; 361: 2005-16.
- [266] Bellosta S, Ferri N, Arnaboldi L, Bernini F, Paoletti R, Corsini A. Pleiotropic effects of statins in atherosclerosis and diabetes. *Diabetes Care* 2000; 23: B72-8.
- [267] Luan Z, Chase AJ, Newby AC. Statins inhibit secretion of metalloproteinases-1, -2, -3, and -9 from vascular smooth muscle cells and macrophages. *Arterioscler Thromb Vasc Biol* 2003; 23: 769-75.
- [268] Song Y, Li C, Cai L. Fluvastatin prevents nephropathy likely through suppression of connective tissue growth factor-mediated extracellular matrix accumulation. *Exp Mol Pathol* 2004; 76: 66-75.
- [269] Furman C, Copin C, Kandoussi M. Rosuvastatin reduces MMP-7 secretion by human monocyte-derived macrophages: potential relevance to atherosclerotic plaque stability. *Atherosclerosis* 2004; 174: 93-8.
- [270] Balk EM, Lau J, Goudas LC. Effects of statins on nonlipid serum markers associated with cardiovascular disease: a systematic review. *Ann Intern Med* 2003; 139: 670-82.
- [271] Sager PT, Melani L, Lipka L. Effect of coadministration of ezetimibe and simvastatin on high-sensitivity C-reactive protein. *Am J Cardiol* 2003; 92: 1414-8.
- [272] Ridker PM, Rifai N, Clearfield M. Measurement of C-reactive protein for the targeting of statin therapy in the primary prevention of acute coronary events. *N Engl J Med* 2001; 344: 1959-65.
- [273] Elisaf M, Mikhailidis DP. Statins and renal function. *Angiology* 2002; 53: 493-502.
- [274] Daskalopoulou SS, Athyros VG, Elisaf M, Mikhailidis DP. Uric acid levels and vascular disease. *Curr Med Res Opin* 2004; 20: 951-4.
- [275] Athyros VG, Elisaf M, Papageorgiou AA. Effect of statins versus untreated dyslipidemia on serum uric acid levels in patients with coronary heart disease: a subgroup analysis of the GREek Atorvastatin and Coronary-heart-disease Evaluation (GREACE) study. *Am J Kidney Dis* 2004; 43: 589-99.

- [276] Francis GA, Annicotte JS, Auwerx J. PPAR agonists in the treatment of atherosclerosis. *Curr Opin Pharmacol* 2003; 3: 186-91.
- [277] Marx N, Schönbeck U, Lazar MA, Libby P, Plutzky J. Peroxisome proliferator-activated receptor gamma activators inhibit gene expression and migration in human vascular smooth muscle cells. *Circ Res* 1998; 83: 1097-103.
- [278] Shu H, Wong B, Zhou G. Activation of PPARalpha or gamma reduces secretion of matrix metalloproteinase 9 but not interleukin 8 from human monocytic THP-1 cells. *Biochem Biophys Res Commun* 2000; 267: 345-9.
- [279] Rizos E, Bairaktari E, Ganotakis E, Tsimihodimos V, Mikhailidis DP, Elisaf M. Effect of ciprofibrate on lipoproteins, fibrinogen, renal function, and hepatic enzymes. *J Cardiovasc Pharmacol Ther* 2002; 7: 219-26.
- [280] Tsimihodimos V, Kostoula A, Kakafika A. Effect of fenofibrate on serum inflammatory markers in patients with high triglyceride values. *J Cardiovasc Pharmacol Ther* 2004; 9: 27-33.
- [281] Wierzbicki AS, Mikhailidis DP, Wray R. Statin-fibrate combination: therapy for hyperlipidemia: a review. *Curr Med Res Opin* 2003; 19: 155-68.
- [282] Kim YS, Joh TH. Microglia, major player in the brain inflammation: their roles in the pathogenesis of Parkinson's disease. *Exp Mol Med* 2006; 38: 333-47.
- [283] Block ML, Zecca L, Hong JS. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nat Rev Neurosci* 2007; 8: 57-69.
- [284] Candelario-Jalil E, Yang Y, Rosenberg GA. Diverse roles of matrix metalloproteinases and tissue inhibitors of metalloproteinases in neuroinflammation and cerebral ischemia. *Neuroscience* 2009; 158: 983-94.
- [285] Rosenberg GA. Matrix metalloproteinases and their multiple roles in neurodegenerative diseases. *Lancet Neurol* 2009; 8: 205-16.
- [286] Yang Y, Hill JW, Rosenberg GA. Multiple roles of metalloproteinases in neurological disorders. *Prog Mol Biol Transl Sci* 2011; 99: 241-63.
- [287] Yang Y, Estrada EY, Thompson JF, Liu W, Rosenberg GA. Matrix metalloproteinase-mediated disruption of tight junction proteins in cerebral vessels is reversed by synthetic matrix metalloproteinase inhibitor in focal ischemia in rat. *J Cereb Blood Flow Metab* 2007; 27: 697-709.
- [288] Lo EH, Dalkara T, Moskowitz MA. Mechanisms, challenges and opportunities in stroke. *Nat Rev Neurosci* 2003; 4: 399-415.
- [289] Asahi M, Wang X, Mori T *et al.* Effects of matrix metalloproteinase-9 gene knock-out on the proteolysis of blood-brain barrier and white matter components after cerebral ischemia. *J Neurosci* 2001; 21: 7724-32.
- [290] Gasche Y, Copin JC, Sugawara T, Fujimura M, Chan PH. Matrix metalloproteinase inhibition prevents oxidative stress-associated blood-brain barrier disruption after transient focal cerebral ischemia. *J Cereb Blood Flow Metab* 2001; 21: 1393-400.
- [291] Chang DI, Hosomi N, Lucero J *et al.* Activation systems for latent matrix metalloproteinase-2 are upregulated immediately after focal cerebral ischemia. *J Cereb Blood Flow Metab* 2003; 23: 1408-19.
- [292] Liu W, Hendren J, Qin XJ, Shen J, Liu KJ. Normobaric hyperoxia attenuates early blood-brain barrier disruption by inhibiting MMP-9-mediated occludin degradation in focal cerebral ischemia. *J Neurochem* 2009; 108: 811-20.
- [293] Gurney KJ, Estrada EY, Rosenberg GA. Blood-brain barrier disruption by stromelysin-1 facilitates neutrophil infiltration in neuroinflammation. *Neurobiol Dis* 2006; 23: 87-96.
- [294] Asahi M, Asahi K, Jung JC, del Zoppo GJ, Fini ME, Lo EH. Role for matrix metalloproteinase 9 after focal cerebral ischemia: effects of gene knockout and enzyme inhibition with BB-94. *J Cereb Blood Flow Metab* 2000; 20: 1681-9.

- [295] Rosenberg GA, Estrada EY, Dencoff JE, Stetler-Stevenson WG. Tumor necrosis factor-alpha-induced gelatinase B causes delayed opening of the blood-brain barrier: an expanded therapeutic window. *Brain Res* 1995; 703: 151-5.
- [296] Candelario-Jalil E, Taheri S, Yang Y *et al.* Cyclooxygenase inhibition limits blood-brain barrier disruption following intracerebral injection of tumor necrosis factor-alpha in the rat. *J Pharmacol Exp Ther* 2007; 323: 488-98.
- [297] Kauppinen TM, Swanson RA. Poly(ADP-ribose) polymerase-1 promotes microglial activation, proliferation, and matrix metalloproteinase-9-mediated neuron death. *J Immunol* 2005; 174: 2288-96.
- [298] del Zoppo GJ, Milner R, Mabuchi T *et al.* Microglial activation and matrix protease generation during focal cerebral ischemia. *Stroke* 2007; 38: 646-51.
- [299] Wang L, Zhang ZG, Zhang RL *et al.* Matrix metalloproteinase 2 (MMP2) and MMP9 secreted by erythropoietin-activated endothelial cells promote neural progenitor cell migration. *J Neurosci* 2006; 26: 5996-6003.
- [300] Lee SR, Kim HY, Rogowska J *et al.* Involvement of matrix metalloproteinase in neuroblast cell migration from the subventricular zone after stroke. *J Neurosci* 2006; 26: 3491-5.
- [301] Harkness KA, Adamson P, Sussman JD, Davies-Jones GA, Greenwood J, Woodroffe MN. Dexamethasone regulation of matrix metalloproteinase expression in CNS vascular endothelium. *Brain* 2000; 123: 698-709.
- [302] Gottschall PE, Yu X. Cytokines regulate gelatinase A and B (matrix metalloproteinase 2 and 9) activity in cultured rat astrocytes. *J Neurochem* 1995; 64: 1513-20.
- [303] Lee WJ, Shin CY, Yoo BK *et al.* Induction of matrix metalloproteinase-9 (MMP-9) in lipopolysaccharide-stimulated primary astrocytes is mediated by extracellular signal-regulated protein kinase 1/2 (Erk1/2). *Glia* 2003; 41: 15-24.
- [304] Hsieh HL, Yen MH, Jou MJ, Yang CM. Intracellular signalings underlying bradykinin-induced matrix metalloproteinase-9 expression in rat brain astrocyte-1. *Cell Signal* 2004; 16: 1163-76.
- [305] Wu CY, Hsieh HL, Jou MJ, Yang CM. Involvement of p42/p44 MAPK, p38 MAPK, JNK and nuclear factor-kappa B in interleukin-1beta-induced matrix metalloproteinase-9 expression in rat brain astrocytes. *J Neurochem* 2004; 90: 1477-88.
- [306] Woo CH, Lim JH, Kim JH. Lipopolysaccharide induces matrix metalloproteinase-9 expression via a mitochondrial reactive oxygen species-p38 kinase-activator protein-1 pathway in Raw 264.7 cells. *J Immunol* 2004; 173: 6973-80.
- [307] Rangaswami H, Bulbule A, Kundu GC. JNK1 differentially regulates osteopontin-induced nuclear factor-inducing kinase/MEKK1-dependent activating protein-1-mediated promatrix metalloproteinase-9 activation. *J Biol Chem* 2005; 280: 19381-92.
- [308] Han S, Ritzenthaler JD, Sitaraman SV, Roman J. Fibronectin increases matrix metalloproteinase 9 expression through activation of c-Fos via extracellular-regulated kinase and phosphatidylinositol 3-kinase pathways in human lung carcinoma cells. *J Biol Chem* 2006; 281: 29614-24.
- [309] Wang HH, Hsieh HL, Wu CY, Sun CC, Yang CM. Oxidized low-density lipoprotein induces matrix metalloproteinase-9 expression via a p42/p44 and JNK-dependent AP-1 pathway in brain astrocytes. *Glia* 2009; 57: 24-38.
- [310] Keller JN, Hanni KB, Markesbery WR. Oxidized low-density lipoprotein induces neuronal death: implications for calcium, reactive oxygen species, and caspases. *J Neurochem* 1999; 72: 2601-9.
- [311] Kim JA, Tran ND, Berliner JA, Fisher MJ. Minimally oxidized low-density lipoprotein regulates hemostasis factors of brain capillary endothelial cells. *J Neurol Sci* 2004; 217: 135-41.

- [312] Lupo G, Nicotra A, Giurdanella G *et al.* Activation of phospholipase A(2) and MAP kinases by oxidized low-density lipoproteins in immortalized GP8.39 endothelial cells. *Biochim Biophys Acta* 2005; 1735: 135-50.
- [313] Vos CM, Sjulson L, Nath A *et al.* Cytotoxicity by matrix metalloproteinase-1 in organotypic spinal cord and dissociated neuronal cultures. *Exp Neurol* 2000; 163: 324-30.
- [314] Deb S, Gottschall PE. Increased production of matrix metalloproteinases in enriched astrocyte and mixed hippocampal cultures treated with beta-amyloid peptides. *J Neurochem* 1996; 66: 1641-7.
- [315] Rosenberg GA, Cunningham LA, Wallace J *et al.* Immunohistochemistry of matrix metalloproteinases in reperfusion injury to rat brain: activation of MMP-9 linked to stromelysin-1 and microglia in cell cultures. *Brain Res* 2001; 893: 104-12.
- [316] Kim KS, Kim HY, Joe EH, Jou I. Matrix metalloproteinase-3 induction in rat brain astrocytes: focus on the role of two AP-1 elements. *Biochem J* 2008; 410: 605-11.
- [317] Crocker SJ, Milner R, Pham-Mitchell N, Campbell IL. Cell and agonist-specific regulation of genes for matrix metalloproteinases and their tissue inhibitors by primary glial cells. *J Neurochem* 2006; 98: 812-23.
- [318] Wells GM, Catlin G, Cossins JA *et al.* Quantitation of matrix metalloproteinases in cultured rat astrocytes using the polymerase chain reaction with a multi-competitor cDNA standard. *Glia* 1996; 18: 332-40.
- [319] Anthony DC, Ferguson B, Matyzak MK, Miller KM, Esiri MM, Perry VH. Differential matrix metalloproteinase expression in cases of multiple sclerosis and stroke. *Neuropathol Appl Neurobiol* 1997; 23: 406-15.
- [320] Rosenberg GA, Sullivan N, Esiri MM. White matter damage is associated with matrix metalloproteinases in vascular dementia. *Stroke* 2001; 32: 1162-8.
- [321] Kim YS, Kim SS, Cho JJ *et al.* Matrix metalloproteinase-3: a novel signaling proteinase from apoptotic neuronal cells that activates microglia. *J Neurosci* 2005; 25: 3701-11.
- [322] Kim YS, Choi DH, Block ML *et al.* A pivotal role of matrix metalloproteinase-3 activity in dopaminergic neuronal degeneration via microglial activation. *FASEB J* 2007; 21: 179-87.
- [323] Woo MS, Park JS, Choi IY, Kim WK, Kim HS. Inhibition of MMP-3 or -9 suppresses lipopolysaccharide-induced expression of proinflammatory cytokines and iNOS in microglia. *J Neurochem* 2008; 106: 770-80.
- [324] Kim EM, Hwang O. Role of matrix metalloproteinase-3 in neurodegeneration. *J Neurochem* 2011; 116: 22-32.
- [325] Kim EM, Shin EJ, Choi JH *et al.* Matrix metalloproteinase-3 is increased and participates in neuronal apoptotic signaling downstream of caspase-12 during endoplasmic reticulum stress. *J Biol Chem* 2010; 285: 16444-52.
- [326] Tan HK, Heywood D, Ralph GS, Bienemann A, Baker AH, Uney JB. Tissue inhibitor of metalloproteinase 1 inhibits excitotoxic cell death in neurons. *Mol Cell Neurosci* 2003; 22: 98-106.
- [327] Tejima E, Guo S, Murata Y *et al.* Neuroprotective effects of overexpressing tissue inhibitor of metalloproteinase TIMP-1. *J Neurotrauma* 2009; 26: 1935-41.
- [328] Rosenberg GA. Inflammation and white matter damage in vascular cognitive impairment. *Stroke* 2009; 40: S20-3.
- [329] Semenza GL. Vasculogenesis, angiogenesis, and arteriogenesis: mechanisms of blood vessel formation and remodeling. *J Cell Biochem* 2007; 102: 840-7.
- [330] Nakaji K, Ihara M, Takahashi C *et al.* Matrix metalloproteinase-2 plays a critical role in the pathogenesis of white matter lesions after chronic cerebral hypoperfusion in rodents. *Stroke* 2006; 37: 2816-23.
- [331] Chandler S, Coates R, Gearing A, Lury J, Wells G, Bone E. Matrix metalloproteinases degrade myelin basic protein. *Neurosci Lett* 1995; 201: 223-6.
- [332] Forsythe JA, Jiang BH, Iyer NV *et al.* Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol Cell Biol* 1996; 16: 4604-4613.

- [333] Huang RQ, Cheng HL, Zhao XD *et al.* Preliminary study on the effect of trauma-induced secondary cellular hypoxia in brain injury. *Neurosci Lett* 2010; 473: 22-7.
- [334] Ihara M, Tomimoto H, Kinoshita M *et al.* Chronic cerebral hypoperfusion induces MMP-2 but not MMP-9 expression in the microglia and vascular endothelium of white matter. *J Cereb Blood Flow Metab* 2001; 21: 828-34.
- [335] Adair JC, Charlie J, Dencoff JE *et al.* Measurement of gelatinase B (MMP-9) in the cerebrospinal fluid of patients with vascular dementia and Alzheimer disease. *Stroke* 2004; 35: e159-62.
- [336] Fernandez-Patron C, Radomski MW, Davidge ST. Vascular matrix metalloproteinase-2 cleaves big endothelin-1 yielding a novel vasoconstrictor. *Circ Res* 1999; 85: 906-11.
- [337] He S, Prasanna G, Yorio T. Endothelin-1-mediated signaling in the expression of matrix metalloproteinases and tissue inhibitors of metalloproteinases in astrocytes. *Invest Ophthalmol Vis Sci* 2007; 48: 3737-45.
- [338] Zhang WW, Badonic T, Höög A *et al.* Structural and vasoactive factors influencing intracerebral arterioles in cases of vascular dementia and other cerebrovascular disease: a review. *Immunohistochemical studies on expression of collagens, basal lamina components and endothelin-1. Dementia* 1994; 5: 153-62.
- [339] Fritsch T, Smyth KA, Wallendal MS, Hyde T, Leo G, Geldmacher DS. Parkinson disease: research update and clinical management. *South Med J* 2012; 105: 650-6.
- [340] Sung JY, Park SM, Lee CH *et al.* Proteolytic cleavage of extracellular secreted {alpha}-synuclein via matrix metalloproteinases. *J Biol Chem* 2005; 280: 25216-24.
- [341] McClain JA, Phillips LL, Fillmore HL. Increased MMP-3 and CTGF expression during lipopolysaccharide-induced dopaminergic neurodegeneration. *Neurosci Lett* 2009; 460: 27-31.
- [342] Choi DH, Kim EM, Son HJ *et al.* A novel intracellular role of matrix metalloproteinase-3 during apoptosis of dopaminergic cells. *J Neurochem* 2008; 106: 405-15.
- [343] Levin J, Giese A, Boetzel K *et al.* Increased alpha-synuclein aggregation following limited cleavage by certain matrix metalloproteinases. *Exp Neurol* 2009; 215: 201-8.
- [344] Kortekaas R, Leenders KL, van Oostrom JC *et al.* Blood-brain barrier dysfunction in parkinsonian midbrain in vivo. *Ann Neurol* 2005; 57: 176-9.
- [345] Carvey PM, Zhao CH, Hendey B *et al.* 6-Hydroxydopamine-induced alterations in blood-brain barrier permeability. *Eur J Neurosci* 2005; 22: 1158-68.
- [346] Zhao C, Ling Z, Newman MB, Bhatia A, Carvey PM. TNF-alpha knockout and minocycline treatment attenuates blood-brain barrier leakage in MPTP-treated mice. *Neurobiol Dis* 2007; 26: 36-46.
- [347] Ji KA, Eu MY, Kang SH, Gwag BJ, Jou I, Joe EH. Differential neutrophil infiltration contributes to regional differences in brain inflammation in the substantia nigra pars compacta and cortex. *Glia* 2008; 56: 1039-47.
- [348] Selkoe DJ. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* 2001; 81: 741-66.
- [349] Honjo K, Black SE, Verhoeff NP. Alzheimer's disease, cerebrovascular disease, and the beta-amyloid cascade. *Can J Neurol Sci* 2012; 39: 712-28.
- [350] Cavallucci V, D'Amelio M, Cecconi F. Aβ toxicity in Alzheimer's disease. *Mol Neurobiol* 2012; 45: 366-78.
- [351] LaFerla FM, Green KN, Oddo S. Intracellular amyloid-beta in Alzheimer's disease. *Nat Rev Neurosci* 2007; 8: 499-509.
- [352] Ahmad M, Takino T, Miyamori H, Yoshizaki T, Furukawa M, Sato H. Cleavage of amyloid-beta precursor protein (APP) by membrane-type matrix metalloproteinases. *J Biochem* 2006; 139: 517-26.
- [353] Allinson TM, Parkin ET, Turner AJ, Hooper NM. ADAMs family members as amyloid precursor protein alpha-secretases. *J Neurosci Res* 2003; 74: 342-52.

- [354] Endres K, Fahrenholz F. Upregulation of the alpha-secretase ADAM10--risk or reason for hope? *FEBS J* 2010; 277: 1585-96.
- [355] Saftig P, Reiss K. The "A Disintegrin And Metalloproteases" ADAM10 and ADAM17: novel drug targets with therapeutic potential? *Eur J Cell Biol* 2011; 90: 527-35.
- [356] Weskamp G, Cai H, Brodie TA *et al.* Mice lacking the metalloprotease-disintegrin MDC9 (ADAM9) have no evident major abnormalities during development or adult life. *Mol Cell Biol* 2002; 22: 1537-44.
- [357] Cong L, Jia J. Promoter polymorphisms which regulate ADAM9 transcription are protective against sporadic Alzheimer's disease. *Neurobiol Aging* 2011; 32: 54-62.
- [358] Cissé MA, Sunyach C, Lefranc-Jullien S, Postina R, Vincent B, Checler F. The disintegrin ADAM9 indirectly contributes to the physiological processing of cellular prion by modulating ADAM10 activity. *J Biol Chem* 2005; 280: 40624-31.
- [359] Tousseyn T, Thathiah A, Jorissen E *et al.* ADAM10, the rate-limiting protease of regulated intramembrane proteolysis of Notch and other proteins, is processed by ADAMS-9, ADAMS-15, and the gamma-secretase. *J Biol Chem* 2009; 284: 11738-47.
- [360] Parkin E, Harris B. A disintegrin and metalloproteinase (ADAM)-mediated ectodomain shedding of ADAM10. *J Neurochem* 2009; 108: 1464-79.
- [361] Postina R, Schroeder A, Dewachter I *et al.* A disintegrin-metalloproteinase prevents amyloid plaque formation and hippocampal defects in an Alzheimer disease mouse model. *J Clin Invest* 2004; 113: 1456-64.
- [362] Schmitt U, Hiemke C, Fahrenholz F, Schroeder A. Over-expression of two different forms of the alpha-secretase ADAM10 affects learning and memory in mice. *Behav Brain Res* 2006; 175: 278-84.
- [363] Bell KF, Zheng L, Fahrenholz F, Cuello AC. ADAM-10 over-expression increases cortical synaptogenesis. *Neurobiol Aging* 2008; 29: 554-65.
- [364] Schroeder A, Fahrenholz F, Schmitt U. Effect of a dominant-negative form of ADAM10 in a mouse model of Alzheimer's disease. *J Alzheimers Dis* 2009; 16: 309-14.
- [365] Chen YY, Hehr CL, Atkinson-Leadbeater K, Hocking JC, McFarlane S. Targeting of retinal axons requires the metalloproteinase ADAM10. *J Neurosci* 2007; 27: 8448-56.
- [366] Jangouk P, Dehmel T, Meyer Zu Hörste G, Ludwig A, Lehmann HC, Kieseier BC. Involvement of ADAM10 in axonal outgrowth and myelination of the peripheral nerve. *Glia* 2009; 57: 1765-74.
- [367] Hattori M, Osterfield M, Flanagan JG. Regulated cleavage of a contact-mediated axon repellent. *Science* 2000; 289: 1360-5.
- [368] Janes PW, Saha N, Barton WA *et al.* Adam meets Eph: an ADAM substrate recognition module acts as a molecular switch for ephrin cleavage in trans. *Cell* 2005; 123: 291-304.
- [369] Buxbaum JD, Liu KN, Luo Y *et al.* Evidence that tumor necrosis factor alpha converting enzyme is involved in regulated alpha-secretase cleavage of the Alzheimer amyloid protein precursor. *J Biol Chem* 1998; 273: 27765-7.
- [370] Merlos-Suárez A, Fernández-Larrea J, Reddy P, Baselga J, Arribas J. Pro-tumor necrosis factor-alpha processing activity is tightly controlled by a component that does not affect notch processing. *J Biol Chem* 1998; 273: 24955-62.
- [371] Slack BE, Ma LK, Seah CC. Constitutive shedding of the amyloid precursor protein ectodomain is up-regulated by tumour necrosis factor-alpha converting enzyme. *Biochem J* 2001; 357: 787-94.
- [372] Endres K, Anders A, Kojro E, Gilbert S, Fahrenholz F, Postina R. Tumor necrosis factor-alpha converting enzyme is processed by proprotein-convertases to its mature form which is degraded upon phorbol ester stimulation. *Eur J Biochem* 2003; 270: 2386-93.

- [373] Skovronsky DM, Fath S, Lee VM, Milla ME. Neuronal localization of the TNF α converting enzyme (TACE) in brain tissue and its correlation to amyloid plaques. *J Neurobiol* 2001; 49: 40-6.
- [374] Gottschall PE, Deb S. Regulation of matrix metalloproteinase expressions in astrocytes, microglia and neurons. *Neuroimmunomodulation* 1996; 3: 69-75.
- [375] Oh LY, Larsen PH, Krekoski CA *et al.* Matrix metalloproteinase-9/gelatinase B is required for process outgrowth by oligodendrocytes. *J Neurosci* 1999; 19: 8464-75.
- [376] Leake A, Morris CM, Whateley J. Brain matrix metalloproteinase 1 levels are elevated in Alzheimer's disease. *Neurosci Lett* 2000; 291: 201-3.
- [377] Yin KJ, Cirrito JR, Yan P *et al.* Matrix metalloproteinases expressed by astrocytes mediate extracellular amyloid-beta peptide catabolism. *J Neurosci* 2006; 26: 10939-48.
- [378] Walsh DM, Minogue AM, Sala Frigerio C, Fadeeva JV, Wasco W, Selkoe DJ. The APP family of proteins: similarities and differences. *Biochem Soc Trans* 2007; 35: 416-20.
- [379] Deb S, Gottschall PE. Increased production of matrix metalloproteinases in enriched astrocyte and mixed hippocampal cultures treated with beta-amyloid peptides. *J Neurochem* 1996; 66: 1641-7.
- [380] Yan P, Hu X, Song H *et al.* Matrix metalloproteinase-9 degrades amyloid-beta fibrils in vitro and compact plaques in situ. *J Biol Chem* 2006; 281: 24566-74.
- [381] Ito S, Kimura K, Haneda M, Ishida Y, Sawada M, Isobe K. Induction of matrix metalloproteinases (MMP3, MMP12 and MMP13) expression in the microglia by amyloid-beta stimulation via the PI3K/Akt pathway. *Exp Gerontol* 2007; 42: 532-7.
- [382] Talamagas AA, Efthimiopoulos S, Tsilibary EC, Figueiredo-Pereira ME, Tzinia AK. A β (1-40)-induced secretion of matrix metalloproteinase-9 results in sAPP α release by association with cell surface APP. *Neurobiol Dis* 2007; 28: 304-15.
- [383] White AR, Du T, Loughton KM *et al.* Degradation of the Alzheimer disease amyloid beta-peptide by metal-dependent up-regulation of metalloprotease activity. *J Biol Chem* 2006; 281: 17670-80.
- [384] Caragounis A, Du T, Filiz G *et al.* Differential modulation of Alzheimer's disease amyloid beta-peptide accumulation by diverse classes of metal ligands. *Biochem J* 2007; 407: 435-50.
- [385] Filiz G, Price KA, Caragounis A, Du T, Crouch PJ, White AR. The role of metals in modulating metalloprotease activity in the AD brain. *Eur Biophys J* 2008; 37: 315-21.
- [386] Selkoe DJ. Biochemistry and molecular biology of amyloid beta-protein and the mechanism of Alzheimer's disease. *Handb Clin Neurol* 2008; 89: 245-60.
- [387] Yoshiyama Y, Asahina M, Hattori T. Selective distribution of matrix metalloproteinase-3 (MMP-3) in Alzheimer's disease brain. *Acta Neuropathol* 2000; 99: 91-5.
- [388] Horstmann S, Budig L, Gardner H *et al.* Matrix metalloproteinases in peripheral blood and cerebrospinal fluid in patients with Alzheimer's disease. *Int Psychogeriatr* 2010; 22: 966-72.
- [389] Stomrud E, Björkqvist M, Janciauskiene S, Minthon L, Hansson O. Alterations of matrix metalloproteinases in the healthy elderly with increased risk of prodromal Alzheimer's disease. *Alzheimers Res Ther* 2010; 2: 20.
- [390] Backstrom JR, Lim GP, Cullen MJ, Tökés ZA. Matrix metalloproteinase-9 (MMP-9) is synthesized in neurons of the human hippocampus and is capable of degrading the amyloid-beta peptide (1-40). *J Neurosci* 1996; 16: 7910-9.
- [391] Lorenzl S, Albers DS, Relkin N *et al.* Increased plasma levels of matrix metalloproteinase-9 in patients with Alzheimer's disease. *Neurochem Int* 2003; 43: 191-6.

- [392] Brown PD. Synthetic inhibitors of matrix metalloproteinases. In: Parks WC, Mecham RP. Eds. Matrix metalloproteinases. San Diego, CA: Academic Press 1998; pp. 243-262.
- [393] Rosenberg GA, Kornfeld M, Estrada E, Kelley RO, Liotta LA, Stetler-Stevenson WG. TIMP-2 reduces proteolytic opening of blood-brain barrier by type IV collagenase. *Brain Res* 1992; 576: 203-7.
- [394] Gijbels K, Galardy RE, Steinman L. Reversal of experimental autoimmune encephalomyelitis with a hydroxamate inhibitor of matrix metalloproteinases. *J Clin Invest* 1994; 94: 2177-82.
- [395] Hewson AK, Smith T, Leonard JP, Cuzner ML. Suppression of experimental allergic encephalomyelitis in the Lewis rat by the matrix metalloproteinase inhibitor Ro31-9790. *Inflamm Res* 1995; 44: 345-9.
- [396] Clements JM, Cossins JA, Wells GM *et al.* Matrix metalloproteinase expression during experimental autoimmune encephalomyelitis and effects of a combined matrix metalloproteinase and tumour necrosis factor-alpha inhibitor. *J Neuroimmunol* 1997; 74: 85-94.
- [397] Mun-Bryce S, Rosenberg GA. Gelatinase B modulates selective opening of the blood-brain barrier during inflammation. *Am J Physiol* 1998; 274: R1203-11.
- [398] Sood RR, Taheri S, Candelario-Jalil E, Estrada EY, Rosenberg GA. Early beneficial effect of matrix metalloproteinase inhibition on blood-brain barrier permeability as measured by magnetic resonance imaging countered by impaired long-term recovery after stroke in rat brain. *J Cereb Blood Flow Metab* 2008; 28: 431-8.
- [399] Leib SL, Clements JM, Lindberg RL *et al.* Inhibition of matrix metalloproteinases and tumour necrosis factor alpha converting enzyme as adjuvant therapy in pneumococcal meningitis. *Brain* 2001; 124: 1734-42.
- [400] Gu Z, Cui J, Brown S *et al.* A highly specific inhibitor of matrix metalloproteinase-9 rescues laminin from proteolysis and neurons from apoptosis in transient focal cerebral ischemia. *J Neurosci* 2005; 25: 6401-8.
- [401] Lapchak PA, Chapman DF, Zivin JA. Metalloproteinase inhibition reduces thrombolytic (tissue plasminogen activator)-induced hemorrhage after thromboembolic stroke. *Stroke* 2000; 31: 3034-40.
- [402] Pfefferkorn T, Rosenberg GA. Closure of the blood-brain barrier by matrix metalloproteinase inhibition reduces rtPA-mediated mortality in cerebral ischemia with delayed reperfusion. *Stroke* 2003; 34: 2025-30.
- [403] Fisher JF, Mobashery S. Recent advances in MMP inhibitor design. *Cancer Metastasis Rev* 2006; 25: 115-36.
- [404] Puerta DT, Mongan J, Tran BL, McCammon JA, Cohen SM. Potent, selective pyrone-based inhibitors of stromelysin-1. *J Am Chem Soc* 2005; 127: 14148-9.
- [405] Corbitt CA, Lin J, Lindsey ML. Mechanisms to inhibit matrix metalloproteinase activity: where are we in the development of clinically relevant inhibitors? *Recent Pat Anticancer Drug Discov* 2007; 2: 135-42.
- [406] Coussens LM, Fingleton B, Matrisian LM. Matrix metalloproteinase inhibitors and cancer: trials and tribulations. *Science* 2002; 295: 2387-92.
- [407] Cho Y, Son HJ, Kim EM *et al.* Doxycycline is neuroprotective against nigral dopaminergic degeneration by a dual mechanism involving MMP-3. *Neurotox Res* 2009; 16: 361-71.
- [408] Du Y, Ma Z, Lin S *et al.* Minocycline prevents nigrostriatal dopaminergic neurodegeneration in the MPTP model of Parkinson's disease. *Proc Natl Acad Sci USA* 2001; 98: 14669-74.
- [409] Tikka T, Fiebich BL, Goldsteins G, Keinanen R, Koistinaho J. Minocycline, a tetracycline derivative, is neuroprotective against excitotoxicity by inhibiting activation and proliferation of microglia. *J Neurosci* 2001; 21: 2580-8.
- [410] Wu DC, Jackson-Lewis V, Vila M *et al.* Blockade of microglial activation is neuroprotective in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson disease. *J Neurosci* 2002; 22: 1763-71.

- [411] Matsukawa N, Yasuhara T, Hara K *et al.* Therapeutic targets and limits of minocycline neuroprotection in experimental ischemic stroke. *BMC Neurosci* 2009; 10: 126.
- [412] Moon M, Kim HG, Hwang L *et al.* Neuroprotective effect of ghrelin in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson's disease by blocking microglial activation. *Neurotox Res* 2009; 15: 332-47.
- [413] Lee J, Lim E, Kim Y, Li E, Park S. Ghrelin attenuates kainic acid-induced neuronal cell death in the mouse hippocampus. *J Endocrinol* 2010; 205: 263-70.
- [414] Lee EJ, Kim SY, Hyun JW, Min SW, Kim DH, Kim HS. Glycitein inhibits glioma cell invasion through down-regulation of MMP-3 and MMP-9 gene expression. *Chem Biol Interact* 2010; 185: 18-24.
- [415] Kim S, Moon M, Park S. Exendin-4 protects dopaminergic neurons by inhibition of microglial activation and matrix metalloproteinase-3 expression in an animal model of Parkinson's disease. *J Endocrinol* 2009; 202: 431-9.
- [416] Kojro E, Gimpl G, Lammich S, Marz W, Fahrenholz F. Low cholesterol stimulates the nonamyloidogenic pathway by its effect on the alpha -secretase ADAM 10. *Proc Natl Acad Sci USA* 2001; 98: 5815-20.
- [417] Matthews V, Schuster B, Schütze S *et al.* Cellular cholesterol depletion triggers shedding of the human interleukin-6 receptor by ADAM10 and ADAM17 (TACE). *J Biol Chem* 2003; 278: 38829-39.
- [418] Felson DT. Developments in the clinical understanding of osteoarthritis. *Arthritis Res Ther* 2009; 11: 203.
- [419] Goldring MB, Goldring SR. Articular cartilage and subchondral bone in the pathogenesis of osteoarthritis. *Ann NY Acad Sci* 2010; 1192: 230-7.
- [420] Blom AB, van Lent PL, Holthuysen AE *et al.* Synovial lining macrophages mediate osteophyte formation during experimental osteoarthritis. *Osteoarthritis Cartilage* 2004; 12: 627-35.
- [421] Goldring MB, Goldring SR. Osteoarthritis. *J Cell Physiol* 2007; 213: 626-34.
- [422] Heinegård D, Saxne T. The role of the cartilage matrix in osteoarthritis. *Nat Rev Rheumatol* 2011; 7: 50-6.
- [423] Nagase H, Kashiwagi M. Aggrecanases and cartilage matrix degradation. *Arthritis Res Ther* 2003; 5: 94-103.
- [424] Billingham RC, Dahlberg L, Ionescu M *et al.* Enhanced cleavage of type II collagen by collagenases in osteoarthritic articular cartilage. *J Clin Invest* 1997; 99: 1534-45.
- [425] Lohmander LS, Atley LM, Pietka TA, Eyre DR. The release of crosslinked peptides from type II collagen into human synovial fluid is increased soon after joint injury and in osteoarthritis. *Arthritis Rheum* 2003; 48: 3130-9.
- [426] Lee JH, Fitzgerald JB, Dimicco MA, Grodzinsky AJ. Mechanical injury of cartilage explants causes specific time-dependent changes in chondrocyte gene expression. *Arthritis Rheum* 2005; 52: 2386-95.
- [427] Guo D, Tan W, Wang F *et al.* Proteomic analysis of human articular cartilage: identification of differentially expressed proteins in knee osteoarthritis. *Joint Bone Spine* 2008; 75: 439-44.
- [428] Kuno K, Okada Y, Kawashima H *et al.* ADAMTS-1 cleaves a cartilage proteoglycan, aggrecan. *FEBS Lett* 2000; 478: 241-5.
- [429] Somerville RP, Longpre JM, Jungers KA *et al.* Characterization of ADAMTS-9 and ADAMTS-20 as a distinct ADAMTS subfamily related to *Caenorhabditis elegans* GON-1. *J Biol Chem* 2003; 278: 9503-13.
- [430] Collins-Racie LA, Flannery CR, Zeng W *et al.* ADAMTS-8 exhibits aggrecanase activity and is expressed in human articular cartilage. *Matrix Biol* 2004; 23: 219-30.
- [431] Zeng W, Corcoran C, Collins-Racie LA, Lavallie ER, Morris EA, Flannery CR. Glycosaminoglycan-binding properties and aggrecanase activities of truncated ADAMTSs: comparative analyses with ADAMTS-5, -9, -16 and -18. *Biochim Biophys Acta* 2006; 1760: 517-24.

- [432] Gendron C, Kashiwagi M, Lim NH *et al.* Proteolytic activities of human ADAMTS-5: comparative studies with ADAMTS-4. *J Biol Chem* 2007; 282: 18294-306.
- [433] Sandy JD, Verscharen C. Analysis of aggrecan in human knee cartilage and synovial fluid indicates that aggrecanase (ADAMTS) activity is responsible for the catabolic turnover and loss of whole aggrecan whereas other protease activity is required for C-terminal processing in vivo. *Biochem J* 2001; 358: 615-26.
- [434] Sandy JD. A contentious issue finds some clarity: on the independent and complementary roles of aggrecanase activity and MMP activity in human joint aggrecanolytic activity. *Osteoarthritis Cartilage* 2006; 14: 95-100.
- [435] Little CB, Meeker CT, Golub SB *et al.* Blocking aggrecanase cleavage in the aggrecan interglobular domain abrogates cartilage erosion and promotes cartilage repair. *J Clin Invest* 2007; 117: 1627-36.
- [436] Bau B, Gebhard PM, Haag J, Knorr T, Bartnik E, Aigner T. Relative messenger RNA expression profiling of collagenases and aggrecanases in human articular chondrocytes in vivo and in vitro. *Arthritis Rheum* 2002; 46: 2648-57.
- [437] Bondeson J, Wainwright SD, Lauder S, Amos N, Hughes CE. The role of synovial macrophages and macrophage-produced cytokines in driving aggrecanases, matrix metalloproteinases, and other destructive and inflammatory responses in osteoarthritis. *Arthritis Res Ther* 2006; 8: R187.
- [438] Glasson SS, Askew R, Sheppard B *et al.* Deletion of active ADAMTS5 prevents cartilage degradation in a murine model of osteoarthritis. *Nature* 2005; 434: 644-8.
- [439] Stanton H, Rogerson FM, East CJ *et al.* ADAMTS5 is the major aggrecanase in mouse cartilage in vivo and in vitro. *Nature* 2005; 434: 648-52.
- [440] Glasson SS, Askew R, Sheppard B *et al.* Characterization of and osteoarthritis susceptibility in ADAMTS-4-knockout mice. *Arthritis Rheum* 2004; 50: 2547-58.
- [441] Song RH, Tortorella MD, Malfait AM *et al.* Aggrecan degradation in human articular cartilage explants is mediated by both ADAMTS-4 and ADAMTS-5. *Arthritis Rheum* 2007; 56: 575-85.
- [442] Naito S, Shiomi T, Okada A *et al.* Expression of ADAMTS4 (aggrecanase-1) in human osteoarthritic cartilage. *Pathol Int* 2007; 57: 703-11.
- [443] Pratta MA, Scherle PA, Yang G, Liu RQ, Newton RC. Induction of aggrecanase 1 (ADAM-TS4) by interleukin-1 occurs through activation of constitutively produced protein. *Arthritis Rheum* 2003; 48: 119-33.
- [444] Rogerson FM, Stanton H, East CJ *et al.* Evidence of a novel aggrecan-degrading activity in cartilage: Studies of mice deficient in both ADAMTS-4 and ADAMTS-5. *Arthritis Rheum* 2008; 58: 1664-73.
- [445] Mitchell PG, Magna HA, Reeves LM *et al.* Cloning, expression, and type II collagenolytic activity of matrix metalloproteinase-13 from human osteoarthritic cartilage. *J Clin Invest* 1996; 97: 761-8.
- [446] Kevorkian L, Young DA, Darrah C *et al.* Expression profiling of metalloproteinases and their inhibitors in cartilage. *Arthritis Rheum* 2004; 50: 131-41.
- [447] Little CB, Barai A, Burkhardt D *et al.* Matrix metalloproteinase 13-deficient mice are resistant to osteoarthritic cartilage erosion but not chondrocyte hypertrophy or osteophyte development. *Arthritis Rheum* 2009; 60: 3723-33.
- [448] Inada M, Wang Y, Byrne MH *et al.* Critical roles for collagenase-3 (Mmp13) in development of growth plate cartilage and in endochondral ossification. *Proc Natl Acad Sci USA* 2004; 101: 17192-7.
- [449] Stickens D, Behonick DJ, Ortega N *et al.* Altered endochondral bone development in matrix metalloproteinase 13-deficient mice. *Development* 2004; 131: 5883-95.
- [450] Zack MD, Malfait AM, Skepner AP *et al.* ADAM-8 isolated from human osteoarthritic chondrocytes cleaves fibronectin at Ala(271). *Arthritis Rheum* 2009; 60: 2704-13.

- [451] van Meurs J, van Lent P, Stoop R *et al.* Cleavage of aggrecan at the Asn341-Phe342 site coincides with the initiation of collagen damage in murine antigen-induced arthritis: a pivotal role for stromelysin 1 in matrix metalloproteinase activity. *Arthritis Rheum* 1999; 42: 2074-84.
- [452] Kubota E, Imamura H, Kubota T, Shibata T, Murakami K. Interleukin 1 beta and stromelysin (MMP3) activity of synovial fluid as possible markers of osteoarthritis in the temporomandibular joint. *J Oral Maxillofac Surg* 1997; 55: 20-7.
- [453] Garner BC, Stoker AM, Kuroki K, Evans R, Cook CR, Cook JL. Using animal models in osteoarthritis biomarker research. *J Knee Surg* 2011; 24: 251-64.
- [454] Tetlow LC, Adlam DJ, Woolley DE. Matrix metalloproteinase and proinflammatory cytokine production by chondrocytes of human osteoarthritic cartilage: associations with degenerative changes. *Arthritis Rheum* 2001; 44: 585-94.
- [455] Sofat N. Analysing the role of endogenous matrix molecules in the development of osteoarthritis. *Int J Exp Pathol* 2009; 90: 463-79.
- [456] Klatt AR, Paul-Klausch B, Klinger G *et al.* A critical role for collagen II in cartilage matrix degradation: collagen II induces pro-inflammatory cytokines and MMPs in primary human chondrocytes. *J Orthop Res* 2009; 27: 65-70.
- [457] Fichter M, Körner U, Schömburg J, Jennings L, Cole AA, Mollenhauer J. Collagen degradation products modulate matrix metalloproteinase expression in cultured articular chondrocytes. *J Orthop Res* 2006; 24: 63-70.
- [458] Jennings L, Wu L, King KB, Hämmerle H, Cs-Szabo G, Mollenhauer J. The effects of collagen fragments on the extracellular matrix metabolism of bovine and human chondrocytes. *Connect Tissue Res* 2001; 42: 71-86.
- [459] Pelletier JP, Martel-Pelletier J, Abramson SB. Osteoarthritis, an inflammatory disease: potential implication for the selection of new therapeutic targets. *Arthritis Rheum* 2001; 44: 1237-47.
- [460] Benito MJ, Veale DJ, FitzGerald O, van den Berg WB, Bresnihan B. Synovial tissue inflammation in early and late osteoarthritis. *Ann Rheum Dis* 2005; 64: 1263-7.
- [461] Goldring MB, Otero M, Tsuchimochi K, Ijiri K, Li Y. Defining the roles of inflammatory and anabolic cytokines in cartilage metabolism. *Ann Rheum Dis* 2008; 67: 75-82.
- [462] Heinegård D, Saxne T. The role of the cartilage matrix in osteoarthritis. *Nat Rev Rheumatol* 2011; 7: 50-6.
- [463] Kapoor M, Martel-Pelletier J, Lajeunesse D, Pelletier JP, Fahmi H. Role of proinflammatory cytokines in the pathophysiology of osteoarthritis. *Nat Rev Rheumatol* 2011; 7: 33-42.
- [464] Haynes MK, Hume EL, Smith JB. Phenotypic characterization of inflammatory cells from osteoarthritic synovium and synovial fluids. *Clin Immunol* 2002; 105: 315-25.
- [465] Sawaji Y, Hynes J, Vincent T, Saklatvala J. Fibroblast growth factor 2 inhibits induction of aggrecanase activity in human articular cartilage. *Arthritis Rheum* 2008; 58: 3498-509.
- [466] Tortorella MD, Malfait AM, Deccico C, Arner E. The role of ADAM-TS4 (aggrecanase-1) and ADAM-TS5 (aggrecanase-2) in a model of cartilage degradation. *Osteoarthritis Cartilage* 2001; 9: 539-52.
- [467] Yamanishi Y, Boyle DL, Clark M *et al.* Expression and regulation of aggrecanase in arthritis: the role of TGF-beta. *J Immunol* 2002; 168: 1405-12.
- [468] Koshy PJ, Lundy CJ, Rowan AD *et al.* The modulation of matrix metalloproteinase and ADAM gene expression in human chondrocytes by interleukin-1 and oncostatin M: a time-course study using real-time quantitative reverse transcription-polymerase chain reaction. *Arthritis Rheum* 2002; 46: 961-7.
- [469] Zwerina J, Redlich K, Polzer K *et al.* TNF-induced structural joint damage is mediated by IL-1. *Proc Natl Acad Sci USA* 2007; 104: 11742-7.

- [470] Séguin CA, Bojarski M, Pilliar RM, Roughley PJ, Kandel RA. Differential regulation of matrix degrading enzymes in a TNF α -induced model of nucleus pulposus tissue degeneration. *Matrix Biol* 2006; 25: 409-18.
- [471] Kobayashi M, Squires GR, Mousa A *et al.* Role of interleukin-1 and tumor necrosis factor alpha in matrix degradation of human osteoarthritic cartilage. *Arthritis Rheum* 2005; 52: 128-35.
- [472] Page Thomas DP, King B, Stephens T, Dingle JT. In vivo studies of cartilage regeneration after damage induced by catabolin/interleukin-1. *Ann Rheum Dis* 1991; 50: 75-80.
- [473] Barksby HE, Hui W, Wappler I *et al.* Interleukin-1 in combination with oncostatin M up-regulates multiple genes in chondrocytes: implications for cartilage destruction and repair. *Arthritis Rheum* 2006; 54: 540-50.
- [474] Vuolteenaho K, Moilanen T, Jalonen U *et al.* TGF β inhibits IL-1 α -induced iNOS expression and NO production in immortalized chondrocytes. *Inflamm Res* 2005; 54: 420-7.
- [475] Glasson SS. In vivo osteoarthritis target validation utilizing genetically-modified mice. *Curr Drug Targets* 2007; 8: 367-76.
- [476] Van den Berg WB. Lessons from animal models of arthritis. *Curr Rheumatol Rep* 2002; 4: 232-9.
- [477] van der Kraan PM, Blaney Davidson EN, van den Berg WB. A role for age-related changes in TGF β signaling in aberrant chondrocyte differentiation and osteoarthritis. *Arthritis Res Ther* 2010; 12: 201.
- [478] Blaney Davidson EN, Remst DF, Vitters EL *et al.* Increase in ALK1/ALK5 ratio as a cause for elevated MMP-13 expression in osteoarthritis in humans and mice. *J Immunol* 2009; 182: 7937-45.
- [479] Hoff P, Buttgerit F, Burmester GR *et al.* Osteoarthritis synovial fluid activates pro-inflammatory cytokines in primary human chondrocytes. *Int Orthop* 2013; 37: 145-51.
- [480] Pulai JI, Chen H, Im HJ *et al.* NF- κ B mediates the stimulation of cytokine and chemokine expression by human articular chondrocytes in response to fibronectin fragments. *J Immunol* 2005; 174: 5781-8.
- [481] Appleton CT, Pitelka V, Henry J, Beier F. Global analyses of gene expression in early experimental osteoarthritis. *Arthritis Rheum* 2007; 56: 1854-68.
- [482] Brühl H, Mack M, Niedermeier M, Lochbaum D, Schölmerich J, Straub RH. Functional expression of the chemokine receptor CCR7 on fibroblast-like synoviocytes. *Rheumatology (Oxford)* 2008; 47: 1771-4.
- [483] Brew K, Nagase H. The tissue inhibitors of metalloproteinases (TIMPs): an ancient family with structural and functional diversity. *Biochim Biophys Acta* 2010; 1803: 55-71.
- [484] Hashimoto G, Aoki T, Nakamura H, Tanzawa K, Okada Y. Inhibition of ADAMTS4 (aggrecanase-1) by tissue inhibitors of metalloproteinases (TIMP-1, 2, 3 and 4). *FEBS Lett* 2001; 494: 192-5.
- [485] Kashiwagi M, Tortorella M, Nagase H, Brew K. TIMP-3 is a potent inhibitor of aggrecanase 1 (ADAM-TS4) and aggrecanase 2 (ADAM-TS5). *J Biol Chem* 2010; 276: 12501-4.
- [486] Wayne GJ, Deng SJ, Amour A *et al.* TIMP-3 inhibition of ADAMTS-4 (Aggrecanase-1) is modulated by interactions between aggrecan and the C-terminal domain of ADAMTS-4. *J Biol Chem* 2007; 282: 20991-8.
- [487] Gendron C, Kashiwagi M, Hughes C, Caterson B, Nagase H. TIMP-3 inhibits aggrecanase-mediated glycosaminoglycan release from cartilage explants stimulated by catabolic factors. *FEBS Lett* 2003; 555: 431-6.
- [488] Sahebjam S, Khokha R, Mort JS. Increased collagen and aggrecan degradation with age in the joints of Timp3(-/-) mice. *Arthritis Rheum* 2007; 56: 905-9.
- [489] Mahmoodi M, Sahebjam S, Smookler D, Khokha R, Mort JS. Lack of tissue inhibitor of metalloproteinases-3 results in an enhanced inflammatory response in antigen-induced arthritis. *Am J Pathol* 2005; 166: 1733-40.
- [490] Morris KJ, Cs-Szabo G, Cole AA. Characterization of TIMP-3 in human articular talar cartilage. *Connect Tissue Res* 2010; 51: 478-90.

- [491] Burrage PS, Brinckerhoff CE. Molecular targets in osteoarthritis: metalloproteinases and their inhibitors. *Curr Drug Targets* 2007; 8: 293-303.
- [492] Yao W, Wasserman ZR, Chao M *et al.* Design and synthesis of a series of (2R)-N(4)-hydroxy-2-(3-hydroxybenzyl)-N(1)-[(1S,2R)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]butanediamide derivatives as potent, selective, and orally bioavailable aggrecanase inhibitors. *J Med Chem* 2001; 44: 3347-50.
- [493] Cherney RJ, Mo R, Meyer DT *et al.* Potent and selective aggrecanase inhibitors containing cyclic P1 substituents. *Bioorg Med Chem Lett* 2003; 13: 1297-300.
- [494] Gilbert AM, Bursavich MG, Lombardi S *et al.* 5-((1H-pyrazol-4-yl)methylene)-2-thioxothiazolidin-4-one inhibitors of ADAMTS-5. *Bioorg Med Chem Lett* 2007; 17: 1189-92.
- [495] Bursavich MG, Gilbert AM, Lombardi S *et al.* Synthesis and evaluation of aryl thioxothiazolidinone inhibitors of ADAMTS-5 (Aggrecanase-2). *Bioorg Med Chem Lett* 2007; 17: 1185-8.
- [496] Glaser KB, Li J, Aakre ME *et al.* Transforming growth factor beta mimetics: discovery of 7-[4-(4-cyanophenyl)phenoxy]-heptanohydroxamic acid, a biaryl hydroxamate inhibitor of histone deacetylase. *Mol Cancer Ther* 2002; 1: 759-68.
- [497] Zhang H, Zou K, Tesseur I, Wyss-Coray T. Small molecule tgf-beta mimetics as potential neuroprotective factors. *Curr Alzheimer Res* 2005; 2: 183-6.
- [498] Bernard DJ, Lee KB, Santos MM. Activin B can signal through both ALK4 and ALK7 in gonadotrope cells. *Reprod Biol Endocrinol* 2006; 4: 52.
- [499] Bondeson J, Lauder S, Wainwright S *et al.* Adenoviral gene transfer of the endogenous inhibitor IkappaBalpha into human osteoarthritis synovial fibroblasts demonstrates that several matrix metalloproteinases and aggrecanases are nuclear factor-kappaB-dependent. *J Rheumatol* 2007; 34(3): 523-33.
- [500] Bondeson J, Wainwright S, Hughes C, Caterson B. The regulation of the ADAMTS4 and ADAMTS5 aggrecanases in osteoarthritis: a review. *Clin Exp Rheumatol* 2008; 26: 139-45.
- [501] Pattoli MA, MacMaster JF, Gregor KR, Burke JR. Collagen and aggrecan degradation is blocked in interleukin-1-treated cartilage explants by an inhibitor of IkappaB kinase through suppression of metalloproteinase expression. *J Pharmacol Exp Ther* 2005; 315: 382-8.
- [502] Pelletier JP, Martel-Pelletier J, Raynauld JP. Most recent developments in strategies to reduce the progression of structural changes in osteoarthritis: today and tomorrow. *Arthritis Res Ther* 2006; 8: 206.

Figure legend

Figure 1. Several stimuli induce specific activated cells to release metalloproteinases which are involved in the pathogenesis of age-related chronic diseases.

Figure 1.




Table 1. Human ADAM members and suggested functions.

| ADAM | Proteolytic activity | Suggested functions |
|--------------|---|--|
| ADAM-1 | No | Possibly involved in sperm-egg fusion |
| ADAM-2 | No | Possibly involved in sperm-egg fusion |
| ADAM-3A | No | Possibly involved in sperm-egg fusion |
| ADAM-5P | No | Unknown |
| ADAM-6 | No | Unknown |
| ADAM-7 | No | Possibly involved in sperm-egg fusion |
| ADAM-8 | CD23, TNFRI, IL-1RII | Involved in cell adhesion during neurodegeneration; a target for allergic respiratory diseases, including asthma |
| ADAM-9 | APP | Involved in induced ectodomain shedding of membrane-anchored heparin binding-epidermal growth factor (HB-EGF)-like growth factor; suggested to cleave APP at the α -secretase site |
| ADAM-10 | APP, CX3CL1, collagen IV | Involved in shedding of various transmembrane proteins, including cadherins; α -secretase for APP; control wound healing, neurogenesis, skin homeostasis |
| ADAM-11 | No | Candidate tumor suppressor gene for human breast cancer; pain transmission, synaptic modulation |
| ADAM-12 | HB-EGF sheddase | Involved in myogenesis and skeletal muscle regeneration; upregulated in tumor progression; involved in osteoclast formation from mononuclear precursors |
| ADAM-15 | collagen IV, gelatin | Involved in cell adhesion through integrin binding; wound healing, mediates heterotypic T-cell interactions; cleaves E-cadherin; glomerular cell migration and pathological neovascularization; cartilage remodeling |
| ADAM-17 | APP, TNF- α , TNFRI and RII, IL-1RII | Proteolytic release of TNF- α , p75TNF receptor, IL-1R type II, p55 TNFR, TGF- α , L-selectin and APP; activation of Notch pathway |
| ADAM-18 | No | Possibly involved in sperm-egg fusion |
| ADAM-19 | TRANCE | Involved in neurogenesis and synaptogenesis; osteoblast differentiation and/or activity in bone |
| ADAM-20 | Yes | Possibly involved in sperm-egg fusion |
| ADAM-21 | Yes | Possibly involved in sperm-egg fusion and in epithelia functions |
| ADAM-22 | No | Ligand for integrin in the brain; involved in regulation of cell adhesion, inhibition of cell proliferation, neuronal receptor for leucin-rich-glioma inactivated1 (LG11) |
| ADAM-23 | No | Involved in cell-cell and cell-matrix interactions in the brain |
| ADAM-28 | CD40, FasL | Role during lymphocyte migration; shedding of lymphocyte surface target protein; sperm maturation |
| ADAM-29, -32 | No | Involved in spermatogenesis and fertilization |
| ADAM-30 | Yes | Involved in spermatogenesis and fertilization |
| ADAM-33 | Yes | Involved in asthma and bronchial hyperresponsiveness |

Table 2. Human MMP members and principal biological effects.

| Group name | MMP | Substrates | Biological effects |
|--------------------------|--------|--|--|
| I. Collagenase | | | |
| Collagenase-1 | MMP-1 | Collagens I, II, III, VII, X, gelatin, proteoglycan, link protein, entactin, tenascin | Keratinocyte migration and reepithelialization, platelet aggregation, release of bFGF, cell proliferation, pro- and anti-inflammatory, osteoclast activation, cell migration, enhanced collagen affinity, apoptosis, increased bioavailability of TGFβ |
| Collagenase-2 | MMP-8 | Collagens I, II, III, gelatin, proteoglycan, link protein | |
| Collagenase-3 | MMP-13 | Collagens I, II, III, IV, IX, X, XIV, proteoglycan, fibronectin, tenascin | |
| II. Gelatinase | | | |
| Gelatinase A | MMP-2 | Gelatin, collagens IV, V, VII, XI, laminin, fibronectin, elastin, proteoglycan, link protein | Neurite outgrowth, generation of angiostatin-like fragment, enhanced collagen affinity, epithelial cell migration, tumor cell resistance, pro-inflammatory, increased bioavailability of TGFβ, generation of vasoconstrictor, recruitment of osteoclasts |
| Gelatinase B | MMP-9 | Gelatin, collagens III, IV, V, elastin, entactin, link protein | |
| III. Stromelysin | | | |
| Stromelysin-1 | MMP-3 | Proteoglycan, collagens III, IV, IX, X, laminin, fibronectin, gelatin, tenascin, link protein, elastin | Mammary epithelial cell apoptosis and alveolar formation, generation of angiostatin-like fragment, release of bFGF, increased bioavailability of IGF1 and cell proliferation, anti-inflammatory, increased bioavailability of TGFβ, increased cell invasion |
| Stromelysin-2 | MMP-10 | Collagens III, IV, V, fibronectin, laminin, proteoglycan, link protein, elastin | |
| Stromelysin-3 | MMP-11 | Fibronectin, laminin, proteoglycan, gelatin | |
| IV. Membrane-type | | | |
| MT1-MMP | MMP-14 | Collagens I, II, III, gelatin, proteoglycan, fibronectin, laminin | Cell migration, kidney tubulogenesis, epithelial cell migration, reduced cell adhesion and spreading, embryo attachment to uterine epithelia |
| MT2-MMP | MMP-15 | Fibronectin, tenascin, entactin, aggrecan, perlecan, laminin | |
| MT3-MMP | MMP-16 | Collagen III, gelatin, fibronectin | |
| MT4-MMP | MMP-17 | ? | |
| MT5-MMP | MMP-24 | Proteoglycan | |
| MP6-MMP | MMP-25 | Gelatin | |
| V. Others | | | |
| Matrilysin | MMP-7 | Proteoglycan, gelatin, fibronectin, tenascin, elastin, collagen IV, laminin, link protein | Adipocyte differentiation, generation of angiostatin-like fragment, increased bioavailability of IGF1 and cell proliferation, epithelial cell migration, increased bioavailability of TGFβ, disrupted cell aggregation and increased cell invasion, Fas-receptor mediated apoptosis, pro-inflammatory, osteoclast activation, vasoconstriction and cell growth |
| Metalloelastase | MMP-12 | Elastin | |
| Collagenase 4 | MMP-18 | Collagen I | |
| RAS 1-1 | MMP-19 | Tenascin, gelatin, aggrecan | |
| Enamelysin | MMP-20 | Enamel, gelatin | |
| XMMP | MMP-21 | ? | |
| No common name | MMP-23 | ? | |
| Matrilysin 2 | MMP-26 | Collagen IV, fibronectin, gelatin | |
| No common name | MMP-27 | ? | |
| Epilysin | MMP-28 | Casein | |