

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**New genetic maps for globe artichoke and wild cardoon and their alignment with an SSR-based consensus map**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/142100> since 2016-07-14T14:51:03Z

*Published version:*

DOI:10.1007/s11032-013-9861-9

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*Molecular Breeding, Volume 32, Issue 1, 2013,*

*DOI:10.1007/s11032-013-9861-9*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*<http://link.springer.com/article/10.1007%2Fs11032-013-9861-9>*

## New genetic maps for globe artichoke and wild cardoon and their alignment with an SSR-based consensus map

Eugenia MARTIN <sup>1</sup>, Vanina CRAVERO <sup>1</sup>, Ezio PORTIS <sup>2\*</sup>, Davide SCAGLIONE <sup>2</sup>, Esteban ACQUAVIVA <sup>3</sup>, Enrique COINTRY <sup>3</sup>

<sup>1</sup> CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas), Parque J.F. Villarino, CC 14, (S2125ZAA) Zavalla, Argentina

<sup>2</sup> DISAFA (Dipartimento di Scienze Agrarie, Forestali e Alimentari) Plant Genetics and Breeding, University of Torino, via L. da Vinci 44, I-10095 Grugliasco (Torino), Italy

<sup>3</sup> Cátedra de Mejoramiento Vegetal y Producción de Semillas, Facultad de Ciencias Agrarias (UNR) Parque J.F. Villarino, (S2125ZAA) Zavalla, Argentina

\*Corresponding author: [ezio.portis@unito.it](mailto:ezio.portis@unito.it); Tel: 39.011.6708807; Fax: 39.011.2368807

**Abstract:** An F<sub>1</sub> mapping population was bred by crossing an accession of wild cardoon with a single Argentinian globe artichoke plant of the variety “Estrella del Sur FCA” with a view to generate new *C. cardunculus* linkage maps. Genotyping was conducted using a set of 553 SRAP, SSR, AFLP and SNP markers. The 1,465.5 cM map based on the segregation of alleles present in the wild cardoon parent comprised 214 loci distributed across 16 linkage groups (LGs), while the 910.1 cM globe artichoke based map featured 141 loci falling into 12 LGs and the total length. Three of the morphological traits (inflorescence spininess, leaf spininess and head color) for which the parents contrasted were inherited monogenically, and the genes conditioning them were mapped. A set of 48 co-dominant loci was used to align the LGs with those derived from a reference SSR-based consensus map of the species.

**Keywords:** *Cynara cardunculus*, globe artichoke, cardoon, molecular markers, linkage groups, genetic map.

## Introduction

The *Asteraceae* species *Cynara cardunculus* L. ( $2n=2x=34$ ) includes two domesticated taxa, namely the globe artichoke (var. *scolymus*) and the cultivated cardoon (var. *altilis*), as well as the wild cardoon (var. *silvestris*), commonly considered to be the wild ancestor of both domesticated forms (Rottenberg and Zohary 1996; Lanteri *et al.* 2004). The species is highly heterozygous and suffers from inbreeding depression when self-fertilization is enforced (Cravero *et al.* 2002). Since inter-taxon hybrids are viable and fertile, globe artichoke improvement programmes can readily access germplasm across the whole species complex (Portis *et al.* 2009). Despite the impact of inbreeding depression, conventional breeding has succeeded in producing a number of high quality and genetically uniform seed-propagated globe artichoke varieties (López Anido *et al.* 2010; Garcia *et al.* 2006).

Up to now, the mode of inheritance of only a few traits (earliness, color, spininess and tightness of the head) has been investigated in detail (Pécaut 1993; López Anido *et al.* 1998; Mauromicale and Ierna 2000; Cravero *et al.* 2005; Lanteri *et al.* 2006; Portis *et al.* 2012). The development of a genetic linkage map simplifies the genetic analysis of trait variation, particularly for those traits which are polygenically inherited. The definition of trait/marker linkages provides a tool for accelerated selection via marker assisted selection (Young 1999). The first such maps for globe artichoke was developed by Lanteri *et al.* (2006) and was based on 204 loci which fell into 18 linkage groups (LGs) for the female map and 180 loci distributed in 17 LGs for the male one. Since then, marker density of the original maps has been increased (Acquadro *et al.* 2006; 2009) and a number of other maps have been elaborated (Sonnante *et al.* 2011; Portis *et al.* 2012).

Building molecular marker based linkage maps has relied on a number of marker technologies. A particularly flexible and effective one, termed “sequence related amplified polymorphism” (SRAP), has been proposed by Li and Quiros (2001). SRAP advantages include simplicity, robustness, a reasonable throughput capacity and the ready sequencing of specific fragments included in the amplicon. The design of the primers ensures that the majority of polymorphisms arise due to variation in the length of intron, promoter and spacer regions (Lin *et al.* 2003). Approximately half of all of SRAP markers are located within the genic portion of the genome (Lin *et al.* 2003; Sun *et al.* 2007).

Here, we present genetic linkage maps of *C. cardunculus* based on a cross between an accession of wild cardoon and a single plant belonging to the open pollinated Argentinian globe artichoke variety “Estrella del Sur FCA”. Most of the genotypic data needed for linkage map construction were derived from SRAP analysis, although other PCR-based markers were also included to allow cross-referencing with the consensus

map recently produced by Portis *et al.* (2012). The resulting maps were used to identify and locate the major loci controlling three key agronomic traits.

## **Materials and methods**

### **Plant materials and DNA extraction**

A mapping population was generated by pollinating a local accession of wild cardoon with a single plant of “Estrella del Sur FCA” (“Est”). The putative F<sub>1</sub> seeds were sown in a greenhouse, and seedlings at the fourth true leaf stage were transplanted, together with both crossing parents, into a loamy soil field at the Experimental Field Station of the Universidad Nacional de Rosario (33°1’S; 60°53’W). This site experiences a temperate climate and an average annual rainfall of 950 mm, and is typical of the zone in Argentina where most of the country's globe artichoke is produced (Cravero *et al.* 2010). The inter-row spacing was 1.4 m and the spacing between plants within each row was 80 cm. Genomic DNA of the presumptive hybrids and the parental genotypes was extracted from fresh leaf using a DNeasy Plant mini Kit (Qiagen). The hybrid status of each population member was verified with two SSR (simple sequence repeats) markers CELMS-30 and -37 (Acquadro *et al.* 2009) that were polymorphic between parents. This resulted in the identification of a set of 91 true F<sub>1</sub> hybrids.

### **Genotypic analysis**

The mapping population was genotyped using SRAP, SSR, AFLP (amplified fragment length polymorphism) and SNP (single nucleotide polymorphisms) markers. The number of SRAP primer combinations (PCs, Table 1) was 25. We used the protocol described by Cravero *et al.* (2007), except that each reaction only contained 1U Taq polymerase. The resulting amplicons were separated by electrophoresis through 6% (w/v) denaturing polyacrylamide gels, and then visualized by silver staining (Bassam *et al.* 1991). The AFLP procedure followed that of Vos *et al.* (1995), as modified by Lanteri *et al.* (2004). Briefly, the genomic DNA was digested with *EcoRI* and *TaqI* and ligated to a standard adaptor oligomer. The product of the ligation reaction represented the template for a PCR driven by a pair of primers complementary to the adaptor sequences plus one selective nucleotide at their 3' end (*EcoRI*+A and *TaqI*+T). A second PCR followed driven by one of 16 PCs (Table 2). The resulting amplicons were separated by electrophoresis through a 6.5% denaturing polyacrylamide gel, and

detected using a Gene ReadIR 4200 system (LI-COR) (Jackson and Matthews 2000). The set of SSRs comprised 247 loci and were assayed using a combination of CELMS (*Cynara* Enriched Library MicroSatellite; Acquadro *et al.* 2009) and CyEM (*Cynara* Expressed Microsatellite; Scaglione *et al.* 2009) markers. An initial screen was performed using as template DNA from the two parentals and six of the F<sub>1</sub> progeny, in order to identify those markers which were more informative. Only the selected markers were applied to the full mapping population. The PCR conditions applied followed the recommendations of the markers developer and the amplicons were separated in the same way as the AFLPs. SNP markers were directed at seven genes involved in the synthesis of caffeoylquinic acids, namely *HCT*, *HQT*, *C3'H*, *C4H*, *4CL*, *Acyltransf\_1* and *\_2* (Comino *et al.* 2007; 2009; Moglia *et al.* 2009; Menin *et al.* 2010). The assay was based on a tetra-primer ARMS-PCR (amplification refractory mutation system) protocol (Ye *et al.* 2001), with the amplicons being separated by electrophoresis through 2% w/v agarose gels and visualized by ethidium bromide staining.

Fragments generated by the SRAP and AFLP assays were treated as dominant markers, with each marker named according to the PC used to generate it (Tables 1 and 2) and the estimated size of the fragment; for example, locus 1.800.190 is an AFLP fragment of 190-bp length amplified by PC *Eco*+*ACC*/*Taq*+*TTT*. SSR and SNP loci were scored as co-dominant markers and identified by the primer pair used in the assay.

### **Phenotypic analysis**

The parents differ from one another in spininess of the leaf and head bracts, and by their head color: “Est” is non-spiny and forms purple heads, while the wild cardoon accession is spiny and forms green heads. The presence/absence of spines on fully developed leaves and bracts, and head color was scored over two consecutive seasons (2008 and 2009). Goodness of fit between the observed and expected segregation according to a proposed inheritance model was evaluated using the  $\chi^2$  test.

### **Construction of the linkage maps**

The marker loci were either (1) maternal testcross markers which segregated only in the female (wild cardoon) gamete, giving a ratio of 1:1 in the F<sub>1</sub> population; (2) paternal testcross markers which segregated only in the male (globe artichoke) gamete, giving a ratio of 1:1 in the F<sub>1</sub> population; and intercross markers, which segregated in both gametes, producing a segregation ratio of either 3:1 (for the SRAP and AFLP markers) or 1:2:1 (SSR and SNP markers) in the F<sub>1</sub> population. The two-way pseudo-testcross mapping strategy described by Weeden (1994) and Grattapaglia and Sederoff (1994) was used to develop two independent linkage maps,

one for each parental genotype. Goodness-of-fit between the observed and expected segregation ratios was determined by a  $\chi^2$  test. Only markers either fully consistent with monogenic segregation ( $\chi^2 \leq \chi^2_{\alpha=0.1}$ ) or showing only minor distortion ( $\chi^2_{\alpha=0.1} < \chi^2 \leq \chi^2_{\alpha=0.01}$ ) were used for the construction of the maps. Those for which the segregation was highly distorted ( $\chi^2 > \chi^2_{\alpha=0.01}$ ) were included in a second round of mapping only when their presence induced no alteration in local marker order. The LOD threshold for accepting an LG was 5.0, as estimated by JoinMap4.0 software (Van Ooijen 2006). The JoinMap settings applied were Rec=0.40, LOD=1.0 and Jump=5. Recombination values were converted to genetic distances using the Kosambi (1944) mapping function. Intercross markers were used to identify homologous LGs. The LGs derived from the wild cardoon parent have been labeled “Wild” and those from the globe artichoke parent as “Est” (Fig. 1A-C, Table 3). Linkage maps were drawn using MapChart 2.2 software (Voorrips 2002)

#### **Alignment of the “Wild” and “Est” LGs with the SSR-based consensus LGs**

The SSR and SNP markers were used to align the *de novo* LGs with those derived by Sonnante *et al.* (2011) and Portis *et al.* (2012). To achieve this, a consensus map, based exclusively on SSR and SNP markers, was assembled from the SSR and SNP segregation data collected from the “Romanesco C3” (globe artichoke) x “Atilis 41” (cultivated cardoon) mapping population described by Portis *et al.* (2012). This SSR-based consensus map had an overall length of 1068.0 cM, comprising 217 SSR and ten SNP loci arranged into 20 LGs (LOD threshold >6.0). The haploid chromosome complement was recovered by lowering the LOD threshold to 5.0. These 17 SSR-based consensus LGs are hereafter referred to as “SSR-Ref\_LGs”, applying the same I through XVII numbering system employed by Portis *et al.* (2012). When bridge markers linked the “SSR-Ref\_LGs” with the “Wild” and “Est” ones, the latter have been numbered accordingly (Fig. 1 A-C, Tab. 3).

## **Results**

### **Markers segregation**

The SRAP analysis generated 336 informative fragments, representing a mean of 13.4 (range: 4-51) per PC. Of these, 186 segregated as expected in the maternal gametes, 130 as expected in the paternal gametes, while 20 were inferred to have been in the heterozygous state in both parental lines (intercross markers). The mean number of informative AFLP markers per PC was 8.4 (range: 1-27), so that in all, 135 loci segregated, of which

78 segregated only in the female gamete, 47 in the male gamete and the remaining ten were intercross markers. On the whole, of the combined 441 SRAP and AFLP testcross markers, 59.9% segregated in the wild cardoon parental genotype, while 40.1% in “Est”. The segregation of ~24% (112 loci) of the SRAP and AFLP markers was distorted ( $\chi^2 > \chi^2_{\alpha=0.1}$ ).

Of the 61 CELMS SSRs tested, 28 were informative between the parents; of these 14 segregated only among the female gametes, 13 among the male, and only one was an intercross marker. Similarly, of the 186 CyEM SSRs tested, 25 segregated only among the female gametes, 17 among the male, and seven were intercross markers. Two of the SSR loci suffered a minor degree of segregation distortion ( $\chi^2_{\alpha=0.05} < \chi^2 \leq \chi^2_{\alpha=0.01}$ ). Five, out of the seven SNP loci genotyped, segregated in the mapping population. The *HCT* marker segregated only among the female gametes, whereas those for *HQT*, *C3'H*, *4CL* and *Acytranf\_1* segregated only among the male gametes.

### **Phenotyping**

The three morphological traits investigated segregated consistently with a 1:1 ratio, suggesting that, in the analyzed progeny, all these traits are controlled by single genes with alternative alleles in each of the parents. The  $\chi^2$  values associated with these ratios were, respectively, 5.13 for the presence/absence spines on the heads, 2.78 for the presence/absence spines on the leaf, and 1.53 for purple-green vs purple heads ( $\chi^2 \leq \chi^2_{\alpha=0.01}=6.63$ ).

### **Linkage maps construction and their alignment**

Five hundred fifty-six markers were used for maps construction (553 molecular and 3 phenotypic markers). The genetic map constructed from the female gametes (the “Wild” map) was based on segregation at 344 loci, while that from the male gametes (the “Est” map) was based on 250 loci. The number of intercross markers was 38. The linkage analysis resulted in the placement of 345 loci across both maps, comprising 187 SRAP markers, 89 AFLP markers, 64 SSR loci, three SNP loci and three phenotypic traits.

The “Wild” map featured 214 loci distributed over 16 LGs, each defined by between five and 34 loci (mean 13.4), their length ranging from 30.1 to 153.1 cM (mean 91.6cM). The overall map length was 1,465.5 cM and the mean inter-marker distance was 7.4 cM (range 3.1–15.0 cM). The placement of one SNP and 28 SSR loci allowed the alignment of the 16 LGs with 13 consensus SSR-based map LGs (Table 3; Fig. 1A-C); three pairs of “Wild” LGs (Ia and Ib, Va and Vb, XVa and XVb) each corresponded with a single consensus

SSR-based one. Among the loci showing segregation distortion, 28 were successfully integrated within ten of the LGs. Clusters of distorted loci were detected on Wild\_VIII (four loci) and Wild\_XII (ten loci) (Fig. 1 B-C).

The “Est” map comprised 141 loci grouped into 12 LGs, which covered a genetic length of 910.1 cM. The individual LGs varied in length from 12.5 to 153.7 cM (mean 75.9 cM), containing between three and 23 (mean 11.8) loci. The mean inter-locus distance was 7.1 cM (range 3.7–19.3 cM). The placement of 19 SSR and two SNP loci allowed 11 of these LGs to be aligned with nine of the reference LGs (Table 3; Fig. 1A-C). Two pairs of “Est” LGs (VIIIa and VIIIb, Xa and Xb) each corresponded with a single consensus SSR-based one. The remaining LG was labeled “Est\_n.a.” (n.a.= not aligned) in Table 3 and Fig. 1C. Among the loci showing segregation distortion, 29 were successfully integrated within eight of the LGs. Two clusters of distorted loci were detected on Est\_Xb (one comprising two and the other three loci), along with one cluster on Est\_XII (ten loci) and a further one on Est\_XVII (three loci) (Fig. 1B-C).

The genes underlying the presence/absence of head spines (*Sp<sub>Head</sub>*), the presence/absence of leaf spines (*Sp<sub>Leaf</sub>*) were represented on the “Est” map (LG VIIIa), and the one responsible for head color (*ColorHead*) was located on LG Wild\_Va.

Ten of the 38 intercross markers (four SRAP, four AFLP and two SSR) were assignable to a specific LG in both maps (three each on IX and XII and four on XI), which aided the identification of homology between the “Wild” and “Est” LGs. Three other pairs (II, III and VIII) were identified as being homologous, based on their alignment with the reference SSR-based map (Tab. 3, Fig. 1 A-C). The distribution across 12 “Wild” and nine “Est” LGs of the 31 markers shared with the map generated by Sonnante et al. (2011) is also shown in Table 3.

## Discussion

A population arising from the cross between wild cardoon and a globe artichoke variety was generated with the view to construct a linkage map and locating the genes underlying the presence/absence of spines on the heads and leaf, and pigmentation of the heads. The high level of inbreeding depression experienced by *C. cardunculus* (Cravero *et al.* 2002) prevents either backcross, F<sub>2</sub> or recombinant inbred line populations being used for mapping quantitative traits, so instead the double pseudo-testcross approach was taken, as also successfully used by Lanteri *et al.* (2006), Portis *et al.* (2009) and Sonnante *et al.* (2011). Mapping population individuals obtained from other globe artichoke x wild cardoon crosses have been shown to vary widely with respect to both

quantitative and qualitative characters (Sonnante *et al.* 2011; Lanteri *et al.* 2012), confirming that a high level of heterozygosity has been retained in both wild and cultivated germplasm (Portis *et al.* 2005a, 2005b; Mauro *et al.* 2009, 2012).

The present linkage maps were based on a variety of molecular marker types. The choice of SRAP PCs reflected prior experience (Cravero *et al.* 2007) and, as SRAP genotyping generated most of the markers applied, these loci constituted the major backbone of the two maps. Since its development some ten years ago, SRAP genotyping has been deployed in a range of plant species for estimating levels of genetic diversity (Cravero *et al.* 2007; Esposito *et al.* 2007; Aneja *et al.* 2012), gene tagging (Martin *et al.* 2008; Zhang *et al.* 2011) and map construction (Lin *et al.* 2003; Sun *et al.* 2007; Xue *et al.* 2010). However, here we report its first usage for a genetic mapping in *C. cardunculus*.

A comparison of heterozygosity between the mapping parents used by Portis *et al.* (2009) showed that fewer loci were informative in the cultivated cardoon than in the globe artichoke variety “Romanesco C3”. An explanation for this difference reflected the fact that while the latter is propagated vegetatively, the former is a seed-propagated variety. Clonal propagation allows the maintenance of high heterozygosity levels, but seed propagation can introduce an element of purifying selection to stabilize production. In contrast, the parental lines used by Sonnante *et al.* (2011) (one a globe artichoke, the other a wild cardoon) produced a pair of maps in which the cultivated parent map include more loci than the wild parent one. In the present case, the wild cardoon-based map was based on a higher number of loci than the globe artichoke one (214 vs 141 markers) and was around 50% longer (1,465.5 cM vs 910.1 cM), implying that the level of heterozygosity in the wild cardoon parent is higher than in the domesticated type. Indeed, the latter was selected from an open pollinated population on the basis of some key commercial traits (Garcia *et al.* 2006), and it was expected that this selection would have reduced its global level of heterozygosity.

Segregation distortion is commonplace in mapping populations, including an intra-taxon one in *C. cardunculus* where it affected ~10% of the loci (Lanteri *et al.* 2006). Here, the segregation of some 19% of mapped loci was distorted, and this higher level may simply reflect the greater genetic separation between the parents of the population (Grandillo and Tanksley 1996; Verde *et al.* 2005). Clusters of distorted markers were detected in both the “Wild” and the “Est” maps. Since the direction of segregation bias within each cluster was unidirectional, the suggestion is that the basis of the phenomenon is biological, rather than it being due to scoring error or chance (Fishman *et al.* 2001).

The mapping of the three morphological traits considered was consistent with each being determined by a single gene. The spininess of the head trait has been known for some time to be determined by the gene *Sp*, where the dominant allele determines lack of spininess (Pochard *et al.* 1969; Basnizki and Zohary 1994). The wild cardoon parent used here was therefore presumably of genotype *spsp*, while that of “Est” was *Spsp*. The homozygous state prevented its location in the “Wild” map, but it was successfully located to LG VIIIa on the “Est” map. The gene underlying the presence/absence of leaf spines (*Sp<sub>Leaf</sub>*) appeared to be tightly linked to *Sp*. The presence of recombinants in the F<sub>1</sub> progeny indicates that this trait is therefore controlled by a locus distinct from *Sp*. The non spiny type is conditioned by the presence of at least one dominant allele (*Sp<sub>Leaf</sub><sup>-</sup>*), while the recessive homozygote *sp<sub>Leaf</sub>sp<sub>Leaf</sub>* produces spiny leaves. At the genetic level therefore, we propose that the wild cardoon accession was *spsp/sp<sub>Leaf</sub>sp<sub>Leaf</sub>* and “Est” was *Spsp/ Sp<sub>Leaf</sub>sp<sub>Leaf</sub>*. With respect to head color, Cravero *et al.* (2005) showed that within globe artichoke, two independent genes act epistatically to determine the trait; in the absence of dominant alleles at *P* the inflorescence bracts remain green (*pp*), while the genotype *P-U* develops purple-green bracts and *P-uu* uniformly purple bracts. The segregation for head color observed in the present mapping population was consistent with homozygosity composition in both parents at locus *P*, which determines the presence/absence of anthocyanic pigments (*pp* in “Wild” and *PP* in “Est”), while “Wild” was *Uu* and “Est” was *uu*; suggesting that the second locus, which determinates anthocyanin distribution, segregated. The *U* locus mapped on the maternal map to Wild\_Va.

The alignment and integration of independently generated linkage maps is desirable, but problematical where different marker types and/or population types are involved (Sun *et al.* 2007). Here, we relied on a set of 48 co-dominant markers for the alignment with the Portis *et al.* (2012) “Romanesco C3” x “Atilis 41” consensus map. This allowed 16 of the “Wild” and 11 of the “Est” LGs to be matched with 16 out of the 17 consensus LGs. Six of the reference LGs could be simultaneously aligned with both the “Wild” and “Est” LGs, leaving seven aligned only with the “Wild” map and four with only the “Est” map. The number of bridging markers per LG required for this exercise varied from one to four, and their linear order was mostly conserved. The exceptions related to three of the CELMS and four of the CyEM SSRs, which appeared to identify small inversions within LGs III, VIII and XV. Such minor differences may be due to mapping imprecisions (Lombard and Delourme 2001), to differences in recombination frequencies of marker pairs in different populations (Studer *et al.* 2010) or might be attributed to different population sizes (Spiller *et al.* 2011). Other potential sources of variation are genotyping errors, an excess of missing values and the mapping of distorted markers (Hackett and Broadfoot 2003). Given the probabilistic nature of genetic maps, the re-ordering of tightly linked

markers will generally produce a version of the map only marginally less probable than the most probable one, so this practice is quite frequently followed to improve alignment quality (Cervera *et al.* 2001; Jeuken *et al.* 2001; Lespinasse *et al.* 2000; Lombard and Delourme 2001; Sebastian *et al.* 2000). The further alignment of the “Wild” and “Est” LGs with 15 of the Sonnante *et al.* (2011) globe artichoke x wild cardoon LGs using 31 common markers evidenced the assignment of 14 SSR loci (ten CyEM, four CELMS) to a specific LG for the first time in a *C. cardunculus* map.

Given that the parents of the present mapping population differ widely with respect to a number of both qualitative and quantitative traits such as shape and weight of the heads; number of head per plant; height and shape of the plant; days to first harvest, opportunities are now available to investigate the inheritance of some important agronomic characteristics. Moreover, these are the first linkage maps of *C. cardunculus* developed from Argentinean genotypes of wild cardoon and globe artichoke. The enrichment of these maps with additional markers, like CAPS and SNPs recently developed by Scaglione *et al.* (2012a, 2012b), will improve their utility for the localization of major genes and quantitative trait loci.

### **Acknowledgments**

This research was supported by Agencia Nacional de Investigaciones Científicas y Técnicas (ANCYT, Argentina), Comisión Nacional de Ciencia y Tecnología (CONICET, Argentina) and by MIPAAF (Ministero delle Politiche Agricole, Alimentari e Forestali - Italy) through the “CYNERGIA” project .

## References

- Acquadro A, Portis E, Moglia A, Magurno F, Lanteri S (2006) Retrotransposon-based S-SAP as a platform for the analysis of genetic variation and linkage in globe artichoke. *Genome* 49:1149-1159
- Acquadro A, Lanteri S, Scaglione D, Arens P, Vosman B, Portis E (2009) Genetic mapping and annotation of genomic microsatellites isolated from globe artichoke. *Theor Appl Genet* 118(8):1573-1587
- Aneja B, Yadav N, Chawla V, Yadav R (2012) Sequence-related amplified polymorphism (SRAP) molecular marker system and its applications in crop improvement. *Mol Breeding* 30:1635-1648
- Basnitzki J, Zohary D (1994) Breeding of seed planted artichoke. *Plant Breed Rev* 12:253-269
- Bassam BJ, Caetano Anolles G, Gresshoff PM (1991) Fast and sensitive silver staining of DNA in polyacrylamide gels. *Analytical Biochemistry* 196:80-83
- Cervera MT, Storme V, Ivens B, Gusmao J, Liu BH, Hostyn V, Van Slycken J, Van Montagu M, Boerjan W (2001) Dense genetic linkage maps of three *Populus* species (*Populus deltoides*, *P-nigra* and *P-trichocarpa*) based on AFLP and microsatellite markers. *Genetics* 158:787-809
- Comino C, Lanteri S, Portis E, Acquadro A, Romani A, Hehn A, Larbat R, Bourgaud F (2007) Isolation and functional characterization of a cDNA coding a hydroxycinnamoyltransferase involved in phenylpropanoid biosynthesis in *Cynara cardunculus* L. *BMC Plant Biol* 7:14
- Comino C, Hehn A, Moglia A, Menin B, Bourgaud F, Lanteri S, Portis E (2009) The isolation and mapping of a novel hydroxycinnamoyltransferase in the globe artichoke chlorogenic acid pathway. *BMC Plant Biol* 9:30
- Cravero V, Lopez Anido FS, Cointry EL (2002) Efecto de una generación de endocría sobre caracteres vegetativos y productivos en dos estructuras familiares de alcaucil. *Horticultura Brasileira*, 20(3):459-464
- Cravero V, Picardi L, Cointry E (2005) An approach for understanding the heredity of two quality traits (head color and tightness) in globe artichoke (*Cynara scolymus* L.). *Genet Mol Biol* 28:431-434
- Cravero V, Martin E, Cointry E (2007) Genetic Diversity in *Cynara cardunculus* determined by Sequence-related Amplified Polymorphism markers. *J Amer Soc Hort Sci* 132(2):1-5
- Cravero V, Martin E, Lopez Anido F, Cointry E (2010) Stability through years in a non-balanced trial of globe artichoke varietal types. *Scientia Horticulturae* 126:73-79
- Esposito MA, Martin EA, Cravero VP, Cointry E (2007) Characterization of pea accessions by SRAP's markers. *SciHortic* 113:329-335
- Fishman L, Kelly AJ, Morgan E, Willis JH (2001) A genetic map in the *Mimulus guttatus* species complex reveals transmission ratio distortion due to heterospecific interactions. *Genetics* 159:1701-1716
- Garcia SM, Cointry EL, Lopez Anido FS, Firpo IT, Cravero V, Muñoz S (2006) Nueva generación de alcaucil multiplicable por semilla. *Horticultura Argentina* 25(59):152
- Grandillo S, Tanksley SD (1996) Genetic analysis of RFLPs, GATA microsatellites and RAPDs in a cross between *L. esculentum* and *L. pimpinellifolium*. *Theor Appl Genet* 92:957-965
- Grattapaglia D, Sederoff R (1994) Genetic linkage maps of *Eucalyptus grandis* and *Eucalyptus urophylla* using a pseudotest-cross: mapping strategy and RAPD markers. *Genetics* 137:1121-1137
- Hackett CA, Broadfoot LB (2003) Effects of genotyping errors, missing values and segregation distortion in molecular marker data on the construction of linkage maps. *Heredity* 90:33-38
- Jackson JA, Matthews D (2000) Modified inter-simple sequence repeat PCR protocol for use in conjunction with the LI-COR gene ImagIR(2) DNA analyzer. *Biotechniques* 28:914-916
- Jeuken M, Van Wijk R, Peleman J, Lindhout, P (2001) An integrated interspecific AFLP map of lettuce (*Lactuca*) based on two *L. sativa* x *L. saligna* F2 populations. *Theor Appl Genet* 103:638-647
- Kosambi D (1994) The estimation of map distances from recombination values. *Ann Eugen* 12:171-175
- Lanteri S, Saba E, Cadinu M, Mallica G, Baghino L, Portis E (2004) Amplified fragment length polymorphism for genetic diversity assessment in globe artichoke. *Theor Appl Genet* 108:1534-1544

- Lanteri S, Acquadro A, Comino C, Mauro R, Mauromicale G, Portis E (2006) A first linkage map of globe artichoke (*Cynara cardunculus* var. *scolymus* L.) based on AFLP, S-SAP, M-AFLP and microsatellite markers. *Theor Appl Genet* 112:1532–1542
- Lanteri S, Portis E, Acquadro A, Mauro RP, Mauromicale G (2012) Morphology and SSR fingerprinting of newly developed *Cynara cardunculus* genotypes exploitable as ornamentals. *Euphytica* 184:311–321
- Lespinasse D, Rodier-Goud M, Grivet L, Leconte A, Legnate H, Seguin M (2000) A saturated genetic linkage map of rubber tree (*Hevea* spp.) based on RFLP, AFLP, microsatellite, and isozyme markers. *Theor Appl Genet* 100:127-138
- Li, G. and C. Quiros. 2001. Sequence-related amplified polymorphism (SRAP), a new marker system based on a simple PCR reaction: Its application to mapping and gene tagging in Brassica. *Theor Appl Genet* 103:455–461
- Lin Z, Zhang X, Yichun N, Daohua H, Maoqing W (2003) Construction of a genetic linkage map for cotton based on SRAP. *Chinese Science Bulletin*, 48(19):2064-2068
- Lombard V, Delourne R (2001) A consensus linkage map for rapessed (*Brassica napus* L.): construction and integration of three individual maps from DH populations. *Theor Appl Genet* 103:491-507
- Lopez Anido F, Firpo I, Garcia S, Cointry E (1998) Estimation of genetic parameters for yield traits in globe artichoke (*Cynara scolymus* L.). *Euphytica* 103:61–66
- Lopez Anido F, Cravero V, Martin E, Crippa I, Cointry, E (2010) Differential expression of the genetic variance in F<sub>2</sub> populations of reciprocal crosses of artichoke. *Span J Agric Res* 8(3):679-685
- Martin E, Cravero V, Esposito M, Lopez anido F, Milanese L, Cointry E (2008) Identification of markers linked to agronomic traits in globe artichoke. *Australian Journal of Crop Science* 1(2):43-46
- Mauro R, Portis E, Acquadro A, Lombardo S, Mauromicale G, Lanteri S (2009) Genetic diversity of globe artichoke landraces from Sicilian small-holdings: implications for evolution and domestication of the species. *Conserv Genet* 10:431-440
- Mauro R, Portis E, Lanteri S, Mauromicale G (2012) Genotypic and bio-agronomical characterization of an early Sicilian landraces of globe artichoke. *Euphytica* 186:357-366
- Mauromicale G, Morello N, Santoiemma G, Ierna A (2000) Nuove varietà per migliorare la cinaricoltura siciliana. *Inf Agr* 26:47-51
- Menin B, Comino C, Moglia A, Dolzhenko Y, Portis E, Lanteri S (2010) Identification and mapping of genes related to caffeoylquinic acid synthesis in *Cynara cardunculus* L. *Plant Sci* 178:338–347
- Moglia A, Comino C, Portis E, Acquadro A, De Vos R, Beekwilder J, Lanteri S (2009) Isolation and mapping of a C3'H gene (CYP98A49) from globe artichoke, and its expression upon UV-C stress. *Plant Cell Rep* 28:963-974
- Pecaut P (1993) Globe artichoke *Cynara scolymus* L. In: Kallo, Berg (ed) Genetic improvement of vegetables crops. Pergamon, Oxford, pp 737-746
- Pochard E, Foury C, Chambonet D (1969) Il miglioramento genetico del carciofo. Proceedings the 1 Congresso Internazionale sul carciofo. Bari, Italy, pp 117-155
- Portis E, Acquadro A, Comino C, Mauromicale G, Saba E, Lanteri S (2005a) Genetic structure of island populations of wild cardoon [*Cynara cardunculus* L. var. *sylvestris* (Lamk) Fiori] detected by AFLPs and SSRs. *Plant Science* 169:199-210
- Portis E, Barchi L, Acquadro A, Macua J, Lanteri S (2005b) Genetic diversity assessment in cultivated cardoon by AFLP (amplified fragment length polymorphism) and microsatellite markers. *Plant Breeding* 124:299-304
- Portis E, Mauromicale G, Mauro R, Acquadro A, Scaglione D, Lanteri S (2009) Construction of a reference molecular linkage map of globe artichoke (*Cynara cardunculus* var. *scolymus*). *Theor Appl Genet* 120(1):59-70
- Portis E, Scaglione D, Acquadro A, Mauromicale G, Mauro R, Knapp S, Lanteri S (2012) Genetic mapping and identification of QTL for earliness in the globe artichoke/cultivated cardoon complex. *BMC Research Notes* 5:252

- Rottenberg A, Zohary D (1996) The wild ancestry of the cultivated artichoke. *Genet Resour Crop Evol* 43:53–58
- Scaglione D, Acquadro A, Portis E, Taylor CA, Lanteri S, Knapp SJ (2009) Ontology and diversity of transcript-associated microsatellites mined from globe artichoke EST database. *BMC Genomics* 10:454
- Scaglione D, Acquadro A, Portis E, Tirone M, Knapp SL, Lanteri S (2012a) RAD tag sequencing as a source of SNP markers in *Cynara cardunculus* L. *BMC Genomics* 13:3
- Scaglione D, Lanteri S, Acquadro A, Lai Z, Knapp SJ, Rieseberg L, Portis E (2012b) Large-scale transcriptome characterization and mass discovery of SNPs in globe artichoke and its related taxa. *Plant Biotechnology Journal* 10: 956-69
- Sebastian RL, Howell EC, King GJ, Marshall DF, Kearsey MJ (2000) An integrated AFLP and RFLP *Brassica oleracea* linkage map from two morphologically distinct doubled-haploid mapping populations. *Theor Appl Genet* 100:75-81
- Sonnante G, Gatto A, Morgese A, Montemurro F, Sarli G, Blanco E, Pignone D (2011) Genetic map of artichoke x wild cardoon: toward a consensus map for *Cynara cardunculus*. *Theor Appl Genet* 123(7):1215–1229
- Spiller M, Linde M, Hibrand-Saint Oyant L, Tsai C, Byrne D, Smulders M, Foucher F, Debener T (2011) Towards a unified genetic linkage map for diploid roses. *Theor Appl Genet* 122:489-500
- Studer B, Kölliker R, Muylle H, Asp T, Frei U, Roldán-Ruiz I, Barre P, Tomaszewski C, Meally H, Barth S, Skot L, Armstead I, Dolstra O, Lübberstedt T (2010) EST-derived SSR markers used as anchor loci for the construction of a consensus linkage map in ryegrass (*Lolium* spp.). *BMC Plant Biology* 10:177
- Sun Z, Wang Z, Tu J, Zhang J, Yu F, McVetty P, Li G (2007) An ultradense genetic recombination map for *Brassica napus*, consisting of 13551 SRAP markers. *Theor Appl Genet* 114:1305-1317
- Van Ooijen JW (2006) JoinMap® v.4, Software for the calculation of genetic linkage maps in experimental populations. Wageningen, Netherlands: Kyazma BV
- Verde I, Lauria M, Dettori M, Vendramin E, Balconi C, Micali S, Wang Y, Marrazzo MT, Cipriani G, Hartings H, Testolin R, Abbott AG, Motto M, Quarta R (2005) Microsatellite and AFLP markers in the *Prunus persica* [L. (Batsch)] x *P-ferganensis* BC(1) linkage map: saturation and coverage improvement. *Theor Appl Genet* 111:1013-1021
- Voorrips R (2002) MapChart: Software for the graphical presentation of linkage maps and QTLs. *Journal of Heredity* 93(1):77–78.
- Vos P, Hogers R, Bleeker M, Reijans M, Van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 23:4407–4414
- Weeden N (1994) Approaches to mapping in horticultural crops. Gressho P (ed) *Plant genome analysis*. CRC Press, Boca Raton, pp 57–68
- Xue D, Feng S, Zhao H, Jiang H, Shen B, Shi N, Lu J, Liu J, Wang H (2010) The linkage maps of *Dendrobium* species based on RAPD and SRAP markers. *J Genetics Genomics* 37:197–204
- Ye S, Dhillon S, Ke X, Collins A, Day I (2001) An efficient procedure for genotyping single nucleotide polymorphisms. *Nucleic Acid Res* 29:E88-8
- Young ND (1999) A cautiously optimistic vision for marker-assisted breeding. *Molecular Breeding* 5: 505-510.
- Zamir D (2001) Improving plant breeding with exotic libraries. *Nat Rev Genet* 2:983–989
- Zhang F, Chen S, Chen F, Fang W, Deng Y, Chang Q, Liu P (2011) Genetic analysis and associated SRAP markers for flowering traits of chrysanthemum (*Chrysanthemum morifolium*). *Euphytica* 177:15–24

## Tables

**Table 1** SRAP primer sequences used for genotyping.

<b>Forward Primer</b>		<b>Reverse Primer</b>	
<b>Me1</b>	5'-TGAGTCCAAACCGGATA-3'	<b>Em1</b>	5'-GACTGCGTACGAATTAAT-3'
<b>Me2</b>	5'-TGAGTCCAAACCGGAGC-3'	<b>Em2</b>	5'-GACTGCGTACGAATTTGC-3'
<b>Me3</b>	5'-TGAGTCCAAACCGGAAT-3'	<b>Em3</b>	5'-GACTGCGTACGAATTGAC-3'
<b>Me4</b>	5'-TGAGTCCAAACCGGACC-3'	<b>Em4</b>	5'-GACTGCGTACGAATTTGA-3'
<b>Me5</b>	5'-TGAGTCCAAACCGGAAG-3'	<b>Em5</b>	5'-GACTGCGTACGAATTAAC-3'

**Table 2** AFLP primer combinations used for genotyping.

<b>EcoRI/TaqI template</b>	
<b>Primer combination</b>	<b>Code</b>
E + ACC/T + TTT	1.800
E + ACG/T + TTT	1.700
E + ACC/T + TAC	2.800
E + ACG/T + TAC	2.700
E + ACC/T + TGT	3.800
E + ACG/T + TGT	3.700
E + ACC/T + TAG	4.800
E + ACG/T + TAG	4.700
E + ACC/T + TAT	5.800
E + ACG/T + TAT	5.700
E + ACC/T + TCC	6.800
E + ACG/T + TCC	6.700
E + ACC/T + TAA	7.800
E + ACG/T + TAA	7.700
E + ACC/T + TTA	8.800
E + ACG/T + TTA	8.700


**Table 3** Key characteristics of the wild cardoon (“Wild”) and globe artichoke (“Est”) linkage maps and their alignment.

LG name		Size (cM) Wild	Size (cM) Est	Total markers	Marker density	Portis <i>et al.</i> 2012		Sonnante <i>et al.</i> 2011	
Wild cardoon	Estrella del Sur					Aligned LGs	Shared markers	Aligned LGs	Shared markers
Wild_Ia		56,9		8	8,1	I	2	I	2
Wild_Ib		41,8		6	8,4		1		0
Wild_II		75,3		7	12,6	II	3	II	2
	Est_II		96,5	6	19,3		4		1
Wild_III		120,4		15	8,6	III	3	III	1
	Est_III		153,7	14	11,8		4		0
	Est_IV		106,2	23	4,8	IV	2	VI	2
Wild_Va		119,4		17	7,5	V	2	V	2
Wild_Vb		93,4		10	10,4		1	X	1
Wild_VI		119,1		10	13,2	VI	2		0
Wild_VII		105,7		16	7,0	VII	2	XVIII	2
Wild_VIII		153,1		34	4,6	VIII	2	XV	1
	Est_VIIIa		120,3	23	5,5		3	VIII	1
	Est_VIIIb		18,5	4	6,2		1	VIII	2
Wild_IX		59,9		5	15,0	IX	2	IX	2
	Est_IX		31,4	4	10,5		2		2
	Est_Xa		12,4	3	6,2	X	3	XI	1
	Est_Xb		70,6	20	3,7		1		1
Wild_XI		108,4		10	12,0	XI	2		0
	Est_XI		36,1	9	4,5		0		0
Wild_XII		67,4		23	3,1	XII	1	IV	1
	Est_XII		92,8	18	5,5		0		1
Wild_XIII		125,8		16	8,4	XIII	1	XIV	1
Wild_XIV		136,4		26	5,5	XIV	1	XIII	1
Wild_XVa		52,4		6	10,5	XV	1	XVI	1
Wild_XVb		30,1		5	7,5		3		3
	Est_XVII		121,1	12	11,0	XVII	1		0
	Est_n.a.		50,5	5	12,6		0		0


## Figure caption

**Figure 1 (A, B, C)** Genetic linkage maps of wild cardoon (blue LGs, shown on the left) and globe artichoke (yellow LGs, on the right) aligned with the reference SSR-based consensus map (green LGs in the center). Shared SSR and SNP loci are shown in bold, and their positions are connected by a line. Marker names are shown to the right of each LG, with map distances (in cM) to the left. Markers showing significant levels of segregation distortion are indicated by asterisks (\*:  $0.1 > P < 0.05$ , \*\*:  $0.05 > P < 0.01$ , \*\*\*  $P > 0.01$ ).


### Wild\_la




### SSR-Ref LGI




### Wild\_lb




### Wild\_III




### SSR-Ref LGIII




### Est\_III




### Wild\_Va




### SSR-Ref LGV




### Wild\_Vb




### Wild\_II




### SSR-Ref LGII




### Est\_II









### SSR-Ref LGIV




### Est\_IV







### Wild\_XII




### SSR-Ref LGXII




### Est\_XII




### Wild\_XIII




### SSR-Ref LGXIII




### Wild\_XIV




### SSR-Ref LGXIV




### Wild\_XVa




### SSR-Ref LGXV




### Wild\_XVb



### SSR-Ref LGXVII



### Est\_XVII



### Est\_n.a.

