

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Impact of control strategies on Thrips tabaci and its predator Aeolothrips intermedius on onion crops

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/138620> since 2016-01-07T17:53:19Z

Published version:

DOI:10.1007/s12600-013-0335-8

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Mautino G.C., Bosco L., Tavella L.
Impact of control strategies on *Thrips tabaci* and its predator *Aeolothrips*
intermedius on onion crops
PHYTOPARASITICA (2014) 42 (1)
DOI: 10.1007/s12600-013-0335-8

The definitive version is available at:

<http://link.springer.com/content/pdf/10.1007/s12600-013-0335-8>

1 **Impact of control strategies on *Thrips tabaci* and its predator *Aeolothrips intermedius* on onion**
2 **crops**

3

4 Giulia C. Mautino – Lara Bosco – Luciana Tavella*

5

6 Dipartimento di Scienze Agrarie, Forestali e Alimentari (DISAFA), University of Torino, 10095

7 Grugliasco (TO), Italy

8

9 *Corresponding author e-mail: luciana.tavella@unito.it

10

11

12 Received: 29 March 2013 / Accepted: 4 July 2013

13

14

15 **Abstract**

16 *Thrips tabaci* (Thysanoptera: Thripidae) is a major pest of onion worldwide. In 2011, research was
17 conducted in a commercial onion field in northwestern Italy to: (i) assess the presence of
18 autochthonous onion thrips predators on the crop; and (ii) evaluate the impact of the commonly
19 used insecticides and alternative pest management strategies on onion thrips and its autochthonous
20 predators. Toxicity of the active ingredients on local populations of onion thrips and its predatory
21 thrips was also evaluated in laboratory bioassays. During field surveys, the highest and lowest
22 thrips infestations were observed in plots treated with lambda-cyhalothrin and spinosad,
23 respectively. The effectiveness of spinosad on *T. tabaci* was also confirmed in laboratory bioassays.
24 The dominant zoophagous species *Aeolothrips intermedius* (Thysanoptera: Aeolothripidae) was
25 more adversely affected by treatment with lambda-cyhalothrin, confirmed by a decrease in
26 predator/prey ratios. The use of spinosad and acibenzolar-S-methyl is suggested as an alternative to
27 conventional insecticides for the preservation of *A. intermedius*, which proved to be a potential
28 biological control agent of *T. tabaci*.

29

30 **Keywords** Acibenzolar-S-methyl – Lambda-cyhalothrin – Onion thrips – Predator thrips –
31 Spinosad – Thysanoptera –

32

33

34 **Introduction**

35 *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) is one of the most serious pests of onion (*Allium*
36 *cepa* L.) and other *Allium* spp. in many parts of the world (MacIntyre Allen *et al.* 2005; Martin *et*
37 *al.* 2003; Nault & Shelton 2010). Moreover, *T. tabaci* is the vector of *Iris yellow spot virus* (IYSV),
38 genus *Tospovirus* (Bunyaviridae), a severe and widespread disease infecting onion, leek, iris and
39 wild *Allium* species (Gent *et al.* 2006; Nagata *et al.* 1999; Pappu *et al.* 2009). Chemical treatments
40 are the main method used by onion growers for thrips control. However, in Italy there is a restricted
41 range of authorized products, predominantly pyrethroids (*e.g.* lambda-cyhalothrin), to which onion
42 thrips can develop high levels of resistance (Martin *et al.* 2003; Shelton *et al.* 2006). Additionally,
43 some of these insecticides have side effects, affecting biological control agents either directly (*i.e.*,
44 physiological or behavioral effects) or indirectly (*e.g.* habitat destruction, oviposition, resting,
45 mating sites) (Desneux *et al.* 2007). Therefore, it is important to develop new reduced-risk
46 insecticides to use in Integrated Pest Management (IPM) programs with minimal impact on
47 beneficial arthropods (Jones *et al.* 2005). In this regard, spinosad is known to be effective against *T.*
48 *tabaci* (Gent *et al.* 2006; Shelton *et al.* 2006; Yarahmadi *et al.* 2009), with a low to moderate
49 toxicity to thrips' common natural enemies (Jones *et al.* 2005; Ludwig & Oetting 2001; Workman
50 & Martin 2002).

51 Pesticide impact on human and environmental health begs for alternative pest management
52 approaches, such as the use of straw mulch (Gent *et al.* 2006; Larentzaki *et al.* 2008; Momol *et al.*
53 2004), of intercrops (Trdan *et al.* 2006), and of thrips-resistant onion cultivars (Diaz-Montano *et al.*
54 2010), or the evaluation of new bio-insecticides (Patil *et al.* 2009) and biologically active plant
55 volatiles (Koschier *et al.* 2002) against *T. tabaci*. Moreover, activators of natural systemic acquired
56 resistance (SAR), such as acibenzolar-S-methyl, appear promising for reducing the use of
57 conventional insecticides (Mautino *et al.* 2012). Acibenzolar-S-methyl was used as an effective
58 alternative to many bactericides and fungicides (Gent *et al.* 2006; Momol *et al.* 2004; Pappu *et al.*

59 2000), and for the control of tospoviruses and their thrips vectors (Gent *et al* 2006; Momol *et al.*
60 2004).

61 Furthermore, the conservation and augmentation of predators feeding on thrips are important
62 control strategies. Different generalist predators are known to be effective biocontrol agents, and
63 they are now commercially available. These predators include anthocorids of the genus *Orius*
64 (Hemiptera: Anthocoridae) (Bosco *et al.* 2008; Funderburk *et al.* 2000; Riudavets 1995), mirids of
65 the genera *Dicyphus* and *Macrolophus* (Hemiptera: Miridae) (Gabarra *et al.* 1995; Riudavets *et al.*
66 1993), and mites of the genus *Amblyseius* (Acarina: Phytoseiidae) (Wimmer *et al.* 2008). There are
67 comparatively few data on the predatory efficiency of the autochthonous thrips species belonging to
68 the genera *Aeolothrips*, *Haplothrips* and *Franklinothrips* (Thysanoptera: Aeolothripidae), that are
69 potential biological control agents of *T. tabaci* (Cox *et al.* 2006; Fathi *et al.* 2008; Kakimoto *et al.*
70 2006; Trdan *et al.* 2005a).

71 Integration of chemical, biological, agronomic and physical control options is increasingly
72 necessary in order to maintain pest populations below economic damage thresholds (Cox *et al.*
73 2006), particularly upon consideration of the weak efficacy of pyrethroids (especially lambda-
74 cyhalothrin) that are widely used on onion (Herron *et al.* 2008; MacIntyre Allen *et al.* 2005;
75 Mautino *et al.* 2012). Moreover, the possible impact on beneficial arthropods should be carefully
76 monitored to preserve the presence and abundance of the natural populations in the field.
77 Independent of the type of insecticide employed, chemical control still remains the primary
78 strategy for pest control on onion crops; therefore, supervised control based on action thresholds or
79 tolerance levels consistent with IPM programs is always recommended for growers (Nault &
80 Shelton 2010).

81 Consequently, this study focused on: (i) assessing the entity of autochthonous predators of *T.*
82 *tabaci* on the onion crop; and (ii) evaluating the impact of the commonly used insecticides and
83 alternative pest management strategies on onion thrips and its autochthonous predators.

84

85 **Materials and methods**

86 *Field experiments* Research was conducted in 2011, in a commercial onion field of
87 approximately 8 ha located in Castellazzo Bormida (province of Alessandria, Piedmont, 44°50'45"
88 N, 8°34'41" E, 105 m a.s.l.). The experimental site was flanked on all sides by at least 2 m of
89 insecticide-free onions within the grower's field. Experimental plots (10.5 m² each) consisted of six
90 onion rows 7 m in length, with rows spaced every 0.2 m. Plots were separated within rows by 1.0
91 m and spaced every 1.5 m. Onions of the golden onion cultivar 'Derek' were seeded in April;
92 diseases and weeds were controlled using products recommended for onion production
93 (pendimethanil, ioxynil, metalaxyl-M + copper, dimethomorph + pyraclostrobin, cyprodinil +
94 fludioxonil).

95 The trial was arranged in a randomized complete block design with four replicates for each
96 of six treatments (24 plots). The treatments consisted of one untreated control (T1); one treatment
97 based on four repeated pyrethroid applications (T2); one treatment based on one pyrethroid
98 application at the action threshold (*i.e.*, when mean number of thrips sampled by plant beating
99 exceeded two thrips per plant) (T3); one treatment based on two spinosad applications (T4); one
100 treatment based on four applications of the plant activator acibenzolar-S-methyl (T5); one treatment
101 based on two acibenzolar-S-methyl applications, followed by two spinosad applications (T6) (Table
102 1). The active ingredients and formulations that we tested were: acibenzolar-S-methyl 500 g a.i. kg⁻¹
103 (Bion[®] 50 WG, Syngenta Crop Protection, Milano, Italy); lambda-cyhalothrin 15 g a.i. l⁻¹ (Karate[®]
104 Zeon 1.5, Syngenta Crop Protection); spinosad 120 g a.i. l⁻¹ (Success[®] SC, Dow AgroSciences,
105 Milano, Italy). Chemicals were applied at the manufacturer's recommended field rates with a
106 precision shoulder sprayer, using 600 l of solution ha⁻¹ of onion crop and producing a fine mist to
107 ensure effective coverage. The delivery pressure at the nozzle was 300 kPa. Rate and timing of
108 applications are listed in Table 1.

109 Commencing in May, onion plots were surveyed weekly for the presence of *T. tabaci* and
110 their predators. At the first occurrence of onion thrips infestation on the crop (pre-sampling, *i.e.*,

111 pre-S, June 15), insecticide applications were sprayed, in relation to the treatment, on June 19, June
112 29, July 9 and July 19. Sampling was carried out 3 days after each cluster of sprays (*i.e.*, S1, June
113 23; S2, July 4; S3, July 13; S4, July 22). *Thrips tabaci*, thrips predators including anthocorid, mirid
114 bugs (Hemiptera: Anthocoridae, Miridae), and predatory thrips (Thysanoptera: Aeolothripidae)
115 were detected during field surveys. Plant beating was chosen as the sampling method for its high
116 feasibility, and because a strong relationship between this method and visual inspection has
117 previously been observed, and also for the larval population, generally underestimated with the
118 beating sampling method (Mautino *et al.* 2012). Five plants were randomly selected at three points
119 in each plot (15 plants per plot), and these were beaten over a plastic tray (350 × 250 mm). Thrips
120 adults and larvae, and their predators, were counted, collected with a mouth aspirator and
121 transferred to the laboratory. Subsequently, adult onion thrips were observed under a
122 stereomicroscope at 160× magnification and identified to the species level according to Mound *et*
123 *al.* (1976). For predatory thrips, 10% to 20% of total adults sampled on each sampling date, and in
124 each treatment, were mounted on microscope slides and identified under a compound microscope
125 according to Schliephake & Klimt (1979). Thrips larvae were observed under a stereomicroscope at
126 160× magnification and attributed to the family Aeolothripidae or Thripidae, according to
127 Vierbergen *et al.* (2010).

128 Data on local weather conditions during field experiments were provided by Rete
129 Agrometeorologica, Regione Piemonte, Settore Fitosanitario. In particular, the following average
130 temperatures were recorded: T mean 18.7°C, T max 27.4°C, T min 10.8°C in May; T mean 20.8°C,
131 T max 27.3°C, T min 15.5°C in June; T mean 22.2°C, T max 29.1°C, T min 16.1°C in July.
132 Rainfall was 34.2, 102.6 and 33.8 mm in May, June and July, respectively.

133 *Laboratory bioassays* Adults of *T. tabaci* and predatory thrips collected in untreated plots on
134 July 13 and July 22 (S3 and S4) were tested in laboratory bioassays. Field-collected thrips were
135 temporarily transferred to 1 l gauze-covered glass jars (approximately 200 thrips per jar), with
136 corrugated cardboard on the bottom to provide pupation sites, and paper to avoid humidity. To

137 provide food sources and oviposition sites, jars were supplied with pollen and green bean pods
138 [*Phaseolus vulgaris* L. (Fabaceae)] for *T. tabaci*, and with leek leaves [*Allium porrum* L.
139 (Alliaceae)], previously infested by *T. tabaci* (providing live prey), for predatory thrips. Mass
140 rearing was conducted in growth chambers at $25 \pm 1^\circ\text{C}$, $65 \pm 5\%$ r.h. and a 16h:8h L:D cycle
141 (Tedeschi *et al.* 2001).

142 The toxicity of tested products was evaluated on *T. tabaci* and predatory thrips using the vial
143 bioassay method described by Mautino *et al.* (2012), which is an adaptation of the thrips
144 insecticides bioassay system (TIBS) described by Rueda & Shelton (2003). Thrips were collected
145 from rearing jars and placed into a plastic microcentrifuge tube previously treated with the product
146 being tested; the tube cap contained a small well with 10% sugar-water solution. The solution was
147 sealed into the well with a small piece of stretched parafilm through which thrips could feed on the
148 sugar solution. The tube, but not the cap, was treated with the product (or water for the untreated
149 control), and after 4 h the chemical solution (or water) was poured out and the tube was allowed to
150 dry overnight. Specifically, ten *T. tabaci* females and five predatory thrips females were introduced
151 separately into each treated tube. Five replicates were used for each of the four treatments: untreated
152 control, lambda-cyhalothrin, spinosad and acibenzolar-S-methyl. The active ingredients were used
153 in the same formulations and doses as those adopted in the field experiment. Acibenzolar-S-methyl
154 is a plant defense activator and it is supposed to have no effect by contact or fumigation on insects;
155 nonetheless, it was tested in the vial bioassay to evaluate a potential side effect (*e.g.* presence of
156 adjuvant compounds) on thrips, even in the absence of the plant. Thrips survival was assessed after
157 24 h and 48 h with the use of a stereomicroscope: thrips which did not move after 2 min of
158 observation were considered dead. The vial bioassay was replicated three times for both onion
159 thrips and predatory thrips.

160 *Statistical analyses* For the field data, the mean numbers of total (adults plus larvae) *T. tabaci* and
161 predatory thrips per plant were log-transformed to achieve homogeneity of variance (Levene) and

162 normality (Shapiro-Wilk), and analyzed by Univariate Analysis of Variance (ANOVA) for
163 randomized blocks (treatments and blocks were the factors).

164 To describe the effect of treatments on the relationship between phytophagous and predatory
165 thrips, the predator/prey ratios were calculated from the mean number of total thrips per plant for
166 each sampling date. Ratio values were log-transformed to achieve homogeneity of variance
167 (Levene) and normality (Shapiro-Wilk), and analyzed by ANOVA and Tukey's *post hoc* test.

168 Percentage survival data obtained in the laboratory bioassays for predatory thrips and *T.*
169 *tabaci* were separately transformed to arcsine square-root values before analysis; the non-
170 parametric Kruskal-Wallis was chosen since data were non-homogeneous, and means were
171 compared using the Mann-Whitney U test. For each treatment, differences between the survival
172 data of *T. tabaci* and predatory thrips were analyzed by ANOVA after tests of homogeneity of
173 variance (Levene) and normality (Shapiro-Wilk).

174 All statistical analyses were performed using SPSS statistical package (version 17.0; SPSS,
175 Chicago, IL, USA).

176

177 **Results**

178 Field experiments

179 *Treatment efficacy against onion thrips* *Thrips tabaci* was the dominant phytophagous species
180 collected on onions, with populations composed of both adults and larval stages. Over 96% of the
181 total sampled adult thrips (n = 2,787) belonged to this species, while the remaining 23 and 79 adult
182 thrips belonged to *Frankliniella intonsa* Trybom and to *Haplothrips* spp., respectively. Overall,
183 larval stages were 13% of the total thrips sampled by plant beating (n = 3,208); among the
184 treatments, percentages of Thripidae larvae varied from 7.6% in T6 to 19.0% in T2. Moreover, the
185 average percentages of larvae varied throughout the growing season: 1.6%, 6.0%, 31.6%, 6.7%, and
186 20.8% of total specimens sampled on June 15, June 23, July 4, July 13 and July 22, respectively.

187 The mean numbers of total (adults plus larvae) onion thrips collected in the plots of each
188 treatment are reported in Table 2. On June 15 (Pre-S), before the first chemical application, mean
189 numbers per plant beating ranged between 1.0 and 1.5 thrips without any significant differences
190 between the treatments (ANOVA: $df = 5, 63; F = 1.182; P = 0.327; n = 12$). On June 23 (S1), after
191 the first chemical application of lambda-cyhalothrin (T2) and acibenzolar-S-methyl (T5 and T6),
192 independently of the product, numbers of *T. tabaci* per plant in the sprayed plots (*i.e.*, T2, T5, T6)
193 were significantly lower than in the unsprayed plots (*i.e.*, T1, T3, T4) (ANOVA: $df = 5, 63; F =$
194 $20.438; P < 0.001; n = 12$). On July 4 (S2), thrips populations naturally decreased in the untreated
195 control (T1) and in all of the unsprayed plots (T3, T4), whereas in the sprayed plots (T2, T5, T6)
196 populations increased despite the second chemical application of lambda-cyhalothrin (T2) and
197 acibenzolar-S-methyl (T5 and T6). Nevertheless, on this sampling date no significant differences
198 between the treatments were recorded (ANOVA: $df = 5, 63; F = 1.347; P = 0.256; n = 12$) (Table
199 2). On July 13 (S3) in treatments T3, T5 and T6, and on July 22 (S4) in the other treatments, thrips
200 populations started to increase and reached maximum infestation levels. On July 13 (S3), the
201 maximum numbers of *T. tabaci* per plant beating were recorded after the third application of
202 acibenzolar-S-methyl in treatment T5; mean numbers were significantly higher than in the untreated
203 control (T1), and in the treatment sprayed with spinosad (T4) (ANOVA: $df = 5, 63; F = 5.512; P <$
204 $0.001; n = 12$). In particular, the first application of spinosad (T4) maintained infestation levels at
205 the lowest values recorded in all treatments (Table 2). On July 22 (S4), at the end of the growing
206 season, thrips populations generally reached maximum values. In treatment T2, after the fourth
207 application of lambda-cyhalothrin, mean numbers of *T. tabaci* per plant beating were significantly
208 higher than in all other treatments, except T5 where acibenzolar-S-methyl was applied for the fourth
209 time (ANOVA: $df = 5, 63; F = 14.438; P < 0.001; n = 12$). Conversely, both the single application
210 of lambda-cyhalothrin (T3) and the second application of spinosad, after acibenzolar-S-methyl (T6),
211 maintained thrips populations at the lowest values recorded in all treatments, reducing infestation
212 levels observed on the previous sampling date (Table 2).

213 *Side effects on predatory thrips* *Aeolothrips intermedius* Bagnall (Thysanoptera: Aeolothripidae)
214 was the dominant zoophagous species collected on onions, with populations composed of both
215 adults (females and males) and larval stages. Overall, 1,492 adult predatory thrips were sampled by
216 plant beating, and all the identified adults ($n = 230$) belonged to this species. Larval stages were 6%
217 of the total predatory thrips sampled by plant beating ($n = 1,595$); among the treatments,
218 percentages of Aeolothripidae larvae varied in the treatments from 3.7% (T1) to 9.5% (T2).
219 Moreover, average percentages were variable throughout the growing season: 2.5%, 3.3%, 23.0%,
220 13.0%, and 3.7% of total specimens sampled on June 15, June 23, July 4, July 13 and July 22,
221 respectively.

222 The mean numbers of *A. intermedius* (adults plus larvae) collected in the plots of each
223 treatment, on five sampling dates, are reported in Table 3. On June 15 (Pre-S), before the first
224 chemical application, mean numbers per plant beating ranged between 1.4 and 1.9 predatory thrips
225 without any significant differences between the treatments (ANOVA: $df = 5, 63$; $F = 0.644$; $P =$
226 0.667 ; $n = 12$). On June 23 (S1), the predatory thrips population increased and reached maximum
227 levels in the unsprayed plots (*i.e.*, T1, T3, T4); the maximum mean number per plant beating (2.3
228 thrips) was observed in treatment T1 (Table 3). In contrast, after the first chemical application of
229 lambda-cyhalothrin (T2) and acibenzolar-S-methyl (T5 and T6), the mean numbers of *A.*
230 *intermedius* per plant beating decreased significantly (ANOVA: $df = 5, 63$; $F = 47.541$; $P < 0.001$; n
231 $= 12$), as previously observed for onion thrips. This side effect was confirmed in particular for
232 lambda-cyhalothrin; in fact, significantly lower numbers of predatory thrips per plant beating were
233 observed in T2 after the second application on July 4 (ANOVA: $df = 5, 63$; $F = 5.234$; $P = 0.001$; n
234 $= 12$), and after the fourth application on July 22 (ANOVA: $df = 5, 63$; $F = 11.500$; $P < 0.001$; $n =$
235 12), and also in T3 on July 22, after the single application at the action threshold (Table 3). By
236 contrast, on July 13 (ANOVA: $df = 5, 63$; $F = 2.367$; $P = 0.049$; $n = 12$) and on July 22, after both
237 applications of spinosad (T4), the numbers of predatory thrips per plant beating were lower but not
238 significantly different from the untreated control (Table 3).

239 *Predator/prey ratio* Population abundances of *A. intermedius* and *T. tabaci*, and therefore the
240 predator/prey ratios, were variable during the field surveys (Fig. 1). On June 15 (Pre-S), predatory
241 thrips were more abundant than onion thrips in all treatments (>1.1 predator/prey). On June 23 (S1),
242 the ratio exceeded 1.3 in T1, T3 and T4, where no chemicals were applied. By contrast, ratios
243 drastically decreased in treatment T2, and especially in T5 and T6, where lambda-cyhalothrin (T2)
244 and acibenzolar-S-methyl (T5, T6) were sprayed for the first time, with mean values significantly
245 different from those recorded in the unsprayed plots (ANOVA: $df = 5, 66; F = 14.145; P < 0.0001;$
246 $n = 12$). After S1, overall ratios started to decline, due to both the increase of *T. tabaci* and the
247 decrease of *A. intermedius*. In treatment T2, where lambda-cyhalothrin was applied four times, *T.*
248 *tabaci* increased more distinctly, whereas *A. intermedius* decreased almost to the point of
249 disappearance. Consequently, the lowest predator/prey ratios were recorded in T2 and T3 on July
250 22, after the first application of lambda-cyhalothrin. On the last sampling date (S4), the ratios
251 recorded in T2 were significantly lower than those in the untreated control (T1) and plots sprayed
252 with spinosad (T4) (ANOVA: $df = 5, 66; F = 6.978; P < 0.0001; n = 12$). After the first application
253 of acibenzolar-S-methyl (S1) in both T5 and T6, predator/prey ratios were drastically reduced;
254 however, after the second application (S2), the ratios were similar to those observed in the other
255 treatments (except the lambda-cyhalothrin treatment), and the same trend was recorded in T5 after
256 the third and fourth applications (Fig. 1).

257 Laboratory bioassays

258 The percentages of onion thrips and predatory thrips alive in the vial bioassays after 24 h and 48 h
259 are shown in Figure 2. Significant differences between the treatments were found for onion thrips
260 after both 24 h (Kruskal-Wallis: $df = 3; \chi^2 = 52.329; P < 0.001; n = 7$) and 48 h (Kruskal-Wallis: df
261 $= 3; \chi^2 = 22.979; P < 0.001; n = 7$), and for predatory thrips after both 24 h (Kruskal-Wallis: $df = 3;$
262 $\chi^2 = 23.007; P < 0.001; n = 15$) and 48 h (Kruskal-Wallis: $df = 3; \chi^2 = 48.498; P < 0.001; n = 15$).

263 After 24 h, adult mortality of onion thrips and predatory thrips in the untreated control never
264 reached 5%. Spinosad was the most effective active ingredient; in fact, there were no adults of *T.*

265 *tabaci* or of *A. intermedius* alive after only 24 h. With lambda-cyhalothrin, 10% of onion thrips and
266 1% of predatory thrips survived, but these percentages were not statistically different from those
267 with spinosad. On the contrary, with acibenzolar-S-methyl, the percentages of live adults of onion
268 thrips and predatory thrips were statistically the same as the control (Fig. 2).

269 After 48 h, survival of *T. tabaci* and *A. intermedius* adults was statistically the same in
270 acibenzolar-S-methyl and control treatments. In the lambda-cyhalothrin treatment, although 6% of
271 onion thrips survived, there were no statistically significant differences between this active
272 ingredient and spinosad for either onion thrips or predatory thrips (Fig. 2).

273 Statistically significant differences were found between the survival data of onion thrips and
274 predatory thrips in certain treatments. *A. intermedius* adults were negatively affected by the TIBS
275 method more than *T. tabaci* adults. Even in the absence of any treatment exposure (*i.e.*, control),
276 after 48 h of isolation in the vials, under the same experimental conditions, mortality of predatory
277 thrips (40%) was significantly higher than mortality of onion thrips (7%) (ANOVA: $df = 1, 20$; $F =$
278 24.288 ; $P < 0.0001$; $n = 15, 7$), showing a greater sensitivity of the former species compared with
279 the latter, at least at the adopted experimental conditions. Exposure to lambda-cyhalothrin resulted
280 in significantly higher mortality of *A. intermedius* than *T. tabaci* after both 24 h and 48 h (ANOVA:
281 $df = 1, 20$; $F = 11.738; 9.679$; $P = 0.003; 0.006$; $n = 15, 7$). No differences between mortality of
282 predatory and onion thrips were recorded with spinosad (where no thrips survived in any of the
283 cases) or acibenzolar-S-methyl (ANOVA: $df = 1, 20$; $F = 0.028; 1.074$; $P = 0.868; 0.312$; $n = 15, 7$).

284

285 **Discussion**

286 During field surveys, two thrips species were dominant on onion, the phytophagous *T. tabaci* and
287 the zoophagous *A. intermedius*. In fact, the most numerous species was *T. tabaci*, independent of
288 the treatments, with population levels increasing from late June–early July, as observed in other
289 countries (Larentzaki *et al.* 2008; Torres-Vila *et al.* 1994). Nevertheless, population levels were
290 very low throughout the season, ranging from 0.3 to 4.7 thrips per plant by plant beating, probably

291 due to unfavorable climatic conditions. In 2011, temperatures in July were on average lower than in
292 2010 (*i.e.*, maximum, minimum and mean of 4.3°, 1.6° and 2.7 °C, respectively). Additionally,
293 rainfall in June (38.4 mm) and in July (30.0 mm) was more abundant than in 2010. It is well known
294 that heavy *T. tabaci* infestations occur mainly under hot and dry conditions (Theunissen &
295 Schelling 1997; Torres-Vila *et al.* 1994; Trdan *et al.* 2005b); thus the weather conditions probably
296 played an important role in the low population levels, compared with those observed in the growing
297 season of 2010 (Mautino *et al.* 2012). Moreover, during June, the frequent rainfalls (13 rainy days)
298 delayed the first chemical application, and consequently the low chemical pressure allowed
299 predatory thrips to migrate and establish in the field. Using the regression equation previously
300 developed to adjust the number of thrips per plant recorded with the beating method into the visual
301 method (Mautino *et al.* 2012), a mean seasonal value of around five thrips per plant (corresponding
302 to 1.5 thrips per plant by beating) overall was detected, and on average around seven thrips per
303 plant (corresponding to 2.3 thrips per plant by beating) were observed on the last sampling date
304 (July 22).

305 Among zoophagous thrips belonging to the family Aeolothripidae that feed on
306 phytophagous thrips (Bournier *et al.* 1979; Yano 2004; Zegula *et al.* 2003), *A. intermedius* is
307 considered to be a potentially important autochthonous facultative predator in Europe (Bournier *et*
308 *al.* 1978; Franco *et al.* 1999; Torres-Vila *et al.* 1994; Trdan *et al.* 2005a). The coexistence of *A.*
309 *intermedius* with the onion thrips, and also with *F. intonsa*, has already been observed in Italy
310 (Bournier *et al.* 1978, 1979; Conti 2009), but it has never been investigated thoroughly. The highest
311 predator population on the onion crop was detected in mid–late June, similar to populations in
312 France and Tuscany (central Italy) (Bournier *et al.* 1978; Conti 2009), and ranged from 0.03 to 2.3
313 predatory thrips per plant in relation to the treatment; these values were higher than those observed
314 on leek in Piedmont, where *Aeolothrips* sp. numbered on average 0.1 and 0.2 predatory thrips per
315 plant (Bosco & Tavella 2010).

316 During the field experiments in this study, the untreated plots did not exhibit the highest
317 onion thrips levels, as had been observed previously (Mautino *et al.* 2012). In this instance of low
318 thrips infestation, the untreated plots presented the best solution; this strongly supports the
319 importance of a supervised control based on pest monitoring before spray applications. On the
320 contrary, lambda-cyhalothrin applications were followed by the highest infestation levels of *T.*
321 *tabaci*. Resistance of onion thrips to pyrethroids (including lambda-cyhalothrin) has been reported
322 worldwide (Foster *et al.* 2010; Herron *et al.* 2008; MacIntyre Allen *et al.* 2005; Martin *et al.* 2003),
323 and in laboratory bioassays performed with the TIBS method (Rueda & Shelton 2003; Shelton *et al.*
324 2006). Nevertheless, in our laboratory bioassays by TIBS the efficacy of lambda-cyhalothrin was
325 high, especially after 48 h of exposure.

326 In the field, the failure of repeated lambda-cyhalothrin applications against onion thrips has
327 previously been observed (Nault & Shelton 2010). In our field experiments, failure was more likely
328 linked to factors other than the inefficacy of this active ingredient on the onion thrips population
329 itself, for example a side effect on autochthonous predators. Little information is available on the
330 direct (*i.e.*, physiological or behavioral) and indirect (*e.g.* habitat destruction, oviposition, resting,
331 mating sites) effects of this chemical on non-target organisms, including predatory thrips (Desneux
332 *et al.* 2007; Li *et al.* 2006; Mori & Gotoh 2001). In the field experiments, the lowest population
333 levels of *A. intermedius* were generally observed in the treatment with four applications of lambda-
334 cyhalothrin. Additionally, commencing from the second consecutive application, the lowest
335 predator/prey ratio was also found in this treatment, confirming that *T. tabaci* increased noticeably
336 whereas *A. intermedius* decreased almost to its disappearance. Moreover, in the laboratory
337 bioassays the sensitivity of *A. intermedius* to lambda-cyhalothrin was significantly higher than that
338 registered for onion thrips both after 24 h and 48 h of exposure (even if after 48 h the TIBS method
339 in itself negatively affected predatory thrips). Until now, no specific data existed regarding *A.*
340 *intermedius* sensitivity to pyrethroids used in IPM programs against *T. tabaci* (Bosco & Tavella
341 2010; Harper 1978), and the result obtained may be corroborated by further laboratory research

342 with *T. tabaci*, and its predatory thrips, under different experimental conditions and concentrations
343 of lambda-cyhalothrin.

344 Spinosad is well known to be one of the most effective insecticides against *T. tabaci* (Gent
345 *et al.* 2006; Shelton *et al.* 2006; Yarahmadi *et al.* 2009), and a reduced-risk insecticide for many
346 useful arthropods, if used properly (Funderburk *et al.* 2000; Jones *et al.* 2005; Ludwig & Oetting
347 2001; Workman & Martin 2002). In our study, the efficacy of spinosad against *T. tabaci* was
348 confirmed both under field and laboratory conditions, where it was the most insect-toxic among the
349 tested products. The same toxicity of this product was recorded in *A. intermedius* in the laboratory
350 bioassays; onion thrips and predatory thrips did not survive beyond 24 h of exposure to the active
351 ingredient at field concentrations. However, in the field experiments spinosad was less insect-toxic
352 than lambda-cyhalothrin to predatory thrips, as already observed on leek in northwestern Italy
353 (Bosco & Tavella 2010). The predator/prey ratios recorded after spinosad applications were similar
354 to ratios recorded in the other treatments, including the untreated control, unlike the observations
355 for lambda-cyhalothrin. Spinosad applications likely equally affected both thrips populations
356 without heavily impacting the predator/prey balance, thereby enabling the re-establishment of
357 predator activity and colonization of the crop. Thus, for the *Aeolothrips* genus, spinosad could
358 represent a reduced threat when used at the recommended timing and number of applications;
359 otherwise the active ingredient could prevent the development of predatory thrips (Workman &
360 Martin 2002) and other natural enemies useful for onion thrips control (Biondi *et al.* 2012).

361 The potential value of SAR compounds against several insect pests Alcantra *et al.* 2010;
362 (Correa *et al.* 2005; Costa *et al.* 2007; Tomquelski *et al.* 2007), and more specifically against
363 tospovirus and in thrips vector control (Gent *et al.* 2006; Momol *et al.* 2004; Pappu *et al.* 2000),
364 has been shown. Thrips feeding induces the expression of gene markers for the jasmonic and
365 salicylic acid pathways (JA and SA, respectively) involved in the basic plant defense response (Abe
366 *et al.* 2008). Acibenzolar-S-methyl is the synthetic functional analog of SA inducing the
367 corresponding response pathway (Gorlach *et al.* 1996). Moreover, a substantial co-regulation or

368 cross-talk among the salicylate and jasmonate plant defense pathways has been demonstrated
369 (Schenk *et al.* 2000); however, it is not fully understood how this signal interaction affects plant
370 response to thrips damage (Abe *et al.* 2012; Thaler *et al.* 2002). In the field experiments,
371 acibenzolar-S-methyl ensured low infestation levels of thrips on June 23 and July 4, as already
372 observed with high infestation levels (Mautino *et al.* 2012); nonetheless, acibenzolar-S-methyl was
373 unable adequately to control the *T. tabaci* population at the following sampling dates when plants
374 started to wither. Therefore, its effects on herbivores could be mediated by plant phenology. On the
375 other hand, where the SAR activator was sprayed only two times, followed by spinosad (T6), onion
376 thrips was effectively controlled. Unlike antiherbivore effects on insect pests, until now no
377 information was available about the effect of acibenzolar-S-methyl on *A. intermedius*. Acibenzolar-
378 S-methyl showed a similar side effect on predatory thrips, when it was applied singly and associated
379 with spinosad. In the laboratory bioassays, acibenzolar-S-methyl showed no effect on either *T.*
380 *tabaci* or *A. intermedius*. Adult survival in vials treated with this active ingredient was not
381 significantly different from that observed in the untreated control. Therefore, in the absence of the
382 plant, there is no side effect of acibenzolar-S-methyl on thrips, as previously observed in the vial
383 bioassay under the same experimental conditions (Mautino *et al.* 2012).

384 Spinosad appears to be the most effective control strategy against onion thrips, and the most
385 low-risk insecticide for *A. intermedius* if applied at the economic threshold, limiting the number of
386 applications, and in alternation with pesticides of a different mode of action (Biondi *et al.* 2012). As
387 an alternative control strategy, the addition of acibenzolar-S-methyl at the beginning of the growing
388 season should be considered also based on its minor impact on the overall predator/prey balance.
389 An environmentally friendly control strategy can maintain populations of the autochthonous *A.*
390 *intermedius* at reasonable levels in the open field. This species is naturally able to reduce *T. tabaci*
391 populations; therefore, its potential economic importance in IPM programs for onion is relevant.
392 Although in the field experiments densities of onion thrips were probably too low to comprehend
393 fully the direct effects of *A. intermedius* predatory activity, predator/prey ratios starting from

394 approximately 0.2 seem to be adequate to contain infestation levels of *T. tabaci*. Moreover,
395 supplemental techniques that are useful in improving thrips control are crucial and thus require
396 further investigation. In particular, methods to increase colonization by autochthonous predators in
397 the onion fields, such as programmed releases of combined beneficial arthropods (Fathi *et al.* 2008)
398 and intercropping with forage plants, should be developed (Bán *et al.* 2010; Theunissen & Schelling
399 1997; Trdan *et al.* 2006).

400

401 **Acknowledgments**

402 Special thanks are expressed to Ivo Rigamonti (Dipartimento di Scienze per gli Alimenti, la
403 Nutrizione e l'Ambiente, University of Milano) for confirming the thrips species identification. We
404 also thank Michela Basso, Sandro Frati and Michele Baudino of CReSO of Boves (province of Cuneo,
405 Italy) and Alessandro Costanzo of Cadir Lab of Quargnento (province of Alessandria, Italy) for
406 technical assistance in the field trials. Research was supported by a grant from Regione Piemonte.

407

408 **References** ‘*et al.*’ should be in italics

- 409 Abe, H., Ohnishi, J., Narusaka, M., Seo, S., Narusaka, Y., Tsuda, S., *et al.* (2008). Function of
410 jasmonate in response and tolerance of *Arabidopsis* to thrips feeding. *Plant Cell Physiology*, *49*, 68-
411 80.
- 412 Abe, H., Yasuhiro, T., Shimoda, T., Seo, S., Sakurai, T., Kugimiya, S., *et al.* (2012). Antagonistic
413 plant defense system regulated by phytohormones assists interactions among vector insect, thrips
414 and a tospovirus. *Plant Cell Physiology*, *53*, 204-212.
- 415 Alcantra, E., Moraes, J. C., & Antonio, A. (2010). Efeito de indutores da resistência e cultivares de
416 algodão no comportamento de *Aphis gossypii*. *Revista Ciência Agronômica*, *41*, 619-624.
- 417 Bán, G., Fetykó, K., & Tóth, F. (2010). Predatory arthropod assemblages of alfalfa and stinging
418 nettle as potential biological control agents of greenhouse pests. *Acta Phytopathologica et*
419 *Entomologica Hungarica*, *45*, 159-172.
- 420 Biondi, A., Mommaerts, V., Smagghe, G., Vinuela, E., Zappala, L., & Desneux, N. (2012). The
421 non-target impact of spinosyns on beneficial arthropods. *Pest Management Science*, *68*, 1523-1536.
- 422 Bosco, L., Giacometto, E., & Tavella, L. (2008). Colonization and predation of thrips
423 (Thysanoptera: Thripidae) by *Orius* spp. (Heteroptera: Anthocoridae) in sweet pepper greenhouses
424 in northwest Italy. *Biological Control*, *44*, 331-340.
- 425 Bosco, L., & Tavella, L. (2010). Population dynamics and integrated pest management of *Thrips*
426 *tabaci* on leek under field conditions in northwest Italy. *Entomologia Experimentalis et Applicata*,
427 *135*, 276-287.

428 Bournier, A., Lacasa, A., & Pivot, Y. (1978). Biologie d'un thrips prédateur *Aeolothrips*
429 *intermedius* [Thys.: Aeolothripidae]. *Entomophaga*, 23, 403-410.

430 Bournier, A., Lacasa, A., & Pivot, Y. (1979). Régime alimentaire d'un thrips prédateur *Aeolothrips*
431 *intermedius* [Thys.: Aeolothripidae]. *Entomophaga*, 24, 353-361.

432 Conti, B. (2009). Notes on the presence of *Aeolothrips intermedius* in northwestern Tuscany and on
433 its development under laboratory conditions. *Bulletin of Insectology*, 62, 107-112.

434 Correa, R. S. B., Moraes, J. C., Auad, A. M., & Carvalho, G. A. (2005). Silicon and acibenzolar-S-
435 methyl as resistance inducers in cucumber, against the whitefly *Bemisia tabaci* (Gennadius)
436 (Hemiptera: Aleyrodidae) biotype B. *Neotropical Entomology*, 34, 429-433.

437 Costa, R. R., Moraes, J. C., & Antunes, C. S. (2007). Resistência induzida em trigo ao pulgão
438 *Schizaphis graminum* (Rondani, 1852) (Hemiptera: Aphididae) por silício e acibenzolar-S-methyl.
439 *Ciência e Agrotecnologia, Lavras*, 31, 393-397.

440 Cox, P. D., Matthews, L., Jacobson, R. J., Cannon, R., MacLeod, A., & Walters, K. F. A. (2006).
441 Potential for the use of biological agents for the control of *Thrips palmi* (Thysanoptera: Thripidae)
442 outbreaks. *Biocontrol Science and Technology*, 16, 871-891.

443 Desneux, N., Decourtye, A., & Delpuech, J. M. (2007). The sublethal effects of pesticides on
444 beneficial arthropods. *Annual Review of Entomology*, 52, 81-106.

445 Diaz-Montano, J., Fuchs, M., Nault, B. A., & Shelton, A. M. (2010). Evaluation of onion cultivars
446 for resistance to onion thrips (Thysanoptera: Thripidae) and *Iris yellow spot virus*. *Journal of*
447 *Economic Entomology*, 103, 925-937.

448 Fathi, S. A. A., Asghari, A., & Sedghi, M. (2008). Interaction of *Aeolothrips intermedius* and *Orius*
449 *niger* in controlling *Thrips tabaci* on potato. *International Journal of Agriculture and Biology*, 10,
450 521-525.

451 Foster, S. P., Gorman, K., & Denholm, I. (2010). English field samples of *Thrips tabaci* show
452 strong and ubiquitous resistance to deltamethrin. *Pest Management Science*, 66, 861-864.

453 Franco, S., Beignet, P., Rat, E., & Thibout, E. (1999). The effect of thrips on wild and cultivated
454 alliaceous plants in France. *Phytoma*, 514, 41-44.

455 Funderburk, J. E., Stavisky, J., & Olson, S. M. (2000). Predation of *Frankliniella occidentalis*
456 (Thysanoptera: Thripidae) in field peppers by *Orius insidiosus* (Hemiptera: Anthocoridae).
457 *Environmental Entomology*, 29, 376-382.

458 Gabarra, R., Castañé, C., & Albajes, R. (1995). The mirid bug *Dicyphus tamanii* as a greenhouse
459 whitefly and western flower thrips predator on cucumber. *Biocontrol Science and Technology*, 5,
460 475-488.

461 Gent, D. H., du Toit, L. J., Fichtner, S. F., Mohan, S. K., Pappu, H. R., & Schwartz, H. F. (2006).
462 *Iris yellow spot virus*: an emerging threat to onion bulb and seed production. *Plant Disease*, 90,
463 1468-1480.

464 Gorlach, J., Volrath, S., Knauf-Beiter, G., Hengy, G., Beckhove, U., Kogel, K. H., *et al.* (1996).
465 Benzothiadiazole, a novel class of inducers of systemic acquired resistance, activates gene
466 expression and disease resistance in wheat. *The Plant Cell*, 8, 629-643.

467 Harper, A. M. (1978). Effect of insecticides on the pea aphid, *Acyrtosiphon pisum* (Hemiptera:
468 Aphididae), and associated fauna on alfalfa. *The Canadian Entomologist*, 110, 891-894.

469 Herron, G. A., James, T. M., Rophail, J., & Mo, J. (2008). Australian populations of onion thrips,
470 *Thrips tabaci* Lindeman (Thysanoptera: Thripidae), are resistant to some insecticides used for their
471 control. *Australian Journal of Entomology*, 47, 361-364.

472 Jones, T., Scott-Dupree, C., Harris, R., Shipp, L., & Harris, B. (2005). The efficacy of spinosad
473 against the western flower thrips, *Frankliniella occidentalis*, and its impact on associated biological
474 control agents on greenhouse cucumbers in southern Ontario. *Pest Management Science*, *61*, 179-
475 185.

476 Kakimoto, K., Inoue, H., Hinomoto, N., Noda, T., Hirano, K., Kashio, T., *et al.* (2006). Potential of
477 *Haplothrips brevitubus* (Karny) (Thysanoptera: Phlaeothripidae) as a predator of mulberry thrips
478 *Pseudodendrothrips mori* (Niwa) (Thysanoptera: Thripidae). *Biological Control*, *37*, 314-319.

479 Koschier, E. H., Sedy, K. A., & Novak, J. (2002). Influence of plant volatiles on feeding damage
480 caused by the onion thrips *Thrips tabaci*. *Crop Protection*, *21*, 419-425.

481 Larentzaki, E., Plate, J., Nault, B. A., & Shelton, M. (2008). Impact of straw mulch on populations
482 of onion thrips (Thysanoptera: Thripidae) in onion. *Journal of Economic Entomology*, *101*, 1317-
483 1324.

484 Li, D. X., Tian, J., & Shen, Z. R. (2006). Effects of pesticides on the functional response of
485 predatory thrips, *Scolothrips takahashii* to *Tetranychus viennensis*. *Journal of Applied*
486 *Entomology*, *130*, 314-322.

487 Ludwig, S., & Oetting, R. (2001). Effect of spinosad on *Orius insidiosus* (Hemiptera:
488 Anthocoridae) when used for *Frankliniella occidentalis* (Thysanoptera: Thripidae) control on
489 greenhouse pot chrysanthemums. *The Florida Entomologist*, *84*, 311-313.

490 MacIntyre Allen, J. K., Scott-Dupree, C. D., Tolman, J. H., & Harris, C. R. (2005). Resistance of
491 *Thrips tabaci* to pyrethroid and organophosphorus insecticides in Ontario, Canada. *Pest*
492 *Management Science*, *61*, 809-815.

493 Martin, N. A., Workman, P. J., & Butler, R. C. (2003). Insecticide resistance in onion thrips (*Thrips*
494 *tabaci*) (Thysanoptera: Thripidae). *New Zealand Journal of Crop and Horticultural Science*, *31*, 99-
495 106.

496 Mautino, G. C., Bosco, L., & Tavella, L. (2012). Integrated management of *Thrips tabaci*
497 (Thysanoptera: Thripidae) on onion in north-western Italy: basic approaches for supervised control.
498 *Pest Management Science*, *68*, 185-193.

499 Momol, M. T., Olson, S. M., Funderburk, J. E., Stavisky, J., & Marois, J. J. (2004). Integrated
500 management of tomato spotted wilt on field-grown tomatoes. *Plant Disease*, *88*, 882-890.

501 Mori, K., & Gotoh, T. (2001). Effects of pesticides on the spider mite predators, *Scolothrips*
502 *takahashii* (Thysanoptera: Thripidae) and *Stethorus japonicus* (Coleoptera: Coccinellidae).
503 *International Journal of Acarology*, *27*, 299-302.

504 Mound, L. A., Morison, G. D., Pitkin, B. R., & Palmer, J. M. (1976). *Thysanoptera. Handbooks for*
505 *the identification of British insects. vol. I, part 11*. London, UK: Royal Entomological Society of
506 London.

507 Nagata, T., Almeida, A. C. L., Resende, R. O., & de Ávila, A. C. (1999). The identification of the
508 vector species of Iris yellow spot tospovirus occurring on onion in Brazil. *Plant Disease*, *83*, 399.

509 Nault, B. A., & Shelton, A. M. (2010). Impact of insecticide efficacy on developing action
510 thresholds for pest management: a case study of onion thrips (Thysanoptera: Thripidae) on onion.
511 *Journal of Economic Entomology*, *103*, 1315-1326.

512 Pappu, H. R., Csinos, A. S., McPherson, R. M., Jones, D. C., & Stephenson, M. G. (2000). Effect of
513 acibenzolar-S-methyl and imidacloprid on suppression of tomato spotted wilt *Tospovirus* in flue-
514 cured tobacco. *Crop Protection*, *19*, 349-354.

515 Pappu, H. R., Jones, R. A. C., & Jain, R. K. (2009). Global status of tospovirus epidemics in diverse
516 cropping systems: successes achieved and challenges ahead. *Virus Research*, *141*, 219-236.

517 Patil, S. D., Chandele, A. G., Wayal, C. B., & Game, B. C. (2009). Efficacy of different newer
518 chemicals and bio-insecticides against onion thrips in *kharif* season. *International Journal of Plant*
519 *Protection*, 2, 227-230.

520 Riudavets, J. (1995). Predators of *Frankliniella occidentalis* and *Thrips tabaci*. in:
521 A. J. M. Loomans, J. C. van Lenteren, M. G. Tommasini, S. Maini, & J. Riudavets (Eds.)
522 *Biological control of thrips pests. Wageningen Agricultural University Papers 95-1*. pp. 47-87.
523 Wageningen, the Netherlands: Veenman Drukkers.

524 Riudavets, J., Gabarra, R., & Castañé, C. (1993). *Frankliniella occidentalis* predation by native
525 natural enemies. *IOBC/WPRS Bulletin*, 16, 137-140.

526 Rueda, A., & Shelton, A. M. (2003). Development of a bioassay system for monitoring
527 susceptibility in *Thrips tabaci*. *Pest Management Science*, 59, 553-558.

528 Schenk, P. M., Kazan, K., Wilson, I., Anderson, J. P., Richmond, T., Somerville, S. C. *et al.*
529 (2000). Coordinated plant defense responses in *Arabidopsis* revealed by microarray analysis. *PNAS*,
530 97, 11655-11660.

531 Schliephake, G., & Klimt, K. (1979). *Thysanoptera, Fransenflügler. Die Tierwelt Deutschlands*, 66.
532 Jena, Germany: VEB Gustav Fisher Verlag.

533 Shelton, A. M., Zhao, J-Z., Nault, B. A., Plate, J., Musser, F. R., & Larentzaki, E. (2006). Patterns
534 of insecticide resistance in onion thrips (Thysanoptera: Thripidae), in onion fields in New York.
535 *Journal of Economic Entomology*, 99, 1798-1804.

536 Tedeschi, R., Ciuffo, M., Mason, G., Roggero, P., & Tavella, L. (2001). Transmissibility of four
537 tospoviruses by a thelytokous population of *Thrips tabaci* from Liguria, northwestern Italy.
538 *Phytoparasitica*, 29, 37-45.

539 Thaler, J. S., Karban, R., Ullman, D. E., Boege, K., & Bostock, R. M. (2002). Cross-talk between
540 jasmonate and salicylate plant defense pathways: effects on several plant parasites. *Oecologia*, 131,
541 227-235.

542 Theunissen, J., & Schelling, G. (1997). Damage threshold for *Thrips tabaci* (Thysanoptera:
543 Thripidae) in monocropped and intercropped leek. *European Journal of Entomology*, 94, 253-261.

544 Tomquelski, G. V., Martins, G. L. M., & Papa, G. (2007). Efeito dos indutores de resistência
545 acibenzolar-S- metil e silício na biologia de *Alabama argillacea* (Lepidoptera: Noctuidae) em
546 algodoeiro. *Revista de Agricultura*, 82, 170-175.

547 Torres-Vila, L. M., Lacasa, A., Bielza, P., & Meco, R. (1994). Dinámica poblacional de *Thrips*
548 *tabaci* Lind. (Thysanoptera: Thripidae) sobre liliáceas hortícolas en Castilla-La Mancha. *Boletín de*
549 *Sanidad Vegetal, Plagas*, 20, 661-677.

550 Trdan, S., Andjus, L., Raspudic, E., & Kac, M. (2005a). Distribution of *Aeolothrips intermedius*
551 Bagnall (Thysanoptera: Aeolothripidae) and its potential prey Thysanoptera species on different
552 cultivated host plants. *Journal of Pest Science*, 78, 217-226.

553 Trdan, S., Rifelj, M., & Valic, N. (2005b). Population dynamics of banded thrips (*Aeolothrips*
554 *intermedius* Bagnall, Thysanoptera, Aeolothripidae) and its potential prey Thysanoptera species on
555 white clover. *Communications in Agricultural and Applied Biological Sciences*, 70, 753-758.

556 Trdan, S., Žnidarčič, D., Valič, N., Rozman, L., & Vidrih, M. (2006). Intercropping against onion
557 thrips, *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) in onion production: on the suitability of
558 orchard grass, lacy phacelia, and buckwheat as alternatives for white clover. *Journal of Plant*
559 *Diseases and Protection*, 113, 24-30.

560 Vierbergen, G. B., Kucharczyk, H., & Kirk, W. D. J. (2010). A key to the second instar larvae of
561 the Thripidae of the Western Palaearctic region (Thysanoptera). *Tijdschrift voor Entomologie*, *153*,
562 99-160.

563 Wimmer, D., Hoffmann, D., & Schausberger, P. (2008). Prey suitability of western flower thrips,
564 *Frankliniella occidentalis*, and onion thrips, *Thrips tabaci*, for the predatory mite *Amblyseius*
565 *swirskii*. *Biocontrol Science and Technology*, *18*, 533-542.

566 Workman, P. J., & Martin, N. A. (2002). Towards integrated pest management of *Thrips tabaci* in
567 onions. *New Zealand Plant Protection*, *55*, 188-192.

568 Yano, E. (2004). Recent development of biological control and IPM in greenhouses in Japan.
569 *Journal of Asia-Pacific Entomology*, *7*, 5-11.

570 Yarahmadi, F., Moassadegh, M. S., Soleymannejadian, E., Saber, M., & Shishehbor, P. (2009).
571 Assessment of acute toxicity of abamectin, spinosad and chlorpyrifos to *Thrips tabaci* Lindeman
572 (Thysanoptera: Thripidae) on sweet pepper by using two bioassay techniques. *Asian Journal of*
573 *Biological Sciences*, *2*, 81-87.

574 Zegula, T., Sengonca, C., & Blaeser, P. (2003). Entwicklung, reproduktion und Prädationsleistung
575 von zwei Raubthrips-arten *Aeolothrips intermedius* Bagnall und *Franklinothrips vespiformis*
576 Crawford (Thysanoptera: Aeolothripidae) mit ernährung zweier natürlicher beutearten. *Gesunde*
577 *Pflanzen*, *55*, 169-174 (German, with English summary).

578

579 Table 1. Active ingredients, rates, number and timing of applications of the products sprayed in the
 580 experimental plots of the onion field during the 2011 growing season

Treatment	Active ingredient	Rate ($l\ ha^{-1}$)	No. ^z	Timing
T1 ^y	-	-	-	-
T2	Lambda-cyhalothrin	1.3	4	19 June; 29 June; 9 July; 19 July
T3	Lambda-cyhalothrin	1.3	1 ^x	19 July
T4	Spinosad	0.8	2	9 July; 19 July
T5	Acibenzolar-S-methyl	0.2	4	19 June; 29 June; 9 July; 19 July
T6	Acibenzolar-S-methyl +Spinosad	0.2 0.8	2 2	19 June; 29 June 9 July; 19 July

581 ^zNumber of applications

582 ^yUntreated control

583 ^xPyrethroid application at the action threshold of two thrips per plant by plant beating

584

585 Table 2. Mean numbers (\pm SE) of *Thrips tabaci* (adults plus larvae) per plant sampled by plant
 586 beating in the six tested treatments during field surveys in 2011 (Statistical analyses were performed on
 587 log-transformed data which are not shown)

Treatment	Pre-S ^z 15 June	S1 ^z 23 June	S2 ^z 04 July	S3 ^z 13 July	S4 ^z 22 July
T1	1.03 \pm 0.14	1.60 \pm 0.14 a ^y	1.13 \pm 0.23	1.38 \pm 0.25 bc	2.52 \pm 0.40 b
T2	1.07 \pm 0.11	0.27 \pm 0.10 b	1.22 \pm 0.19	2.68 \pm 0.46 ab	4.70 \pm 0.63 a
T3	0.98 \pm 0.15	1.20 \pm 0.15 a	0.98 \pm 0.29	2.05 \pm 0.31 abc	1.08 \pm 0.25 c
T4	1.45 \pm 0.17	1.52 \pm 1.27 a	1.35 \pm 0.27	1.25 \pm 0.20 c	1.67 \pm 0.20 bc
T5	1.02 \pm 0.14	0.30 \pm 0.09 b	0.87 \pm 0.18	3.23 \pm 0.33 a	2.87 \pm 0.42 ab
T6	1.13 \pm 0.17	0.47 \pm 0.08 b	0.67 \pm 0.14	1.92 \pm 0.26 abc	1.10 \pm 0.18 c
<i>P</i>	0.327	< 0.001	0.256	< 0.001	< 0.001
<i>F</i> _{5,63}	1.182	20.438	1.347	5.512	14.438
SED ^x	0.101	0.103	0.152	0.147	0.146

588

589 ^zPre-S represents the sampling before chemical applications; S1, S2, S3, and S4 represent sampling after the
 590 1st, 2nd, 3rd and 4th applications, respectively

591 ^yWithin columns, means followed by a common letter do not differ significantly ($P < 0.05$, Tukey's test
 592 following ANOVA). ANOVA results (P and F values, $df = 5, 63, n = 12$) are reported

593 ^xStandard errors of the difference values

594

595 Table 3. Mean numbers (\pm SE) of *Aeolothrips intermedius* (adults plus larvae) per plant sampled by
 596 plant beating in the six tested treatments during field surveys in 2011 (Statistical analyses were
 597 performed on log-transformed data which are not shown)

Treatment	Pre-S ^z 15 June	S1 ^z 23 June	S2 ^z 04 July	S3 ^z 13 July	S4 ^z 22 July
T1	1.63 \pm 0.24	2.32 \pm 0.25 a ^y	0.65 \pm 0.10 a	0.25 \pm 0.09 ab	0.87 \pm 0.15 a
T2	1.78 \pm 0.21	0.18 \pm 0.13 b	0.15 \pm 0.06 b	0.07 \pm 0.04 b	0.10 \pm 0.04 c
T3	1.80 \pm 0.23	1.88 \pm 0.28 a	0.73 \pm 0.18 a	0.28 \pm 0.09 ab	0.10 \pm 0.06 c
T4	1.63 \pm 0.28	2.10 \pm 0.30 a	0.78 \pm 0.15 a	0.17 \pm 0.05 ab	0.48 \pm 0.08 ab
T5	1.87 \pm 0.20	0.03 \pm 0.02 b	0.45 \pm 0.13 ab	0.40 \pm 0.07 a	0.47 \pm 0.13 abc
T6	1.43 \pm 0.21	0.12 \pm 0.05 b	0.35 \pm 0.11 ab	0.28 \pm 0.14 ab	0.15 \pm 0.03 bc
P	0.667	< 0.001	0.001	0.049	< 0.001
F _{5,63}	0.644	47.541	5.234	2.367	11.500
SED ^x	0.120	0.113	0.099	0.081	0.085

598

599 ^zPre-S represents the sampling before chemical applications; S1, S2, S3, and S4 represent sampling after the
 600 1st, 2nd, 3rd and 4th applications, respectively

601 ^yWithin columns, means followed by a common letter do not differ significantly ($P < 0.05$, Tukey's test
 602 following ANOVA). ANOVA results (P and F values, $df = 5, 63$, $n = 12$) are reported

603 ^xStandard errors of the difference values

604

605 **Figure captions**

606 **Fig. 1** Mean ratios of predator/prey (*Aeolothrips intermedius*/*Thrips tabaci*) per plant in the six
607 tested treatments during field surveys. Within the same sampling date, bars labeled with different
608 letters are significantly different (Tukey's test following ANOVA, $P < 0.01$). Statistical analyses
609 were performed on log-transformed data which are not shown. Pre-S represents the sampling before
610 chemical applications; S1, S2, S3, and S4 represent sampling after the 1st, 2nd, 3rd and 4th applications,
611 respectively

612

613

614 **Fig. 2** Mean (\pm SE) survival percentages of adult *Thrips tabaci* (**a**) and *Aeolothrips intermedius* (**b**)
615 after 24 h and 48 h in vials treated with the tested products at field concentrations. Within thrips
616 treatments, bars labeled with different letters (lower case and capital, for survivor thrips at 24 h and
617 48 h, respectively) are significantly different ($P < 0.05$, Kruskal-Wallis, $df = 3$)




Fig. 1




Figure 2