

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Biliary tract cancer in male printers and typesetters in the European rare cancer case-control study.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/155441> since

Published version:

DOI:10.1136/oemed-2014-102322

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Ahrens W, Merletti F, Mirabelli D
Biliary tract cancer in male printers and typesetters in the European rare
cancer case-control study.
OCCUPATIONAL AND ENVIRONMENTAL MEDICINE (2014) 71
DOI: 10.1136/oemed-2014-102322

The definitive version is available at:

<http://oem.bmj.com/cgi/doi/10.1136/oemed-2014-102322>

Biliary tract cancer in male printers and typesetters in the European rare cancer case-control study

Kumagai et al recently observed a striking increase in mortality from cholangiocarcinoma (CC) among workers in a colour Occup Environ Med August 2014 Vol 71 No 8 591 proof-printing shop in Japan, based on six deaths among 11 CC cases in a group of 62 workers employed between 1991 and 2006. Exposure to 1,2-dichloropropane and dichloromethane was reported.¹ Kubo et al extended the observation including 111 workers employed from 1981 to 2012, and reported 17 CC cases.² The Nordic Occupational Cancer Study (NOCCA) was used to explore the generalisability of the Japanese cluster findings to the printing industry and related occupations at large.³ An overall standardised incidence ratio (SIR) of 2.34 (95% CI 1.45 to 3.57) was found for intrahepatic CC, based on 21 cases in men. Women had an SIR of 1.95 (95% CI 0.84 to 3.85) with eight cases. Printers and lithographers had the highest SIRs and both occupations were recognised as entailing exposures to cleaning agents and being possibly related with the use of chlorinated solvents. A European multicentric case-control study was conducted between 1995 and 1997 on rare cancers of unknown aetiology, including gallbladder and extrahepatic tract cancers (ECC). We used the study dataset to assess the association of ECC with printing jobs in men. The study design has been fully reported.⁴ In brief, nine countries were involved. In six of them population-based cases and controls were recruited, while centres from three countries recruited hospital case-series and hospital recruitment-area controls. Study subjects were administered a standard questionnaire during face-to-face or telephone interviews. Surrogate interviews with a next-of kin (NoK) were performed if the index subject was too ill or had died before contact was established. All jobs lasting at least 6 months were recorded with the start and end years, working hours, materials handled, and chemical exposures. Jobs were coded according to the 1968 International Standard Classifications of Occupations (ISCO) on a four-digit level.⁵ Analyses were restricted to the pooled European dataset with available population controls, using 153 cases considered as definite or possible after pathological review and 1421 population controls. Printers were defined as individuals ever employed in an occupation corresponding to the ISCO code 92*** (ie, the first two ISCO digits), and typesetters as ISCO code 921**. The reference group was formed by all participants who were never employed as printers or typesetters. Results are presented as ORs with 95% CIs (95%-CI). Risk estimates were calculated by unconditional logistic regression models. Logistic regression models included country (one dummy variable for each country) and year of birth as a continuous variable (OR₁). Further models adjusted for history of gallstones (self-reported as being confirmed or treated by a physician) and NoK status (OR₂). Analyses were performed with SAS (V.8.2). As shown in table 1, ORs were increased for both printers and typesetters. CIs were wide due to the rarity of the cancers and occupations of interest, but our results lend support to the hypothesis of an increased risk for ECC in printing occupations, and are in agreement with the fact that cases of extrahepatic CC were observed along with those of intrahepatic CC in Japanese printers. ^{1 2} Out of five cases among printing workers, two were from Sweden—the only Nordic country involved in our casecontrol study—and both were typesetters. We were unable to identify candidate agents to explain this association because no information on specific chemicals used in the printing industry was available in our study, differently from other exposures.^{6–8} It is clear, however, that if the association is causal in nature, the causative agent(s) must have been present in printing rooms for a long time.

Wolfgang Ahrens,^{1,2} Franco Merletti,³ Dario Mirabelli³ ¹Department of Epidemiological Methods and Etiologic Research, Leibniz Institute for Prevention Research and Epidemiology—BIPS, Bremen, Germany ²Faculty of Mathematics and Computer Science, Institute for Statistics, University of Bremen, Bremen, Germany ³Unit of Cancer Epidemiology, Department of Medical Sciences, University of Turin and Città della Scienza e della Salute Hospital, Turin, Italy Correspondence to Professor Franco Merletti, Epidemiologia dei Tumori, via Santena 7, Torino 10126, Italy; franco.merletti@unito.it Contributors WA conducted the analyses, all authors designed the study, assessed results and wrote the letter. Competing interests None. Patient consent Obtained. Ethics approval Multicentric study: all participating countries and centres obtained approval by the appropriate independent ethical Committee. Provenance and peer review Not commissioned; internally peer reviewed. To cite Ahrens W, Merletti F, Mirabelli D. Occup Environ Med 2014;71:591–592. Received 9 May 2014 Accepted 27 May 2014 Published Online First 10 June 2014 Occup Environ Med 2014;71:591–592. doi:10.1136/oemed-2014-102322 REFERENCES 1 Kumagai S, Kurumatani N, Arimoto A, et al. Cholangiocarcinoma among offset colour proof-printing workers exposed to 1,2-dichloropropane and/or dichloromethane. Occup Environ Med 2013;70:508–10. 2 Kubo S, Nakanuma Y, Takemura S, et al. Case series of 17 patients with cholangiocarcinoma among young adult workers of a printing company in Japan. J Hepatobiliary Pancreat Sci 13 Jan 2014. doi:10.1002/jhbp.86. 3 Vlaanderen J1, Straif K, Martinsen JI, et al. Cholangiocarcinoma among workers in the printing industry: using the NOCCA database to elucidate the generalisability of a cluster report from Japan. Occup Environ Med 2013;70:828–30. 4 Lynge E, Afonso N, Kaerlev L, et al. European multi-centre case-control study on risk factors for rare cancers of unknown aetiology. Eur J Cancer 2005;41:601–12. 5 International Labour Office. International standard classification of occupations. Geneva, Switzerland: International Labour Office, 1968. 6 Schmeisser N, Kaerlev L, Bourdon-Raverdy N, et al. Occupational exposure to pesticides and bile tract carcinoma in men: results from a European multicenter case-control study. Cancer Causes Control 2010;21:1493–502. 7 Ahrens W, Timmer A, Vyberg M, et al. Risk factors for extrahepatic biliary tract carcinoma in men: medical conditions and lifestyle: results from a European multicentre case-control study. Eur J Gastroenterol Hepatol 2007;19:623–30. 8 Ahrens W1, Mambetova C, Bourdon-Raverdy N, et al. Occupational exposure to endocrine-disrupting compounds and biliary tract cancer among men. Scand J Work Environ Health 2007;33:387–96.

Table 1 Number of printers and typesetters among cases and controls and corresponding ORs in the European case-control study of rare cancers

Job category	Controls n (%)	Cases n (%)	OR ₁ (95% CI)	OR ₂ (95% CI)
Printing worker (ISCO 92***)				
No	1396 (98.2)	148 (96.7)	1.00 (–)	1.00 (–)
Yes	25 (1.8)	5 (3.3)	1.88 (0.69 to 5.17)	2.42 (0.81 to 7.24)
Typesetter (ISCO 921**)				
No	1411 (99.3)	150 (98.0)	1.00 (–)	1.00 (–)
Yes	10 (0.7)	3 (2.0)	3.26 (0.82 to 12.99)	5.78 (1.43 to 23.29)