

To the editor:

Can the revised IPSS predict response to erythropoietic-stimulating agents in patients with classical IPSS low or intermediate-1 MDS?

The “classical” International Prognostic Scoring System (IPSS), based on cytogenetics, marrow blast percentage, and number of cytopenias, has played a major role in prognosis assessment in myelodysplastic syndromes (MDS).¹ The recently published revised IPSS (IPSS-R), using the same parameters, but with 5 rather than 3 cytogenetic subgroups and new cutoff values for cytopenias and marrow blast percentages, refines the original IPSS prognostic value.^{2,3} However, its prognostic value for response to erythropoiesis-stimulating agents (ESA) has not been assessed. We analyzed it retrospectively in 456 IPSS low/intermediate (Int)-1–risk MDS patients treated with ESA in France, Germany, and Italy.

Those 456 patients had serum erythropoietin (EPO) <500 mU/mL and hemoglobin (Hb) ≤10 g/dL and had received ESA (EPO alfa or β 40 000-60 000 IU/week, or darbepoetin 150-300 μg/week) for at least 12 weeks. In addition to IPSS-R parameters, age, sex, serum EPO level, serum ferritin (SF), red blood cell (RBC) transfusion requirement before ESA onset were assessed for response to ESA (based on International Working Group 2006 criteria), and overall survival (OS) from ESA onset. Characteristics of the 456 patients at ESA onset are listed in Table 1. Seventy-one percent of the patients had never received RBC transfusions, and their median Hb level was 9.3g/dL (range 7.0-10); 29% of patients had received at least 4 RBC concentrates/8 weeks before ESA onset (with a maximum of 12 concentrates). Median SF was 357 ng/mL and serum EPO was 60 mU/mL (range 6-483). IPSS was low in 55% and Int-1 in 45% of patients. IPSS-R was very low in 15%, low in 61%, intermediate in 19%, and high in 4% of the patients.

A total of 303 (61%) patients had an erythroid response, including 72% and 52% of low and Int-1 risk patients, respectively ($P = .001$). Using IPSS-R, 85%, 68%, 48%, and 31% of patients had erythroid response in the very low, low, intermediate, and high-risk groups, respectively ($P < .0001$).

Other prognostic factors of erythroid response, in univariate analysis, included individual IPSS-R parameters analyzed according to IPSS-R thresholds (Hb level, platelet count, absolute neutrophil count, marrow blast %, cytogenetics), serum EPO level, SF (variables tested as continuous variables), and previous RBC transfusions. In multivariate analysis, IPSS-R, serum EPO, and SF remained significantly associated with erythroid response ($P < .0001$, $P < .0001$, and $P = .002$, respectively).

Applying 1 point to each of the following unfavorable variables of response to ESA, serum EPO >200 mU/mL (=1), SF >350 ng/mL (=1), and IPSS-R (very low = 0, low = 1, intermediate = 2, and high = 3) yielded a score ranging from 0 to 4, with response rates of 85%, 80%, 64%, 40%, and 20%, respectively. As expected, IPSS-R also had strong prognostic value for OS (not shown).

Thus, in this patient cohort with overall favorable prognostic factors of response to ESA according to the Nordic score⁴ (ie, serum EPO <500 mU/mL and no or limited transfusion dependence), IPSS-R alone, and even better, a score ≥3 (using IPSS-R, serum EPO, and SF) proved useful to identifying patients with low response to ESA who also have worse OS⁵ and may require alternative treatments.

Jennifer Schemenau

Heinrich-Heine University,
Düsseldorf, Germany

Alessandro Levis

ASO Santi Antonio e Biagio,
Düsseldorf, Germany

Enrico Balleari

University of Genoa,
Genoa, Italy

Rosa Sapena

Groupe Francophone des Myélodysplasies,
Paris, France

Lionel Adès

Hopital Avicenne,
Paris, France

Agnès Guerci

Centre Hospitalo Universitaire,
Nancy, France

Odile Beyne-Rauzy

Centre Hospitalo Universitaire, Toulouse Purpan,
Toulouse, France

Marie-Pierre Gourin

Groupe Francophone des Myélodysplasies,
Limoges, France

Stephane Cheze

Groupe Francophone des Myélodysplasies,
Caen, France

Aspasia Stamatoullas

Groupe Francophone des Myélodysplasies,
Rouen, France

Alessandro Sanna

Fundazion Italiana delle sindromi mielodisplastiche,
Florence, Italy

Daniela Gioia

Fundazion Italiana delle sindromi mielodisplastiche,
Alessandria, Italy

Gianni Cametti

Fundazion Italiana delle sindromi mielodisplastiche,
Chieri, Italy

Dario Ferrero

Fundazion Italiana delle sindromi mielodisplastiche,
Torino, Italy

Emmanuel Raffoux

Hospital Saint Louis,
Paris, France

Christian Rose

Hopital Saint Vincent de Paul, Université Catholique de Lille,
Lille, France

Antonella Poloni

Universita Politecnica delle Marche,
Ancona, Italy

Thomas Prebet

Groupe Francophone des Myélodysplasies,
Marseille, France

Valeria Santini

Fundazion Italiana delle sindromi mielodisplastiche,
Florence, Italy

Table 1. Baseline patient characteristics

	n = 456	Patients, %	Erythroid response (IWG 2006 criteria)	Univariate analysis P	Multivariate analysis P
Hb, g/dL					
<8	52	11	33		
8-10	292	64	66		
≥10	112	25	78	<.0001	
ANC					
<0.8	42	9	55		
≥0.8	414	91	66	.16	
Platelet count*					
<50	17	3	37		
50-100	58	13	57		
>100	381	84	67	.01	
Marrow blasts, %					
≤2	238	52	72		
<2-<5	118	26	61		
5-10	100	22	53	.0052	
IPSS-R karyotype					
Very good	21	5	71		
Good	379	83	65		
Intermediate	45	10	75		
Poor-very poor	11	2	20	.008	
IPSS					
Low	256	55	72		
Int-1	210	45	52	.001	
IPSS-R					
Very low	69	15	85		
Low	281	61	68		
Intermediate	87	19	48		
High	19	4	31	<.0001	<.0001
Sex					
Male	250	55	62		
Female	206	45	68	.14	
Age, years					
≤60	29	6	55		
>60	427	94	66	.23	
EPO, mU/mL					
≤100	306	67	75		
>100	150	33	45	<.0002	
≤200	393	86	75		
>200	63	13	31	<.0001	<.0001
Serum ferritin, ng/mL					
≤350	224	49	72		
>350	232	51	58	.001	.002
Previous RBC transfusions					
No	323	71	73		
Yes	133	29	55	.0005	
WHO classification					
RA	110	24	73		
RAEB-1	69	15	50		
RARS	114	25	70		
RARS-T	2	0.3	100		
RCMD	124	27	62		
MDS WITH DEL 5q	21	5	62		
MDS-U	16	4	55	.03	.71

ANC, absolute neutrophil count; IWG, International Working Group; MDS-U, MDS-unclassified; MDS WITH DEL, MDS associated with isolated del (5q); RA, refractory anemia; RAEB-1, refractory anemia with excess blasts-1; RARS, refractory anemia with ring sideroblasts; RARS-T, refractory anemia with ring sideroblasts associated with marked thrombocytosis; RCMD, refractory cytopenia with multilineage dysplasia; WHO, World Health Organization.

Univariate and multivariate analyses for response (IWG 2006 criteria). The variables with significant prognostic value in univariate analysis were integrated in the multivariate analysis, excepting Hb, platelet count, IPSS-R karyotype, and % of blasts, because IPSS-R integrates already those variables. For EPO level, the 200 mU/mL threshold was chosen because it was the most significant in univariate analysis. Previous transfusions were not integrated in the model because they "duplicate" the Hb variable.

*10⁹/L.

Laurence Legros

Groupe Francophone des Myélodysplasies,
Nice, France

Shanti Natarajan-Amé

Hopital Civil, Centre Hospitalo Universitaire,
Strasbourg, France

Pierre Fenaux

Hopital Avicenne, Paris 13 University,
Paris, France

Ulrich Germing

Heinrich-Heine University,
Düsseldorf, Germany

François Dreyfus

Hopital Cochin, Assistance Publique des Hôpitaux de Paris,
Paris, France

Sophie Park

Centre Hospitalo Universitaire,
Grenoble, France

Contribution: V.S., J.S., A.L., E.B., R.S., L.A., A.G., O.B.-R., M.-P.G., S.C., A. Stamatoulas, A. Sanna, D.G., G.C., D.F., E.R., C.R., A.P., T.P., L.L., S.N.-A., P.F., U.G., F.D., and S.P. collected and analyzed the data; V.S., U.G., P.F., F.D., and S.P. designed the study and wrote the manuscript; and S.P. performed the statistical analysis.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Sophie Park, Centre Hospitalo Universitaire (CHU) of Grenoble, Clinique Universitaire d'hématologie, 6eA, CS 10217, 38043 Grenoble, France; e-mail: spark@chu-grenoble.fr.

References

- Greenberg P, Cox C, LeBeau MM, et al. International scoring system for evaluating prognosis in myelodysplastic syndromes. *Blood*. 1997;89(6):2079-2088.
- Greenberg PL, Tuechler H, Schanz J, et al. Revised international prognostic scoring system for myelodysplastic syndromes. *Blood*. 2012;120(12):2454-2465.
- Schanz J, Tüchler H, Solé F, et al. New comprehensive cytogenetic scoring system for primary myelodysplastic syndromes (MDS) and oligoblastic acute myeloid leukemia after MDS derived from an international database merge. *J Clin Oncol*. 2012;30(8):820-829.
- Hellström-Lindberg E, Gulbrandsen N, Lindberg G, et al; Scandinavian MDS Group. A validated decision model for treating the anaemia of myelodysplastic syndromes with erythropoietin + granulocyte colony-stimulating factor: significant effects on quality of life. *Br J Haematol*. 2003;120(6):1037-1046.
- Kelaidi C, Park S, Sapena R, et al. Long-term outcome of anemic lower-risk myelodysplastic syndromes without 5q deletion refractory to or relapsing after erythropoiesis-stimulating agents. *Leukemia*. 2013;27(6):1283-1290.

© 2013 by The American Society of Hematology

To the editor:**Impact of lead intoxication in children with iron deficiency anemia in low- and middle-income countries**

Recently, Pasricha et al¹ reported on the control of iron deficiency anemia in low- and middle-income countries.

The article discusses common contributory etiologies ranging from nutritional to infectious causes for iron deficiency anemia in the pediatric age group. Given that lead ingestion exacerbates iron deficiency,²⁻⁴ may accelerate the cognitive injuries of iron deficiency,⁵ and may also confound the diagnosis of microcytic anemia, we were surprised that the authors did not mention lead in their article.

Clune et al⁶ found that lead intoxication is remarkably common in many areas in the world including those described by the authors. Our own pilot study, conducted in a semiurban Indian setup, indicated strikingly high prevalence rates of lead intoxication. Fifty-six percent of the children in the age group of 0 to 6 years had a mean blood lead level (BLL) above 5 mcg/dL, with close to 10% of children in this age group having BLL dangerously above 15 mcg/dL.

Similar results were reported^{7,8} in separate moderate to large population-based studies from different parts of India with an average BLL in toxic ranges, well above the upper limit set by the World Health Organization and the Centers for Disease Control and Prevention for acceptable BLLs.⁹

Increased existence of hotspots of lead exposure in the South Asian region exist, but the lack of universal lead-screening programs in the developing world underrecognizes lead poisoning as a significant health hazard. Given the geographical overlap, it is likely that many of the patients in the study of Pasricha et al had concomitant lead poisoning.

Given the value of the authors' discussion in creating awareness and perhaps initiating legislation on the global level, we think the likely contribution of lead intoxication should be brought to the attention of investigators and public health officials.

Akshat Jain

Pediatric Hematology and Oncology,
Steven and Alexandra Cohen Children's Medical Center of New York,
New Hyde Park, NY

Lawrence C. Wolfe

Pediatric Hematology and Oncology,
Steven and Alexandra Cohen Children's Medical Center of New York,
New Hyde Park, NY

Ginee Jain

Department of General Dentistry,
Boston University School of Dentistry,
Boston, MA

Contribution: A.J. reviewed the literature and provided original research input; A.J. and L.C.W. wrote the manuscript; and G.J. provided statistics for original research.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Akshat Jain, Pediatric Hematology and Oncology, Steven and Alexandra Cohen Children's Medical Center of New York, 26901 76th Ave, New Hyde Park, NY 11040; e-mail: ajain12@nshs.edu.

References

- Pasricha SR, Drakesmith H, Black J, Hipgrave D, Biggs BA. Control of iron deficiency anemia in low- and middle-income countries. *Blood*. 2013;121(14):2607-2617.
- Clark M, Royal J, Seeler R. Interaction of iron deficiency and lead and the hematologic findings in children with severe lead poisoning. *Pediatrics*. 1988;81(2):247-254.
- Zimmermann MB, Muthayya S, Moretti D, Kurpad A, Hurrell RF. Iron fortification reduces blood lead levels in children in Bangalore, India. *Pediatrics*. 2006;117(6):2014-2021.
- Choi JW, Kim SK. Association between blood lead concentrations and body iron status in children. *Arch Dis Child*. 2003;88(9):791-792.



blood

2013 122: 2286-2288
doi:10.1182/blood-2013-07-512442

Can the revised IPSS predict response to erythropoietic-stimulating agents in patients with classical IPSS low or intermediate-1 MDS?

Valeria Santini, Jennifer Schemenau, Alessandro Levis, Enrico Balleari, Rosa Sapena, Lionel Adès, Agnès Guerci, Odile Beyne-Rauzy, Marie-Pierre Gourin, Stephane Cheze, Aspasia Stamatoullas, Alessandro Sanna, Daniela Gioia, Gianni Cametti, Dario Ferrero, Emmanuel Raffoux, Christian Rose, Antonella Poloni, Thomas Prebet, Laurence Legros, Shanti Natarajan-Amé, Pierre Fenaux, Ulrich Germing, François Dreyfus and Sophie Park

Updated information and services can be found at:
<http://www.bloodjournal.org/content/122/13/2286.full.html>

Articles on similar topics can be found in the following Blood collections
[Myeloid Neoplasia](#) (1303 articles)

Information about reproducing this article in parts or in its entirety may be found online at:
http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:
<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:
<http://www.bloodjournal.org/site/subscriptions/index.xhtml>