

Mosaic genome of endobacteria in arbuscular mycorrhizal fungi: Transkingdom gene transfer in an ancient mycoplasma-fungus association

Gloria Torres-Cortés^{a,1}, Stefano Ghignone^b, Paola Bonfante^c, and Arthur Schüßler^a

^aDepartment of Biology, Ludwig-Maximilians-University Munich, 82152 Martinsried, Germany; ^bInstitute for Sustainable Plant Protection, Torino Unit, National Research Council, 10125 Torino, Italy; ^cDepartment of Life Sciences and Systems Biology, University of Torino, 10125 Torino, Italy

Edited by Nancy A. Moran, University of Texas, Austin, TX, and approved April 14, 2015 (received for review January 27, 2015)

For more than 450 million years, arbuscular mycorrhizal fungi (AMF) have formed intimate, mutualistic symbioses with the vast majority of land plants and are major drivers in almost all terrestrial ecosystems. The obligate plant-symbiotic AMF host additional symbionts, so-called *Mollicutes*-related endobacteria (MRE). To uncover putative functional roles of these widespread but yet enigmatic MRE, we sequenced the genome of *DhMRE* living in the AMF *Dentiscutata heterogama*. Multilocus phylogenetic analyses showed that MRE form a previously unidentified lineage sister to the hominis group of *Mycoplasma* species. *DhMRE* possesses a strongly reduced metabolic capacity with 55% of the proteins having unknown function, which reflects unique adaptations to an intracellular lifestyle. We found evidence for transkingdom gene transfer between MRE and their AMF host. At least 27 annotated *DhMRE* proteins show similarities to nuclear-encoded proteins of the AMF *Rhizophagus irregularis*, which itself lacks MRE. Nuclear-encoded homologs could moreover be identified for another AMF, *Gigaspora margarita*, and surprisingly, also the non-AMF *Mortierella verticillata*. Our data indicate a possible origin of the MRE-fungus association in ancestors of the *Glomeromycota* and *Mucoromycotina*. The *DhMRE* genome encodes an arsenal of putative regulatory proteins with eukaryotic-like domains, some of them encoded in putative genomic islands. MRE are highly interesting candidates to study the evolution and interactions between an ancient, obligate endosymbiotic prokaryote with its obligate plant-symbiotic fungal host. Our data moreover may be used for further targeted searches for ancient effector-like proteins that may be key components in the regulation of the arbuscular mycorrhiza symbiosis.

arbuscular mycorrhizal fungi | endobacteria | intracellular symbiont | *Mycoplasma* | horizontal gene transfer

Soil fungi from the *Glomeromycota*, which form arbuscular mycorrhiza (AM) symbioses with the vast majority of land plants, are major players in terrestrial ecosystems (1). This symbiosis originated more than 450 million years ago (2) and represents an impressive example of an ancient and stable mutualistic association: AM fungi (AMF) efficiently explore the soil with their fine mycelium and supply plants with inorganic nutrients and water, whereas the plants provide carbohydrates derived from photosynthesis in above-ground organs.

Interestingly, AMF, as biotrophic and obligate plant symbionts, themselves host additional endosymbionts in their cytoplasm, biotrophic endobacteria (3, 4). These endobacteria were electron-microscopically described in the 1970s as bacterium-like organisms (5). Two types were later defined, the first one being found only in members of the family *Gigasporaceae*. This Gram-negative bacterium was named *Candidatus Glomeribacter gigasporarum* (*CaGg*) and is related to *Burkholderia* (4). The second and much more widespread type represents the only known fungal endobacteria belonging to the *Mollicutes* (“*Mollicutes*-related endobacteria”; MRE), although related endobacteria (e.g., *Mycoplasma* and *Phytoplasma* species) are widespread as biotrophic parasites of humans, mammals, reptiles, fishes, arthropods, or plants. MRE

were frequently detected in the intraradical and extraradical mycelium and in spores of AMF; however, they could never be detected free-living (3). Strikingly, MRE have recently been demonstrated to also occur in several non-AMF species from the genus *Endogone* (*Mucoromycotina*), where some members are also plant symbionts (6).

MRE are associated with all major phylogenetic lineages of AMF studied so far and, thus, indirectly also with more than 80% of all land plants. They are coccoid, located in the cytoplasm without a surrounding host-membrane, and appear to possess a Gram-positive cell wall, which is surprising because of the phylogenetic relationship with cell wall-lacking *Mollicutes* (3). During their long-lasting coevolution, MRE have formed distinct, monophyletic evolutionary lineages within their fungal hosts, with a 16S rRNA gene (16S) sequence divergence of up to 20% (3, 7).

We hypothesized that MRE play an important biological role in AM, consistent with the observation that they have been maintained as ancient endosymbionts in major evolutionary AMF lineages that separated hundreds of million years ago. To obtain hints for such roles and to shed light on MRE evolution, we

Significance

Obligate plant-symbiotic, arbuscular mycorrhizal fungi (AMF) are major drivers of terrestrial ecosystems and host enigmatic *Mollicutes*-related endobacteria (MRE) in their cytoplasm. The genome analysis of a MRE living in the AMF *Dentiscutata heterogama* revealed it to represent a previously unidentified bacterial lineage of *Mycoplasma*-related species. *DhMRE* shows strongly reduced metabolic capacity and underwent transkingdom gene transfer: its genome codes for an arsenal of eukaryotic-like putative effector proteins, with nuclear encoded homologues in AMF and *Mortierella*. The MRE-fungus (-plant) association probably evolved in ancestors of *Glomeromycota* and *Mucoromycotina*. This calls for a targeted search for ancient effector proteins that play crucial roles in the MRE interaction with fungal hosts, and putatively also with plants.

Author contributions: G.T.-C., S.G., P.B., and A.S. designed research; G.T.-C. performed research; A.S. contributed new reagents/analytic tools; G.T.-C., S.G., P.B., and A.S. analyzed data; and G.T.-C. and A.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The *DhMRE* genome assembly and annotation were deposited at the EMBL-EBI European Nucleotide Archive (accession no. PRJEB8356). Alignments and phylogenetic trees were deposited at TreeBase under submission ID 17164 for the multilocus analysis (Fig. 1), 17266 for the protein kinase analysis (Fig. 2), and 17268 for the AIG domain protein analysis (SI Appendix, Fig. S6). *Gigaspora margarita* proteins used for phylogenetic analyses are part of the *G. margarita* BEG34 Transcriptome Shotgun Assembly project (BioProject PRJNA267628; Biosamples SAMN03216569-SAMN03216586), which was deposited at DDBJ/EMBL/GenBank (accession no. GBYF00000000). The sequences used in the analyses are GBYF01079662, GBYF01002907, GBYF01035393, GBYF01077659, GBYF01035394, GBYF01010756, GBYF01025162, and GBYF01016789 and belong to the first TSA version, GBYF01000000.

¹To whom correspondence should be addressed. Email: gloria.torres@bio.lmu.de.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1501540112/-DCSupplemental.

analyzed the genome of a MRE colonizing the AMF *Denticutata heterogama* FL654 (syn.: *Scutellospora heterogama*; *Gigasporaceae*), termed *DhMRE*. Only three genomes of fungal endobacteria have been published so far: from *CaGg*, which colonize some AMF from the *Gigasporaceae* (8); from *Burkholderia rhizoxinica*, the endobacterium of *Rhizopus microsporus*, which participates in the production of rhizoxin (a potent antimitotic agent that acts as a virulence factor) (9); and from a betaproteobacterial endosymbiont associated with *Mortierella elongata* (10).

We present the annotation of a MRE genome draft, the phylogenetic placement of MRE based on a multilocus analysis, the evolutionary implications about the origin of MRE-fungus (-plant) associations, and evidence for transkingdom horizontal gene transfer (HGT). Moreover, the genome presents genes coding for proteins with likely regulatory functions in the interaction between this obligate symbiotic prokaryote and its obligate plant-symbiotic fungal host.

Results

General Features of the *DhMRE* Genome. To study the genomic organization and gene repertoire of an MRE living in AMF, we isolated the DNA of the endobacterial population from spores of *D. heterogama* FL654, which harbors MRE but not *CaGg* (*SI Appendix*, Fig. S1). Following Illumina sequencing and PCR based gap-closing approaches, three *DhMRE* genomic scaffolds could be defined. One is circular, with a size of 648,879 bp and 34.1% GC content (scaffold A), the second (scaffold B) is 60,375 bp in size with a GC content of 32.3%, and the third (scaffold C) is a putative plasmid of 3,779 bp and 34.1% GC content (*SI Appendix*, Fig. S2 and Tables S1 and S2). Tetranucleotide frequencies analyses validated the origin of scaffolds A and B (*SI Appendix*, Table S3). The total size of the *DhMRE* genome draft assembly is 713,033 bp, with an estimated sequence coverage of approximately 130-fold.

Most of the studied AMF possess two major phylotypes of MRE 16S rRNA genes co-occurring in the same spore (3, 7). The analysis of 16S rRNA gene PCR-amplicon derived clone libraries from *D. heterogama* also revealed these two major phylotypes (*SI Appendix*, Fig. S3). Phylotype I was more abundant (83%) than phylotype II (17%). The dominance of phylotype I in clone libraries rose to 96% after applying the genomic DNA extraction protocol developed for *DhMRE* (*SI Appendix*, Table S4), explaining why phylotype I was almost exclusively identified in the Illumina dataset (*SI Appendix*, Fig. S4). We speculate that the phylotype I hosting MRE might possess a less rigid cell envelope leading to digestion of its DNA in the DNase treatment applied. To ensure that reads from different phylotypes were not co-assembled, we performed a QualitySNPng analysis (11) on the contig with the 16S rRNA gene (which is organized in a split rRNA operon). Only when relaxing the parameters (“minimal percentage of reads per allele” set to 15%), four haplotypes were discovered, but their very low read composition does not support the existence of a second phylotype in the assembled sequences.

Thus, the genome assembly presented here carries only one of the two major MRE 16S phylotypes and, therefore, putatively only one of the corresponding MRE genomes found in *D. heterogama*. This reduction in complexity enabled us to close most of the genome assembly.

Evolutionary Placement. To determine the phylogenetic placement of *DhMRE* as a representative of the monophyletic MRE clade (3), a maximum likelihood phylogenetic analysis of amino acid sequences from 10 housekeeping genes from 38 bacteria was conducted (listed in *SI Appendix*, Table S5). The MRE lineage was firmly placed in the *Mycoplasmatales* (Fig. 1), most likely being closest related to the hominis group of *Mycoplasma* species.

Transkingdom Horizontal Gene Transfer Between *DhMRE* and the Fungal Host. The analysis of the annotated genes has revealed that several *DhMRE* proteins share similarity with proteins coded by

the nuclear genome of *Rhizophagus irregularis*, which is the only AMF with a sequenced genome (12, 13). All evaluations to date indicate that this fungus does not host endobacteria and separated evolutionary from *D. heterogama* more than 200 million years ago (3, 14). In total, we identified 17 proteins in scaffold A and 10 proteins in scaffold B showing best BLAST hits with an identity higher than 30% and query coverage above 35% against *R. irregularis* proteins. Among the 27 proteins, 21 (78%) have predicted functional domains, which could be related to bacteria-host association: eight are nonspecific tyrosine-kinases like proteins (TKLs; two of them located in putative genomic islands), seven contain leucine rich repeats (LRR) domains normally functioning in protein-protein interactions, and six contain an AIG1 domain, usually found in plant proteins thought to be involved in plant resistance to bacteria (15) (*SI Appendix*, Table S6).

We performed phylogenetic analyses to identify the putative origin of the *DhMRE* proteins sharing similarity with nuclear-encoded *R. irregularis* proteins, including homologous nuclear-encoded proteins from *Gigaspora margarita* BEG34, an AMF not hosting MRE (Fig. 2 and *SI Appendix*, Fig. S5). The phylogenetic analysis of the *DhMRE* TKLs show them to robustly cluster together with protein kinases from the AMF *R. irregularis*, *G. margarita*, and also from the non-AMF *Mortierella verticillata* (*Mortierellales*). This clade is clearly separated from other prokaryotic and eukaryotic protein kinases (Fig. 2 and *SI Appendix*, Fig. S5).

The other main group of *DhMRE* proteins similar to nuclear-encoded proteins of *R. irregularis* possesses an AIG1 domain. *DhMRE*, *Rhizophagus* and *Gigaspora* AIG1 domains also form a monophyletic clade. The only other AIG1 domains that also clustered in this clade are from the soil amoeba *Dictyostelium fasciculatum*. According to the phylogenetic analysis, the ancestor of these domains is eukaryotic, but not fungal (*SI Appendix*, Fig. S6).

Proteins with Eukaryotic Domains. Analysis of the *DhMRE* genome revealed an exceptionally large arsenal of proteins (including some with homologs to nuclear-encoded *R. irregularis* proteins) with eukaryotic-like domains that are likely to interfere with eukaryotic cellular functions. Several of the coding genes are located in putative genomic islands and/or near transposases. At least 29 proteins carry domains predominantly found in eukaryotic proteins (4% of the total proteome). The most abundant eukaryotic domains are LRR (*SI Appendix*, Fig. S7), which are versatile binding motifs involved in protein-protein interactions and in the interaction of microbes with host cells (16). One gene in scaffold B codes for a protein with a heterokaryon incompatibility domain (HET domain), which is part of the nonself recognition systems in ascomycetes (17).

The *DhMRE* proteome also contains proteins related to the eukaryotic ubiquitination system that controls the degradation of proteins by the proteasome and may interfere with host transcriptional gene activation or repression, nuclear transport and enzyme activity (*SI Appendix*, Fig. S7). We found three proteases with a small ubiquitin-like modifier (SUMO)/sentrin domain, reversibly regulating the SUMO modification pathway in eukaryotes (18), one of them predicted to be secreted. Another protein predicted to be secreted carries an ovarian tumor (OTU)-like cysteine protease domain, which is also putatively involved in the ubiquitination system.

Overall, *DhMRE* encodes a large repertoire of proteins with eukaryotic domains, some of them already characterized in prokaryotic effector proteins. Moreover, a subset of them has probably gone through HGT with the host (*SI Appendix*, Table S6).

Functional Annotation of Predicted Protein-Coding Genes. The gene repertoire of *DhMRE* comprises 90 of the 100 essential cluster of orthologous groups (COG) conserved in 99% of bacteria (*SI Appendix*, Table S7) (19). The COG functional categories with




Fig. 2. Maximum likelihood phylogenetic reconstruction of the protein kinases candidates for horizontal gene transfer (A) and taxonomic composition of the *DhMRE* scaffolds according to sequence similarity (B). BLASTP-based affiliation was based on the best-hit with a cutoff value of e^{-03} . For each protein, BLASTP hits in the nonredundant database were recovered by using the program Blast2GO, and the classification of the corresponding organism was extracted according to NCBI taxonomy. SA, scaffold A; SB, scaffold B.

and interactions with their fungal host. The *DhMRE* genome revealed HGT events between the fungal host and MRE, an arsenal of proteins with eukaryotic domains that are likely involved in host–cell interaction and, despite the much-reduced genome, a large repertoire of unknown, unique proteins. Furthermore, *DhMRE* has extremely reduced metabolic capacities. This set of biotrophic adaptations raises new questions about the role of these endobacteria in the evolution of the fungal host and about transkingdom interactions of symbionts.

Evolution of *DhMRE*. *DhMRE* belongs to a bacterial lineage closely related to the pneumoniae and hominis mycoplasma groups, which mainly occur as biotrophic extracellular parasites of animals (23). The *DhMRE* gene homologs we detected in the nuclear genome of *M. verticillata* (*Mortierellales*), together with the recent findings of MRE in *Endogone* (*Endogonales*) species (24), indicate that the MRE association with fungi was already present in a common ancestor of the *Glomeromycota* and *Mucoromycotina* (25).

The phylogenetic position of *DhMRE* in-between mycoplasma groups indicates that its ancestor had shifted from a eukaryotic, probably animal, host to a common ancestor of the *Glomeromycota* and *Mucoromycotina*, resulting in the ancient MRE–fungus association. Host shifts are not uncommon for *Mycoplasma* species,

e.g., causing the *Mycoplasma gallisepticum* outbreak in wild house finches (26).

Although MRE are phylogenetically embedded in the *Mycoplasmatales* and neither bacteria from *Mycoplasmatales* nor from lineages branching earlier (*Spiroplasma*, *Phytoplasma*, *Acholeplasma*) possess a cell wall, MRE surprisingly present a structure that was interpreted as being a Gram-positive cell wall (3). However, the nature of the MRE cell wall was never experimentally confirmed and the genome data presented here indicate that this structure does not represent a prokaryotic murein sacculus.

Mechanisms Promoting *DhMRE* Evolution and Diversification. The *DhMRE* draft genome lacks some genes for DNA repair, for example *ung*, coding for uracil-DNA glycosylase, which is in line with the suggestion that species of the *Mycoplasmatales* are constitutive mutators because of the lack of some components of the SOS response and DNA repair machinery (27, 28). High mutation rates also play an important role in adaptation processes and may have helped the last common ancestor of the MRE in its shift to a fungal host.

Another important mechanism for rapid evolution and adaptation in *Mycoplasmatales* is HGT (29, 30). It appears to be also a major driving force in MRE evolution. Transkingdom HGT events between MRE and AMF could have helped MRE to use acquired eukaryotic genes to exploit fungal metabolism or other functions, but it could also have been a key element enabling the fungi to control the plant host, as indicated by the fact of such genes being kept in the MRE-free *R. irregularis*. Horizontal gene transfer and evolution of eukaryotic-like proteins to effector proteins have been proposed to enable the intracellular life of *L. pneumophila* (31). A similar scenario may have happened in the ancestral MRE–AMF association, or in even more ancestral fungi, like such from the *Mucoromycotina*.

A major group of genes that underwent HGT between AMF and *DhMRE* encode proteins related to signal transduction, in




Fig. 3. Functional classification of annotated protein-coding genes for the scaffolds A (A) and B (B). The functional categorization of each protein-coding gene was classified according to the cluster of orthologs groups (COG assignments). SA, scaffold A; SB, scaffold B.

particular TKLs. The origin of these TKLs is uncertain, but the sequencing of genomes from more AMF and *Endogone*- and *Mortierella*-related fungi and their MRE will clarify this point. The nuclear genome of *R. irregularis* showed TKL gene family expansions, and several TKL-containing proteins are strongly expressed in germinating spores and intraradical mycelia, indicating their functional relevance (12). The roles that the protein tyrosine kinases that underwent HGT, and which have homologs in the nuclear genomes of species from the *Glomeromycota* and *Mortierellales*, might play in signal transduction in early diverging fungal lineages becomes a highly interesting question.

The other major group of genes likely to have undergone HGT between AMF and *DhMRE* encode proteins with an AIG1 domain, which appears to be involved in plant resistance to bacteria. The surprising finding of AIG1 homologs existing in the soil-living amoeba *D. fasciculatum* makes it tempting to speculate about an early association between the ancestors of MRE and amoeba. In this context, it may be noted that it was suggested that *L. pneumophila* evolved some of its host-interaction systems in associations with amoeba (32).

Phylogenetic analyses indicated different possible scenarios for MRE HGT and evolution. Perhaps the most parsimonious scenario that we can envisage is an ancestral HGT from animals to biotrophic MRE ancestors, with subsequent HGT to the AMF host in the AM symbiosis, followed by a loss of the MRE symbionts in some AMF, such as certain members of the genera *Rhizophagus* and *Gigaspora*. The alternative of an ancestral HGT from animals or amoeba to the fungi followed by HGT from the fungi to the bacteria appears less likely. The sequencing of more MRE and AMF genomes will shed more light on this open question.

DhMRE also possess several genes encoding eukaryotic-like proteins carrying domains known from effector proteins. For example, SUMO and OTU proteases that could interfere with the host ubiquitination system. Different microbial effector proteins targeting the host ubiquitination system have been characterized, e.g., the effectors XopD, AvrXv4, and AvrBsT from *Xanthomonas*, and YopJ from *Yersinia* mimic SUMO protease activity, disrupting the regulation of the SUMO pathway and altering the activity or stability of the host proteins (33). *DhMRE*, or their fungal host, might use such proteins with eukaryotic domains to communicate with or regulate host processes.

In addition to the proteins described above, the proteome of *DhMRE* also contains a high proportion (1.5%) of proteins with LRR domains, similar to intracellular bacteria associated with amoebae (1.3%) (34), but higher than in *Wolbachia* or *Rickettsia* endobacteria (usually below 0.25%). These proteins are also candidates for playing a role in the interaction between *DhMRE* and the fungal host.

Taking all together, *DhMRE* genome sequencing has highlighted its plasticity, with both localized (mutations) and broad (genome reduction, HGT) processes altering its genetic content, in close coevolution with its hosts.

Overall Traits of the Small, *Mycoplasma*-Like *DhMRE* Genome. The *DhMRE* genome shares a set of characteristics with small genomes of endobacteria, reflecting a high level of adaptation to an obligate intracellular lifestyle. For example, the absence of glycolysis, the tricarboxylic acid cycle, and a functional respiratory chain together indicate that *DhMRE* could generate energy only by substrate-level phosphorylation through the degradation of substrates obtained from its host, as the amoeba symbiont *Candidatus* Amoebophilus asiaticus (34). Another trait occasionally associated to intracellular symbionts (35) is the absence of the *dnaA* gene, which has been considered as an indication of a more direct control of DNA replication and regulation of endosymbionts propagation by the host.

In conclusion, the *DhMRE* genome possesses typical features of members of the *Mollicutes*, such as low GC content, a reduced

genome size, high gene density, and 55% of its genes coding for PUF, which is even more than normally found in other species from the *Mollicutes* (36, 37). In our opinion, the occurrence of MRE in evolutionary lineages of AMF that separated more than 400 million years ago points to a mutualistic association, but it remains unknown how the MRE impact the AMF or, for that matter, the plant host. The large number of PUF as yet may hide important functions of the MRE in the symbiosis. We found transkingdom HGT of putatively regulatory genes of which homologs are retained in MRE-free AMF and, in general, a large arsenal of proteins with eukaryotic domains. Thus, our study may open the way for a more targeted search for effectors playing roles in AM. Taking the ancestral nature of the MRE–fungus association into account, the analysis of putative effector proteins with eukaryotic like domains and derived from HGT might lead to the identification of crucial effectors in bacteria–fungus–plant associations.

Material and Methods

Biological Material. *D. heterogama* FL654 spores hosting MRE but not CaGg were used for genome sequencing. FL654 was produced in vitro in monoxenic culture on root organs by The Energy and Resources Institute. The spores were handled in a sterile manner under a laminar flow hood.

Endobacteria DNA Extraction for Sequencing. For *DhMRE* genomic DNA preparation, a new protocol was established, attempting to enrich the *DhMRE* fraction and to remove AMF nuclear DNA (protocol details in *SI Appendix, SI Materials and Methods*). To obtain 50 ng of DNA, ~300,000 spores from *D. heterogama* were used. The noted DNA isolation procedure was specifically designed with the aim to isolate intact small and rigid appearing endobacteria. Shearing and osmotic forces were used to rupture nuclei and mitochondria, the DNA of which was then removed by Dnase digestion. In this process, we seem to have selectively enriched only one of the two MRE phylotypes known from previous Sanger sequencing approaches. In a semiquantitative approach to study this aspect, we constructed and analyzed clone libraries (details in *SI Appendix, SI Materials and Methods*).

DNA Sequencing and Sequence Assembly. Illumina sequencing was performed by using the Illumina MiSeq platform at the Genomics Service Unit of the Ludwig-Maximilian-University Munich Biocenter, generating 41×10^6 paired end 150-bp raw reads. The paired-end reads were quality trimmed by using CLC workbench v5 (CLC Bio), under the default parameters. Trimmed reads were mapped against the three main bacterial contaminant (*Delftia acidovorans*, *Stenotrophomonas maltophilia*, *Achromobacter xylosoxidans*) genomes to remove contaminant reads. The remaining, cleaned reads were assembled by using the CLC de novo assembly algorithm, identifying 24 *DhMRE* contigs (details in *SI Appendix, SI Materials and Methods*). After assembly, the 24 remaining gaps were closed by using conventional directed PCR and primer-walking approaches followed by Sanger sequencing.

Annotation and Metabolic Reconstruction. The *DhMRE* genome was integrated into the MicroScope platform to perform automatic and manual annotation and comparative analysis with other species from *Mycoplasmatales* and *Firmicutes*. AMIGene software was used to predict coding DNA sequences (CDSs). More information on the syntactic and functional annotation process is given in ref. 38.

Phylogenetic Analyses. For the multilocus phylogenetic reconstruction, 10 housekeeping genes (single loci genes coding for eight ribosomal proteins, the translation initiation factor IF-2, and the DNA-directed RNA polymerase subunit beta) often used for multilocus analyses (8) were selected from 38 bacteria (listed in *SI Appendix, Table S5*). The sequences were manually concatenated. The final length of the concatenated sequences was 2,900 aa after removing ambiguously aligned positions.

To study *DhMRE* proteins candidate for HGT by phylogenetic analyses, homologs were selected from the NCBI database (details in *SI Appendix, SI Materials and Methods*). For all phylogenetic analyses, protein sequences were aligned by using MAFFT under default parameters (39). JalView (40) was used to manually remove ambiguously aligned positions and regions. Highly divergent sequences were also removed from the alignment. Only the conserved parts of the alignment were selected for the maximum likelihood analysis. Best-fitting evolutionary models for the alignment were determined by its Bayesian information

criterion values with ProtTest version 3.4. Maximum likelihood phylogenetic analyses were computed through the CIPRES web portal with RAxML-HPC2 on XSEDE (41) with the best-fit model by using 100 bootstraps.

Accession Numbers and Alignments. Alignments and phylogenetic trees were deposited at TreeBase under submission ID 17164 for the multilocus analysis (Fig. 1), 17266 for the protein kinase analysis (Fig. 2), and 17268 for the AIG domain protein analysis (SI Appendix, Fig. S6).

The *DhMRE* genome assembly and annotation were deposited at the European Molecular Biology Laboratory (EMBL)-European Bioinformatics (EBI) European Nucleotide Archive under accession no. PRJEB8356.

- van der Heijden MGA, Bardgett RD, van Straalen NM (2008) The unseen majority: Soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol Lett* 11(3):296–310.
- Schüßler A, Walker C (2011) Evolution of the 'Plant-Symbiotic' Fungal Phylum, *Glomeromycota*. *Evolution of Fungi and Fungal-Like Organisms, The Mycota*, eds Pöggeler S, Wöstemeyer J (Springer, Berlin), pp 163–185.
- Naumann M, Schüßler A, Bonfante P (2010) The obligate endobacteria of arbuscular mycorrhizal fungi are ancient heritable components related to the *Mollicutes*. *ISME J* 4(7):862–871.
- Bianciotto V, Lumini E, Bonfante P, Vandamme P (2003) 'Candidatus glomeribacter gigasporarum' gen. nov., sp. nov., an endosymbiont of arbuscular mycorrhizal fungi. *Int J Syst Evol Microbiol* 53(Pt 1):121–124.
- Mosse B (1970) Honey-coloured, sessile *Endogone* spores: II. Changes in fine structure during spore development. *Arch Microbiol* 74(2):129–145.
- Desirò A, Faccio A, Kaech A, Bidartondo MI, Bonfante P (2015) *Endogone*, one of the oldest plant-associated fungi, host unique *Mollicutes*-related endobacteria. *New Phytol* 205(4):1464–1472.
- Desirò A, et al. (2014) Detection of a novel intracellular microbiome hosted in arbuscular mycorrhizal fungi. *ISME J* 8(2):257–270.
- Ghignone S, et al. (2012) The genome of the obligate endobacterium of an AM fungus reveals an interphylum network of nutritional interactions. *ISME J* 6(1):136–145.
- Partida-Martinez LP, Hertweck C (2005) Pathogenic fungus harbours endosymbiotic bacteria for toxin production. *Nature* 437(7060):884–888.
- Fujimura R, et al. (2014) Draft genome sequence of the betaproteobacterial endosymbiont associated with the fungus *Mortierella elongata* FMR23-6. *Genome Announc* 2(6):e01272-14.
- Nijveen H, van Kaauwen M, Esselink DG, Hoegen B, Vosman B (2013) QualitySNPng: A user-friendly SNP detection and visualization tool. *Nucleic Acids Res* 41(web server issue, W1):W587–590.
- Tisserant E, et al. (2013) Genome of an arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis. *Proc Natl Acad Sci USA* 110(50):20117–20122.
- Lin K, et al. (2014) Single nucleus genome sequencing reveals high similarity among nuclei of an endomycorrhizal fungus. *PLoS Genet* 10(1):e1004078.
- Redecker D, Kodner R, Graham LE (2000) Glomalean fungi from the Ordovician. *Science* 289(5486):1920–1921.
- Reuber TL, Ausubel FM (1996) Isolation of *Arabidopsis* genes that differentiate between resistance responses mediated by the *RPS2* and *RPM1* disease resistance genes. *Plant Cell* 8(2):241–249.
- Marino M, Braun L, Cossart P, Ghosh P (2000) A framework for interpreting the leucine-rich repeats of the *Listeria internalins*. *Proc Natl Acad Sci USA* 97(16):8784–8788.
- Glass NL, Kaneko I (2003) Fatal attraction: Nonself recognition and heterokaryon incompatibility in filamentous fungi. *Eukaryot Cell* 2(1):1–8.
- Hickey CM, Wilson NR, Hochstrasser M (2012) Function and regulation of SUMO proteases. *Nat Rev Mol Cell Biol* 13(12):755–766.
- Merhej V, Royer-Carenzi M, Pontarotti P, Raoult D (2009) Massive comparative genomic analysis reveals convergent evolution of specialized bacteria. *Biol Direct* 4(1):13.
- Finn RD, Clements J, Eddy SR (2011) HMMER web server: Interactive sequence similarity searching. *Nucleic Acids Res* 39(web server issue, suppl 2):W29–W37.
- Nishino K, Yamaguchi A (2008) Role of xenobiotic transporters in bacterial drug resistance and virulence. *IUBMB Life* 60(9):569–574.
- Tian C, et al. (2010) Regulation of the nitrogen transfer pathway in the arbuscular mycorrhizal symbiosis: Gene characterization and the coordination of expression with nitrogen flux. *Plant Physiol* 153(3):1175–1187.
- Rosengarten R, et al. (2000) Host-pathogen interactions in mycoplasma pathogenesis: Virulence and survival strategies of minimalist prokaryotes. *Int J Med Microbiol* 290(1):15–25.
- Nadimi M, Beaudet D, Forget L, Hijri M, Lang BF (2012) Group I intron-mediated trans-splicing in mitochondria of *Gigaspora rosea* and a robust phylogenetic affiliation of arbuscular mycorrhizal fungi with *Mortierellales*. *Mol Biol Evol* 29(9):2199–2210.
- Hibbett DS, et al. (2007) A higher-level phylogenetic classification of the *Fungi*. *Mycol Res* 111(Pt 5):509–547.
- Delaney NF, et al. (2012) Ultrafast evolution and loss of CRISPRs following a host shift in a novel wildlife pathogen, *Mycoplasma gallisepticum*. *PLoS Genet* 8(2):e1002511.
- Carvalho FM, et al. (2005) DNA repair in reduced genome: The *Mycoplasma* model. *Gene* 360(2):111–119.
- Citti C, Blanchard A (2013) Mycoplasmas and their host: Emerging and re-emerging minimal pathogens. *Trends Microbiol* 21(4):196–203.
- Sirand-Pugnet P, et al. (2007) Being pathogenic, plastic, and sexual while living with a nearly minimal bacterial genome. *PLoS Genet* 3(5):e75.
- Sirand-Pugnet P, Citti C, Barré A, Blanchard A (2007) Evolution of mollicutes: Down a bumpy road with twists and turns. *Res Microbiol* 158(10):754–766.
- Al-Quadani T, Price CT, Abu Kwaik Y (2012) Exploitation of evolutionarily conserved amoeba and mammalian processes by *Legionella*. *Trends Microbiol* 20(6):299–306.
- Molmeret M, Horn M, Wagner M, Santic M, Abu Kwaik Y (2005) Amoebae as training grounds for intracellular bacterial pathogens. *Appl Environ Microbiol* 71(1):20–28.
- Hotson A, Mudgett MB (2004) Cysteine proteases in phytopathogenic bacteria: Identification of plant targets and activation of innate immunity. *Curr Opin Plant Biol* 7(4):384–390.
- Schmitz-Esser S, et al. (2010) The genome of the amoeba symbiont "Candidatus Amoebophilus asiaticus" reveals common mechanisms for host cell interaction among amoeba-associated bacteria. *J Bacteriol* 192(4):1045–1057.
- Nakabachi A, et al. (2006) The 160-kilobase genome of the bacterial endosymbiont *Carsonella*. *Science* 314(5797):267.
- Chambaud I, et al. (2001) The complete genome sequence of the murine respiratory pathogen *Mycoplasma pulmonis*. *Nucleic Acids Res* 29(10):2145–2153.
- Tran-Nguyen LTT, Kube M, Schneider B, Reinhardt R, Gibb KS (2008) Comparative genome analysis of "Candidatus Phytoplasma australiense" (subgroup tuf-Australia I; rp-A) and "Ca. Phytoplasma asteris" Strains OY-M and AY-WB. *J Bacteriol* 190(11):3979–3991.
- Vallenet D, et al. (2013) MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res* 41(Database issue):D636–D647.
- Katoh K, Kuma K, Toh H, Miyata T (2005) MAFFT version 5: Improvement in accuracy of multiple sequence alignment. *Nucleic Acids Res* 33(2):511–518.
- Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ (2009) Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25(9):1189–1191.
- Stamatakis A (2006) RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22(21):2688–2690.