

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Prospective molecular monitoring of minimal residual disease after nonmyeloablative allografting in newly diagnosed multiple myeloma

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1507008> since 2015-08-26T07:06:09Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Prospective molecular monitoring of minimal residual disease after nonmyeloablative allografting in newly diagnosed multiple myeloma

Benedetto Bruno, MD, PhD¹, Simone Ferrero, MD¹, Daniela Drandi, PhD¹, Moreno Festuccia, MD¹, Francesca Patriarca, MD², Nicola Mordini, MD³, Silvia Cena, PhD¹, Daniela Barbero, PhD¹, Luigia Monitillo, PhD¹, Federica Ferrando MD¹, Luisa Giaccone, MD, PhD¹, Renato Fanin, MD², Roberto Passera, PhD⁴, Antonio Palumbo, MD¹, Mario Boccadoro, MD¹, Paola Omedè, PhD¹ and Marco Ladetto, MD¹

¹Division of Hematology, University of Torino, A.O.U. Città della Salute e della Scienza, Torino, Italy

²Haematological Clinic, Azienda Ospedaliera Universitaria, Udine, Italy

³Hematology, Azienda Ospedaliera S. Croce e Carle, Cuneo, Italy

⁴Division of Nuclear Medicine, Statistical Consultant, University of Torino, Torino, Italy

Data on minimal residual disease (MRD) after tandem autologous-nonmyeloablative allografting (auto-allo) are lacking

Aim of the study. To carry out MRD analyses by nested qualitative PCR and real time quantitative (RQ) PCR on newly diagnosed MM patients treated with auto-allo

Methods. Twenty-six patients with a diagnostic bone marrow (BM) specimen suitable for immunoglobulin heavy-chain gene rearrangement (IGH) sequencing were evaluated for MRD by PCR methods. Auto-allo consisted of an autograft followed by 200 cGy TBI and an allograft. BM samples were collected at diagnosis, after the autograft, at month 1, 3, 6 after the allograft and then every 6 months. Nested-PCR and RQ-PCR analyses were carried out using patient-specific primers. FullMR and StandardMR indicated MRD negativity on two consecutive samples by nested-PCR or RQ-PCR respectively

Results. In 19/26 patients had a molecular marker. At a median follow-up of 10 years (4.4-12) from diagnosis and 8.9 years (3.5-11) from the allograft, overall survival (OS) was 61% and median time-to-progression (TTP) 5.6 years. TRM was 16%. MRD studies showed that after the autograft 3/19 patients (16%) were negative by nested-PCR. After the allograft PCR-negativity rates gradually increased to 4/18 (22%) at 1 and 3 months, 7/17 (41%) at 6 months and 8/15 (53%) at 1 year post-transplant. Overall, 8 patients achieved FullMR at a median time from allograft of 6 months (1-12) and for a median duration of 33 months (6-102). Overall, 8 relapses occurred, 6 in the 11 patients who never achieved FullMR and 2 in patients who reached FullMR: one showed molecular relapse, with persistent CCR, after 3 years and the other after 6 months from the last PCR-negative sample. Patients in FullMR had better median TTP (not reached vs 1.6 years, $p=0.043$) and OS (not reached vs 3.3 years, $p=0.008$) than patients who did not achieve FullMR. StandardMR occurred in 12/19 patients (63%) during the first 24 months post-transplant, at a median time of 2 months (1-18) and for a median duration of 27 months (3-102). Patients in StandardMR showed better median TTP (not reached vs 1 year, $p=0.005$) and OS (not reached vs 3.3 years, $p=0.031$) as compared to patients with positive PCR. There was no correlation with chronic graft-vs-host disease suggesting specific graft-vs-myeloma

Conclusions. Auto-allo induces high molecular remission rates, significantly associated with better TTP and OS, that indicate potentially curative graft-vs-myeloma