



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

# Chemical characterisation and cytotoxic effects in A549 cells of urban-air PM10 collected in Torino, Italy.

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/73135 since
Published version:
DOI:10.1016/j.etap.2009.12.005
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in:

# [Environmental toxicology and pharmacology, 29, issue 2, 2010, DOI: 10.1016/j.etap.2009.12.005]

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en), [+ Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect<sup>®</sup> platform]

## Elsevier Editorial System(tm) for Environmental Toxicology and Pharmacology Manuscript Draft

Manuscript Number:

# Title: CYTOTOXIC EFFECTS INDUCED ON A549 CELLS AND CHEMICAL CHARACTERIZATION OF URBAN-AIR PM10 COLLECTED IN TORINO (ITALY).

Article Type: Research Paper

Keywords: Particulate Matter; Cell Proliferation; Cytotoxicity; LDH; Cytokine.

Corresponding Author: Dr. Tiziana Schilirò,

Corresponding Author's Institution: University of Torino

First Author: Tiziana Schilirò

Order of Authors: Tiziana Schilirò; Luca Alessandria; Raffaella Degan; Deborah Traversi; Giorgio Gilli

Abstract: Human type II alveolar cell line (A549) was exposed to aqueous and organic solvent PM10 extracts in order to evaluate cell proliferation, proinflammatory cytokines and cytotoxicity (lactate dehydrogenase, LDH). PM10 samples, collected in Torino (north-west Italy), were analyzed for inorganic chemical species (bioavailable iron and secondary particulates) and endotoxins, potentially inflammatory promoters to human airways.

Meanly, in the considered period, PM10 concentration was  $55.4 \pm 39.1 \mu g/m3$ , iron concentration was  $0.078 \pm 0.095$  and secondary particles constituted  $42 \pm 9$  % of the PM10 total mass. PM10 inhibits cell proliferation and induces both IL-6 and LDH release in a dose and time dependent manner, with a seasonal trend. The different roles of aqueous and organic solvent PM10 extracts demonstrate the importance of particle composition for the biological effects. Nevertheless few significant correlations were found between the biological effects and PM10 components evaluated in this methods development study.

Suggested Reviewers: Ann E. Aust Department of Chemistry and Biochemistry., Utah State University, Logan, UT 84322. AAUST@cc.usu.edu Extensive experience in research of cytotoxic effects of air pollution particulate matter in vitro.

Elisabetta Carraro Università Piemonte Orientale "A. Avogadro" elisabetta.carraro@mfn.unipmn.it Great knowledge of the evaluation of cytotoxicity and genotoxicity in environmental matrices.

Roberto Bono University of Torino roberto.bono@unito.it Great experience in the field of the evaluation of particulate matter genotoxicity.

Eudes Lanciotti University of Florence eudes.lanciotti@unifi.it Great interests in the monitoring and risk assessment of the environmental pollutants.

**Opposed Reviewers:** 

Dear Editor,

We are sending the manuscript *"Cytotoxic effects induced on A549 cells and chemical characterization of urban-air PM10 collected in Torino (Italy)"* that we submit for possible publication on *Environmental Toxicology and Pharmacology.* 

PM10 samples were collected in an urban site of Torino, an industrial north-western Italian city, with high levels of particulate matter among European cities. PM10 samples were analyzed for the amount of inorganic chemical species (bioavailable iron and secondary particulates: sulfates and nitrates) and endotoxins, described as potentially inflammatory promoters to human airways. In order to better understand how ambient particulate matter affects lung health, A549 cells were exposed to aqueous and organic solvent PM10 extracts. Effects on cell proliferation, on the release of proinflammatory cytokines and cytotoxicity, measuring lactate dehydrogenase (LDH) release in the extracellular medium, were evaluated.

In general PM10 inhibits cell proliferation and induces both IL-6 and LDH release in a dose and time dependent manner and these biological effects have a seasonal trend. Few significant correlations were found between the biological effects and PM10 components evaluated in this methods development study. Nevertheless the different roles of aqueous and organic solvent PM10 extracts demonstrate the importance of particle composition for the biological effects.

Best regards

Tiziana Schilirò

1	1	CYTOTOXIC EFFECTS INDUCED ON A549 CELLS AND CHEMICAL
2 3 4	2	CHARACTERIZATION OF URBAN-AIR PM10 COLLECTED IN TORINO (ITALY).
5 6	3	
7 8 0	4	Tiziana Schilirò <sup>1</sup> *, Luca Alessandria <sup>1</sup> , Raffaella Degan <sup>1</sup> , Deborah Traversi <sup>1</sup> , Giorgio
9 10 11	5	Gilli <sup>1</sup>
12 13	6	
14 15 16	7	<sup>1</sup> Department of Public Health and Microbiology, University of Torino, Via Santena 5bis,
10 17 18	8	10126 Torino, Italy.
19 20	9	
21 22	10	* <u>Corresponding Author:</u>
23 24 25	11	Tiziana Schilirò
25 26 27	12	Department of Public Health and Microbiology, University of Torino.
28 29	13	Via Santena, 5bis - 10126 Torino – ITALY
30 31	14	tel. + 39 011 670 5810
32 33 24	15	Fax + 39 011 236 5810
34 35 36	16	e-mail: <u>tiziana.schiliro@unito.it</u>
37 38	17	
39 40	18	
4⊥ 42 43	19	Running head: Biological effects of PM10 induced on A549 cells.
44 45	20	
46 47	21	
48 49	22	
50 51 52	23	
53 54	24	
55 56	25	
57 58	26	
59 60 61	21	
62 63		1
64 65		1

28 Abstract

 Human type II alveolar cell line (A549) was exposed to aqueous and organic solvent PM10 extracts in order to evaluate cell proliferation, proinflammatory cytokines and cytotoxicity (lactate dehydrogenase , LDH). PM10 samples, collected in Torino (northwest Italy), were analyzed for inorganic chemical species (bioavailable iron and secondary particulates) and endotoxins, potentially inflammatory promoters to human airways.

Meanly, in the considered period, PM10 concentration was 55.4  $\pm$  39.1  $\mu$ g/m<sup>3</sup>, iron concentration was  $0.078 \pm 0.095$  and secondary particles constituted  $42 \pm 9$  % of the PM10 total mass. PM10 inhibits cell proliferation and induces both IL-6 and LDH release in a dose and time dependent manner, with a seasonal trend. The different roles of aqueous and organic solvent PM10 extracts demonstrate the importance of particle composition for the biological effects. Nevertheless few significant correlations were found between the biological effects and PM10 components evaluated in this methods development study.

**Keywords:** Particulate Matter, Cell Proliferation, Cytotoxicity, LDH, Cytokine.

#### **1. Introduction**

The health burden due to particulate matter (PM) air pollution is one of the biggest environmental health concerns in the World Health Organization (WHO) European Region and all around the world. PM is a complex and heterogeneous mixture, whose composition (particle size distribution and chemical characteristics) changes in time, space and depends on emissions from various sources, atmospheric chemistry and weather conditions (WHO, 2007).

The breathable fraction of ambient PM is often referred to PM10, defined as particles with a median aerodynamic diameter less than 10 µm, that is a quality air indicator widely measured (WHO, 2006). PM10 may further be divided into two main size fractions, a "coarse" (2.5-10 µm) and a "fine" (0.1-2.5 µm). The coarse fraction is dominated by natural sources (geological material: fugitive and resuspended dust; biological material: pollen, endotoxin) and its composition changes depending on the geology of the site considered. The fine fraction is dominated by anthropogenic emissions: mixture of carbon particles from combustion processes and secondary particles produced by photochemical reactions in the atmosphere (sulfates and nitrates). The carbonaceous fraction consists of aggregates of organic and inorganic carbon on which are adsorbed transition metals (Pb, Cd, V, Ni, Cu, Zn, Mn, Fe), organic compounds and biological constituents (US EPA, 1996).

Epidemiological and experimental studies have shown that particulate air pollution may induce and aggravate respiratory and cardiovascular diseases. Significant associations between exposure to ambient air particles and increased morbidity and mortality have been demonstrated (Englert 2004; Katsouyanni et al. 2001; Krewski and Rainham 2007; Pope 2007). Most of the available studies do not attribute the observed health effects to a particular characteristic of PM (WHO, 2007). Moreover the exact physiochemical mechanism by which PM produces adverse effects is still poorly

known; one of the hypotheses considered on PM's mechanisms of action is the
oxidative potential of the particles or specific components.

Several in vitro studies with collected particulate matter fractions indicate that particles with a higher oxidative potential have a greater ability to deplete antioxidant defences and to induce airway inflammation (Ghio et al. 2002; Moller et al. 2008; Schins et al. 2004). In this process transition metals such as Fe (Carter et al. 1997; Hutchison et al. 2005; Schins et al. 2004), organics such as polycyclic aromatic hydrocarbons (PAHs) (Billet et al. 2008) and endotoxins (Becker et al. 2003; Oberdorster et al. 2000) in PM seem to be involved (Donaldson et al. 2003; Schwarze et al. 2006; Sorensen et al. 2003). Moreover, although secondary inorganic aerosols have less toxic activity when tested under controlled laboratory conditions, epidemiological studies show significant associations between sulfates and nitrates and various health outcomes. In ambient air, this fraction may act as a carrier for other components or as a surrogate for PM emitted from the combustion of sulfur-containing fuels (Schwarze et al. 2006).

Recent studies have emphatised the ability of particles to induce cellular responses in different types of lung cells, as well as inflammation and toxic effects in the whole lung (Calcabrini et al. 2004; Churg and Brauer 2000; de Kok et al. 2006; Gualtieri et al. 2008; Hetland et al. 2004; Monn and Becker 1999; Shi et al. 2006). In vitro studies on the toxicity of PM10 and its fractions were made by exposing cultures of macrophages and pulmonary epithelial cells to suspension of particles (Becker et al. 2005; Churg and Brauer 2000). The results of these studies show that exposure to environmental particles induces cytotoxicity, increase in transcription factors expression (NFkB) and release in the culture medium of cytokines (such as IL-1, IL- 6, IL-8, MIP-2, GM CSF, TNF) and reactive oxygen species (Becker et al. 2005; Huang et al. 2002; Monn and Becker 1999; Ning et al. 2000). Other in vitro studies show that particles can induce damage in plasma membrane and release of cytosolic enzyme lactate dehydrogenase

(LDH) (Billet et al. 2007; Garcon et al. 2006; Geng et al. 2006; Molinelli et al. 2002); LDH is considered a marker for acute cytotoxicity (Seagrave and Nikula 2000). The aim of the present study was to investigate the biological in vitro effect of urban-air PM10 collected in Torino, a north-western Italian city, during two years (2005-2006). PM10 samples were analyzed for the amount of inorganic species (bioavailable iron and secondary particulates: sulfates and nitrates) and endotoxins, described as potentially toxic to human airways. Adherent A549 cells were exposed to PM10 aqueous and organic solvent extracts at different concentrations. A549 cells constituted a useful in vitro model for studying human respiratory epithelial cell biology as they present characteristics similar to human alveolar type II cells (Hatch 1992; Li et al. 2003; Shi et al. 2006; Veronesi et al. 2002) . This study investigated whether different PM10 extracts could 1) affect the cell proliferation, 2) increase the release of proinflammatory cytokines e.g. IL-6, and 3) induce cytotoxicity by measuring LDH release in culture medium. The association of the chemical and in vitro toxicological characterization of urban-air PM10 in one of the most industrialized areas of Italy, with high levels of particulate matter among European cities (Hazenkamp-Von Arx et al. 2004; Marcazzan et al. 2003) may provide scientific evidences for environmental and sanitary purposes. 2. Materials and Methods 

127 2.1 PM10 sampling

 PM10 (PM passing through a size-selective inlet with a 50% efficiency cut-off at 10  $\mu$ m aerodynamic diameter) was sampled on glass microfiber filters (Type A/E, 8" x 10", Gelman Sciences, Michigan, USA), with a Sierra Andersen High Volume Sampler 1200/VFC (Andersen Samplers, Atlanta, Georgia, USA) using a flow of 1160 L/min. Sample duration was controlled by a timer accurate to ± 15 min over a 24 hr sample period. The exact flow was calculated daily, corrected for variation in atmospheric

pressure and actual differential pressure across the filter. The filters were pre- and post- conditioned by moving them to a dry and dark environment for 48h, and they were weighed in a room with controlled temperature and humidity. Procedures were conducted according to the European Committee for Standardization (CEN, 1998). The PM10 concentration, C ( $\mu$ g/m<sup>3</sup>), in the air volume sampled, V (L), was calculated as follows:

140 C =  $[(W2-W1)-(B2-B1)]*10^{3}/V$ 

141 where  $W_1$  is the mean of three tare weights of the same filter before sampling (mg),  $W_2$ 142 is mean of three post-sampling weights of the same sample-containing filter (mg),  $B_1$  is 143 mean tare weight of blank filters (mg).  $B_2$  is mean post-sampling weight of blank filters 144 (mg) and V is volume as sampled at the nominal flow rate.

Samplings were carried out from January 2005 to December 2006 (2 filters every
month, one sampled at the half and one at the end of every month, without considering
week-ends) in a meteorological-chemical background station located in the urban
centre of Torino, an industrial north-western Italian city (Gilli et al. 2007b).

## 2.2 Particles extractions

Each PM10 filter (typically 20.32 cm by 25.40 cm) was treated individually: two different 6.35 cm by 20.32 cm wide strips were cut from the same filters and each portion has been used with different extraction medium: acetone and the Dulbecco's Modified Eagle's Medium (DMEM) culture medium without fetal calf serum (FCS). Acetone was chosen as solvent to extract organic-extractable compounds (Claxton et al. 2004). DMEM without FCS was chosen to extract water-soluble components, hypothetically comparable to the extraction at the lung cells (Calcabrini et al. 2004; Hetland et al. 2004). Each portion of the filter has been cut into little strips and placed in 50 mL polypropylene sterile tube with 15 mL of the extraction medium. The tubes were placed in an ultrasonic water bath for 10 min followed by 1 min of vortexing. This procedure was repeated 3 times. The samples were centrifugated at 5000 rpm for 10 min to 

remove the filter material and the supernatant was collected. Acetone extracts were evaporated with a rotary evaporator and re-suspended in culture medium. Aqueous extracts were directly assayed. It was assumed that particles were totally removed from the filters through extraction procedures (Hutchison et al. 2005). Unless specified otherwise, all chemicals were purchased from Sigma, USA.

#### 2.3 Iron and bioavailable iron determinations

167 Iron determination was performed according to the procedure of Gilli et al. (2007b).
168 Briefly, the metals were extracted from filter strips (3.18 cm by 20.32 cm) by a nitric
169 acid solution. After cooling, the sample was mixed and centrifuged and the trace
170 element concentrations in each sample were determined by atomic absorption
171 spectrometry, Varian GTA-96.

The bioavailable iron from urban particulates was determined as previously described (Lund and Aust 1990), with some modifications as reported by (Smith and Aust 1997) and Gilli et al. (2007b). Briefly, filter strips (3.18 cm by 20.32 cm) were suspended in NaCl 50nM, mixed and pH adjusted to 7.5. Citrate was then added to samples to obtain a final concentration of 1 mM and all samples were placed on a wrist-action shaker in the dark for 24 h. One mL samples were withdrawn and centrifuged to remove the particulates. The amount of iron mobilized as the citrate: Fe complex in the supernatant was determined using a spectrophotometric total non-heme iron assay (Brumby and Massey, 1967). The concentration of iron mobilized by citrate was expressed as µg of  $Fe/m^3$  and as ng of  $Fe/\mu g$  of particulate.

#### 2.4 Sulfates and nitrates determination

Sulfates and nitrates determination was performed according to the procedure of Gilli et al. (2007a, 2007b). Briefly each filter strip (3.18 cm by 20.32 cm) was extracted in 15 mL in distilled deionized waters, via 30 min sonication, 30 min agitation and an overnight refrigeration at 4°C. Prior to the analysis, the samples were centrifuged to remove particles. Ion chromatography (IC) was used to determine the soluble ion

188 content sulfates (SO<sub>4</sub><sup>=</sup>) and nitrates (NO<sub>3</sub><sup>-</sup>), applying a Dionex DX-100 ion 189 chromatograph with NaHCO<sub>3</sub> 0.3 mM and Na<sub>2</sub>CO<sub>3</sub> 2.7 mM for eluent and IonPac 190 analytical column S12A for anions. Both the utilized standards, sodium sulfate and 191 sodium nitrate (71959 and 71759 FLUKA respectively) ranged from 0.1  $\mu$ g/mL to 100 192  $\mu$ g/mL. The ions were identified by their elution/retention times (about 8.75 min and 193 12.5 min for nitrates and sulfates respectively) and quantified by the conductivity peak 194 area or peak height (300A method, US EPA, 1996).

#### 

## 2.5 Endotoxins determination

Endotoxin was assayed with an end-point chromogenic Limulus amebocyte lysate (LAL) method (QLC-1000 n° 50-648U, Cambrex, Walkersville, MD, USA) at 37°C with an automated micro-plate reader (ELX 800 UV, Bio-Tek Instruments, Inc) following manufacturer's instructions. Escherichia coli 0111:B4 endotoxin was used as standard endotoxin. Sample concentration were reported as endotoxin units (EU) per millilitre of eluant, EU per milligram of PM10, and EU per cubic meter of air collected. The limit of detection (LOD) was 0.01 EU/mL. To control for the potential interference of samples substances with the LAL assay, all samples were spiked with a known amount of endotoxin (0.4 EU/mL). The spiked solution was assayed along with the unspiked samples and their respective endotoxin concentrations are determined. The difference between these two calculated endotoxin values should equal the known concentration of the spike  $\pm$  25% to accept the results of the determinations.

<sup>45</sup><sub>46</sub> 208 2.6 Cell culture

The human lung epithelial cell line A549 (non-small cell lung cancer) from Interlab Cell Line Collection (Genova, IT) was used as a model for human epithelial lung cells. Cells were grown as a monolayer, maintained and treated in DMEM supplemented with 10% FCS, 2% L-glutamine 200mM, 2% HEPES 1M, 1% sodium pyruvate 100 mM and 1% penicillin/streptomycin 10mg/mL, at 37°C in an humidified atmosphere containing 5%  $CO_2$ . 

 

#### 2.7 Cell proliferation

Cell proliferation was evaluated using crystal violet method (Kueng et al. 1989). Cells were seeded in 24-well plates at a density of 4\*10<sup>4</sup> cell/well and exposed to particle concentration of 50 and 100 µg/mL (30 and 60 µg/cm<sup>2</sup>), (Alfaro-Moreno et al. 2002; Calcabrini et al. 2004; Veronesi et al. 2002). Cell proliferation was determined at 24h, 48h and 72h of exposure measuring the residual cell number by crystal violet by determining the absorbance at 595 nm with a micro-plate reader (ELX 800 UV, Bio-Tek Instruments, Inc). At the same time blank filters were treated in the same way. All experiments were performed in triplicate. Percentage cell proliferation was calculated comparing the absorbance of exposed cultures with the absorbance of non-exposed cultures.

#### 2.8 Proinflammatory cytochine in cell supernatant

IL-6 is a multifactorial cytokine protein that plays a major role in the mediation of airways inflammation associated with infection or toxic insult. A549 cells were exposed to particle concentration of 50 and 100 µg/mL (30 and 60 µg/cm<sup>2</sup>) and after 24, 48 and 72h of treatment, supernatant aliquots were removed for the cytochine assay. IL-6 was measured in the culture supernatant by enzime-lynked immunoabsorbent assay (ELISA) kits (PeliPair reagent set, Sanguin, Netherlands) with an automated micro-plate reader (ELX 800 UV, Bio-Tek Instruments, Inc) according to the manufacturer's recommendations. Results were calculated by expressing samples IL-6 increment as a percentage of IL-6 expression from non exposed cells.

2.9 LDH assay

To evaluate PM10 cytotoxicity, released lactate dehydrogenase (LDH) activities, from damaged cells, were measured in cell-free culture supernatants. Enzymatic activities, measured spectrophotometrically as absorbance variation at 340 nm (37°C), were expressed as nmol NADH oxidized/min in the conversion of pyruvate to lactate (Golladay et al. 1997; Kinnula et al. 1994; Riganti et al. 2002). A549 cells were seeded

in 6 well plates at a density of 1\*10<sup>6</sup> cell/well and exposed to PM10 extracts containing 50, 100 and 200 µg/mL of particles (30, 60 and 110 µg/cm<sup>2</sup>), (Billet et al. 2007; Geng et al. 2006; Hetland et al. 2000). At the same time blank filters were treated in the same manner. At 24h, 48h and 72h LDH activity was measured in supernatant and cell lysate. LDH activity was calculated as ratio between the extracellular LDH (measured in the supernatant) and total LDH (expressed as sum of LDH measured in supernatant and cell lysate). To obtain cell lysate, cells were washed with PBS, detached with trypsin-EDTA, resuspended with 1 mL of TRAP (82.3 mM triethanolamine hydrochloride, pH 7.6) and sonicated for 10". Then LDH was measured by adding 250 µl of mix containing NADH 0.25 mM and pyruvate 0.5 mM and consumption of NADH was measured as absorbance in a microplate reader (Benchmark Plus Microplate Reader, Biorad) at 340 nm. All experiments were performed in triplicate. Results were calculated by expressing samples LDH activity as a percentage of LDH activity from non exposed cells.

<sup>32</sup> <sub>33</sub> 256

#### 2.10 Statistical analysis

257 Statistical analyses were performed using the SPSS Package, version 14.0 for 258 Windows. T-test analysis was used to compare means and Spearman correlation 259 coefficient was used to assess relationship between variables. The mean difference 260 and correlation were considered significant at p<0.05.

- **3. Results**

#### 263 3.1 PM10 concentration

A total of 46 PM10 filters were analyzed during 2005 and 2006 without considering week-ends. The mean PM10 concentration during the whole sampling period was 55.4  $\pm$  39.1 µg/m<sup>3</sup>. The highest value was observed in winter (198.5 µg/m<sup>3</sup>), while the lowest was observed in summer (12.4 µg/m<sup>3</sup>); the T-test analysis show significant (p < 0.01)

difference for the seasonal (autumn/winter vs spring/summer) PM10 concentrations (Table 1).

#### 

6

#### 3.2 Iron and bioavailable iron concentrations

The total iron mean concentration was  $1.035 \pm 0.811 \,\mu g/m^3$  and, in relation to PM10, 19.43  $\pm$  11.64 ng/µg particles. The highest value was observed in winter (3.751 µg/m<sup>3</sup>), while the lowest was observed in summer (0.099 µg/m<sup>3</sup>). A statistically significant difference was found for the seasonal concentrations (autumn/winter vs spring/summer) expressed as  $\mu g/m^3$  (t-test, p<0.01) but not as ng/ $\mu g$  particles (Table 2). A significant positive correlation was found with PM10 concentrations (Spearman correlation: r=0.800, p<0.01).

Table 2 shows also the mean concentrations of bioavailable iron,  $6.9 \pm 4.5\%$  of total iron was bioavailable. No statistically significant differences were found for the mean seasonal concentrations expressed both as  $\mu g/m^3$  (t-test, p<0.01) and as ng/ $\mu g$ particles. A positive significant correlation was found with PM10 concentrations (Spearman correlation: r=0.670, p<0.01). 

#### 3.3 Sulfates and nitrates concentrations

The sulfates mean concentration was  $10.5 \pm 5.7 \ \mu g/m^3$  and, in relation to PM10, 216.7  $\pm$  72.7 ng/µg particles, such that sulfates represented 21.2  $\pm$  6.9 % of total PM10 mass. The nitrates mean concentration was  $11.7 \pm 9.2 \,\mu g/m^3$  and, in relation to PM10, 216.3  $\pm$  80.5 ng/µg particles, such that nitrates represented 20.6  $\pm$  6.1% of total PM10 mass. Considering the sum of sulfates and nitrates as principal components of secondary particulate, it was found that they meanly constituted  $42 \pm 9$  % of the PM10 total mass (Table 1). A statistically significant difference was found for the seasonal concentrations for both sulfates and nitrates expressed as  $\mu q/m^3$  (t-test, p < 0.05 and p < 0.01, respectively). Considering the concentrations of the two species in relation to PM10 (percentage), it was found that in the hot season the sulphate amount (23.6  $\pm$ 5.6 %) was more elevated than in the cold season (19.2  $\pm$  7.3 %) and this trend was 

significant (t-test, p < 0.01), while for nitrates the amount was similar in the two seasons. A significant positive correlation was found with PM10 concentrations both for sulfates (Spearman correlation: r=0.740, p<0.01) and for nitrates (Spearman correlation: r=0.900, p<0.01) and also a significant positive correlation between sulfates and bioavailable iron (Spearman correlation: r=0.490, p<0.01).

## 3.4 Endotoxins concentrations

Endotoxin mean concentration during the whole sampling period was  $0.19 \pm 0.07$ EU/m<sup>3</sup> and in relation to PM10 3.70  $\pm$  0.95 EU/mg. A statistically significant difference was found for the seasonal concentrations (autumn/winter vs spring/summer) expressed as EU/m<sup>3</sup> (0.24  $\pm$  0.07 vs 0.14  $\pm$  0.01 respectively) (t-test, p<0.05) but not as EU/mg particles. Endotoxins, EU/mg, were inversely correlated with PM10 concentrations (Spearman correlation: r = -0.710, p < 0.05).

3.5 A549 cell proliferation

Mean absorbance values at 595 nm of the negative control at 24, 48 and 72 h were respectively 0.120  $\pm$  0.010, 0.225  $\pm$  0.010 and 0.410  $\pm$  0.015. Extracts from blank filters had no significant effect on cell proliferation. Figure 1 shows effects of organic solvent (1a) and aqueous (1b) PM10 extracts on cell proliferation. PM10 inhibits cell proliferation in a dose and time dependent manner. Aqueous extracts seem to have stronger effects than organic solvent extracts on cell proliferation, this trend was confirmed by T-tests for inhibition at 48 and 72 h (p < 0.05) but not at 24 h, for both the tested concentrations. The maximum inhibition of cell proliferation,  $25.3 \pm 5.8$  %, was achieved at 72h by the 100 µg/mL aqueous extract. 

Proliferation inhibition of A549 cells due to organic solvent extract, in some cases (50 µg/mL at 72h, 100 µg/mL at 48 and 72h) was correlated with PM10 concentrations (Spearman correlations: r=0.860, p<0.01; r=0.810, p<0.01 and r=0.620, p<0.05 

320 respectively), no correlations were found with any other parameters or for the aqueous321 extracts.

Figure 3(a) shows different seasonal (autumn/winter *vs* spring/summer) proliferation inhibition of A549 cells at 72h after their incubation in the continuous presence of two concentrations of particle matter (50 and 100  $\mu$ g/mL). Despite PM10 organic solvent extracts and PM10 aqueous extracts always had higher effect in the cold months these differences were not significant (T-tests p > 0.05).

#### 3.6 IL-6 release in A549 cells

In order to investigate the proinflammatory potential of the particulate, the production of IL-6 in the presence of two different extracts of PM10 was analyzed. Mean IL-6 release in the negative control at 24, 48 and 72h range from 0.2 to 1.0 pg/mL. PM10 extracts induce IL-6 release in a dose and time dependent manner. As shown in figure 2a, the organic solvent extract induce significant release of IL-6 in relation to the control at 48h and 72h of exposure (T-test p < 0.05) for both the concentrations, the lowest concentration at 24h of exposure has no significant increase of IL-6 release in relation to the control (6.2  $\pm$  3.5%). The two tested concentrations exhibited similar expression of cytokine without significant difference (t-test p > 0.05).

<sup>2</sup> 337 In figure 2b, the aqueous extract induce significant release of IL-6 in relation to the <sup>1</sup> 338 control at 24h, 48h and 72h of exposure (T-test p < 0.05) for cells exposed to 100 <sup>3</sup> 339 µg/mL while for cells exposed to 50 µg/ml only at 72h.

340 Despite the aqueous extract exhibit higher mean levels of IL-6 expression the 341 difference between the production of IL-6 in the presence of organic solvent extract or 342 aqueous extract was not statistically significant (T-test p>0.05) for any concentrations 343 or times. The maximum increment of IL-6 release,  $108.3 \pm 39.1$  %, was obtained at 72h 344 by the 100 µg/mL aqueous extract. IL-6 release by A549 due to organic solvent extracts (50 and 100  $\mu$ g/mL at 72h) was correlated with PM10 concentrations (Spearman correlations: r=0.680, p<0.05 and r=0.730, p<0.05 respectively). No correlations were found with any other parameters. IL-6 release by A549 due to aqueous extracts was always correlated with PM10, sulfate, nitrate and Fe concentrations for both the extract concentrations and at each time of exposure in a range of Spearman correlation between r = 0.720 to r = 0.920, p<0.05.

Figure 3(b) shows different seasonal (autumn/winter *vs* spring/summer) IL-6 release at 72h after their incubation in the continuous presence of two concentrations of particle matter (50 and 100  $\mu$ g/mL). PM10 organic solvent extracts and PM10 aqueous extracts always had higher effect in the cold months and these differences were statistically significant (t-test, p < 0.01) except for one case (50 $\mu$ g/mL organic solvent extract).

#### 

#### 3.7 LDH release in A549 cells

An increase of extracellular LDH enzyme activity reflects an increase in the amount of membrane-damage cells (Lobner 2000). The release of the cytoplasmatic enzyme LDH into the culture supernatant was used to measure cell cytotoxicity. Extracts from blank filters had no significant effect on LDH release. Incubation of A549 cells with particles at a concentration of 50 µg/mL had no effect on LDH release as compared to control cells (data not shown). Results are shown in table 3: for organic solvent extract and for both the concentrations the highest and significant release of LDH was obtained at 72h of exposure (T-tests, p < 0.05), while at 24h and 48h no significant effects were observed in relation to the control (T-tests, p > 0.05). For aqueous extracts exposure to 100 µg/mL of particles increase LDH release by approximately 30-35% for all incubation time. Statistically significant increase of extracellular LDH activities were observed at 24h, 48h and 72h after cell exposure at 200 µg/mL of particles concentration (T-tests, p < 0.05). The maximum increment of LDH release, 87.6 ± 41.9 

371 % and 83.7  $\pm$  44.1 %, was respectively obtained at 48h by the 200 µg/mL aqueous 372 extract and at 72h by the 200 µg/mL organic solvent extract .

LDH increase by A549 due to both organic solvent extracts and aqueous extracts was
never correlated with PM10 concentrations or with any other parameters, with the
exception of the aqueous 200 µg/mL extract at 72h *vs* PM10 concentrations.

Figure 3(c) shows different seasonal (autumn/winter *vs* spring/summer) release of LDH activity from A549 cells at 72h after their incubation in the continuous presence of two concentrations of particle matter (100 and 200 µg/mL). PM10 organic solvent extracts had similar effect in the two seasons while PM10 aqueous extracts always had higher effect in the cold months, this difference was significant only for the lowest extract concentration.

 

#### 4. Discussion and Conclusions

The PM10 monitoring in Torino has showed concentrations of PM10 often over the daily (50 µg/m<sup>3</sup>) and yearly (40 µg/m<sup>3</sup>) quality targets (Air Quality Directive, 2008/50/CE). High pollution episodes are frequent during winter time in Northern Italy. This is due to the regional unique climate and topography which hamper particles dispersion. Human alveolar epithelial cells were exposed to concentration of particles comparable to 100 µg/m<sup>3</sup> such as those found in winter season in Torino, indeed the averages were 93.5  $\pm$  54.1  $\mu$ g/m<sup>3</sup> for 2005 and 78.4  $\pm$  53.1  $\mu$ g/m<sup>3</sup> for 2006 (Oberdörster and Yu 1999).

In urban sites like that utilized in this study, significant seasonal differences in total Fe concentrations occur probably due to minor presence of anthropogenic sources (for example domestic and industrial heating) in the warm season (Gilli et al. 2007b; Ziemacki et al. 2003). A small amount of iron is presumably bioavailable (about 6% of total Fe) and this form is capable of generating ROS and induces inflammation in 

 cellular systems (Aust et al. 2002). In this study this bioavailable Fe has not a
significant seasonal trend and it is less correlated with PM10 concentration than total
Fe.

In relation to secondary PM10 components, the seasonal trend of the two species is normally due to the photochemical reactions that, in warm season occur more frequently. It was found that the secondary airborne particulate matter (as sum of sulfates and nitrates) mass concentration was about 42% of the total PM10, and this value was confirmed by other studies conducted in the Po valley of Northern Italy (Gilli et al. 2007b; Marcazzan et al. 2003).

406 Among relevant air pollution components in causing toxicity there are also endotoxins; 407 in this study the mean endotoxins concentration was low and quite comparable with 408 those present in literature (Heinrich et al. 2003; Morgenstern et al. 2005; Mueller-409 Anneling et al. 2004; Solomon et al. 2006).

410 Many studies have investigated the toxicity and mechanism of PM in the airway 411 epithelial cells (Scapellato and Lotti 2007); PM induces different biologic effects 412 depending on sampling site (Rosas-Pérez et al., 2007), size fraction (Alfaro-Moreno et 413 al. 2002; Hetland et al. 2004; Osornio-Vargas et al. 2003), sampling time (Frampton et 414 al. 1999) and contaminants adsorbed on the particles (Baulig et al. 2003; Billet et al. 415 2007; Calcabrini et al. 2004; Muller et al. 2006). Moreover all in vitro studies with PM10 416 usually used high concentration of particles so that a potential confounding effect of 417 high doses need to be taken into account when extrapolating to *in vivo* conditions 418 (Oberdorster and Yu 1999).

In this study PM10 inhibits cell proliferation and induces both IL-6 and LDH release in a
dose and time dependent manner according to other studies on epithelial cells (AlfaroMoreno et al. 2002; Osornio-Vargas et al. 2003). These biological effects have a
seasonal trend as well as other biological effects such as mutagenicity evaluated in a
study conducted in the Torino city (Gilli et al. 2007a).

In general the maximum biological effect on A549 cells was obtained in term of IL-6 release (108.3  $\pm$  39.1 %), then LDH production (87.6  $\pm$  41.9 %) and in the end inhibition of cell proliferation (25.3  $\pm$  5.8 %). Indeed other studies have shown that PM10 induce - first the production of reactive oxygen species and inflammatory mediators such as interleukin IL-6 - than acute citotoxicity as release of LDH and in the end - an alteration of the cellular function highlighted by inhibition of the cellular proliferation (Jalava et al. 2006). It is also important to consider that these outcomes might occur independently or in association with each others (Seagrave and Nikula 2000).

In our investigation aqueous extracts seem to induce a greater IL-6 release and a greater proliferation inhibition than organic solvent (acetone) ones, while, on the contrary, organic solvent extract induce a greater release of LDH. Probably ,although both the two PM10 extracts produce effects on cells, water-soluble components, such as transition metals, could rapidly cross the plasma membrane and induce the inflammatory response, while polar compounds, from organic solvent extracts, cross the plasma membrane more slowly and are able to directly interact with DNA and induce mutations (Gilli et al. 2007a; Gilli et al. 2007b).

In general the biological effects are frequently attributed to the intracellular generation of ROS, usually correlated with aqueous PM extracts, that may damage protein structure, plasmatic membrane associated lipids and other cellular compartments (Knaapen et al. 2004; Sevanian and Ursini 2000; Sorensen et al. 2003). Soukup and Becker (Soukup and Becker 2001) found that expression of cytokines by human alveolar macrophages incubated with ambient insoluble PM10 was more than 50 times higher than that caused by the water-soluble fraction. Another study, conducted by Javala et al. (2008), is focused on the stronger cytotoxic effects of insoluble (organic) than water-soluble fraction of PM. In another investigation, mRNA levels of IL-1 $\alpha$  and IL-8 were significantly more expressed by human bronchial epithelial cells when

450 incubated with the water-soluble fraction of ambient PM10 than with the corresponding
451 suspensions (10–500 µg/mL, up to 24 h) (Fujii et al. 2001). To our knowledge, few
452 studies in literature compare the different effects of urban PM10 extracts.

Few significant correlations were found between the biological effects and PM10 components (bioavailable iron, secondary particulates and endotoxins) evaluated in this methods development study and it is important to note that the correlations have become significant only after some days of exposure (72h). These results confirmed other recent studies in which, for in vitro lung cells exposure, there were inconsistent associations between both cell proliferation and IL-6 release and secondary particulates or metals (Happo et al. 2008). One possible explanation for the presence of few correlations could be that the biological end-points and the PM10 components were quantified in chemically separate fractions. Furthermore the aqueous extracts did not reflected the activity of any particular PM10 chemicals components but only highlighted the activity of physiologic-soluble components.

Nevertheless the significant difference in biological effects induced by the two different extracts (aqueous and organic solvent) and the observed seasonal trend, support the need of further studies, in order to determine particle components responsible for biological effects associated both to PM10 and fine particulate matter.

41 468 The identification of PM components carrying the relevant burden in causing
43 469 inflammation remains unsettled, and it was said to represent one of the biggest gaps in
45 460 our knowledge (Pope and Dockery 2006).

48 471 In the future, better understanding of the relative toxicity and health effects of particles
 49
 50 50 51
 51 472 from various sources could facilitate targeted abatement policies and more effective
 52 53
 53 473 control measures to reduce the burden of disease due to air pollution.

59 476

1	477	Acknowledgments
2 3 4	478	This study was financed by a Piedmont Regional Grant (C.I.P.E. 2004). The authors
5 6	479	kindly thank Drs. M. Sacco, Drs. M. Maringo and Dr. M. Grosa of Environmental
7 8 0	480	Protection Regional Agency (Piedmont A.R.P.A.).
9 10 11	481	
12 13	482	
14 15	483	
16 17	484	
18 19	485	
20 21 22	486	
23 24	487	
25 26	488	
27 28	489	
29 30 31	490	
32 33	491	
34 35	492	
36 37	493	
38 39 40	494	
41 42	495	
43 44	496	
45 46	497	
47 48	498	
49 50 51	499	
52 53	500	
54 55	501	
56 57	502	
58 59 60	503	
61 62		
63 64		19
65		

5. References Ambient air guality and cleaner air for Europe. Directive 2008/50/CE. European Parliament and Council, 2008 б Alfaro-Moreno E, Martinez L, Garcia-Cuellar C, Bonner JC, Murray JC, Rosas I, Rosales SP, Osornio-Vargas AR. 2002. Biologic effects induced in vitro by PM10 from three different zones of Mexico City. Environ Health Perspect 110(7):715-20. Aust AE, Ball JC, Hu AA, Lighty JS, Smith KR, Straccia AM, Veranth JM, Young WC. 2002. Particle characteristics responsible for effects on human lung epithelial cells. Res Rep Health Eff Inst(110):1-65; discussion 67-76. Baulig A, Sourdeval M, Meyer M, Marano F, Baeza-Squiban A. 2003. Biological effects of atmospheric particles on human bronchial epithelial cells. Comparison with diesel exhaust particles. Toxicol In Vitro 17(5-6):567-73. Becker S, Mundandhara S, Devlin RB, Madden M. 2005. Regulation of cytokine production in human alveolar macrophages and airway epithelial cells in response to ambient air pollution particles: Further mechanistic studies. Toxicol Appl Pharmacol 207(2 Suppl):269-75. Becker S, Soukup JM, Sioutas C, Cassee FR. 2003. Response of human alveolar macrophages to ultrafine, fine, and coarse urban air pollution particles. Exp Lung Res 29(1):29-44. Billet S, Abbas I, Goff JL, Verdin A, Andre V, Lafargue PE, Hachimi A, Cazier F, Sichel F, Shirali P and others. 2008. Genotoxic potential of Polycyclic Aromatic Hydrocarbons-coated onto airborne Particulate Matter (PM(2.5)) in human lung epithelial A549 cells. Cancer Lett. Billet S, Garcon G, Dagher Z, Verdin A, Ledoux F, Cazier F, Courcot D, Aboukais A, Shirali P. 2007. Ambient particulate matter (PM(2.5)): Physicochemical 

1	530	characterization and metabolic activation of the organic fraction in human lung
2 3 4	531	epithelial cells (A549). Environ Res 105(2):212-223.
5 6	532	Brumby, P.E.; Massey, V.; 1967. Determination of nonheme iron, total iron, and
7 8	533	copper. In: Estabrook, R.W.; Pullman, M.E.; eds. Methods in Enzymology, vol.
9 10 11	534	10. New York: AcademicPress;. p. 463–74.
11 12 13	535	Calcabrini A, Meschini S, Marra M, Falzano L, Colone M, De Berardis B, Paoletti L,
14 15	536	Arancia G, Fiorentini C. 2004. Fine environmental particulate engenders
16 17	537	alterations in human lung epithelial A549 cells. Environ Res 95(1):82-91.
18 19	538	Carter JD, Ghio AJ, Samet JM, Devlin RB. 1997. Cytokine production by human airway
20 21 22	539	epithelial cells after exposure to an air pollution particle is metal-dependent.
23 24	540	Toxicol Appl Pharmacol 146(2):180-8.
25 26	541	CEN. Air quality - determination of the PM10 fraction of suspended particulate matter –
27 28	542	reference method and field test procedure to demonstrate reference
29 30 31 32 33	543	equivalence of measurement methods. Brussels: European Committee for
	544	Standardization; (European Standard EN 12341) 1998.
34 35	545	Churg A, Brauer M. 2000. Ambient atmospheric particles in the airways of human
36 37	546	lungs. Ultrastruct Pathol 24(6):353-61.
38 39	547	Claxton LD, Matthews PP, Warren SH. 2004. The genotoxicity of ambient outdoor air, a
40 41 42	548	review: Salmonella mutagenicity. Mutat Res 567(2-3):347-99.
43 44	549	de Kok TM, Driece HA, Hogervorst JG, Briede JJ. 2006. Toxicological assessment of
45 46	550	ambient and traffic-related particulate matter: a review of recent studies. Mutat
47 48	551	Res 613(2-3):103-22.
49 50 51	552	Donaldson K, Stone V, Borm PJ, Jimenez LA, Gilmour PS, Schins RP, Knaapen AM,
52 53	553	Rahman I, Faux SP, Brown DM and others. 2003. Oxidative stress and calcium
54 55	554	signaling in the adverse effects of environmental particles (PM10). Free Radic
56 57	555	Biol Med 34(11):1369-82.
58 59 60		
61 62		
63 64		21

1556Englert N. 2004. Fine particles and human health--a review of epidemiological studies.23557Toxicol Lett 149(1-3):235-42.

56
 558 Frampton MW, Ghio AJ, Samet JM, Carson JL, Carter JD, Devlin RB. 1999. Effects of aqueous extracts of PM(10) filters from the Utah valley on human airway
 560 epithelial cells. Am J Physiol 277(5 Pt 1):L960-7.

- 12<br/>13561Fujii T, Hayashi S, Hogg JC, Vincent R, Van Eeden SF. 2001. Particulate matter14<br/>15562induces cytokine expression in human bronchial epithelial cells. Am J Respir16<br/>17563Cell Mol Biol 25(3):265-71.
- <sup>18</sup> 19 564 Garcon G, Dagher Z, Zerimech F, Ledoux F, Courcot D, Aboukais A, Puskaric E,
  - 565 Shirali P. 2006. Dunkerque City air pollution particulate matter-induced
- 23<br/>24566cytotoxicity, oxidative stress and inflammation in human epithelial lung cells25<br/>26567(L132) in culture. Toxicol In Vitro 20(4):519-28.
- 568 Geng H, Meng Z, Zhang Q. 2006. In vitro responses of rat alveolar macrophages to
   569 particle suspensions and water-soluble components of dust storm PM(2.5).
   570 Toxicol In Vitro 20(5):575-84
- <sup>32</sup> 570 Toxicol In Vitro 20(5):575-84.
- <sup>34</sup> 571 Ghio AJ, Silbajoris R, Carson JL, Samet JM. 2002. Biologic effects of oil fly ash.
   <sup>36</sup> 572 Environ Health Perspect 110 Suppl 1:89-94.
- Gilli G, Pignata C, Schiliro T, Bono R, La Rosa A, Traversi D. 2007a. The mutagenic
   hazards of environmental PM2.5 in Turin. Environ Res 103(2):168-75.
- 43
   44
   575 Gilli G, Traversi D, Rovere R, Pignata C, Schiliro T. 2007b. Chemical characteristics
   576 and mutagenic activity of PM10 in Torino, a northern Italian city. Sci Total
   577 Environ 385(1-3):97-107.

# 0578Golladay SA, Park SH, Aust AE. 1997. Efflux of reduced glutathione after exposure of2579human lung epithelial cells to crocidolite asbestos. Environ Health Perspect 1054580Suppl 5:1273-7.

 $3 \frac{1}{9} \frac{1}{50} \frac{1}{52} \frac{1}{53} \frac{1}{55} \frac{1}{55}$ 

11

20 21

22

- 62 63
- 64 65

- Gualtieri M, Mantecca P, Cetta F, Camatini M. 2008. Organic compounds in tire particle induce reactive oxygen species and heat-shock proteins in the human alveolar cell line A549. Environ Int 34(4):437-42. Happo MS, Hirvonen MR, Halinen AI, Jalava PI, Pennanen AS, Sillanpaa M, Hillamo R, Salonen RO. 2008. Chemical compositions responsible for inflammation and tissue damage in the mouse lung by coarse and fine particulate samples from contrasting air pollution in europe. Inhal Toxicol 20(14):1215-31. Hatch G. 1992. Comparative biochemistry of airway lining fluid. In: Parent RA, editor. Comparative Biology of the Normal Lung. Boca Ranton, FL: CRC Press. p 617-632. Hazenkamp-Von Arx ME, Gotschi T, Ackermann-Liebrich U, Bono R, Burney P, Cyrys J, Jarvis D, Lillienberg L, Luczynska C, Maldonado JA and others. 2004. PM2.5 and NO2 assessment in 21 European study centres of ECRHS II: annual means and seasonal differences. Atmospheric Environment 38(13):1943-1953. Heinrich J, Pitz M, Bischof W, Krug N, Borm PJA. 2003. Endotoxin in fine (PM2.5) and coarse (PM2.5-10) particle mass of ambient aerosols. A temporo-spatial analysis. Epidemiology 14(5):S62-S62. Hetland RB, Cassee FR, Refsnes M, Schwarze PE, Lag M, Boere AJ, Dybing E. 2004. Release of inflammatory cytokines, cell toxicity and apoptosis in epithelial lung cells after exposure to ambient air particles of different size fractions. Toxicol In Vitro 18(2):203-12. Hetland RB, Refsnes M, Myran T, Johansen BV, Uthus N, Schwarze PE. 2000. Mineral and/or metal content as critical determinants of particle-induced release of IL-6 and IL-8 from A549 cells. J Toxicol Environ Health A 60(1):47-65. Huang SL, Cheng WL, Lee CT, Huang HC, Chan CC. 2002. Contribution of endotoxin in macrophage cytokine response to ambient particles in vitro. J Toxicol Environ Health A 65(17):1261-72.

1	608	Hutchison GR, Brown DM, Hibbs LR, Heal MR, Donaldson K, Maynard RL, Monaghan
2 3 4	609	M, Nicholl A, Stone V. 2005. The effect of refurbishing a UK steel plant on
5 6	610	PM10 metal composition and ability to induce inflammation. Respir Res 6:43.
7 8	611	Jalava PI, Salonen RO, Halinen AI, Penttinen P, Pennanen AS, Sillanpaa M, Sandell E,
9 10	612	Hillamo R, Hirvonen MR. 2006. In vitro inflammatory and cytotoxic effects of
11 12 13	613	size-segregated particulate samples collected during long-range transport of
14 15	614	wildfire smoke to Helsinki. Toxicol Appl Pharmacol 215(3):341-53.
16 17	615	Katsouyanni K, Touloumi G, Samoli E, Gryparis A, Le Tertre A, Monopolis Y, Rossi G,
18 19	616	Zmirou D, Ballester F, Boumghar A and others. 2001. Confounding and effect
20 21 22	617	modification in the short-term effects of ambient particles on total mortality:
22 23 24	618	results from 29 European cities within the APHEA2 project. Epidemiology
25 26	619	12(5):521-31.
27 28	620	Kinnula VL, Aalto K, Raivio KO, Walles S, Linnainmaa K. 1994. Cytotoxicity of oxidants
29 30	621	and asbestos fibers in cultured human mesothelial cells. Free Radic Biol Med
32 33	622	16(2):169-76.
34 35	623	Knaapen AM, Borm PJ, Albrecht C, Schins RP. 2004. Inhaled particles and lung
36 37	624	cancer. Part A: Mechanisms. Int J Cancer 109(6):799-809.
38 39	625	Krewski D, Rainham D. 2007. Ambient air pollution and population health: overview. J
40 41 42	626	Toxicol Environ Health A 70(3-4):275-83.
43 44	627	Kueng W, Silber E, Eppenberger U. 1989. Quantification of cells cultured on 96-well
45 46	628	plates. Anal Biochem 182(1):16-9.
47 48	629	Li N, Hao M, Phalen RF, Hinds WC, Nel AE. 2003. Particulate air pollutants and
49 50	630	asthma. A paradigm for the role of oxidative stress in PM-induced adverse
51 52 53	631	health effects. Clin Immunol 109(3):250-65.
54 55	632	Lobner D. 2000. Comparison of the LDH and MTT assays for quantifying cell death:
56 57	633	validity for neuronal apoptosis? J Neurosci Methods 96(2):147-52.
58 59		
6U 61 62		
63		

Lund LG, Aust AE. 1990. Iron mobilization from asbestos by chelators and ascorbic acid. Arch Biochem Biophys 278(1):61-4. Marcazzan GM, Ceriani M, Valli G, Vecchi R. 2003. Source apportionment of PM10 б and PM2.5 in Milan (Italy) using receptor modelling. Sci Total Environ 317(1-3):137-47. Molinelli AR, Madden MC, McGee JK, Stonehuerner JG, Ghio AJ. 2002. Effect of metal removal on the toxicity of airborne particulate matter from the Utah Valley. Inhal Toxicol 14(10):1069-86. Moller P, Folkmann JK, Forchhammer L, Brauner EV, Danielsen PH, Risom L, Loft S. 2008. Air pollution, oxidative damage to DNA, and carcinogenesis. Cancer Lett 266(1):84-97. Monn C, Becker S. 1999. Cytotoxicity and induction of proinflammatory cytokines from human monocytes exposed to fine (PM2.5) and coarse particles (PM10-2.5) in outdoor and indoor air. Toxicol Appl Pharmacol 155(3):245-52. Morgenstern V, Carty CL, Gehring U, Cyrys J, Bischof W, Heinrich J. 2005. Lack of spatial variation of endotoxin in ambient particulate matter across a German metropolitan area. Atmospheric Environment 39(36):6931-6941. Mueller-Anneling L, Avol E, Peters JM, Thorne PS. 2004. Ambient endotoxin concentrations in PM10 from Southern California. Environ Health Perspect 112(5):583-8. Muller A, Wichmann G, Massolo L, Rehwagen M, Grabsch C, Loffhagen N, Herbarth O, Ronco A. 2006. Cytotoxicity and oxidative stress caused by chemicals adsorbed on particulate matter. Environ Toxicol 21(5):457-63. Ning Y, Imrich A, Goldsmith CA, Qin G, Kobzik L. 2000. Alveolar macrophage cytokine production in response to air particles in vitro: role of endotoxin. J Toxicol Environ Health A 59(3):165-80. 

1	660	Oberdorster G, Finkelstein JN, Johnston C, Gelein R, Cox C, Baggs R, Elder AC. 2000.
2 3 4	661	Acute pulmonary effects of ultrafine particles in rats and mice. Res Rep Health
5 6	662	Eff Inst(96):5-74; disc 75-86.
7 8	663	Oberdorster G, Yu CP. 1999. Lung dosimetryconsiderations for noninhalation studies.
9 10	664	Exp Lung Res 25(1):1-6.
11 12 13	665	Osornio-Vargas AR, Bonner JC, Alfaro-Moreno E, Martinez L, Garcia-Cuellar C,
14 15	666	Ponce-de-Leon Rosales S, Miranda J, Rosas I. 2003. Proinflammatory and
16 17	667	cytotoxic effects of Mexico City air pollution particulate matter in vitro are
18 19	668	dependent on particle size and composition. Environ Health Perspect
20 21 22	669	111(10):1289-93.
22 23 24	670	Pope CA, 3rd. 2007. Mortality effects of longer term exposures to fine particulate air
25 26	671	pollution: review of recent epidemiological evidence. Inhal Toxicol 19 Suppl
27 28	672	1:33-8.
29 30 21	673	Pope CA, 3rd, Dockery DW. 2006. Health effects of fine particulate air pollution: lines
3⊥ 32 33	674	that connect. J Air Waste Manag Assoc 56(6):709-42.
34 35	675	Riganti C, Aldieri E, Bergandi L, Fenoglio I, Costamagna C, Fubini B, Bosia A, Ghigo
36 37	676	D. 2002. Crocidolite asbestos inhibits pentose phosphate oxidative pathway
38 39	677	and glucose 6-phosphate dehydrogenase activity in human lung epithelial cells.
40 41 42	678	Free Radic Biol Med 32(9):938-49.
43 44	679	Scapellato ML, Lotti M. 2007. Short-term effects of particulate matter: an inflammatory
45 46	680	mechanism? Crit Rev Toxicol 37(6):461-87.
47 48	681	Schins RP, Lightbody JH, Borm PJ, Shi T, Donaldson K, Stone V. 2004. Inflammatory
49 50 51	682	effects of coarse and fine particulate matter in relation to chemical and
51 52 53	683	biological constituents. Toxicol Appl Pharmacol 195(1):1-11.
54 55	684	Schwarze PE, Ovrevik J, Lag M, Refsnes M, Nafstad P, Hetland RB, Dybing E. 2006.
56 57	685	Particulate matter properties and health effects: consistency of epidemiological
58 59 60 61 62	686	and toxicological studies. Hum Exp Toxicol 25(10):559-79.
63		26

1	687	Seagrave JC, Nikula KJ. 2000. Multiple modes of responses to air pollution particulate
2 3 4	688	materials in A549 alveolar type II cells. Inhal Toxicol 12 Suppl 4:247-60.
5 6	689	Sevanian A, Ursini F. 2000. Lipid peroxidation in membranes and low-density
7 8	690	lipoproteins: similarities and differences. Free Radic Biol Med 29(3-4):306-11.
9 10	691	Shi T, Duffin R, Borm PJ, Li H, Weishaupt C, Schins RP. 2006. Hydroxyl-radical-
11 12 13	692	dependent DNA damage by ambient particulate matter from contrasting
14 15	693	sampling locations. Environ Res 101(1):18-24.
16 17	694	Smith KR, Aust AE. 1997. Mobilization of iron from urban particulates leads to
18 19	695	generation of reactive oxygen species in vitro and induction of ferritin synthesis
20 21 22	696	in human lung epithelial cells. Chem Res Toxicol 10(7):828-34.
23 24	697	Solomon GM, Hjelmroos-Koski M, Rotkin-Ellman M, Hammond SK. 2006. Airborne
25 26	698	mold and endotoxin concentrations in New Orleans, Louisiana, after flooding,
27 28	699	October through November 2005. Environmental Health Perspectives
29 30 21	700	114(9):1381-1386.
32 33	701	Sorensen M, Autrup H, Moller P, Hertel O, Jensen SS, Vinzents P, Knudsen LE, Loft S.
34 35	702	2003. Linking exposure to environmental pollutants with biological effects. Mutat
36 37	703	Res 544(2-3):255-71.
38 39 40	704	Soukup JM, Becker S. 2001. Human alveolar macrophage responses to air pollution
40 41 42	705	particulates are associated with insoluble components of coarse material,
43 44	706	including particulate endotoxin. Toxicol Appl Pharmacol 171(1):20-6.
45 46	707	U.S. EPA. Air quality criteria for particulate matter. vol 1-3. EPA/600/P-95/001a. 1996.
47 48 40	708	Veronesi B, de Haar C, Lee L, Oortgiesen M. 2002. The surface charge of visible
49 50 51	709	particulate matter predicts biological activation in human bronchial epithelial
52 53	710	cells. Toxicol Appl Pharmacol 178(3):144-54.
54 55	711	World Health Organization Europe. 2006. Air Quality Guidelines — Global Update
56 57	712	2005.
58 59 60		
61 62		
63 64		27

1	713	World Health Organization Regional Office for Europe, 2007. Health relevance of
3 4	714	particulate matter from various sources. Report on a WHO Workshop, Bonn,
5 6 7 8 9 10	715	Germany, 26-27 March 2007.
	716	Ziemacki G, Cattani G, Cusano MC, Stacchini G, Marconi A. 2003. [Occurrence of
	717	metals in various size fractions of particulate matter]. Ann Ist Super Sanita
12 13	718	39(3):371-9.
14 15	719	
16 17	720	Figure Captions
18 19	721	
20 21 22	722	Figure 1. Proliferation inhibition of A549 cells after their incubation in the continuous
23 24	723	presence of increasing concentrations of particulate matter (50 and 100 $\mu\text{g/mL})$ from
25 26	724	PM10 organic solvent extracts (a) and PM10 aqueous extracts (b). Inhibition of cell
27 28	725	proliferation was calculated comparing the absorbance of exposed cultures with the
29 30 31	726	absorbance of non-exposed cultures. Asterisks indicate statistically significant
32 33	727	differences from the control , p< 0.05 (T-test).
34 35	728	
36 37	729	Figure 2. IL-6 release by A549 cells after their incubation in the continuous presence of
38 39 40	730	increasing concentrations of particulate matter (50 and 100 $\mu\text{g/mL})$ from PM10 organic
41 42	731	solvent extracts (a) and PM10 aqueous extracts (b). Results were calculated by
43 44	732	expressing samples IL-6 relies as a percentage of IL-6 release from non exposed cells.
45 46	733	Asterisks indicate statistically significant differences from the control , p< 0.05 (T-test).
47 48	734	
49 50 51	735	Figure 3. Different seasonal (autumn/winter vs spring/summer) proliferation inhibitions
52 53	736	3(a), IL-6 releases 3(b) and LDH releases 3(c) from A549 cells at 72h after their
54 55	737	incubation in the continuous presence of PM10 organic solvent extracts and PM10
56 57	738	aqueous extracts (50, 100 $\mu$ g/mL or 200 $\mu$ g/mL). Asterisks indicates statistically
58 59 60	739	significant differences for the two seasons, p<0.05 (T-test).
61 62		
63 64		28
65		

#### \* Conflict of Interest Statement Click here to download high resolution image



#### Environmental Toxicology and Pharmacology Conflict of Interest Policy

Manuscript number (if applicable): Article Title: Cytotoxic effects induced on a549 cells and chemical characterization of urban-air PM10 collected in Torino (Italy). Author name: Tiziana Schilirò, Luca Alessandria, Raffaella Degan, Deborah Traversi, Giorgio Gilli

#### Declarations

Environmental Toxicology and Pharmacology requires that all authors sign a declaration of conflicting interests. If you have nothing to declare in any of these categories then this should be stated.

#### **Conflict of Interest**

A conflicting interest exists when professional judgement concerning a primary interest (such as patient's welfare or the validity of research) may be influenced by a secondary interest (such as financial gain or personal rivalry). It may arise for the authors when they have financial interest that may influence their interpretation of their results or those of others. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding.

Please state any competing interests

No conflict of Interest

#### Funding Source

All sources of funding should also be acknowledged and you should declare any involvement of study sponsors in the study design; collection, analysis and interpretation of data; the writing of the manuscript; the decision to submit the manuscript for publication. If the study sponsors had no such involvement, this should be stated.

Please state any sources of funding for your research

Piedmont Regional Grant (C.I.P.E. 2004).

Signature (a scanned signature is acceptable, but each author must sign)

in Slille 1. -

Print name

Tiziana Schilirò, Luca Alessandria Raffaella Degan, Deborah Traversi Giorgio Gilli **Table 1.** Means and standard deviations of PM10, sulfates and nitrates concentrations ( $\mu$ g/m<sup>3</sup>) of the whole sampling period (2005/2006) and divided by seasons.

	PM10	SO₄ <sup>⁼</sup>	NO <sub>3</sub> <sup>-</sup>	SO₄⁼	NO <sub>3</sub> <sup>-</sup>
	(µg/m³)	(µg/m³)	(µg/m³)	(% of PM10	(% of PM10
				mass)	mass)
Whole period	55.4 ± 39.1	$10.5\pm5.7$	11.7 ± 9.2	$21.2\pm 6.9$	$20.6\pm6.1$
Winter/Autumn	71.3 ± 43.1**	$12.4\pm6.4^{\star}$	14.9 ± 10.1**	$19.2\pm7.3^{*}$	$20.8\pm6.0$
Summer/Spring	37.4 ± 24.1**	8.3 ± 5.7*	$8.0\pm6.3^{\star\star}$	$23.6\pm5.6^{\ast}$	$20.3\pm6.3$

\*\* statistically significant differences, (autumn/winter vs spring/summer) p < 0.01 (T-test).

\* statistically significant differences, (autumn/winter vs spring/summer) p < 0.05 (T-test).

**Table 2.** Means and standard deviations of Fe concentrations ( $\mu$ g/m<sup>3</sup>) of the whole sampling period (2005/2006) and divided by seasons.

	Fe total	Fe bio	% Fe bio	
	(µg/m³)	(µg/m³)	(of total)	
Whole sampling period	$1.035\pm0.811$	$0.078\pm0.095$	$6.9\pm4.5$	
Winter/Autumn	1.355 ± 0.919**	$0.097\pm0.089$	$7.5\pm4.8$	
Summer/Spring	$0.672 \pm 0.463^{**}$	$0.056\pm0.099$	$\textbf{6.2}\pm\textbf{4.1}$	

\*\* statistically significant differences, (autumn/winter vs spring/summer) p < 0.01 (T-test).

**Table 3.** Released lactate dehydrogenase (LDH) activity from A549 cell 24h, 48h and 72h after their incubation in the continuous presence of increasing concentrations of particle matter (100 and 200  $\mu$ g/mL) from PM10 organic solvent extracts and from PM10 aqueous extracts. Results were calculated by expressing samples LDH activity as a percentage of LDH activity from non exposed cells.

LDH increase %		24 h	48 h	72 h
PM10 organic solvent	100 µg/mL	-2.7 ± 7.9	3.9 ± 7.1	64.0 ± 22.5*
extracts	200 µg/mL	-7.3 ± 3.1	$2.9\pm5.5$	83.7 ± 44.1*
PM10 aqueous extracts	100 µg/mL	$30.2\pm24.3$	33.5 ± 23.0*	35.6 ± 18.3*
	200 µg/mL	78.1 ± 34.1*	87.6 ± 41.9*	53.6 ± 24.6*

\* statistically significant differences in relation to the control, p < 0.05 (T-test).



Figure 1b Click here to download high resolution image



Figure 2a Click here to download high resolution image



Figure 2b Click here to download high resolution image





3c