

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Three-dimensional laser scanner evaluation of facial soft tissue changes after Le Fort I advancement and rhinoplasty surgery: patients with cleft lip and palate vs patients with nonclefted maxillary retrognathic dysplasia (control group)

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/157585> since 2015-09-09T08:46:08Z

Published version:

DOI:10.1016/j.oooo.2013.12.406

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in:

[Oral surgery, oral medicine, oral pathology and oral radiology, Volume 117, Issue 4, April 2014, DOI: 10.1016/j.oooo.2013.12.406]

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect® platform]

Manuscript Number: TRIPLEO-D-13-00740R2

Title: THREE-DIMENSIONAL LASER SCANNER EVALUATION OF FACIAL SOFT-TISSUE CHANGES AFTER LEFORT I ADVANCEMENT AND RHINOPLASTY SURGERY: CLEFT LIP AND PALATE PATIENTS VS NON-CLEFTED MAXILLARY RETROGNATHIC PATIENTS (CONTROL GROUP)

Article Type: Original Research Article

Keywords: Cleft lip and palate, LeFort I osteotomy and secondary RSP, soft -tissue, three-dimensional surface laser scanner.

Corresponding Author: Dr. Laura Verzé,

Corresponding Author's Institution: Università di Torino

First Author: Laura Verzé

Order of Authors: Laura Verzé; Francesca Antonella Bianchi; Guglielmo Ramieri

Abstract: Objective: The aim of this study was to analyse the differences in facial soft-tissue changes, despite the same extent of upper jaw forward movement, between uCLP patients compared with non-CLP retruded maxilla patients, after LeFort I osteotomy and secondary Rhinoplasty. Study design: Twelve patients with maxillary retrognathic dysplasia and nose deformity were divided in two groups: A (uCLP) and B (control) and compared on the basis of the same maxillary advancement. Cephalometry and 3D mean facial model of A and B was obtained before and after surgery. Linear/angular measurements were calculated. Results: Upper vermilion and alar base remained unchanged in A but increased in B. In both groups, symmetry of the nasal base was improved and an increase of the sagittal projection of the lips was observed. Conclusions: 3D analysis showed that surgical procedures, in uCLP, can provide a satisfactory aesthetic outcome but some differences are evident in comparison to control group.

Journal of Oral Surgery Oral Medicine Oral Pathology Oral Radiology.

Editor-in-Chief Mark W. Lingen, DDS, PhD , Chicago, IL

Manuscript submission

I have enclosed the manuscript entitled: “Three dimensional laser scanner evaluation of facial soft tissue changes after Lefort I advancement and Rhinoplasty surgery: cleft lip and palate patients vs retrognathic-(control group)” for publication on the *Journal of Oral Surgery Oral Medicine Oral Pathology Oral Radiology*.

The present study performed a 3D analysis of the differences in the facial soft-tissue changes after surgical treatment (LeFort I osteotomy and secondary Rhinoplasty surgery) in unilateral cleft lip and palate (uCLP) patients vs retrognathic patients with the same amount of bone correction.

3D analysis showed that surgical procedures, in uCLP adult patients, can provide a satisfactory aesthetic outcome but some differences are evident in comparison to control group. Therefore, further technical improvements are still possible.

Hereby I certify that the authors have participated actively in the writing of the manuscript and they approve to submit this version of the manuscript to the *Journal of Oral Surgery Oral Medicine Oral Pathology Oral Radiology*.

This paper is unpublished and it is not under current consideration elsewhere.

There is no conflict of interest directly or indirectly for either authors.

Best regards.

Laura Verzé

June, 3, 2013

Laura Verzé, MD

Department of Public and Pediatric Health Sciences, Legal Medicine Section, University of Turin,
Turin, Italy.

Corso Galileo Galilei 22, 10126 Turin, Italy.

telephone: 00390116705684; fax: 00390116705934; E-mail: laura.verze@unito.it

LIST OF POTENTIAL REVIEWERS

Takako OKAWACHI, Associate Researcher, Kagoshima University

Chiarella SFORZA, Full Professor, Università di Milano

Turin, 11/11/2013

Ms. Ref. No.: TRIPLEO-D-13-00740R1

Title: THREE-DIMENSIONAL LASER SCANNER EVALUATION OF FACIAL SOFT-TISSUE CHANGES AFTER LEFORT I ADVANCEMENT AND RHINOPLASTY SURGERY: CLEFT LIP AND PALATE PATIENTS VS NON CLEFT RETRUDED MAXILLA PATIENTS (CONTROL GROUP).

Dr Mark W. Lingen, DDS, PhD , Chicago, IL

Editor-in-Chief. *Journal of Oral Surgery Oral Medicine Oral Pathology Oral Radiology*.

Dear Professor Lingen,

we thank the reviewers for their comments about our study that will help us to improve the level of our scientific communication.

We really appreciate the hard work of the reviewers and we agree that our paper needed some relatively minor recommendations.

We shall try to answer to reviewer's # 1 comments:

1. **“Non-cleft retruded maxilla patients”**: This term was replaced with “non-clefted maxillary retrognathic patients”, accepting your suggestion.
2. **“No primary rhinoplasty is performed”**: the authors treated all the patients with current standard of care techniques. The phrase “no primary rhinoplasty is performed” intended that no further corrective rhinoplasty was performed. All patients underwent a primary cheilorhinoplasty therefore we edited the phrase in the Material and Methods Section P2 Lines 29-32.
3. **Discussion P2 Lines 53-54, P5 Lines 36-37**: we changed “defects” to “deformities”.

4. **Discussion P3 Lines 38-39:** the sentence “the alar base width in CLP was bigger than in control pre-operatively” means that the alar base was wide and in the next paragraph we explained the reason of that “the nasal alar widening seems to be due to the release of the soft tissue attachment and muscle insertion”. The term “bigger” was replaced with “wider and vertically increased” because we agree that this second statement is more understandable.
5. **Discussion P5 Lines 43-44:** the authors wrote the sentence “CLP patients had partially edentulous premaxilla...” because in four out six patients one or two frontal teeth were missing at the CLP side but we think that this should not altered the 3D facial morph. The sentence could be misunderstood so we decided to replaced it with “In addition, in some of the CLP patients, one or two frontal teeth were missing at the cleft side and this fact may contribute to reduce the upper lip support”.

We hope that these changes will fit the request of the reviewer and that our paper will be accepted for publication in the *Journal of Oral Surgery Oral Medicine Oral Pathology Oral Radiology*.

With regards.

Sincerely yours,

Laura Verzé, M.D.

Statement of Clinical Relevance. 3D analysis showed that surgical procedures, in uCLP adult patients, can provide a symmetric nasal base and a satisfactory facial profile but some differences remain in the post-operative frontal and profile views in uCLP patients, in comparison with the control group, indicating further technical requirement.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

THREE-DIMENSIONAL LASER SCANNER EVALUATION OF FACIAL SOFT-TISSUE CHANGES AFTER LEFORT I ADVANCEMENT AND RHINOPLASTY SURGERY: CLEFT LIP AND PALATE PATIENTS VS NON-CLEFT RETRUDED MAXILLA PATIENTS — NON-CLEFTED MAXILLARY RETROGNATHIC PATIENTS (CONTROL GROUP).

Laura Verzé, MD^{*}, Francesca Antonella Bianchi, MD[^], Guglielmo Ramieri, MD, DDS[^]

^{*} Department of Public and Pediatric Health Sciences, Legal Medicine Section, University of Turin, Turin, Italy.

[^] Department of Surgical Sciences, Maxillofacial Surgery Section, San Giovanni Battista Hospital, University of Turin, Turin, Italy.

Corresponding Author:

Laura Verzé, MD; Department of Science of Public Health and Pediatrics, Legal Medicine Section, University of Turin, Turin, Italy.

Corso Galileo Galilei 22, 10126 Turin, Italy.

telephone : 00390116705684; fax: 00390116705934; e-mail: laura.verze@unito.it

Disclosure. Hereby I certify that there is no conflict of interest directly or indirectly for either authors.

WORD COUNT ABSTRACT: 150

COMPLETE MANUSCRIPT WORD COUNT: 3363

NUMBER OF REFERENCES: 39

NUMBER OF FIGURES: 4

NUMBER OF TABLES: 2

ABSTRACT

Objective: The aim of this study was to analyse the differences in facial soft-tissue changes, despite the same extent of upper jaw forward movement, between uCLP patients compared with non-CLP patients, after LeFort I osteotomy and secondary Rhinoplasty.

Study design: Twelve patients with maxillary retrognathic dysplasia and nose deformity were divided in two groups: A (uCLP) and B (control) and compared on the basis of the same maxillary advancement. Cephalometry and 3D mean facial model of A and B was obtained before and after surgery. Linear/angular measurements were calculated.

Results: Upper vermilion and alar base remained unchanged in A but increased in B. In both groups, symmetry of the nasal base was improved and an increase of the sagittal projection of the lips was observed.

Conclusions: 3D analysis showed that surgical procedures, in uCLP, can provide a satisfactory aesthetic outcome but some differences are evident in comparison to control group.

Keywords. Cleft lip and palate, LeFort I osteotomy and secondary RSP, soft -tissue, three-dimensional surface laser scanner.

Statement of Clinical Relevance. 3D analysis showed that surgical procedures, in uCLP adult patients, can provide a symmetric nasal base and a satisfactory facial profile but some differences remain in the post-operative frontal and profile views in uCLP patients, in comparison with the control group, indicating further technical requirement.

INTRODUCTION

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Improvement of facial aesthetics is one of the primary objectives of modern orthognatic surgery; attractiveness is a major component of the self concept. The appearance of the face has been found to influence the social acceptance and psychological well being of the individual. In the literature, it has been reported that symmetrical body shape is a central cue for attractiveness.^{1,2,3}

This fact allows the assumption of a potential disadvantage in the visual perception of patient with cleft lip and palate (CLP).⁴

The cleft malformation shows a variety of inter-individual shapes. Even when surgery was completed early in infancy and followed by therapeutic rehabilitation, adult patients with CLP show secondary deformities in the maxillary and nasal regions. These deformities may consist of defects that are unrepaired in primary surgery and distortions that develop through growth or caused by residual scars. Clinical examination usually revealed upper lip scars from previous corrective plastic surgery, maxillary hypoplasia, difference in lip length and nasal deformities which can vary from almost invisible to catastrophic, mostly dependent on the severity and type of cleft⁵ and on ability of cleft surgeon. Nose distortions include deviated columella, a depressed and deviated nasal tip, dislocation of the alar cartilage, webbing at the alar rim, flat and V- shaped nostrils, and scarring or fistulae of the nostril floor. These abnormalities are involved in all components of the nose, such as the facial skeleton, cartilage, muscle, skin, subcutaneous tissue and mucosal lining.⁶⁻⁹

There are many well-established surgical techniques to repair residual maxillary, lip and nose deformities in CLP adult patients. In all cases the aim of the therapy is to reach normal anatomy with symmetrical relations between the cleft and non-cleft sides.

Several cephalometric studies in cleft patients after surgery were conducted. The results of soft tissue changes have so far been interpreted with the aid of lateral

1 cephalograms.¹⁰ However in the lateral view structural comparison is limited in the medial
2 plane and the asymmetries were not quantified.¹¹
3

4 Analytical and objective three-dimensional (3D) laser scanner evaluation could help to
5 quantify the post-operative soft-tissue changes.
6
7

8
9 To the best of the author's knowledge, no 3D laser scanner studies have been
10 performed on the volumetric 3D soft-tissue changes after LeFort I osteotomy and secondary
11 Rhinoplasty surgery (RSP) in unilateral CLP (uCLP) patients compared with a control group
12 (~~non-cleft retruded maxilla~~ non-clefted maxillary retrognathic patients: non-CLP).
13
14
15
16
17
18

19 The aim of this study was therefore to analyse the differences in facial soft-tissue
20 changes in uCLP patients who underwent LeFort I osteotomy and secondary RSP surgery
21 compared with non-CLP patients who underwent the same extent of upper jaw forward
22 movement and secondary RSP surgery.
23
24
25
26
27
28

29 30 31 **PATIENTS AND METHODS** 32

33 From January 2010 to December 2011, 53 adults Caucasian patients with maxillary
34 retrognathic dysplasia and varying degrees of nasal deformity underwent LeFort I osteotomy
35 and secondary Rhinoplasty surgery (RSP) at the Division of Maxillofacial Surgery, San
36 Giovanni Battista Hospital, University of Turin, Turin, Italy.
37
38
39
40
41
42

43 The criteria for inclusion in the present study were as follows: white males or females,
44 adult age (> 18 years), skeletal class III (with maxillary retrognathic dysplasia, SNA<
45 82°±2°) and nasal deformity. All the patients underwent a similar surgical procedure, by the
46 same surgeon, consisting of a maxillary advancement (standardized surgical treatment
47 consisting of a LeFort I osteotomy) and RSP with or without grafts. Following maxillary
48 stabilisation, a resorbable (2.0 Vicryl) oblique suture, performed intraorally with two
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 symmetrical tension stitches between levator labii superioris alaeque nasi (LLSAN) of the
3 right side and LLSAN of the left side, allowed a good control of alar flaring.

4
5 Patients with congenital syndromes or previous facial injuries, with incomplete
6 clinical and radiologic records, and those who had not completed their post-operative follow-
7 up were excluded from the study.
8
9

10
11
12 Twelve patients fulfilled inclusion criteria for the study and they were divided
13 in two groups (Table I). Variables examined include: age, sex and deformity of the nose.

14
15
16 Group A: included 6 adult patients (4 women, 2 men) with complete uCLP (with no
17 other associated malformations or distinctive features in the face such as piercing or tattoos);
18 mean age 28.5 years, range 18-39 years. Three patients had a uCLP on the right side; the
19 others (1 woman, 2 men) had a uCLP on the left side. ~~Primary closure of the lip~~
20 Cheilorhinoplasty was conducted between the 6 and 9 months of age. ~~No primary rhinoplasty~~
21 ~~was performed.~~ Closure of the hard and soft palates was done at 12 to 18 months of age.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

999
1000 Pre-operative patients' features and description of the surgical procedures are shown
1001 in Table I. The costal and auricular cartilage grafts were used to support the nasal tip
1002 projection. The quadrangular cartilage graft was used in patient no. 6 to reconstruct the left
1003 nasal alae.

1 Lateral (L) cephalometry and 3D facial surface data were obtained before (T0) and
2 one year (T1) after surgery.
3

4 Informed consent was obtained from all participants. This study was performed in
5 agreement with local institutional review board. We followed the Helsinki Declaration
6 guidelines.
7
8
9

10 ***Cephalometric measurements***

11 Lateral cephalograms were traced by one examiner using the software Dolphin 11.0
12 Premium (Dolphin Imaging, Chatsworth, CA, USA). Only SNA measurements were obtained
13 to assess sagittal skeletal movements.
14
15
16
17
18
19
20

21 A subsample of 20 randomly selected radiographs were retraced and digitized 1
22 month later to calculate the systematic errors. All the measurements were compared between
23 the two time sets by the paired t test. All the measurements presented no significant
24 difference at retracing.
25
26
27
28
29
30

31 ***Facial scan and data processing***

32 3D images of all subjects have been achieved at T0 and at T1. Surface data were
33 acquired using a Head and Face Colour 3D Scanner (3030RGB; Cyberware, Inc., Monterey,
34 California). Subjects were registered with the head in natural position (nhp), the eyes closed
35 and teeth in occlusion. The acquired data was transferred to a graphics workstation for
36 viewing and elaboration with Cyberware Echo software (Cyberware Inc., Monterey,
37 California). The scanning method and the detailed protocol regarding how to reduce the
38 artifacts was previously described.¹²
39
40
41
42
43
44
45
46
47
48
49
50

51 Scanned data arrays of the facial area were then firstly restricted and then reduced
52 from around 160.000 to 30.000 points. Facial surface reconstruction, multiple scan alignment
53 and measurements were carried out using Rapid Form 2004 software (INUS Technologies
54 Inc., Seoul, South Korea).
55
56
57
58
59
60
61
62
63
64
65

1 Facial scans at T0 and T1 were pooled together by electronic surface averaging to
2 obtain the mean facial model of uCLP patients (A) and control group (B), before (T0) and
3 after treatment (T1); the rater was L.V. (Fig. 1). 3D average surfaces were constructed using
4 the software Morphostudio (Biomodelling Solutions, UK) and a mesh framework algorithm
5 based on nine anatomical landmarks.
6
7
8
9
10

11 Reference vertical (Y: midline through glabella; Y1: vertical plane through left
12 endocanthion and perpendicular to X) and horizontal (X: through right and left endocanthion)
13 planes were constructed on models.
14
15
16
17
18

19 Different linear and angular measurements of the mean faces were calculated for
20 comparison of the T0 and T1 models using 10 landmarks taken from classical anthropometry.
21 The landmarks were (Fig. 2-Table II): 1. enr, right endocanthion; 2. enl, left endocanthion; 3.
22 alr, right alar crest point; 4. all, left alar crest point; 5. prn, tip of nose; 6. sn, subnasale; 7.
23 chr, right cheilion; 8. chl, left cheilion; 9. ls, labialis superior; 10. stomation, sto. Differences
24 of linear measurement greater than one millimeter and angular measurements superior to 3
25 grades were considered significant.
26
27
28
29
30
31
32
33
34
35

36 Axial cross sections through prn, sn and ls and sagittal cross sections were also
37 obtained (Fig. 3-4).
38
39
40

41 **RESULTS**

42 *Cephalometric measurements*

43 The L Cephalometric measurements showed a maxillary advancement between 5 to 8
44 mm (mean 6.33 mm) in each group (Table I).
45
46
47
48
49
50

51 *Facial scan and data processing*

52 The comparison between A and B revealed that uCLP patients had a shorter and more
53 round face than non-CLP subjects, in both T0 and T1 (Fig. 1).
54
55
56
57
58
59
60
61
62
63
64
65

1
2 In frontal view, a vertical increase of the upper vermilion and lengthening of the alar
3 base width were evident in B. At T0, the alar base width in A was bigger than in B and was
4 unchanged at T1; in A also the upper vermilion remained unchanged. In both groups,
5 symmetry of the alar forms were improved at T1 (Fig. 1).
6
7

8
9 In profile view, both in A and B, an improvement of the orbito-maxillary-zygomatic
10 sulcus and increase of the sagittal projection of the lips were observed; at T1, a successfully
11 projected nasal tip was noticed more in B than in A (Fig. 1).
12
13
14
15

16 Table II showed different linear and angular measurements of the mean faces at T0
17 and T1. Measurements at T0 and T1 documented that the major post-surgical changes in A
18 and B were in upper lip and nose. At T0, $al_r - al_l$ in A was greater than in B, at T1 this
19 measurement significantly decreased in A. Normalization of the nasal alae were evident in B,
20 reduced but still noticeable asymmetry of the alar was observed in A ($al_{r-l} - prn$). At T0, the al
21 point of the two sides demonstrated vertical asymmetry, significantly improved at T1; $al_{r-l} - X$
22 was greater in B than in A both at T0 and T1. $Prn - Y$ in A showed that surgery allowed an
23 improvement of deviation of the tip of nose on the symmetry axis. After treatment, the
24 distance of the ls from sto was increased in B, demonstrating lengthening of the upper
25 vermilion.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 In A $prn - sn - ls^\circ$ significantly increased and was larger than in B either at T0 or T1. The
42 distance between prn and the horizontal axis ($prn - X$) is longer in B vs A and was unchanged
43 at T1.
44
45
46
47

48 Increase of the sagittal projection of the lips was also evident, in particular in B, with
49 lengthening of the $ls - Y_1$.
50
51

52 Axial cross sections through prn, sn and ls clearly demonstrated a post-op
53 displacement of facial soft tissue in B greater than in A; reduced, but still noticeable deviation
54 of the tip of nose on the symmetry axis (Y) was observed in A (Fig. 3). Sagittal sections
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

illustrated a refinement of the naso-labial sulcus in A and increase sustain of the upper lip in B (Fig. 4).

DISCUSSION

The impairment of maxillary growth resulting in retrusion of the maxilla is a frequent finding in CLP adult patients. To correct these dento-facial deformities, orthognathic surgery may therefore be indicated.¹³

A maxillary advancement with a Le Fort I osteotomy is the most common orthognathic procedure. In the literature, the frequency of indications for a Le Fort I osteotomy in uCLP patients varies from 22% to 48.3%.¹⁴⁻¹⁷ CLP children often have midfacial growth deficiency, with a a characteristic concave profile. This generally increases during adolescence.¹⁸ For a few authors, these growth disturbances are intrinsic to the cleft itself, as it was observed in children who were never operated on for their cleft.¹⁹⁻²¹ Instead, for many authors, maxillary growth deficiency is mainly iatrogenic in nature and a consequence of the primary surgical repair of the palate.²²⁻²³

CLP adult patients can also show secondary nasal deformities.

Recently, primary rhinoplasty has been highlighted for the management of patients with uCLP and these techniques have been shown to clearly improve the results of the nasal deformity and overall symmetry.²⁴⁻²⁶

However, definitive rhinoplasty may still be necessary as the child grows.

The goal of the secondary treatment of uCLP deformities is to achieve naturally balanced nasal forms with an adequately projected nasal tip and a repositioning of retrognathic maxilla. Because a uCLP involved more or less inherent tissue ~~defects~~ deformities in the lip and nose, the secondary correction does not always achieve the level of the healthy frontal/profile configuration. Many rating systems for nasal deformity have been established and used worldwide.^{27, 28}

1 Recent developments in computer technology have facilitated the more accurate and
2 objective 3D characterization of facial forms of CLP patients.^{29, 30}
3

4 Laser surface scanning has been reported as a reliable and accurate method for
5 identifying cranio-facial surface landmarks.^{12, 31, 32}
6

7 Previous studies on the 3D laser scanner analysis of the morphologic changes of the
8 nose and lips after a Le Fort I osteotomy, in non-CLP patients, were conducted. The authors
9 observed that the labial changes were mainly due to the movements of the jawbone. The nasal
10 morphology changes after a Le Fort I osteotomy and consists mainly of widening of the nasal
11 alae caused by the release of the muscle insertion and their retraction. This change resulted
12 not to be influenced by the direction of the maxillary movement.³³
13
14
15
16
17
18
19
20
21
22
23

24 To date, very limited evaluation of facial morphology changes, after orthognathic
25 surgery in CLP adult patients in comparison to non-CLP subjects, were performed with 3D
26 analysis.^{10, 11} McCance et al.¹⁰ measured the 3D soft-tissue changes in a group of adult
27 patients with various forms of clefts following orthognathic surgery to correct jaw
28 disproportion. The cleft patient groups were also compared to a control group of normal
29 adults with skeletal and occlusal Class I relationships and average facial heights, before and
30 after surgery. A previous 3D soft-tissue evaluation of facial morphology changes after rapid
31 maxillary expansion and Delaire Facemask, in CLP children and the class III malocclusion
32 patients among the groups, were described.¹¹
33
34
35
36
37
38
39
40
41
42
43
44
45

46 The present study attempted to determine the soft-tissue changes in uCLP adult
47 patients who underwent LeFort I advancement and Rhinoplasty surgery, by comparing the
48 3D mean facial model (A) at T0 and T1 and also comparing A with a 3D mean facial model
49 B (control group without CLP), who underwent the same bone displacement.
50
51
52
53
54
55

56 Although the results of our 3D analyses should be interpreted with caution because of
57 the relatively small sample size, several conclusions from our data seem to be warranted.
58
59
60
61
62
63
64
65

1 It is well known that the morphology of the soft tissues, such as the nose and lips, as well as
2 the maxilla, changes after a Le Fort I osteotomy. They have been reported as a flattening and
3
4 thinning of the upper lip, expansion of the nose, and antero-superior movement of the nasal
5
6 tip.^{34,35}
7

8
9 Baek et al.³⁶ compared the treatment outcomes and relapse between maxillary
10 advancement surgery with LeFort I osteotomy (group 1) and maxillary distraction
11 osteogenesis (group 2) in 25 patients with cleft lip and palate with maxillary hypoplasia.
12 They founded that the nasal-labial angle increased more in group 2 than in group 1. In
13 addition, the forward movement of the upper lip and nasal tip was significantly greater in
14 group 2. These findings supported the results in the study of Chua and Cheung.³⁷
15
16

17 However, there are few reports that evaluated the form of these soft tissues three-
18 dimensionally.
19
20

21 In our study, the alar base width in CLP ~~was bigger~~ was wider and vertically increased
22 than in control pre-operatively and was reduced at T1 (Fig. 1, Table II). This finding is
23 consistent with previous results in the literature^{7, 38}, instead the upper vermilion remained
24 unchanged. This is probably because it is more difficult to detach tissues when there are
25 previous scars.
26
27

28 Otherwise, lengthening of the alar base width and a vertical increase of the upper
29 vermilion were evident in B (Fig. 1, Table II). The vertical increase of upper lip in B is
30 positive and probably due to the V-Y closure. The nasal alar widening seems to be due to the
31 release of the soft tissue attachment and muscle insertion. Once released from the bone,
32 muscles such as zygomaticus major, levator labii superioris, levator labii superioris alaeque
33 nasi, and nasalis give rise to lateral retraction, thus resulting in alar widening.
34
35
36
37
38
39
40

41 To prevent nasal alar widening and labial flattening after the maxillary osteotomy,
42 alar cinch suture and V-Y closure are conceptually good procedures. However, although
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 these procedures were performed, some reports, as well as this study, state that several
2 millimeters of widening of the nasal alae were still observed.⁷
3

4 The reasons may include insufficient sutures under general anesthesia with nasal intubation
5 and the short duration of tensile strength of the threads. For the aim of tightening up the nasal
6 alae, some tools such as an external fixator may be effective. Subspinal Le Fort I osteotomy
7 may be another solution for this problem.³⁹
8
9

10 An improvement of symmetry of the alar forms, both vertically and horizontally, were
11 observed in both groups. A normalization of the position of the tip of nose in the center of the
12 face was observed in A (Table II). These effects may be attributed to the closed rhinoplasty
13 which allowed the reshaping of the alar cartilage, columella and nasal dorsum.
14

15 In profile views, an increased support of the lips ~~were~~ was shown to a greater degree
16 in B than in A (Fig. 1- 4). Though a significant normalization of soft-tissue profiles was
17 generally observed in A, residual ~~defects~~ deformities were documented in the post-operative
18 upper lip and tip of nose projection (Fig. 1). This could be explained by the fact that patients
19 with CLP had scar contractures which prevented the correct soft-tissues countering. In
20 addition, ~~CLP patients had partially edentulous premaxilla so the upper lip was without~~
21 support from the teeth. in some of the CLP patients, one or two frontal teeth were missing at
22 the cleft side and this fact may contribute to reduce the upper lip support.
23
24

25 The main limit of this study is the small sample. More experience and further long-
26 term follow-up studies are needed to evaluate a much larger patient population with better
27 control over the variables.
28

29 In conclusion, 3D analysis performed in this study showed quantitative outcomes on
30 the secondary treatment of uCLP nose/lip/maxilla deformities. Our surgical procedures can
31 provide a symmetric nasal base and a satisfactory facial profile but some differences remain
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 in the post-operative frontal and profile views in uCLP patients, in comparison with the
2 control group.
3

4 Precise correction of secondary deformities in CLP adult patients still appears a
5 challenging task in maxillofacial surgery, and further technical improvements are still
6 possible.
7
8
9
10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES

1. Rhodes G, Sumich A, Byatt G. Are Average Facial Configurations Attractive Only Because of Their Symmetry? *Psychol Sci* 1999;10:52-58.
2. Meyer-Marcotti P, Alpers GW, Gerdes AB, Stellzig-Eisenhauer A. Impact of facial asymmetry in visual perception: a 3-dimensional data analysis. *Am J Ortod Dentofacial Ortop* 137:168.e1-8; discussion 2010;168-169.
3. Meyer-Marcotti P, Gerdes AB, Reuther T, Stellzig-Eisenhauer A, Alpers GW. Persons with cleft lip and palate are looked at differently. *J Dent Res* 2010;89:400-404.
4. Meyer-Marcotti P, Gerdes AB, Stellzig-Eisenhauer A, Alpers GW. Visual face perception of adults with unilateral cleft lip and palate in comparison to controls- an eye-tracking study. *Cleft Palate Craniofacial J* 2011;48:210-216.
5. van Loon B, Maal TJ, Plooij JM, Ingels KJ, Borstlap WA, Kuijpers-Jagtman AM, Spauwen PH, Bergé SJ. 3D Stereophotogrammetric assessment of pre- and postoperative volumetric changes in the cleft lip and palate nose. *Int J Oral Maxillofac Surg* 2010;39:534-40.
6. Okawachi T, Nozoe E, Nishihara K, Nakamura N. 3-dimensional analyses of outcomes following secondary treatment of unilateral cleft lip nose deformity. *J Oral Maxillofac Surg* 2011;69:322-332.
7. Nakamura N, Okawachi T, Nozoe E, Nishihara K, Matsunaga K. Three-dimensional analyses of nasal forms after secondary treatment of bilateral cleft lip-nose deformity in comparison to those of healthy young adults. *J Oral Maxillofac Surg* 2011;69:e469-e481.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
8. Spira M, Hardy SB, Gerow FJ. Correction of nasal deformities accompanying unilateral cleft lip. *Cleft Palate J* 1970; 7:112-123.
9. Jackson IT, Fasching MC. Secondary deformities of cleft lip, nose and cleft palate. In: McCarthy JG, May JW, Littler JW. Eds. *Plastic Surgery*. 1st edition, vol 4. Philadelphia: W.B. Saunders Company: 1990; p. 2771.
10. McCance AM, Moss JP, Fright WR, Linney AD, James DR. Three-dimensional analysis techniques-Part 1: three-dimensional soft-tissue analysis of 24 adult cleft palate patients following LeFort I maxillary advancement: a preliminary report. *Cleft Palate Craniofac J* 1997;34:36-45.
11. Sade Hoefert C, Bacher M, Herberts T, Krimmel M, Reinert S, Göz G. 3D soft tissue changes in facial morphology in patients with cleft lip and palate and class III malocclusion under therapy with rapid maxillary expansion and delaire facemask. *J Orofac Orthop* 2010;71:136-51.
12. Ramieri GA, Spada C, Nasi A, Tivolaccini A, Vezzetti E, Tornincasa S, Bianchi SD, Verzé L. Reconstruction of facial morphology from laser scanned data – Part I: Reliability of the technique. *Dentomaxillofacial Radiology* 2006;35:158-64.
13. Chigurupati R. Orthognathic surgery for secondary cleft and craniofacial deformities. *Oral Maxillofac Surg Clin North Am* 2005;17:503-517.
14. Ross RB. Treatment variables affecting facial growth in complete unilateral cleft lip and palate. *Cleft Palate J* 1987;24:5-77.
15. DeLuke DM, Marchand A, Robles EC, Fox P. Facial growth and the need for orthognathic surgery after cleft palate repair: literature review and report of 28 cases. *J Oral Maxillofac Surg* 1997;55:694-697.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
16. Good PM, Mulliken JB, Padwa BL. Frequency of Le Fort I osteotomy after repaired cleft lip and palate or cleft palate. *Cleft Palate Craniofac J* 2007;44:396-401.
 17. Daskalogiannakis J, Mehta M. The need for orthognathic surgery in patients with repaired complete unilateral and complete bilateral cleft lip and palate. *Cleft Palate Craniofac J* 2009;46:498-502.
 18. Broome M, Herzog G, Hohlfeld J, de Buys Roessingh A, Jaques B. Influence of the primary cleft palate closure on the future need for orthognathic surgery in unilateral cleft lip and palate patients. *J Craniofac Surg* 2010;21:1615-1618.
 19. Isiekwe MC, Sowemimo GO. Cephalometric findings in a normal Nigerian population sample and adult Nigerians with unrepaired clefts. *Cleft Palate J* 1984;21:323-328.
 20. Bishara SE, Jakobsen JR, Krause JC, Sosa-Martinez R. Cephalometric comparisons of individuals from India and Mexico with unoperated cleft lip and palate. *Cleft Palate J* 1986;23:116-125.
 21. Shetye PR. Facial growth of adults with unoperated clefts. *Clin Plast Surg* , 2004;31:361-371.
 22. Capelozza Filho L, Normando AD, daSilva Filho OG. Isolated influences of lip and palate surgery on facial growth: comparison of operated and unoperated male adults with UCLP. *Cleft Palate Craniofac J* 1996;33:51-56.
 23. Liao YF, Mars M. Long-term effects of palate repair on craniofacial morphology in patients with unilateral cleft lip and palate. *Cleft Palate Craniofac J* 42:594-600, 2005.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
24. Grayson BH, Santiago PE, Brecht LE, Cutting CB. Presurgical nasoalveolar molding in infants with cleft lip and palate. *Cleft Palate Craniofac J* 1999;36:486-498.
 25. Maull DJ, Grayson BH, Cutting CB, Brecht LL, Bookstein FL, Khorrambadi D, Webb JA, Hurwitz DJ. Long-term effects of nasoalveolar molding on three-dimensional nasal shape in unilateral cleft. *Cleft Palate Craniofac J* 1999;36:391-397.
 26. Nakamura N, Sasaguri M, Nozoe E, Nishihara K, Hasegawa H, Nakamura S. Postoperative nasal forms after presurgical nasoalveolar molding followed by medial-upward advancement of the nasolabial components with vestibular expansion for children with unilateral complete cleft lip and palate. *J Oral Maxillofac Surg* 2009;67:2222-2231.
 27. Seidenstricher-Kink LM, Becker DB, Govier DP, DeLeon VB, Lo LJ, Kane AA. Comparative osseous and soft tissue morphology following cleft lip repair. *Cleft Palate Craniofac J* 2008;45:511-517.
 28. Russell KA, Tompson B, Paedo D. Correlation between facial morphology and esthetics in patients with repaired complete unilateral cleft lip and palate. *Cleft Palate Craniofac J* 2009;46:319-325.
 29. Nakamura N, Suzuki A, Takahashi H, Honda Y, Sasaguri M, Ohishi M. A longitudinal study on influence of primary facial deformities on maxillofacial growth in patients with cleft lip and palate. *Cleft Palate Craniofac J* 2005;42:633-640.
 30. Wong JY, Oh AK, Ohta E, Hunt AT, Rogers GF, Mulliken JB, Deutsch CK. Validity and reliability of craniofacial anthropometric measurement of 3D digital photogrammetric images. *Cleft Palate Craniofac J* 2008;45:232-239.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
31. Bush K, Antonyshyn O. Three-dimensional facial anthropometry using a laser surface scanner: validation of the technique. *Plast Reconstr Surg* 1996;98:226-235.
 32. Littlefield TR, Kelly KM, Cherney JC, Beals SP, Pomatto JK. Development of a new three-dimensional cranial imaging system. *J Craniofac Surg* 2004;15:175-181.
 33. Yamada T, Mishima K, Moritani N, Janune D, Matsumura T, Ikeya Y, Yamamoto T. Nasolabial morphologic changes after a LeFort I osteotomy: a three-dimensional anthropometric study. *J Craniofac Surg* 2010;21:1089-1095.
 34. Altman JI, Oeltien JC. Nasal deformities associated with orthognathic surgery: analysis, prevention, and correction. *J Craniofac Surg* 2007;18:734-739.
 35. Mitchell C, Oeltien J, Panthaki Z, Thaller SR. Nasolabial aesthetics. *J Craniofac Surg* 2007;18:756-765.
 36. Baek SH, Lee JK, Lee JH, Kim MJ, Kim JR. Comparison of treatment outcome and stability between distraction osteogenesis and Le Fort I osteotomy in cleft patients with maxillary hypoplasia. *J Craniofac Surg* 2007;18:1209-1215.
 37. Chua HD, Cheung LK. Soft tissue changes from maxillary distraction osteogenesis versus orthognathic surgery in patients with cleft lip and palate-a randomized controlled clinical trial. *J Oral Maxillofac Surg* 2012;70: 1648-1658.
 38. Ferrario VF, Sforza C, Dellavia C, Tartaglia GM, Colombo A, Carù A. A quantitative three-dimensional assessment of soft tissue facial asymmetry of cleft lip and palate adult patients. *J Craniofac Surg* 2003;14:739-46.
 39. Mommaerts MY, Abeloos JV, De Clercq CA, Neyt LF. The effect of the subspinal Le Fort I-type osteotomy on interalar rim width. *Int J Adult Orthodon Orthognath Surg* 1997;12:95-100.

LEGENDS TO ILLUSTRATIONS

1
2
3
4
5 **Figure 1:** Mean facial model of uCLP patients (A) and control group (B) at T0, T1.
6
7

8
9
10 **Figure 2:** Demonstration of 10 facial landmarks taken from classical anthropometry
11 employing 3D analyses.
12
13

14
15
16
17 **Figure 3:** Axial sections of the 2 superimposed shells (T0: green; T1: red) at different levels
18 passing through prn, sn, ls. A: mean facial model of uCLP patients; B: mean facial model of
19 control group.
20
21
22
23

24
25
26
27 **Figure 4:** Sagittal sections of superimposed shells (T0: green; T1: red). A: mean facial model
28 of uCLP patients; B: mean facial model of control group.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure
[Click here to download high resolution image](#)






Figure
[Click here to download high resolution image](#)



Figure
[Click here to download high resolution image](#)



A



B



Table I. Patients involved in the study.

A (uCLP)							B (Control)						
Surgery							Surgery						
Patient	CLP side	Sex	Age	Deformity of the nose	LeFort I (Advancement mm)	RSP	Patient	Angle	Sex	Age	Deformity of the nose	LeFort I (Advancement mm)	RSP
1	right	F	18	b, c	5	B, +	1	III	F	24	b, e	5	A, C, E
2	right	F	35	a, d, e	5	C, E	2	III	F	22	e	5	A, C, +
3	right	F	18	c, e	6	B, D	3	III	F	22	c, e	6	C, E
4	left	F	39	d	7	‡	4	III	F	45	e	7	C, E
5	left	M	20	b, d	7	†	5	III	M	23	e	7	C, D
6	left	M	21	b, d, e	8	A, B, C, E	6	III	M	32	e	8	A, C, E

Abbreviations: **uCLP**: unilateral cleft lip and palate.

Deformity of the nose. **a**: deviated columella; **b**: depressed and/or deviated nasal tip; **c**: wide nasal ala; **d**: flat and v-shaped nostril; **e**: hump.

RSP: Rhinoplasty surgery. †: costal cartilage graft; ‡: auricular cartilage graft; +: quadrangular cartilage graft. **A**: symmetrizing of alar cartilages ;

B: basal osteotomy; **C**: hump; **D**: septum; **E**: tip of the nose (upward rotation).

Table II. Point to point distances of the landmarks considered.

		A	B	
Frontal view	al _r -al _l	T0	35.22	31.10
		T1	33.62	32.06
	al _r -prn	T0	24.19	25.30
		T1	22.32	23.19
	al _l -prn	T0	27.53	22.09
		T1	25.90	22.18
	al _r -X	T0	36.53	40.21
		T1	37.02	44.69
	al _l -X	T0	38.11	44.27
		T1	37.10	45.57
	prn-Y	T0	1.60	0.97
		T1	0.53	0.87
	sn-ls	T0	11.08	12.39
		T1	11.92	12.24
ch _r -ch _l	T0	51.20	52.89	
	T1	50.38	53.01	
ls-sto	T0	3.18	5.64	
	T1	3.27	7.39	
Lateral view	prn-sn-ls °	T0	134.05	133.05
		T1	137.54	133.47
	prn-X	T0	32.98	37.64
		T1	32.10	37.23
	ls-Y ₁	T0	38.64	38.56

T1	39.52	40.08
----	-------	--------------

The values are in millimetres (or degrees for angles).

A: uCLP patients; **B**: control group. **X**: horizontal reference plane; **Y**: vertical reference plane through glabella; **Y1**: vertical plane through left endocanthion and perpendicular to X; **r**: right; **l**: left.

(Differences of linear measurement major of 1 mm and angular measurements superior to 3 grades were considered significant and highlighted in bold).