

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Quality of life among healthcare workers: A multicentre cross-sectional study in Italy.

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/129666> since

Published version:

DOI:10.1016/j.puhe.2012.03.006

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in:

**[Public Health, Vol. 126, issue 7, July 2012,
DOI: 10.1016/j.puhe.2012.03.006]**

<http://www.sciencedirect.com/science/article/pii/S0033350612001084>

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect® platform]

The quality of life in the healthcare professionals: a multicenter study in Italy

Flavia Kheiraoui^{1§*}, Maria Rosaria Gualano^{1*}, Alice Mannocci^{2*}, Walter Ricciardi^{1*}, Antonio Boccia^{2*}, Giuseppe La Torre^{2*}

¹Institute of Hygiene, Catholic University of the Sacred Heart, Largo Francesco Vito 1, 00168 - Rome, Italy

²Clinical Medicine and Public Health Unit, Sapienza University of Rome, Viale Regina Elena, 324 00161 – Rome, Italy

*These authors contributed equally to this work

§Corresponding author:

Flavia Kheiraoui

Email addresses:

FK: flavia.kheiraoui.iphu@rm.unicatt.it

MRG: mar.guala@gmail.com

AM: alice.mannocci@uniroma1.it

WR: wricciardi@rm.unicatt.it

AB: antonio.boccia@uniroma1.it

GLT: giuseppe.latorre@uniroma1.it

Abstract

Background

The study aims to evaluate the quality of life in healthcare professional, physicians, nurses and prevention technicians for health safety at workplaces and environment's safeguard.

Methods

A cross sectional study was carried out in a population of health professionals in ten Italian Regions, to evaluate the health-perceived status related to their role.

Health status of participants was assessed using the Italian version of the SF-36 questionnaire. Differences for quantitative variables between groups were tested using T-test and ANOVA. A multiple linear regression was performed to assess the influence on the SF-36 scales of the variables age, gender, role, socializing time, working time, years spent in the health care system and in the specific department.

Results

Sample was comprised of 324 healthcare workers (57.1% women, mean age 39.04 ± 10.15), of which 52.6% was medical doctors, 36.8% nurses, 10.5% prevention technicians.

Differences in Mental Health score was positively associated with the amount of socializing time. Workers with a job career longer than 15 years achieved a General Health score lower than those spending less time.

The multivariate analysis showed that increasing age is positively associated with Role Emotional level ($\beta = 0.243$; $p = 0.002$), while an inverse relationship appears for General Health ($\beta = -0.218$; $p = 0.007$) and Physical Function ($\beta = -0.246$; $p = 0.001$). For the roles, nurses seem to experience lower values of Bodily Pain ($\beta = -0.214$; $p < 0.001$), Social Function ($\beta = -0.242$; $p = 0.001$), Role Emotional ($\beta = -0.211$; $p = 0.006$) scores if compared with physicians. By comparing the mean scores of the sample with the Italian general population, the healthcare workers achieved higher values of General

Health, Physical Function, Role Physical, Bodily Pain and Mental Health while lower ones resulted for Vitality, Social Function and Role Emotional.

Conclusions

Healthcare professionals have different Health related quality of life results. In particular, nurses is a professional category with low levels of quality of life. These results can be taken into account in order to recognize the roles and the attitudes that could mainly causes frustration, dissatisfaction and emotional stress in the healthcare professionals.

Background

What is currently defined as the Quality of Life (QoL) in bio-medical issues corresponds in reality to what the English authors call “Health Related Quality of Life” and concerns to those aspects of quality of life that are related to disease and health and therefore “adjusted” by the medicine [1].

The concept of Quality of Life requires to add to the biomedical factors other factors such as:

- Individual factors
- Environmental factors
- Economical factors

The perception of health and quality of life in patients has been an interesting target of intensive investigation in recent years, but for the healthcare personnel the literature focusing on their quality of life in Italy is scarce, except for some specific items, and these issues need to be better appreciated. It is important to understand healthcare personnel’s characteristics and explore the relationship of their level of burden with quality-of-life parameters and psychosocial aspects of their job. This will optimize the use of support and interventional measures and help to reduce negative effects on their lives. Minimizing the burden on healthcare personnel will possibly improve the quality of life and medical outcomes of their patients and the relationships with their private life.

There is evidence of elevated rates of psychological stress and stress related disorder do exist among hospital staff [2].

For example, Burnout syndrome (BOS) was identified in the early 1970s in human service professionals, most notably healthcare workers. Burnout syndrome has been described as an inability to cope with emotional stress at work or as excessive use of energy and resources leading to feelings of failure and exhaustion. Although depression affects nearly every aspect of the person's life, symptoms of burnout occur only at work. Wide variations in the prevalence of Burnout in health care professionals have been reported across specialties, both in doctors and in nurses. Workplace climate and workload were determinants of Burnout syndrome [3-6].

Concerning the possible predictors of burnout, Shanafelt et al. reported that physicians who spent less than 20% of time working on the most meaningful activities had a higher risk of burnout syndrome [7].

Hospitals and health prevention unit are characterized by a high level of work-related stress, a factor known to increase the risk of low quality of life [8-11].

So, the working lives of physicians and nurse are often characterized by a high level of work-related stress: tiredness, high turnover, night shift, workload and stressful work environments like intensive care units, severity of illness and conflicts with co-workers or with patients [12].

The workload of Intensive Care Unit physicians, as an example, is physically demanding, allows limited rest and is associated with sleep deprivation and objective markers of physiologic stress (ketonuria, arrhythmia or heart rate abnormalities)[13], that may causes rebounds in their private and social life.

Also the prevention technicians are exposed to high level of stress. They have the civil and legal liability of their role, their workload is often related to the needs of the health service surveillance and they have to guarantee the availability in their activities, so they often have no spare time for the human relationships, except that with their colleagues.

The work environment and psychosocial factors have been considered largely responsible for the health problems observed in hospital's staff.

Psychosocial aspects refer to the interaction between work environment, content and conditions and worker capacity, needs, culture, extra-work personal elements, which may, according to perception and experience, influence health, satisfaction, and work performance [14-16].

The aim of this study was to evaluate the health-perceived status of healthcare professionals, in particular of :

- medical doctors;
- nurses;
- prevention technicians for health safety at workplaces and environment's safeguard.

Moreover, we wanted to compare the results from the above healthcare personnel with those from the general population.

Methods

Population and setting

A pilot cross sectional study was carried out, in a population of health professionals, in the following Italian Regions: Abruzzi, Calabria, Emilia Romagna, Lazio, Liguria, Lombardy, Marche, Sicily, Apulia and Tuscany. The selection of the participants was made recruiting healthcare personal, participating in courses of continual medical education in all the Regions described .

Concerning physicians and nurses, they are representative of the whole activity of the Italian's structures, working in a plurality of wards and contests in the regions that we chose in the Italian territory and for the prevention technicians we selected the staff of the whole activity of the Prevention's Department (safety and prevention at workplace, public health service, veterinary

service and food safety service) in the area of the central Italy, specifically Rome, the Province of Rome and the Province of Latina.

Quality of life measurement

The health status of participants was assessed using the Italian version of the Short-Form 36 item questionnaire. The crude estimates were transformed, using the procedure described by Apolone and Mosconi [17-20].

Statistical analysis

Descriptive analysis

The aim of this analysis was to describe quantitative and categorical variables means and standard deviations (SD) and frequencies were used, respectively.

Univariate and multivariate analysis

Differences between groups in the SF-36 scales were tested using T-test and ANOVA test. Moreover, the means of the variables (SF -36 Scores) of our sample were compared with the means of the Italian sample [21] using the T-test. Furthermore, we conducted a multivariate analysis using a linear regression in order to assess the influence on the SF-36 scales (as dependent variables) of the explanatory variables: age, gender, role, time of socializing in hours per week, time of working in hours per week, years spent in health system, years spent in the specific department. We selected variables that had a $p\text{-value} < 0.25$ at univariate analysis, according with Hosmer and Lemeshow procedure [22].

The level of significance was set at $p \leq 0.05$. Statistical analysis was performed with SPSS 12.0 software for Windows.

Results

The sample was comprised of 324 healthcare workers. Of them, 57.1% (185) were women. The mean age of this sample was 39.04 ± 10.15 . The characteristics of the sample are showed in Table 1.

52.6 % (170) were medical doctors, 36.8% (119) were nurses and 10.5 % (34) were prevention technicians. The mean of years spent in the Health Care Service was 16.64 ± 9.08 .

At the univariate analysis women had Bodily Pain scores lower than men, and this result was statistically significant ($p=0.005$).

We found differences for the outcome Mental Health that results to be positively associated with the amount of socializing time: in fact who spent more time in social activities seems to have a higher Mental Health score ($p=0.009$ at T- test). Moreover, health care workers with a job career longer than 15 years achieved a General Health score lower than those spending less working time in Health System ($p=0.005$).

Moreover, the healthcare workers spending in socializing activities more than 8 hours for week had a Mental Health score higher than others ($p=0.009$), while who worked more than 40 hours for week had a lower Vitality score ($p=0.005$).

All the other results of the univariate analysis are showed in Table 2 and 3.

As shown in Table 4, results of the multivariate analysis showed that increasing age results to be positively associated with Role Emotional level ($\beta= 0.243$; $p=0.002$), while an inverse relationship appears for General Health and Physical Function levels ($\beta=-0.218$; $p=0.007$ and $\beta=-0.246$; $p=0.001$, respectively).

Moreover, nurses seem to experience lower values of Bodily Pain ($\beta=-0.214$; $p<0.001$), Social Function ($\beta=-0.242$; $p=0.001$), Role Emotional ($\beta=-0.211$; $p=0.006$) scores if compared with physicians.

As concerning gender, women achieved lower values of Vitality and Mental Health scores ($\beta=-0.210$; $p=0.007$ and $\beta=-0.168$; $p=0.046$, respectively) than men do.

Additionally in Table 5 findings of comparing the mean scores of the study sample with the Italian general population [21] are reported: the healthcare workers achieved higher values with statistically significant differences for General Health ($p=0.002$), Physical Function ($p<0.0001$), Mental Health ($p=0.042$) while lower ones resulted for Vitality ($p=0.019$), Social Function ($p<0.0001$) and Role Emotional ($p=0.015$).

Discussion

To our knowledge, this is the first Italian study evaluating the quality of life in the healthcare staff and its perception among different categories of health care workers as physicians, nurses and technicians.

Even if a strong association between healthcare working and occupational stress has already been described by several Italian studies, how different categories of health care workers face the potential effect of healthcare working has not yet been fully explored [23-27].

By comparing the mean scores of the study sample with the Italian general population [21], the health care workers achieved higher values of General Health, Physical Function, Role Physical, Bodily Pain and Mental Health while lower ones resulted for Vitality, Social Function and Role Emotional.

Our findings show that health care workers have a more positive attitude towards their health status if compared with the general population, probably due to the type of daily activities closely

connected with physical suffering, pain and emotional distress. Additionally, our study puts in evidence that women seem to be more sensitive to health care work-related stress, probably because to their greater empathy respect to men.

Workers in the health systems are constantly called to handle feelings and “strong emotions”, they bring to the workplace an emotional "baggage” that may undermine the professional skills and their lack of vitality , as our results show, in our opinion, was due to restrictions on their personal time because they spend many hours each day at workplace and to the sense of failure regarding their hopes and expectations at work. We suppose that it could be useful to act in two ways. On one hand, improving all those interventions targeted to training, that is an effective approach that delineates methods for revitalizing, reawakening the ideals of the health profession and increasing energy and enthusiasm for working. On the other hand, giving health care professionals the opportunity to benefit from interventions oriented to cope the sense of failure and the physical stress, for example an individual counselling or continued informal support. The improve of good relationships with colleagues and positive coping mechanisms could be good to prevention depression [28]. May be needed a review of the workload in our Health System, because high turnover, night shifts and workload seems, according with our questionnaires, so hard and tiring and that often causes rebounds in the private and social life [29].

Some limitations and strengths of this study should be reported.

First of all this study has some limitations related to the study design because we performed a cross-sectional study, but we supposed that SF- 36 is the best way to investigate these topics. For the strengths, we considered a sample distributed in different Italian regions, to acquire knowledge and data before missing regarding quality of life in different categories of health care professionals [30-32]. Moreover, to avoid confounding we performed the multivariate analysis so the results were adjusted for possible confounders.

Conclusions

This study put in evidence that the health care professionals have a different perception of quality of life, based on their different role. The nurses and the women seem to experience the worst results related to the emotional factors.

We showed like the health care workers have a better perception of life if compared with the general population, even if they are daily exposed to physical suffering, pain and emotional distress. It is important that as health professionals, we recognize the roles and the attitudes that could mainly causes frustration, dissatisfaction and emotional stress, to prevent the development of these symptoms in order to protect our colleagues and the people, patients or not, that we meet everyday in our job.

Competing interests

The Authors stated they have no conflict of interest.

Authors' contributions

FK administered questionnaires, participated in the design of the study and drafted the manuscript. MRG performed the statistical analysis and helped to draft the manuscript. AM performed the statistical analysis. GLT conceived of the study and designed the study and WR and AB carried out the coordination.

All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Paola Santini for helping the authors in administering the questionnaires to the participants.

References

1. World Health Organization: WHOQoL Study Protocol. 1993.
2. Fichera GP, Sartori S, Costa G: **Post-traumatic stress disorder following robbery at the workplace: a pilot study on 136 pharmacy workers.** *Med Lav.* 2009; **100**(2):97-108.
3. Häggström E, Skovdahl K, Fläckman B, Kihlgren AL, Kihlgren M: **Work satisfaction and dissatisfaction--caregivers' experiences after a two-year intervention in a newly opened nursing home.** *J Clin Nurs.* 2005;**14**(1):9-19.
4. Poncet MC, Toullic P, Papazian L, Kentish-Barnes N, Timsit JF, Pochard F, Chevret S, Schlemmer B, Azoulay E: **Burnout syndrome in critical care nursing staff.** *Am J Respir Crit Care Med* 2007 ;**175**(7):698-704.
5. Embriaco N, Papazian L, Kentish-Barnes N, Pochard F, Azoulay E: **Burnout syndrome among critical care healthcare workers.** *Curr Opin Crit Care.* 2007;**13**(5):482-8
6. Parshuram C, Dhanani S, Kirsh J, Cox P : **Fellowship training, workload, fatigue and physical stress: a prospective observational study.** *CMAJ*; 2004, **170**:965–970 .
7. Shanafelt TD, West CP, Sloan JA, Novotny PJ, Poland GA, Menaker R, Rummans TA, Dyrbye LN: **Career Fit and Burnout Among Academic Faculty.** *Arch Intern Med.* 2009 25; **169**(10):990-5.
8. Chang EM, Bidewell JW, Huntington AD, Daly J, Johnson A, Wilson H, Lambert VA, Lambert CE.: **A survey of role stress, coping and health in Australian and New Zealand hospital nurses.** *Int J Nurs Stud* 2007;**44**(8):1354-62.
9. Sherman AC, Edwards D, Simonton S, Mehta P: **Caregiver stress and burnout in an oncology unit.** *Palliat Support Care.* 2006 ;**4**(1):65-80.

10. Jonsson A, Halabi J: **Work related post-traumatic stress as described by Jordanian emergency nurses.** *Accid Emerg Nurs.* 2006 ;**14**(2):89-96.
11. Hamaideh SH, Mrayyan MT, Mudallal R, Faouri IG, Khasawneh NA : **Jordanian nurses' job stressors and social support .** *Int Nurs Rev* 2008;**55**(1):40-7.
12. de Carvalho EC, Muller M, de Carvalho PB, de Souza Melo A: **Stress in the professional practice of oncology nurses.** *Cancer Nurs.* 2005;**28**(3):187-92.
13. Embriaco N, Azoulay E, Barrau K, Kentish N, Pochard F, Loundou A, Papazian L : **High level of burnout in intensivists: prevalence and associated factors.** *Am J Respir Crit Care Med.* 2007 1;**175**(7):686-92.
14. Klersy C, Callegari A, Martinelli V, Vizzardi V, Navino C, Malberti F, Tarchini R, Montagna G, Guastoni C, Bellazzi R, Rampino T, David S, Barbieri C, Dal Canton A, Politi P: **Working Group on Burnout and Dialysis. Burnout in health care providers of dialysis service in Northern Italy--a multicentre study .** *Nephrol Dial Transplant* 2007;**22**(8):2283-90.
15. International Labour Office: **Psychosocial factors at work: recognition and control.** Geneva: ILO; 1986.
16. L. Luchettiab, E. Uhunmwanghoa, G. Dordonia, A. Loridaab, S. Barbieriab, A.G. Bolognesiab, G. Gobbib, F. Franchib: **The subjective feeling of burden in caregivers of elderly with dementia: how to intervene?** *Arch Gerontol Geriatr.* 2009;**49** Suppl 1:153-61.
17. Apolone G, Mosconi P: **Questionario sullo stato di salute SF-36. Manuale d'uso e guida all'interpretazione dei risultati.** Istituto di Ricerche Farmacologiche "Mario Negri", Milano. Guerini e Associati, Abbiategrosso (MI) 2007.
18. Ware, Snow, Kosinski, & Gandek: **SF-36 user's manuals,** 1993.
19. Ware, Kosinski, & Keller: **SF-36 user's manuals,**1995.
20. Keller SD, Ware JE Jr, Bentler PM, Aaronson NK, Alonso J, Apolone G, Bjorner JB, Brazier J, Bullinger M, Kaasa S, Leplège A, Sullivan M, Gandek B: **Use of structural equation modeling to**

- test the construct validity of the SF-36 Health Survey in ten countries: results from the IQOLA Project. International Quality of Life Assessment.** *J Clin Epidemiol.* 1998 ;**51**(11):1179-88.
21. Nante N, Groth N , Guerrini M, Galeazzi M,. Kodraliu G , Apolone G: **Using the SF-36 in a rural population of elderly in Italy: a pilot study.** *Journal of Preventive Medicine and Hygiene* 1999; **40**: 1-7.
22. Hosmer DW, Lemeshow S: **Applied logistic regression.** New. York, NY: John Wiley & Sons, 1989.
23. Patti F, Amato MP, Battaglia MA, Pitaro M, Russo P, Solaro C, Trojano M : **Caregiver quality of life in multiple sclerosis: a multicentre Italian study.** *Mult Scler.* 2007;1 **3**(3):412-9. Epub 2007 Jan 29.
24. Pozzilli C, Palmisano L, Mainero C, Tomassini V, Marinelli F, Ristori G, Gasperini C, Fabiani M, Battaglia MA: **Relationship between emotional distress in caregivers and health status in persons with multiple sclerosis.** *Mult Scler.* 2004; **10**(4):442-6.
25. Aprile I, Caliandro P, La Torre G , Tonali P, Foschini M, Mondelli M, Bertolini C. Piazzini D.B, Padua L: **Multicenter study of peroneal mononeuropathy: clinical, neurophysiologic, and quality of life assessment.** *J Peripher Nerv Syst.* 2005; **10**(3):259-68.
26. Tabolli S, Baliva G, Lombardo GA, Sampogna F, Di Pietro C, Mannooranparampil TJ, Alvetreti G, Abeni D: **Health related quality of life assessment in the routine clinical practice of a dermatology unit.** *Eur J Dermatol.* 2006 ;**16**(4):409-15.
27. Bressi C, Manenti S, Porcellana M, Cevalles D, Farina L, Felicioni I, Meloni G, Milone G, Miccolis IR, Pavanetto M, Pescador L, Poddigue M, Scotti L, Zambon A, Corrao G, Lambertenghi-Deliliers G, Invernizzi G: **Haemato-oncology and burnout: an Italian survey.** *Br J Cancer.* 2008 ;**98**(6):1046-52.

28. Penson RT, Dignan FL, Canellos GP, Picard CL, Lynch TJ Jr. **Burnout: caring for the caregivers.** *Oncologist.* 2000;**5**(5):425-34.
29. Drach-Zahavy A: **Workplace health friendliness: a cross level model for predicting workers' health.** *J Occup Health Psychol.* 2008;**13**(3):197-213.
30. La Torre G, Moscato U, La Torre F, Ballini P, Marchi S, Ricciardi W: **Environmental noise exposure and population health: a cross sectional study in the Province of Rome.** *J Public Health* 2007; **15**:339-334.
31. de Vet HC, Adèr HJ, Terwee CB, Pouwer F: **Are factor analytical techniques used appropriately in the validation of health status questionnaires? A systematic review on the quality of factor analysis of the SF-36.** *Qual Life Res.* 2005;**14**(5):1203-18.
32. Ware JE Jr, Gandek B, Kosinski M, Aaronson NK, Apolone G, Brazier J, Bullinger M, Kaasa S, Leplège A, Prieto L, Sullivan M, Thunedborg K: **The equivalence of SF-36 summary health scores estimated using standard and country-specific algorithms in 10 countries: results from the IQOLA Project. International Quality of Life Assessment.** *J Clin Epidemiol.* 1998; **51**(11):1167-70.

Table 1. Characteristics of the sample

Characteristics		Frequencies N (%)
Gender	M	133 (42.9)
	F	182 (57.1)
Role	Physicians	166 (52.6)
	Nurses	118 (36.8)
	Technicians	31 (10.5)
Age		Mean±SD
		39.04±10.15
Macro Regions	North -Central	105 (33.1)
	South- Central	212 (66.8)
Working time (hours)	≥ 40	185 (58.4)
	≤ 40	131 (41.4)
Socializing time per week (hours)	≥ 8	176 (55.6)
	≤ 8	140 (44.3)
Years in healthcare system	≤ 15	211 (66.7)
	≥ 15	105 (33.2)

Table 2. Results of univariate analysis for SF-36 scales according to gender and age groups

	General health (GH)	Physical function (PF)	Role physical (RP)	Bodily pain (BP)	Social functioning (SF)	Vitality (VT)	Emotional Role and status (ER)	Mental health (MH)
Variables (N)	Media±SD	Media±SD	Media±SD	Media±SD	Media±SD	Media±SD	Media±SD	Media±SD
Gender								
M (133)	70.43±15.23	93.12±12.62	82.70±29.07	80.24±21.50	69.64±20.94	62.33±17.77	75.06±34.31	71.62±15.48
F (182)	68.23±18.41	92.65±11.00	76.65±32.14	73.09±22.08	66.35±23.22	56.69±19.53	67.86±35.99	67.09±18.15
p	0.19	0.39	0.068	0.005	0.12	0.008	0.084	0.015
Age								
<40 (180)	71.25±17.62	93.97±9.29	77.08±31.37	75.72±23.03	64.65±22.57	56.72±19.42	65.46±36.01	66.75±17.26
≥40 (136)	66.39±16.08	91.42±14.16	82.17±30.87	76.68±21.77	71.70±21.36	62.14±17.93	78.30±33.16	72.07±16.64
p	0.012	0.048	0.10	0.53	0.004	0.007	0.002	0.006

p-value at t-test

Table 3. Results of univariate analysis for SF-36 scales according to professional factors

	General health (GH)	Physical function (PF)	Role physical (RP)	Bodily pain (BP)	Social functioning (SF)	Vitality (VT)	Emotional Role and status (ER)	Mental health (MH)
Variables (N)	Media±SD	Media±SD	Media±SD	Media±SD	Media±SD	Media±SD	Media±SD	Media±SD
Role								
Physicians (166)	71.04±16.84	94.58±9.98	82.83±29.23	80.49±19.90	69.50±21.86	56.87±18.21	73.69±34.38	67.89±16.14
Nurses (118)	66.81±17.69	91.13±10.63	72.45±33.07	69.54±24.16	64.30±23.81	59.95±19.47	66.24±36.62	70.07±18.98
Tecnicos (31)	68.45±15.83	90.16±20.18	86.29±30.84	79.03±22.83	71.37±17.13	68.06±18.33	74.73±35.71	72.25±14.45
p [^]	0.133	0.043	0.011	0.001	0.095	0.008	0.211	0.337
Socializing time per week (hours)								
<8 (140)	68.08±18.11	93.27±10.87	77.05±32.06	74.94±21.77	66.16±22.57	54.83±18.35	68.45±35.07	66.07±16.44
≥8 (176)	69.45±16.34	92.56±12.32	80.68±30.53	77.09±23.04	68.89±22.79	62.41±18.81	73.01±35.62	71.04±17.43
p [°]	0.772	0.773	0.355	0.495	0.347	<0.001	0.243	0.009
Working Time (hours)								
<40 (131)	66.78±17.07	91.05±13.54	79.39±32.00	75.20±22.62	69.46±23.41	61.33±17.63	71.24±36.40	70.09±16.84
≥40 (185)	70.85±17.00	94.16±10.01	79.19±30.72	76.08±22.04	66.41±21.44	57.44±19.72	70.81±34.76	68.29±17.41
p [°]	0.036	0.038	0.86	0.576	0.214	0.050	0.954	0.341
Years in health system								
<15 (211)	71.03±16.85	93.60±9.59	77.61±31.42	78.80±22.77	65.52±22.41	57.69±19.17	68.01±35.55	68.00±17.76
≥15 (105)	65.43±17.13	91.41±14.99	82.62±30.64	76.80±21.95	72.02±21.55	61.08±18.29	76.98±34.47	71.13±15.83
p [°]	0.005	0.097	0.205	0.897	0.016	0.054	0.041	0.141
Years in specific word								
<6 (160)	70.44±17.22	93.11±9.91	77.97±30.63	74.63±21.99	67.27±21.49	56.20±19.14	69.58±34.55	67.13±17.53
≥6 (124)	67.08±17.35	93.19±10.99	79.23±32.15	77.64±22.89	67.44±24.48	60.69±18.06	71.91±36.71	70.94±17.19
p [°]	0.200	0.882	0.823	0.378	0.98	0.48	0.553	0.090

[°] t-test[^] Anova

Table 4. Results of multivariate analysis for each SF-36 Scale (only significant results of linear regressions are shown)

Outcomes	Variables		Coefficiente β	P-value	R²
General health (GH)	Age		-0.218	0.007	0.048
Physical function (PF)	Age		-0.246	0.001	0.061
Social function (SF)	Nurses		-0.243	0.001	0.059
	Physicians*				
Role physical (RP)	Nurses		-0.204	0.006	0.042
	Physicians*				
Bodily pain (BP)	Nurses		-0.214	<0.001	0.046
	Physicians*				
Vitality (VT)	Gender	M*			0.137
		F	-0,210	0.007	
	Socializing Time (hours)		0,151	0.048	
	Working Time (hours)		-0,291	<0.001	
Emotional Role and status (ER)	Age		0.243	0.002	0.124
	Nurses		-0.211	0.006	
	Physicians*				
Mental health (MH)	Gender	M*			0.028
		F	-0.168	0.046	

* Reference group

Table 5 . Comparisons of SF-36 score means between Healthcare professional and the Italian general population

SF 36 scores	Italy 1995 (Mean±SD)	Health professional 2009 (Mean±SD)	p-value
Physical function (PF)	84.46±23.18	92.83± 11.73	< 0.0001
Role physical (RP)	78.21±35.93	79.27±31.06	0.601
Bodily pain (BP)	73.67±27.65	75.92±22.72	0.149
General health (GH)	65.22±22.28	69.14±17.05	0.002
Vitality (VT)	61.89±20.69	59.14±18.91	0.019
Social function (SF)	77.43±23.34	67.69±22.16	< 0.0001
Emotional Role and status (ER)	76.16±37.25	71.04± 35.31	0.015
Mental health (MH)	66.59±20.89	68.99 ±17.13	0.042

