

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Fluorescent in situ hybridization mapping of three fecundity genes on cattle, river buffalo, sheep and goat

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1530131> since 2015-11-28T12:47:36Z

Published version:

DOI:10.1080/00087114.2014.996038

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[Caryologia: International Journal of Cytology, Cytosystematics and Cytogenetics, 2015, 68:1, 9-12, DOI: [10.1080/00087114.2014.996038](https://doi.org/10.1080/00087114.2014.996038)]

ovvero [Alessandra Iannuzzi, Angela Perucatti, Alfredo Pauciullo, Viviana Genuardo, Lisa De Lorenzi, Pietro Parma, Leopoldo Iannuzzi (2015) Fluorescent in situ hybridization mapping of three fecundity genes on cattle, river buffalo, sheep and goat, Caryologia: International Journal of Cytology, Cytosystematics and Cytogenetics, 68:1, 9-12, DOI: [10.1080/00087114.2014.996038](https://doi.org/10.1080/00087114.2014.996038).]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<http://dx.doi.org/10.1080/00087114.2014.996038>]

1 **Fluorescent in situ hybridization mapping of three fecundity genes on cattle, river buffalo,**
2 **sheep and goat**

3
4
5 Alessandra Iannuzzi^{a*}, Angela Perucatti^a, Alfredo Pauciullo^a, Viviana Genualdo^a, Lisa De Lorenzi^b,
6 Pietro Parma^b and Leopoldo Iannuzzi^a

7
8
9 ^aLaboratory of Animal Cytogenetics and Gene Mapping, National Research Council (CNR),
10 Institute of Animal Production Systems in Mediterranean Environments (ISPAAM), Via Argine,
11 1085, 80147, Naples, Italy

12
13 ^bDepartment of Animal Science, Agricultural Faculty of Sciences, Milan, Italy

14
15
16
17
18 *Corresponding author. Email: alessandra.iannuzzi@cnr.it

19

1 **ABSTRACT**

2 One of the goals of molecular cytogenetics applied to livestock is the extension of their genetic
3 physical maps, especially of loci containing genes related to productions. In this study, a
4 comparative fluorescence in situ (FISH) mapping of three genes related to fecundity of cattle, river
5 buffalo, sheep and goat is reported using bovine BAC-clones taking in account the data available on
6 the BovMap database and considering their physical position and the data obtained from banding
7 experiments. The following three gene sequences were mapped: tumor necrosis factor- α (TNF),
8 correlated to male fertility; signal transducer and activator of transcription 5A (STAT5A), important
9 for its influence on milk production and reproduction activity; melatonin receptor 1A (MTNR1A)
10 important for reproductive seasonality. BAC probes containing these gene sequences were assigned
11 by FISH, for the first time, on RB-banded chromosomes of these four important bovids. TNF was
12 assigned to BTA/CHI23q21-22, OAR20q21-22 and BBU 2p21-22; STAT5A was assigned to
13 BTA/CHI19q17-21, OAR11q17-21 and BBU3p15-21; MTNR1A was assigned to BTA/CHI27q14-
14 15, OAR26q14-15 and BBU1p21-22. The three loci were located in homoeologous chromosomes
15 and chromosome bands, underling the high degree of chromosome homologies among Bovids and
16 extending the cytogenetic maps of this economically important species.

17

18 **Keywords:** FISH-mapping, fecundity genes, cattle, sheep, goat, river buffalo

19

1 INTRODUCTION

2 Cytogenetic mapping is a method used to construct physical maps which are useful tools for
3 various applications, especially in animal cytogenetics. Indeed, it allows: (a) a precise physical
4 position on single chromosome bands of both type I and type II loci, especially in bovids using
5 FISH mapping on R-banded chromosome preparations (Di Meo et al. 2007); (b) to confirm
6 chromosomes and chromosome regions involved in chromosome abnormalities by using specific
7 molecular markers (Di Meo et al. 2000; Perucatti et al. 2011; Iannuzzi et al. 2013); (c) to study
8 chromosome aneuploidies in sperms and oocytes (Pauciullo et al. 2011, 2012; Hornak et al. 2011);
9 (d) to precisely anchor linkage and RH maps (Stafuzza et al. 2013), as well as genome sequence
10 contigs to specific chromosome regions (Goldammer et al. 2009).

11 Cattle (*Bos taurus*, $2n = 60$, BTA), river buffalo (*Bubalus bubalis*, $2n = 50$, BBU), sheep
12 (*Ovis aries*, $2n = 54$, OAR) and goats (*Capra hircus*, $2n = 60$, CHI) are very related species from the
13 evolutionary point of view and, also, the four major domestic bovid species of great economic
14 importance. Although the location of a lot of genes in these species was identified by linkage and
15 RH mapping, a small percentage of those loci were physically assigned to the corresponding bands
16 of specific chromosomal location (Iannuzzi et al. 2003a). So far, several studies on the physical
17 gene mapping using FISH methodology were reported for cattle, river buffalo, sheep, goat and other
18 farm animals (Iannuzzi et al. 2003a, 2003b; Di Meo et al. 2007; Schibler et al. 2009).

19 In the present study, three important fecundity genes (TNF, STAT5A and MTNR1A) were
20 comparatively FISH mapped on cattle, sheep, goat and river buffalo R-banded chromosomes for
21 first time extending the cytogenetic maps of these species. Tumor necrosis factor- α (TNF) is
22 correlated to male fertility (Eggert-Kruse et al. 2007; Kocak et al. 2002); signal transducer and
23 activator of transcription 5A (STAT5A) is important for its influence on milk production and
24 reproduction activity (Yang et al. 2000; Homer et al. 2013); melatonin receptor 1A (MTNR1A) is
25 important for reproductive seasonality (Chu et al. 2007; Luridiana et al. 2012).

26

1 MATERIALS AND METHODS

2 Peripheral blood samples from cattle (Agerolese breed), sheep (Laticauda breed), goat
3 (Cilentana breed) and riverbuffalo were cultured and treated for late BrdU and Hoechst 33258
4 incorporation according to Iannuzzi and Di Berardino (2008). The bovine BAC clones overlapping
5 studied genes (Table 1) were screened by database searching and ordered from INRA bovine BAC
6 library (CRB- Biological Resources Centre dedicated to livestock genomics –INRA, Jouy-en Josas,
7 France) (<http://locus.jouy.inra.fr/cgibin/bovmap/intro2.pl>). Extraction of DNA was done using
8 CHORI (Children’s Hospital Oakland Research Institute) recommended protocol. DNA was labeled
9 with biotin and digoxigenin using nick-translation kit (Roche applied science Inc.). Slides were then
10 treated for FISH with BAC clones overnight in presence of bovine COT-1 DNA and sonicated
11 salmon sperm allocated in a moist chamber. After detection steps with FITC-avidin and anti-
12 digoxigenin antibodies, Chromosomes were counterstained with Vectashield DAPI H1500 in
13 Vectashield H 1000 (Vector Lab) antifade solution. Both RB-banding (R-banding by late
14 incorporation of BrdU) metaphases and fluorescence FITC and TRIC signals were separately
15 captured by a CCD-camera (Photometrics, cool SNAP, Nikon) and processed by superimposing
16 FITC and TRIC signals on RB-banding preparations. Chromosome identification and banding
17 followed the standard karyotypes for cattle, sheep and goat (ISCNDB2000 2001) and river buffalo
18 (CSKBB 1994).

19

20 RESULTS AND DISCUSSION

21 Three major fecundity genes (TNF, STAT5A and MTNR1A), were comparatively
22 physically FISH-mapped on cattle, sheep goat and river buffalo R-banded metaphase chromosomes
23 (Figure 1). Loci FISH-mapped with locus name, symbol, clone identification and chromosome
24 localization are reported in Table 1. TNF maps on BTA/CHI23q21-22, OAR20q21-22 and BBU
25 2p21-22; STAT5A maps on BTA/CHI19q17-21, OAR11q17-21 and BBU3p15-21; MTNR1A maps
26 to BTA/CHI27q14-15, OAR11q17-21 and BBU1p21-22. The three loci were located in

1 homoeologous chromosomes and chromosome bands of the four species extending the cytogenetic
2 maps in these three species chromosomes. FISH-mapping of STAT5A agrees with previous
3 localizations performed in BTA/CHI19 and OAR11 by sequential GTG-banding and FISH
4 (Goldammer et al. 1997), while MTNR1A, earlier assigned to BBU1 by RH-mapping (Miziara et al.
5 2007) was now assigned to specific chromosome arms and bands (1p21-22).

6 During the last fifteen years, FISH techniques have been used in domestic animals research
7 mainly to identify chromosomal rearrangements, gene mapping, comparative mapping, and
8 evolutionary chromosome studies. The localization of TNF, STAT5A and MTNR1A on
9 homologous chromosomes and chromosome bands in cattle, sheep, goat and river buffalo (Figure 1;
10 Table 1) confirmed the high conservation of autosomal chromosomes among the bovid species and
11 extended the cytogenetic maps of the four economically important domestic species. However,
12 some discrepancies may exist between the localization of loci reported in a reference genome and
13 the localization obtained by FISH physical mapping, as supported by different papers (De Lorenzi
14 et al. 2010; Partipilo et al. 2011; De Lorenzi et al. 2013). These results clearly indicate the idea that
15 physical localization of genomic elements by FISH can further improve the excellent results
16 obtained by genome sequence projects.

17

18 **ACKNOWLEDGEMENTS**

19 The authors are very grateful to Domenico Incarnato, CNR-ISPAAAM, Naples for excellent
20 technical assistance. The study has been supported in part by CISIA-VARIGEAV project and in
21 part by project RARECa, PSR, Measure 214, e2 of Campania Region.

22

23 **REFERENCES**


24 CSKBB 1994. Standard karyotype of the river buffalo (*Bubalus bubalis* L., 2n=50). Report of the
25 committee for the standardization of banded karyotype of the river buffalo (Iannuzzi L.,
26 coordinator). *Cytogenet Cell Genet.* 67:102–113.

- 1
- 2 Chu MX, He YQ, Cheng DX, Ye SC, Fang L, Wang JY. 2007. Association between expression of
3 reproductive seasonality and alleles of melatonin receptor 1A in goats. *Anim Reprod Sci.*
4 101(3–4):276–84.
- 5
- 6 De Lorenzi L, Molteni L, Parma P. 2010. FISH mapping in cattle (*Bos taurus* L.) is not yet out of
7 fashion. *J Appl Genet.* 51(4):497–499.
- 8
- 9 De Lorenzi L, Genuardo V, Perucatti A, Iannuzzi A, Iannuzzi L, Parma P. 2013. Physical mapping
10 of 20 unmapped fragments of Btau_4.0 genome assembly in cattle, sheep and river buffalo.
11 *Cytogenet Genome Res.* 140:29–35.
- 12
- 13 Di Meo GP, Molteni L, Perucatti A, De Giovanni A, Incarnato D, Succi G, Schibler L, Cribiu EP,
14 Iannuzzi L. 2000. Chromosomal characterization of three centric fusion translocations in
15 cattle using G-, R- and C-banding and FISH technique. *Cariologia* 53(3–4):213–218.
- 16
- 17 Di Meo GP, Perucatti A, Floriot S, Hayes H, Schibler L, Rullo R, Incarnato D, Ferretti L, Cockett N,
18 Cribiu E, Williams JL, Eggen A, Iannuzzi L. 2007. An advanced sheep (*Ovis aries*, 2n = 54)
19 cytogenetic map and assignment of 88 new autosomal loci by fluorescence in situ
20 hybridization and R-banding. *Anim Genet.* 38:233–240.
- 21
- 22 Eggert-Kruse W, Kiefer I, Beck C, Demirakca T, Strowitzki T. 2007. Role for tumor necrosis factor
23 alpha (TNF- α) and interleukin 1-beta (IL-1 β) determination in seminal plasma during
24 infertility investigation. *Fertil Steril.* 87(4):810–23.
- 25

- 1 Goldammer T, Di Meo GP, Lühken G, Drögemüller C, Wu CH, Kijas J, Dalrymple BP, Nicholas
2 FW, Maddox JF, Iannuzzi L, Cockett NE. 2009. Molecular Cytogenetics and Gene Mapping
3 in Sheep (*Ovis aries*, $2n = 54$). *Cytogenet Genome Res.* 126:63–76.
4
- 5 Goldammer T, Meyer L, Seyfert H-M, Brunner RM, Schwerin M. 1997. STAT5A encoding gene
6 maps to Chromosome 19 in cattle and goat and to Chromosome 11 in sheep. *Mamm Genome.*
7 8:705–706.
8
- 9 Homer EM, Derecka K, Webb R, Garnsworthy PC. 2013. Mutations in genes involved in oestrous
10 cycle associated expression of oestrus. *Anim Reprod Sci.* 142(3–4):106–12.
11
- 12 Hornak M, Jeseta M, Musilova P, Pavlok A, Kubelka M, Motlik J, Rubes J, Anger M. 2011.
13 Frequency of aneuploidy related to age in porcine oocytes. *PLoSOne.* 6:e18892.
14
- 15 Iannuzzi L, Perucatti A, Di Meo GP, Schibler L, Incarnato D, Crihiu EP. 2003a. Chromosomal
16 localization of sixty autosomal loci in sheep (*Ovis aries*, $2n = 54$) by fluorescence in situ
17 hybridization and R-banding. *Cytogenet Genome Res.* 103:135–138.
18
- 19 Iannuzzi L, Di Meo GP, Perucatti A, Schibler L, Incarnato D, Gallagher D, Eggen A, Ferretti L,
20 Crihiu EP, Womack J. 2003b. The river buffalo (*Bubalus bubalis*, $2n=50$) cytogenetic map:
21 assignment of 64 loci by fluorescence in situ hybridisation and R-banding. *Cytogenet*
22 *Genome Res.* 102:65–75.
23
- 24 Iannuzzi L, Di Berardino D. 2008. Tools of the trade: diagnostics and research in domestic animal
25 cytogenetics. *J Appl Genet.* 49(4):357–366.
26

1 Iannuzzi A, Perucatti A, Genuardo V, De Lorenzi L, Di Berardino D, Parma P, Iannuzzi L. 2013.
2 Cytogenetic Elaboration of a Novel Reciprocal Translocation in Sheep. *Cytogenet Genome*
3 *Res.* 139:97–101.
4
5 ISCNDB2000. 2001. International System for Chromosome Nomenclature of Domestic Bovids. Di
6 Berardino D, Di Meo GP, Gallagher DS, Haves H., Iannuzzi L. (coordinator), eds. *Cytogenet*
7 *Cell Genet.* 92: 283–99.
8
9 Luridiana S, Mura MC, Pazzola M, Paludo M, Cosso G, Dettori ML, Bua S, Vacca GM, Carcangiu
10 V. 2012. Association between melatonin receptor 1A (MTNR1A) gene polymorphism and the
11 reproductive performance of Mediterranean Italian buffaloes. *Reprod Fertil Dev.* 24 (7):983–
12 87.
13
14 Koçak I, Yenisey C, Dündar M, Okyay P, Serter M. 2002. Relationship between seminal plasma
15 interleukin-6 and tumor necrosis factor α levels with semen parameters in fertile and infertile
16 men. *Urol Res.* 30(4):263–67.
17
18 Miziara MN, Goldammer T, Stafuzza NB, Ianella P, Agarwala R, Schäffer AA, Elliott JS, Riggs PK,
19 Womack JE, Amaral MEJ. 2007. A radiation hybrid map of river buffalo (*Bubalus bubalis*)
20 chromosome 1 (BBU1). *Cytogenet Genome Res.* 119:100–104.
21
22 Partipilo G, D’Addabbo P, Lacalandra GM, Liu GE, Rocchi M. 2011. Refinement of *Bos taurus*
23 sequence assembly based on BAC-FISH experiments. *BMC Genomics.* 12:639.
24

- 1 Perucatti A, Genuardo V, Iannuzzi A, De Lorenzi L, Matassino D, Parma P, Di Berardino D,
2 Iannuzzi L, Di Meo GP. 2011. A new case and unusual reciprocal translocation in cattle:
3 rcp(11;25)(q11;q14-21). *Cytogenet. Genome Res.* 134:96–100.
4
- 5 Pauciullo A, Cosenza G, Peretti V, Iannuzzi A, Di Meo GP, Ramunno L, Iannuzzi L, Rubes J, Di
6 Berardino D. 2011. Incidence of X-Y aneuploidy in sperm of two indigenous Cattle breeds by
7 using dual color fluorescent in situ hybridization (FISH). *Theriogenology* 76:328–333.
8
- 9 Pauciullo A, Nicodemo D, Cosenza G, Peretti V, Iannuzzi A, et al. 2012. Similar rate of
10 chromosomal aberrant secondary oocytes in two indigenous cattle (*Bos taurus*) breeds as
11 determined by dual-color FISH. *Theriogenology* 77:675–683.
12
- 13 Schibler L, Di Meo GP, Cribru EP, Iannuzzi L. 2009. Molecular Cytogenetics and Comparative
14 Mapping in Goats (*Capra hircus*, 2n = 60). *Cytogenet Genome Res.* 126:77–85.
15
- 16 Stafuzza NB, Greco AJ, Grant JR, Abbey CA, Gill CA, Raudsepp T, Skow LC, Womack JE, Riggs
17 PK, Amaral MEJ. 2013. A high-resolution radiation hybrid map of the river buffalo major
18 histocompatibility complex and comparison with BoLA. *Anim Genet.* 44:369–376.
19
- 20 Yang J, Kennelly J, Baracos VE. 2000. Physiological levels of Stat5 DNA binding activity and
21 protein in bovine mammary. *J Anim Sci.* 78:3126–3134.
22



1

2

3 Figure 1. (Color online) Representative FISH results on cattle (BTA), river buffalo (BBU), sheep
 4 (OAR) and goat (CHI) chromosomes, using bovine BAC clones containing genes related to
 5 fecundity (TNF, STAT5A, MTNR1A). FITC and TRIC signals were superimposed on R-banding
 6 chromosomes counterstained with DAPI. For each chromosome, the corresponding standard
 7 ideogram (ISCNDB2000 2001; CSKBB 1994) it is also reported.

8

1 Table 1. BAC-probes, identified DNA sequences of FISH-mapped genes in cattle (BTA), river buffalo (BBU), sheep (OAR), goat (CHI)
 2 chromosomes (ISCNDB2000 2001 and CSKBB 1994), comparison with human (HSA) chromosomes (HGNC).

3

BAC FISH Probe	Identified DNA sequence within BAC and locus symbol (HGNC)	Gene name	Cytogenetic localization on RBPI-bands				
			BTA	BBU	OAR	CHI	HSA
BtINRA-81C03	<i>TNF</i>	<i>tumor necrosis factor-α</i>	23q21-22	2p21-22	20q21-22	23q21-22	6p21.3
BtINRA-243A01	<i>STAT5A</i>	<i>signal transducer and activator of transcription 5A</i>	19q17-21	3p15-21	11q17-21	19q17-21	17q11.2
BtINRA-448A07	<i>MTNRIA</i>	<i>melatonin receptor 1 A</i>	27q14-15	1p21-22	26q14-15	27q14-15	4q35

4

5

6