

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Mediterranean River Buffalo CSN1S1 gene: search for polymorphisms and association studies.**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1507139> since 2015-11-26T10:29:28Z

*Published version:*

DOI:10.1071/AN13438

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

*This is an author version of the contribution published on:*

*Animal Production Science*

*Questa è la versione dell'autore dell'opera:*

*[Animal Production Science, Vol. 55 No. 5 Pages 654 – 660, 2015, DOI:*

*http://dx.doi.org/10.1071/AN13438]*

*ovvero [G. Cosenza, A. Pauciullo, N. P. P. Macciotta, E. Apicella, R. Steri, A. La Battaglia, L. Jemma, A. Coletta, D. Di Bernardino, L. Ramunno, Animal Production Science, 2015, Vol. 55 No. 5 Pages 654 – 660, CSIRO publishing]*

*The definitive version is available at:*

*La versione definitiva è disponibile alla URL:*

*[<http://www.publish.csiro.au/paper/AN13438.htm>]*

22 **Mediterranean River Buffalo *CSN1S1* gene: search for polymorphisms**  
23 **and association studies**

24  
25  
26 **G. Cosenza<sup>A,F</sup>, A. Pauciullo<sup>B,F</sup>, N. P. P. Macciotta<sup>C</sup>, E. Apicella<sup>A</sup>, R. Steri<sup>C</sup>, A. La Battaglia<sup>D</sup>,**  
27 **L. Jemma<sup>A</sup>, A. Coletta<sup>E</sup>, D. Di Bernardino<sup>A</sup>, L. Ramunno<sup>A</sup>**

28  
29 <sup>A</sup>Dipartimento di Agraria, Università degli Studi di Napoli “Federico II”, via Università 100, Portici,  
30 NA 80055 Italy

31 <sup>B</sup>Istituto per il Sistema Produzione Animale in Ambiente Mediterraneo, Consiglio Nazionale delle  
32 Ricerche, via Argine 1085, Napoli, 80147 Italy

33 <sup>C</sup>Dipartimento di Agraria, Università degli Studi di Sassari, via De Nicola 9, Sassari, 07100 Italy

34 <sup>D</sup>Comunità Montana Alto Agri, Villa D’Agri di Marsicovetere (PZ), 85050 Italy

35 <sup>E</sup>Associazione Nazionale Allevatori Specie Bufalina, Via F. Petrarca 42-44 Località Centurano, Caserta,  
36 81100 Italy

37  
38 <sup>F</sup>Corresponding author: Email: giacosen@unina.it; alfredo.pauciullo@cnr.it

39

40 **Abstract.** The aim of the present work was to study the variability at *CSN1S1 locus* of the Italian  
41 Mediterranean river buffalo and to investigate possible allele effects on milk yield and its composition.  
42 Effects of parity, calving season and month of production were also evaluated. Three SNPs were detected.  
43 The first mutation, located at position 89 of 17<sup>th</sup> exon (c.628C>T), is responsible for the amino acid  
44 change p.Ser178 (B allele)/Leu178 (A allele). The other two polymorphisms, detected at the positions  
45 144 (c.882G>A) and 239 (c.977A>G) of 19<sup>th</sup> exon respectively, are silent (3’ UTR).

46 Associations between the *CSN1S1* genotypes and milk production traits were investigated using  
47 4,122 test day records of 503 lactations from 175 buffalo cows. Milk yield, fat and protein percentages

48 were analyzed using a mixed linear model. A significant association between the c.628C>T SNP and the  
49 protein percentage was found. In particular, the CC genotype showed an average value of about 0.04%  
50 higher than the CT and TT genotypes. The allele substitution effect of the cytosine into the thymine was  
51 -0.014, with a quite low (0.3%) protein percentage (PP) contribution on total phenotypic variance. A  
52 large dominance effect was detected.

53 Furthermore, a characterization of the *CSN1S1* transcripts and a method based on *MboI*-ACRS-  
54 PCR for a rapid genotyping of c.628C>T were provided.

55

56 **Additional keywords:** Mediterranean river buffalo, *Bubalus bubalis*, *CSN1S1*, milk protein percentage,  
57 marker assisted selection

58

## 59 **Introduction**

60 Historically domestic water buffalo were divided into swamp and river subspecies that differ in  
61 morphology, behaviour, and chromosome number (2n=48 and 2n=50, respectively) (Ajmone-Masan P.  
62 *et al*, 2013). In particular, the Mediterranean river buffalo, is the only indigenous Italian breed of water  
63 buffalo. It is of the River subtype of water buffalo, and is genetically and ecological similar to the buffalo  
64 breeds of Hungary, Romania and the Balkan countries. Previously considered to belong to the  
65 Mediterranean buffalo group, it was officially recognised as a breed in 2000, based on its long isolation  
66 period and lack of interbreeding with other buffalo breeds.

67 The improvement of animal performances represents a priority for the Italian dairy Buffalo  
68 industry in order to fulfill the increasing market demand for mozzarella cheese. Italian buffalo stock  
69 consists of approximately 344,000 Mediterranean river buffaloes (<http://faostat.fao.org/>). In 2011, the  
70 average milk yield per lactation per buffalo cow (35,963 registered cows in the national herd book) was  
71 kg 2,223 with 8.49 and 4.65 % of fat and protein content, respectively (<http://www.aia.it/aia->

72 website/it/home). The milk is almost completely processed into cheese. A breeding program aimed at  
73 improving buffalo milk yield and composition is currently operating in Italy, however the low efficiency  
74 of the artificial insemination, the difficulties to detect the oestrus and the variability of its length are  
75 among the main causes of a very limited impact on the population (Barile, 2005).

76         Recent advances of molecular genetics offer the possibility to investigate genomic regions that  
77 affect traits of economic importance and to identify genetic polymorphisms useful for marker-assisted  
78 selection (MAS) programs. In the last decades, several association studies between milk production traits  
79 and markers located in milk protein genes have been carried out in cattle, sheep and goat (Ibeagha-  
80 Awemu *et al*, 2008; Martin *et al*, 2002; Mroczkowski *et al*, 2004). In particular, the goat  $\alpha$ 1 encoding  
81 gene (*CSN1S1*) was found to be highly polymorphic with at least 17 alleles associated with qualitative  
82 and quantitative differences for the content of  $\alpha$ 1casein (Ramunno *et al*, 2004; Ramunno *et al*, 2005),  
83 fat (Grosclaude *et al*, 1994; Chilliard *et al*, 2006), urea level (Schmidely *et al*, 2002; Bonanno *et al*, 2007;  
84 Avondo *et al*, 2009), fatty acid profile Chilliard *et al*, 2006) and milk yield (Yue *et al*, 2011). Associations  
85 between alleles at *CSN1S1 locus* and protein content were also observed for other species as for instance  
86 in bovine (Rando *et al*, 1998; Prinzenberg *et al*, 2003; Çardak, 2005) and ovine (Pirisi *et al*, 1999;  
87 Wessels *et al*, 2004).

88         In ruminant species, the *CSN1S1* gene is characterized by an extremely split architecture with 19  
89 exons, many of which (exons 5, 6, 7, 8, 10, 13 and 16) of small size (24 bp) (Ramunno *et al*, 2004;  
90 Koczan *et al*, 1991; Calvo *et al*, 2011). In buffalo, the  $\alpha$ 1-casein gene codes for a precursor of 214 amino  
91 acids with a signal peptide of 15 amino acid residues (Ferranti *et al*, 1998; Sukla *et al*, 2007). Currently,  
92 several partial or complete bubaline *CSN1S1* cDNA sequences are available in EMBL (FJ392261;  
93 AJ005430; AY948385; EF025981; EF025982; EF025983; DQ111783). The similarity between buffalo  
94 and cattle, goat and sheep  $\alpha$ 1-casein mRNA sequence is 97.2, 93 and 92.3%, respectively. A similar  
95 trend was observed comparing amino acid sequences of these species (Sukla *et al*, 2007).

96 Few polymorphisms have been reported for the *CSN1S1* locus in buffalo. The occurrence of a  
97  $\alpha$ s1-casein B genetic variant characterized by a single amino acid substitution (p.Leu178Ser) as  
98 consequence of single nucleotide substitutions was found for the first time in Romanian Buffalo breed  
99 (Balteanu *et al*, 2008) and confirmed at amino acid level in Mediterranean water buffalo (Chianese *et al*,  
100 2009). Furthermore in Indian water buffalo a novel *CNS1S1* allele has been characterized by c.620G>A  
101 substitution. It led to a p.Gly192Glu replacement in the peptide chain (Sukla *et al*, 2007).

102 Recently, additional SNPs were detected: a transition c.136G>A at exon 5, leading to a  
103 p.Val31Met substitution, a transition c.175A>G at exon 7, leading to a p.Ile44Val substitution in the  
104 peptide chain, one SNP (g.218T>C) in intron 5 and three SNPs (g.472G>C; g.547C>T; g.856T>C) in  
105 intron 6. In particular, it was reported that the g.472G>C substitution inactivates the intron 6 splice donor  
106 site promoting the skipping of exon 6 of the buffalo *CSN1S1* mRNA and, as consequence, triggering the  
107 synthesis of a defective protein lacking eight amino acids (Balteanu *et al*, 2013).

108 In recent years, many studies have been carried out for the identification of the genetic  
109 polymorphisms at the *loci* coding for the buffalo milk proteins (Cosenza *et al*, 2009a, b; Masina *et al*,  
110 2007), and candidate genes responsible for the variation of the quali-quantitative characteristics of the  
111 Mediterranean water buffalo milk have been found (Cosenza *et al*, 2007; Pauciullo *et al*, 2010). Efforts  
112 in this direction also allowed to find significant associations with traits of economic interest, as milk  
113 yield (Pauciullo *et al*, 2012a, b) and milk coagulation properties (Bonfatti *et al*, 2012b).

114 The aim of our work was to study the variability at the Italian Mediterranean river buffalo *CSN1S1*  
115 cDNA and to investigate possible associations with milk yield and composition.

116  
117

118 **Materials and methods**

119 *Sampling*

120 Individual blood samples were collected from 175 Italian river buffaloes randomly chosen and belonging  
121 to an experimental herd, located in Salerno province (Southern Italy).

122 In order to characterize the *CSN1S1* transcripts and to detected polymorphisms at this *locus*,  
123 individual milk samples were collected from 10 animals at comparable age, in third calving, at 120 days  
124 in milking and free of clinical mastitis and randomly chosen in different farms located in the province of  
125 Salerno and Caserta (Southern Italy). After collection, milk samples were immediately frozen and kept  
126 at -80 °C until analysis. Sampling was carried out in collaboration with the Italian National Association  
127 of Buffalo Breeds (ANASB).

128

129 *RNA extraction*

130 Total RNA was isolated from somatic cells (SCC range from 10,000 to 12,000/mL) present in individual  
131 milk samples by using NucleoSpin® Extract Kits (Macherey-Nagel). A digestion with 2U of DNase I  
132 (Ambion) in 1X DNase buffer was carried out according to the manufacture guidelines at 37°C for 30  
133 min followed by the enzyme inactivation at 75°C for 5 min. The quantity, quality, purity, and integrity  
134 of RNA after DNase treatment were estimated by means of Nanodrop 2000c spectrophotometer (Thermo  
135 Scientific, Barrington, IL) and by electrophoresis on a denaturing agarose gel.

136

137 *Reverse transcription, PCR and cloning*

138 Total RNA was converted into cDNA by reverse transcription using Improm- II Reverse Transcriptase  
139 (Promega) with a final volume of 20µl. The reaction was performed using cDNA19R (5'-  
140 CAAAATCTGTTACTGCACA - 3'), a reverse primer complementary to nt 327–345 of 19<sup>th</sup> exon of  
141 buffalo *CSN1S1* cDNA sequence (accession number AY948385).

142 The PCR was performed using: cDNA19R and cDNA1F (5'- AACCCAGCTTGCTGCTT - 3'),  
143 a forward primer corresponding to nt 1–17 of partial 1<sup>st</sup> exon of buffalo *CSN1S1* cDNA sequence  
144 (accession number AY948385). The PCR reaction mix comprised 20 µl of RT reaction product, 50 mM  
145 KCl, 10 mM Tris–HCl, 0.1% Triton X-100, 2 mM MgCl<sub>2</sub>, 10 pmol of each primer, dNTPs each at 0.2  
146 mM, 5 U of Taq DNA Polymerase (Promega, Madison, WI), with a final volume of 100 µl.

147 The amplification protocol consisted of 39 cycles: the first cycle involved a denaturation step at  
148 97 °C for 2 min, an annealing step at 57 °C for 30 s and an extension step at 72 °C for 1 min and 30 s.  
149 The next 37 cycles were performed under the following conditions: 94 °C for 30 s, 57 °C for 30 s. and  
150 72 °C for 1 min and 30 s. In the 39<sup>th</sup> cycle, the final extension step was carried out at 72 °C for 10 min.

151 The amplified products were first analyzed by electrophoresis on 3% agarose gel in TBE 1X  
152 buffer (Bio-Rad) and then cloned in pCR2.1-TOPO plasmid by using the TOPO TA cloning kit  
153 (Invitrogen, Pro, Milan, Italy). White recombinant clones were randomly chosen and screened by PCR  
154 using standard vector primers M13. Recombinant clones underwent plasmid purification by PureYield™  
155 Plasmid Midiprep System (Promega, USA) and then sequenced on both strands at CEINGE -  
156 Biotecnologie Avanzate (Naples, Italy).

157

#### 158 *DNA extraction*

159 DNA was extracted from leukocyte, using the procedure described by Gossens and Kan (1981). Briefly,  
160 fresh buffy coat samples were washed twice with distilled water and NaCl 1.8% to remove the excess of  
161 red cells, protein digestion was carried out with 500 µl of proteinase K solution (2 mg/ml of proteinase  
162 K, 1% w/v SDS and 0.02 M EDTA). Proteins were extracted using phenol-chloroform method followed  
163 by DNA precipitation with cold isopropanol. The isolated DNA was then resuspended in 100 µl TE buffer  
164 pH 7.6 (10 mM Tris, 1mM EDTA). DNA concentration and OD<sub>260/280</sub> ratio of the samples were then  
165 measured by the Nanodrop ND-2000C Spectrophotometer (Thermo Scientific).



166

167 *CSN1S1 locus genotyping*

168 In order to genotype 175 individual samples of water buffalo for the c628C>T mutation, a method based  
169 on ACRS-PCR (Amplification Created Restriction Site PCR) was developed according to Lien *et al*  
170 (1992). The ACRS-PCR was performed using: ACRS17F (5'- CAATACCCTGATGCCCCGAT - 3') as  
171 forward and ACRS17R (5'- CACCACAGTGGCATAGTAG - 3') as reverse, corresponding to nt 70–88  
172 and complementary to nt 137–155 of the 17<sup>th</sup> exon of buffalo *CSN1S1* cDNA sequence (EMBL  
173 HE573919), respectively. According to the method, the forward primer was modified by changing C→G  
174 in position 17 in order to provide a restriction site for the *Mbo*I (!GATC) endonuclease.

175 The amplification protocol consisted of 39 cycles: the first cycle involved a denaturation step at  
176 95 °C for 5 min. The next 38 cycles were performed under the following conditions: 95 °C for 45 s, 63.2  
177 °C for 45 s. and 72 °C for 20 s. In the 39<sup>th</sup> cycle, the extension step was carried out at 72 °C for 10 min.  
178 PCR reaction mix comprised: 100 ng of genomic DNA, 50mM KCl, 10mM Tris–HCl, 0.1% Triton X-  
179 100, 3mM MgCl<sub>2</sub>, 5 pmol of each primer, dNTPs each at 400 μM, 2.5U of *Taq* DNA Polymerase  
180 (Promega, Madison, WI), and 0.04% BSA, with a final volume of 25 μl.

181

182 *Digestion and electrophoresis condition*

183 Each PCR amplification product was digested with 10U of *Mbo*I after incubation for 5 h at 37 °C;  
184 following the supplier's guidelines. The restricted fragments were analyzed directly by electrophoresis  
185 on 3% TBE agarose gel in 1 X TBE buffer and stained with ethidium bromide.

186

187 *Association study*

188 Associations between *CSN1S1* genotypes and milk production traits were carried out on 4,122 test day  
189 records of 503 lactations from 175 buffalo cows, supplied by the Italian Association of Buffalo Breeders

190 (ANASB). Milk yield (MY), fat (FP) and protein percentages (PP) were tested with the following mixed  
191 linear model:

192

$$193 \quad y_{ijklmno} = \text{Month}_i + \text{Par}_j + \text{Sea}_K + \text{DIM}_l + \alpha_{s1m} + c_n(\alpha_{s1m}) + e_{ijklmno} \quad [*]$$

194

195 where:  $y_{ijklmno}$  is the test-day record of MY, FP or PP;  $\text{Month}_i$  is the fixed effect of the  $i$ -<sup>th</sup> month of  
196 production (12 levels);  $\text{Par}_j$  is the fixed effect of the  $j$ -th parity (6 levels: 1 to 5, >5);  $\text{Sea}_K$  is the fixed  
197 effect of the  $k$ -<sup>th</sup> calving season (4 levels: autumn, winter, spring, summer);  $\text{DIM}_l$  (Days In Milk) is the  
198 fixed effect of the  $l$ -<sup>th</sup> stage of lactation (30 levels of 10 days each);  $\alpha_{s1m}$  is the fixed effect of the  $m$ -<sup>th</sup>  
199 genotype at the c.628C>T SNP of *CSN1S1* gene (3 levels: CC, CT, TT);  $c_n$  is the random effect of  
200 individual cow (175 levels), nested within  $\alpha_{s1}$  genotype; and  $e_{ijklmno}$  is the random residual. Pairwise  
201 comparisons among different levels of fixed effects included in model were performed using a  
202 Bonferroni adjusted test. (Co)variance matrices of random effects of cow and residual were assumed to  
203 be diagonal,  $\mathbf{I}\sigma^2_c$  and  $\mathbf{I}\sigma^2_e$ , respectively. They allow for the REML estimation of variance components  
204 associated to individual cow ( $\sigma^2_c$ ) and residual ( $\sigma^2_e$ ). Variance component associated to the  $\alpha_{s1}$  locus  
205 ( $\sigma^2_{\alpha_{s1}}$ ) was estimated running a mixed model having the same structure of [\*] but with the  $\alpha_{s1}$  genotype  
206 treated as random. Contributions of  $\alpha_{s1}$  locus ( $r^2_{\alpha_{s1}}$ ) and cow ( $r^2_c$ ) to the total phenotypic variance of the  
207 trait was calculated as the ratio between  $\sigma^2_{\alpha_{s1}}$  and  $\sigma^2_c$ , respectively and the sum of all variance  
208 components (i.e.  $\sigma^2_{\alpha_{s1}} + \sigma^2_c + \sigma^2_e$ ).

209 In order to estimate the average of gene substitution effect ( $\alpha$ ) and a possible dominance effects  
210 (d), gene effect was treated as a covariable, represented by the number of T alleles at the  $\alpha_{s1}$  locus (0, 1,  
211 2). Finally, an interaction between alleles at the SNP locus was considered (Banos *et al*, 2008; Barendse  
212 *et al*, 2008).

213

## 214 **Results and discussion**

215

### 216 *a) Characterization of CSN1S1 transcripts*

#### 217 *Analysis of the cloned RT-PCR fragments*

218 The mRNAs extracted from individual milk samples obtained from 10 Mediterranean river buffaloes,  
219 randomly chosen in the province of Salerno and Caserta, was investigated through the clone analysis. 10  
220 positive clones for each individual were screened. The electrophoretic analysis of the PCR products and  
221 the subsequent sequencing of clones, showed at least two populations of transcripts for each examined  
222 individual. The most represented population (about 90 %) was correctly assembled, followed by that one  
223 deleted of the first triplet of the 11<sup>th</sup> exon. The last event is a constitutive allele independent event which  
224 takes place during the maturation of the pre-mRNA. The first codon of exon 11 (CAG), coding for  
225 glutamine in position 78, is in fact competitively eliminated from the mRNA because it is recognized as  
226 cryptic site of splice instead of the canonic site (AG) located at the end of the 10<sup>th</sup> intron. This feature  
227 was already observed in buffalo (Ferranti *et al*, 1999) and it is common event for the *CSN1S1* gene of  
228 other ruminants, like sheep (Ferranti *et al*, 1998), goat (Ramunno *et al*, 2005) and cattle (Ferranti *et al*,  
229 1999). Furthermore, as already observed in the afore mentioned species, also for the river buffalo it is  
230 reasonable to hypothesize the existence of other transcripts different from those reported in this study.  
231 Their undetectable amounts opens a new opportunity of investigation in the field of the transcript analysis  
232 for buffalo milk protein.

233

#### 234 *Polymorphism detection*

235 In order to detect polymorphisms at the Mediterranean river buffalo *CSN1S1 locus*, the correctly  
236 assembled transcripts were sequenced. The analysis of the sequences showed a transcript of 1083 bp,

237 spanning from the 16<sup>th</sup> nt of the 1<sup>st</sup> exon to the 345<sup>th</sup> nt of the 19<sup>th</sup> exon. The comparison of the obtained  
238 sequences (EMBL HE573919 and HE573920) showed 3 transitions. The first was located at the position  
239 89 of the 17<sup>th</sup> exon (c.628C>T) and the other two at the position 144 and 239 of the 19<sup>th</sup> exon, respectively  
240 (c.882G>A and c.977A>G). The last two mutations are silent because they are located in the 3' UTR,  
241 whereas the first is a miss-sense SNP (p.Ser178Leu). This amino acid change at the buffalo *CSN1S1*  
242 locus was already observed and it characterizes the genetic variants of the  $\alpha$ s1-casein, named B and A,  
243 respectively (Balteanu *et al*, 2008; Chianese *et al*, 2009).

244 Since the presence of the cytosine in position 628 characterizes even other *CSN1S1* sequences of  
245 buffalo (FJ392261, AY948385, AJ005430, DQ111783) and of other ruminants, such as goat  
246 (AJ504710.2), sheep (NM\_001009795), bison (EU862388) and cattle (X59856), its presence might be  
247 indicative of an ancestral condition. According to the sequence analysis of the cDNA samples belonging  
248 to individuals with informative genotypes, the G in position 882 and the A in position 977 are in *cis* with  
249 the T at 89<sup>th</sup> nucleotide of exon 17. Although the complete genomic sequence of the river buffalo *CSN1S1*  
250 is not available, a close distance exists between the exons 17 and 19 in the homologous sequences of the  
251 other ruminants. For instance, this DNA region is only 1962 bp long in the bovine *CSN1S1* gene (EMBL  
252 acc. No. X59856), therefore, a condition of linkage disequilibrium can be assumed for these SNP in river  
253 buffalo.

254  
255 *b) Association of CSN1S1 polymorphism on milk yield and composition*

256 *Genotyping of Mediterranean river buffalo CSN1S1 alleles*

257 The c.628C>T does not alter or create any restriction site, therefore, we established a screening method  
258 based on the *Mbo*I-ACRS-PCR to identify the carriers of this mutation in an easy and rapid way.

259 The amplified fragment includes the last 86 bp of 17<sup>th</sup> exon. Therefore, the digestion with such  
260 endonuclease produces an undigested fragment of 86 bp in individuals homozygous for the thymine and

261 two fragments of 70 and 16 bp (not visible electrophoretically) in buffaloes homozygous for the cytosine.  
262 The heterozygous individuals produce a pattern characterized by all 3 restriction fragments: 86, 70 and  
263 16 bp (Figure 1). The investigated population was in Hardy–Weinberg equilibrium. Genotype  
264 distribution and allele frequencies are reported in Table 1. The frequency of the thymine was 0.33. This  
265 result is in agreement with data reported by Chianese *et al* (2009) and Bonfatti *et al* (2012a) and it  
266 confirms that the most common genetic variant in Mediterranean breed reared in Campania (Italy) is the  
267  $\alpha s1$ -CN B variant.

268

#### 269 *Association study*

270 All environmental factors included in model [\*] affected significantly all the traits considered (Table 2)  
271 except the calving season for FP. The genotype at the *CSN1S1* locus was significantly associated with  
272 protein percentage ( $p < 0.04$ ). In particular (Table 3), the CC genotype showed an average value of about  
273 0.04% higher than the CT and TT genotypes. The allele substitution effect of a cytosine in a thymine was  
274  $-0.014 \pm 0.014$ , with a quite low contribution of the *CSN1S1* locus to the total phenotypic variance of PP  
275 ( $r^2_{\alpha s1} = 0.003$ ) PP. A large effect of dominance ( $-0.028 \pm 0.019$ ) was also observed. Often such effect is not  
276 detected or considered non relevant because numerically much lower than the additive effect. Although  
277 dominance effects are not important in the estimation of breeding values, being not transmitted in the  
278 offspring, they might have an impact on allele substitution effect in the population as recently reported  
279 (Pauciullo *et al*, 2012a, b).

280 Recently, Bonfanti *et al* (2012a) estimated the effects of the *CSN1S1* (B and A alleles, c.628C>T  
281 transition) -CSN3 (k-casein, X1 and X2 alleles) genotypes on milk production traits and milk coagulation  
282 properties in Mediterranean water buffalo. In particular, these authors report that genotypes did not affect  
283 milk protein content, but the composite genotype AB-X1X1, compared to genotype AA-X1X1, was  
284 associated with an increased fat content and they indicate a role for the casein genes in the variation of

285 the coagulation properties of the buffalo milk. However, the same authors (Bonfatti *et al*, 2012b) reported  
286 that the increased proportion of the  $\alpha$ s1-CN on the total casein (TCN) content is associated with  
287 genotypes carrying the allele *CSN1S1* A. On the contrary, genotypes associated with a marked decrease  
288 of the  $\alpha$ s1-CN on the TCN (composite genotypes AB-X1X1 and BB-X1X2) are associated with marked  
289 increases in the proportion of  $\alpha$ s2-CN.

290 Although results of the present work need to be confirmed with large-scale studies, they might be  
291 of great economic interest for the buffalo dairy industry. In fact, increases in average protein content  
292 would lead consequently to an expected increase of mozzarella production.

293

## 294 **Conclusions**

295 The present study reports a characterization of the Mediterranean water buffalo *CSN1S1* transcripts. At  
296 least two populations of transcripts were detected. The most represented population (about 90 %) was  
297 correctly assembled, followed by that one deleted of the first triplet of the 11<sup>th</sup> exon.

298 Also the study confirms the existence of genetic polymorphisms at this *locus* and it offers a  
299 method based on the *Mbo*I-ACRS-PCR for a rapid genotyping at DNA level for the *CSN1S1* A and B  
300 alleles. Furthermore, a significant association between the c.628C>T SNP and the protein percentage  
301 was found. In particular, the CC genotype showed an average value of about 0.04% higher than the CT  
302 and TT genotypes.

303 Therefore, further studies are necessary to better determine the real effects of the transition  
304 c.628C>T on milk composition. Besides, an investigation on larger population are needed in order to  
305 validate its application.

306 In addition, it is necessary to investigate the remaining polymorphisms detected in the 3' un-  
307 translated regions (UTR). In fact, it is well known that the sequences in the 3' UTR can affect the  
308 mechanism of mRNA regulation, such as de-adenylation and degradation. Therefore, it is reasonable to

309 hypothesize that these observed mutations might influence, directly or indirectly, the gene expression  
310 and, consequently, the milk protein composition.

311

## 312 **Acknowledgements**

313 This research was financially supported by the Italian Ministry for Agriculture and Forestry Policy-  
314 MiPAF (INNOVAGEN project).

315

## 316 **References**

- 317 Ajmone-Masan P, Marco Milanesi M, Bombal, Colli L, Bacciu N, Ramunno L, Cosenza G, Coletta A,  
318 Zicarelli L, Fritz E, Koltjes J, Reecy JM, Williams J, Iamartino D, Sonstegard TS, Van Tassell C P,  
319 The International Buffalo Consortium (2013) Biodiversity among Buffalo genomes. *International*  
320 *Plant and Animal Genome Conference Future Dates, XXI Conference*, January 12-16, - San Diego
- 321 Avondo M, Pagano RI, Guastella AM, Criscione A, Di Gloria M, Valenti B, Piccione G, Pennisi P (2009)  
322 Diet selection and milk production and composition in Girgentana goats with different  $\alpha$ s1-casein  
323 genotype. *Journal of Dairy Research* **76**, 202–209. doi: 10.1017/S0022029909003914
- 324 Barile VL (2005) Improving reproductive efficiency in female buffaloes. *Livestock Production Science*  
325 **92**,183-194. doi:10.1016/j.livprodsci.2004.06.014
- 326 Balteanu VA, Carsai TC, Vlaic A (2013) Identification of an intronic regulatory mutation at the buffalo  
327  $\alpha$ s1-casein gene that triggers the skipping of exon 6. *Molecular Biology Reports* **40**, 4311–4316.  
328 doi: 10.1007/s11033-013-2518-2
- 329 Balteanu VA, Pop FD, Vlaic A, Rusu AR (2008) Multiple mutations events are characterizing alphas1  
330 casein Brv and beta casein Crv alleles discovered in Romanian buffalo breed. *Bulletin of University*  
331 *Of Agricultural Sciences And Veterinary Medicine Cluj-Napoca. Animal Science and*  
332 *Biotechnologies* **65**, 478.

- 333 Banos G, Woolliams JA, Woodward BW, Forbes AB, Coffey MP (2008) Impact of single nucleotide  
334 polymorphisms in leptin, leptin receptor, growth hormone receptor, and diacylglycerol  
335 acyltransferase (DGAT1) gene loci on milk production, feed, and body energy traits of UK dairy  
336 cows. *Journal of Dairy Science* **91**, 3190-3200. doi: 10.3168/jds.2007-0930
- 337 Barendse W, Harrison BE, Bunch RJ, Thomas MB (2008) Variation at the Calpain 3 gene is associated  
338 with meat tenderness in zebu and composite breeds of cattle. *BMC Genetics* **9**, 41. doi:  
339 10.1186/1471-2156-9-41
- 340 Bonanno A, Finocchiaro R, Di Grigoli A, Sardina MT, Tornambè D, Gigli I (2007) Energy intake effects  
341 at pasture on milk production and coagulation properties in Girgentana goats with different  $\alpha$ s1-  
342 casein genotypes. *Proceedings Symposium of the International Goat Association (IGA) The quality  
343 of goats products*, CRA-Uze, Bella (PZ), Italy, p. 70.
- 344 Bonfatti V, Giantin M, Gervaso M, Coletta A, Dacasto M, Carnier P (2012a) Effect of CSN1S1-CSN3  
345 ( $\alpha$ (S1)- $\kappa$ -casein) composite genotype on milk production traits and milk coagulation properties in  
346 Mediterranean water buffalo. *Journal of Dairy Science* **95**(6), 3435-3443. doi: 10.3168/jds.2011-  
347 4901
- 348 Bonfatti V, Giantin M, Gervaso M, Rostellato R, Coletta A, Dacasto M, Carnier P (2012b) CSN1S1-  
349 CSN3 ( $\alpha$ s1-k-casein) composite genotypes affect detailed milk protein composition of  
350 Mediterranean water buffalo. *Journal of Dairy Science* **95**, 6801–6805. doi: 10.3168/jds.2012-5601
- 351 Calvo JH, Dervishi E, Sarto P, Berzal B, Serrano M, Joy M (2011) Direct Submission EMBL JN560175.
- 352 Çardak AD (2005) Effects of genetic variants in milk protein on yield and composition from Holstein-  
353 Friesian and Simmentaler cows. *The South African Journal of Animal Science* **35** (1), 41-47.
- 354 Chianese L, Quarto M, Pizzolongo F, Calabrese MG, Caira S, Mauriello R, De Pascale S, Addeo F (2009)  
355 Occurrence of genetic polymorphism at the alpha(s1)-casein locus in Mediterranean water buffalo  
356 milk. *International Dairy Journal* **19** (4), 181–189. doi:10.1016/j.idairyj.2008.09.008



357 Chilliard Y, Rouel J, Leroux C (2006) Goats alpha-s1-casein genotype influences its milk fatty acid  
358 composition and delta-9 desaturation ratios. *Animal Feed Science and Technology* **131**, 474–487.  
359 doi:10.1016/j.anifeedsci.2006.05.025

360 Cosenza G, Pauciullo A, Mancusi A, Nicodemo D, Di Palo R, Zicarelli L, Di Berardino D, Ramunno L  
361 (2007) Mediterranean river buffalo oxytocin-neurophysin I (OXT) gene: structure, promoter  
362 analysis and allele detection. *Italian Journal of Animal Science* **6 (Suppl.2)**, 303-306. doi:  
363 10.4081/ijas.2007.s2.303

364 Cosenza G, Feligini M, Mancusi A, D’Avino A, Coletta A, Di Berardino D, Ramunno L (2009a) Italian  
365 Mediterranean river buffalo CSN2 gene structure and promoter analysis. *Italian Journal of Animal*  
366 *Science* **8 (Suppl.2)**, 57-59. doi: 10.4081/ijas.2009.s2.57

367 Cosenza G, Pauciullo A, Feligini M, Coletta A, Colimoro L, Di Berardino D, Ramunno L (2009b) A  
368 point mutation in the splice donor site of intron 7 in the  $\alpha$ 2-casein encoding gene of the  
369 Mediterranean River buffalo results in an allele-specific exon skipping. *Animal Genetics* **40 (5)**,  
370 791. doi: 10.1111/j.1365-2052.2009.01897.x.

371 Ferranti P, Chianese L, Malorni A, Migliaccio F, Stingo V, Addeo F (1998) Copresence of deleted protein  
372 species generates structural heterogeneity of ovine alpha(s1)-casein. *Journal of Agricultural and*  
373 *Food Chemistry* **46 (2)**, 411-416. doi: 10.1021/jf970609s

374 Ferranti P, Scaloni A, Caira S, Chianese L, Malorni A, Addeo F (1998) The primary structure of water  
375 buffalo alpha(s1)- and beta-casein identification of phosphorylation sites and characterization of a  
376 novel beta-casein variant. *Journal of Protein Chemistry* **17 (8)**, 835-844.

377 Ferranti P, Lilla S, Chianese L, Addeo F (1999). Alternative nonallelic deletion is constitutive of  
378 ruminant alpha(s1)-casein. *Journal of Protein Chemistry* **18 (5)**, 595-602.

379 Gossens M, Kan YW (1981). DNA analysis in the diagnosis of hemoglobin disorders. *Methods in*  
380 *Enzymology* **76**, 805-817.

381 Grosclaude F, Ricordeau G, Martin P, Remeuf F, Vassal L, Bouillon J (1994) From gene to cheese:  
382 caprine alpha s1 casein polymorphism, its effects and its evolution. *INRA Productions Animales* **7**  
383 **(1)**, 3–19.

384 Ibeagha-Awemu EM, Kgwatalala P, Zhao X (2008) A critical analysis of production-associated DNA  
385 polymorphisms in the genes of cattle, goat, sheep and pig. *Mammalian Genome* **19** **(9)**, 591-617.  
386 doi: 10.1007/s00335-008-9141-x

387 Koczan D, Hobom G, Seyfert HM (1991) Genomic organization of the bovine alpha-S1 casein gene.  
388 *Nucleic Acids Research* **19** **(20)**, 5591-5596.

389 Lien S, Alestrom, P., Klungland H., Rogne S. (1992) Detection of multiple beta-casein (CASB) alleles  
390 by amplification created restriction sites (ACRS). *Animal Genetics* **23** **(4)**, 333-338.  
391 doi:10.1111/j.1365-2052.1992.tb00155.x

392 Martin P, Szymanowska M, Zwierzchowski L, Leroux C (2002). The impact of genetic polymorphisms  
393 on the protein composition of ruminant milks. *Reproduction Nutrition Development* **42**, 433-459.  
394 doi: 10.1051/rnd:2002036

395 Masina P, Rando A, Di Gregorio P, Cosenza G, Mancusi A (2007). Water buffalo kappa-casein gene  
396 sequence. *Italian Journal of Animal Science* **6** **(Suppl.2)**, 353-355. doi: 10.4081/ijas.2007.s2.353

397 Mroczkowski S, Korman K, Erhardt G, Piwczyński D, Borys B (2004) Sheep milk protein polymorphism  
398 and its effect on milk performance of Polish Merino. *Archiv Tierzucht, Dummerstorf* **47**, 114-121.

399 Pauciullo A, Cosenza G, D'Avino A, Colimoro L, Nicodemo D, Coletta A, Feligini M, Marchitelli C, Di  
400 Berardino D, Ramunno L (2010) Sequence analysis and genetic variability of stearoyl CoA  
401 desaturase (SCD) gene in the Italian Mediterranean river buffalo. *Molecular and Cellular Probes*  
402 **24** **(6)**, 407-410. doi: 10.1016/j.mcp.2010.07.009

403 Pauciullo A, Cosenza G, Steri R, Coletta A, Jemma L, Feligini M, Di Berardino D, Macciotta NPP,  
404 Ramunno L (2012a) An association between Single Nucleotide Polymorphisms at the Oxytocin

405 locus and milk yield in Italian Mediterranean river buffalo. *Journal of Dairy Research* **79** (2), 150-  
406 156. doi: 10.1017/S0022029911000914

407 Pauciullo A, Cosenza G, Steri R, Coletta A, La Battaglia A, Di Bernardino D, Macciotta NPP, Ramunno  
408 L (2012b) A single nucleotide polymorphism (SNP) in the promoter region of River buffalo  
409 Stearoyl CoA Desaturase gene (SCD) is associated with milk yield. *Journal of Dairy Research* **79**  
410 (4), 429-435. doi: 10.1017/S0022029912000507

411 Pirisi A, Piredda G, Papoff CM, Di Salvo R, Pintus S, Garro G, Ferranti P, Chianese L (1999). Effects  
412 of sheep alpha s1-casein CC, CD and DD genotypes on milk composition and cheesemaking  
413 properties. *Journal of Dairy Research* **66** (3), 409-419.

414 Prinzenberg EM, Weimann C, Brandt H, Bennewitz J, Kalm E, Schwerin M, Erhardt G (2003)  
415 Polymorphism of the Bovine *CSN1S1* Promoter: Linkage Mapping, Intragenic Haplotypes, and  
416 Effects on Milk Production Traits. *Journal of Dairy Science* **86**, 2696–2705.  
417 doi:10.3168/jds.S0022-0302(03)73865-X

418 Rando A, Di Gregorio P, Ramunno L, Mariani P, Fiorella A, Senese C, Marletta D, Masina P (1998)  
419 Characterization of the *CSN1AG* allele of the bovine  $\alpha$ s1-casein locus by the insertion of a relict  
420 of a long interspersed element. *Journal of Dairy Science* **81** (6), 1735–1742.  
421 doi:10.3168/jds.S0022-0302(98)75741-8

422 Ramunno L, Cosenza G, Rando A, Illario R, Gallo D, Di Bernardino D, Masina P (2004) The goat  $\alpha$ s1-  
423 casein gene: gene structure and promoter analysis. *Gene* **334**, 105-111  
424 doi:10.1016/j.gene.2004.03.006

425 Ramunno L, Cosenza G, Rando A, Pauciullo A, Illario R, Gallo D, Di Bernardino D, Masina P (2005)  
426 Comparative analysis of gene sequence of goat *CSN1S1* F and N alleles and characterization of  
427 *CSN1S1* transcript variants in mammary gland. *Gene* **345** (2), 289-299.  
428 doi:10.1016/j.gene.2004.12.003

- 429 Schmidely Ph, Meschy F, Tessier J, Sauvant D (2002) Lactation response and nitrogen, calcium, and  
430 phosphorus utilization of dairy goats differing by the genotype for alpha S1-casein in milk, and fed  
431 diets varying in crude protein concentration. *Journal of Dairy Science* **85** (9), 2299–2307.  
432 doi:10.3168/jds.S0022-0302(02)74310-5
- 433 Sukla S, Bhattacharya TK, Venkatachalapathy RT, Kumar P, Sharma A (2007). Molecular cloning and  
434 characterization of buffalo alpha(s1)-casein gene. *DNA Seq.* **18** (5), 334-340.
- 435 Yue XP, Zhang XM, Wang W, Ma RN, Deng CJ, Lan XY, Chen H, Li F, Xu XR, Ma Y, Lei CZ (2011)  
436 The *CSN1S1* N and F alleles identified by PCR-SSCP and their associations with milk yield and  
437 composition in Chinese dairy goats. *Molecular Biology Reports* **38** (4), 2821–2825. doi:  
438 10.1007/s11033-010-0428-0
- 439 Wessels G, Hamann H, Erhardt G, Distl O (2004) Genotype effects of milk protein polymorphisms on  
440 milk production in East Friesian dairy sheep. *Berliner Munchener Tierarztl Wochenschr* **117** (9-  
441 **10**), 414–419 [Article in German]
- 442

443 **Table 1.** Genotyping data and allele frequency of the c.628C>T SNP at the *CSN1S1* gene in the Italian  
 444 Mediterranean river buffalo population.

	Genotypes			Total	Allelic frequency	
	CC	CT	TT		C	T
<i>Observed</i>	75	84	16	<b>175</b>	0.67	0.33
<i>Expected</i>	78.22	77.55	19.22			

445  $\chi^2=1.20$ -  $p\leq 0.05$  - d.o.f.=1

446  
447

448 **Table 2.** Statistical significance of factors included in model

Effect	P-value		
	Milk yield	Fat	Protein
Genotype	0.63	0.93	0.04
Parity	<.001	<.001	<.001
Month of production	<.001	<.001	<.001
DIM	<.001	0.02	<.001
Season	<.001	0.48	0.02

449  
450

451 **Table 3.** Least squares means of milk yield (kg/d), fat and protein percentage (%) for the genotypes at  
 452 the *locus* c.628C>T of river buffalo *CSN1S1* gene estimated with model [\*].

Genotype	Animals n.	Milk yield (kg/d)	Fat (%)	Protein (%)
CC	75	7.81±0.17	9.22±0.12	4.72±0.016 <sup>a</sup>
CT	84	7.92±0.17	9.27±0.12	4.68±0.016 <sup>b</sup>
TT	16	7.56±0.36	9.20±0.23	4.69±0.029 <sup>ab</sup>

453 <sup>a,b</sup> Means within columns with different superscripts differ (P=0.038)


454  
455  
456  
457  
458  
459  
460  
461  
462

463 **Table 4.** Substitution effect of a cytosine with a thymine at the c.628C>T SNP in the *CSN1S1* gene (mean  
 464  $\pm$  SE) and contribution of the  $\alpha$ 1 polymorphism to the phenotypic variance for protein percentage.

Statistic	Value
$\alpha$	-0.014 $\pm$ 0.014
d	-0.028 $\pm$ 0.019
$\sigma^2_{\alpha 1}$	0.0004
$\sigma^2_c$	0.0034
$\sigma^2_e$	0.1596
$r^2_{\alpha 1}$	0.003
$r^2_c$	0.021

465  $\alpha$ : Substitution effect;  
 466 d: dominance effect;  
 467  $\sigma^2$ : variance components associated to the genotype ( $\alpha$ 1); to the individual buffalo cow (c), to residuals  
 468 (e);  
 469  $r^2$ : contributions of genotype ( $\alpha$ 1) and of individual buffalo cow (c) to the total phenotypic variance.  
 470

471  
 472  
 473  
 474  
 475  
 476  
 477  
 478  
 479  
 480  
 481  
 482  
 483  
 484  
 485  
 486  
 487  
 488  
 489  
 490  
 491



**Figure 1.** Observed genotypes after *Mbo*I digestion of fragments obtained by ACRS-PCR of a DNA region corresponding to the last 86 bp of the 17<sup>th</sup> exon into Mediterranean river buffalo *CSN1S1* gene. M=2-Log DNA ladder (0.1-10.0kb) (New England Biolabs); lane 1: *CSN1S1* C/C; lane 2: *CSN1S1* C/T; lane 3 *CSN1S1* T/T