



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

# $\beta$ -cyclodextrin-based nanosponges improve 1-MCP efficacy in extending the postharvest quality of cut flowers

This is the author's manuscript		
Original Citation:		
Availability:		
This version is available http://hdl.handle.net/2318/136531since 2015-12-09T14:33:51Z		
Published version:		
DOI:10.1016/j.scienta.2013.05.019		
Terms of use:		
Open Access		
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.		

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

*This is an author version of the contribution published on*: Questa è la versione dell'autore dell'opera: Scientia Horticulturae, Volume 159, 30 July 2013, Pages 162-165

*The definitive version is available at:* La versione definitiva è disponibile alla URL: http://www.sciencedirect.com/science/article/pii/S030442381300246X

1	$\beta$ -cyclodextrin-based nanosponges improve 1-MCP efficacy in extending the
2	postharvest quality of cut flowers
3	Ludovica Seglie <sup>a</sup> , Marco Devecchi <sup>a</sup> , Francesco Trotta <sup>b</sup> , Valentina Scariot <sup>a,*</sup>
4	
5	<sup>a</sup> Department of Agriculture, Forest and Food Sciences, University of Torino, Via
6	Leonardo da Vinci 44 - 10095 Grugliasco (TO), Italy.
7	<sup>b</sup> Department of Chemistry, University of Torino, Via Pietro Giuria 7 - 10125 Turin,
8	Italy.
9	
10	*Corresponding author. Tel.: +39 011 6708932, fax: +39 011 6708798, e-mail address:
11	valentina.scariot@unito.it (V. Scariot).
12	
13	
14	Abstract
15	Investigations of the effect of the $\beta$ -cyclodextrin-based nanosponge-1-
16	methylcyclopropene (1-MCP) complex ( $\beta$ -CD-NS complex) on six ethylene-sensitive
17	cut flowers were performed. Anemone coronaria L. multicolor, Ranunculus asiaticus L.
18	'Minou Abrown', Helianthus annuus L. 'Sunrich Orange', Rosa hybrida L. 'Jupiter',
19	Paeonia lactiflora L. 'Sarah Bernhardt', and Papaver nudicaule L. multicolor

20 underwent four different treatments: a vase placement in a suspension of  $\beta$ -CD-NS 21 complex (0.25  $\mu$ L L<sup>-1</sup>), exposure to gaseous 1-MCP (0.25  $\mu$ L L<sup>-1</sup>) for 6 h, exposure to 22 exogenous ethylene (1 ± 0.2  $\mu$ L L<sup>-1</sup>) for 24 h, and placement in tap water.

β-cyclodextrin-based nanosponges proved to enhance 1-MCP efficacy in all the tested
species. According to the species, senescence symptoms were decreased or delayed.
Anemone and poppy had a longer aesthetic flower quality and vase life, and a reduced

petal abscission. In ranunculus, β-CD-NS complex improved longevity. In sunflower, it
also maintained longer chlorophyll leaf content and cut flower ornamental value. Peony
opened more slowly and rose maintained its ornamental quality longer than 1-MCP
treated flowers. This study opens possibilities for commercial use of the β-CD-NS
complex in the floriculture industry.

31

*Keywords:* Ethylene inhibitor; Flower longevity; Nano-carrier; Preservative;
Senescence; Vase life.

34

*Abbreviations:* CD, cyclodextrin; 1-MCP, 1-methylcyclopropene; β-CD-NS, β-CDbased-nanosponge; β-CD-NS complex, 1-MCP included in β-CD-NS

37

#### 38 **1. Introduction**

Postharvest performance is of crucial importance to the value of cut flowers. The life span of flowers is generally determined by the time to abscission of petals that are still turgid, or by the time to petal wilting or withering. In many species, these phenomena are regulated by ethylene (Woltering and Van Doorn, 1988; van Doorn, 2001). Premature senescence and abscission resulting from exposure to exogenous or endogenous ethylene can be controlled several ways including ethylene action inhibitors (Martínez-Romero et al., 2007) quite commonly.

In particular, while 1-methylcyclopropene (1-MCP), has proved itself to be a very
effective preventer of negative ethylene responses, its gaseous nature makes treatment
difficult (Serek and Sisler, 2005; Serek et al., 2006).

Recently, we developed a non-volatile formulation of 1-MCP in a β-cyclodextrinbased nanosponge (β-CD-NS 1:8) structure. Supplied in the conservation solution, this

formulation (β-CD-NS complex) prolonged the vase life of carnation cut flowers (Seglie
et. al., 2011a, b), and controlled *B. cinerea* damage (Seglie et al., 2012) better than
gaseous 1-MCP.

54 While carnation is generally used in studies on postharvest preservation of ethylene-55 sensitive flowers (Woltering and van Doorn, 1988; Serek et al., 1995a, b); it is not a 56 universal paradigm for flower senescence. The response to ethylene varies widely 57 according to species (Reid and Wu, 1992). Therefore, our study evaluated the 58 effectiveness of the  $\beta$ -CD-NS complex to improve cut flower vase life on a number of 59 ethylene-sensitive species.

60

# 61 2. Materials and methods

The flower market of Sanremo (UCFlor, Liguria, Italy) seasonally provided cut
flowers of *Anemone coronaria* L. multicolor, *Ranunculus asiaticus* L. 'Minou Abrown', *Helianthus annuus* L. 'Sunrich Orange', *Rosa hybrida* L. 'Jupiter', *Paeonia lactiflora*Pall. 'Sarah Bernhardt', and *Papaver nudicaule* L. multicolor.

Stems were transferred to the postharvest laboratory of the Department of Agricultural, Forest and Food Sciences at the University of Turin. Experimental evaluations of vase life were performed in an imitated inside environment (IE) maintained at  $20\pm2^{\circ}$  C, 60 % RH, and 12 h daily of 46 µmol m<sup>-2</sup> s<sup>-1</sup> cool white light (meter model HT307; HT, Faenza, Italy).

Cut flowers were re-cut, labeled and re-watered. Each species sample was divided into sub-groupings of six cut stems (each 30 cm long after re-cutting). Each sub-group was then treated according to the following schemes: in vases with an aqueous suspension of  $\beta$ -CD-NS complex to supply 0.25  $\mu$ L L<sup>-1</sup> of active ingredient; in an air tight cabinet (112 L) exposed to equal concentrations of volatile 1-MCP (3.3% a. i.,

SmartFresh<sup>TM</sup>, AgroFresh Inc., USA) for 6 h; in another air tight cabinet exposed to exogenous ethylene  $(1 \pm 0.2 \ \mu L \ L^{-1})$  for 24 h; and in vases of tap water as a control. The experiment was performed twice for each species.

Endogenous ethylene production of the control flowers was measured using a digital Agilent Technologies gas chromatograph, 7890A Network GC system (Santa Clara, California). N<sub>2</sub> at 40 mL min<sup>-1</sup> was used as the gas carrier at a column temperature 60 °C and calibration range between 0.5-3 ppm. For each species, six samples were considered.

Flower postharvest performance was measured and visual checks for symptoms of variation in senescence were performed daily. In particular, we monitored senescence level (1 = initial senescence, 2 = complete senescence; Seglie et al., 2011a), flower opening stage (1 = initial opening, 2 = half opening, 3 = full opening; Guo et al., 2004, modified), and abscission. Chlorophyll content was indirectly measured in leaves through the Chlorophyll Meter SPAD-502 instrument (Konica Minolta Sensing Inc., Osaka, Japan).

Analysis of variance (ANOVA) was performed to assess statistical significance
among mean values using the Ryan–Einot–Gabriel–Welsch's multiple step-down F
(REGW-F) test (p≤0.05) with SPSS Software (Chicago, USA).

94

### 95 3. Results and Discussion

96 Senescence performance and ethylene sensitivity are highly species related (van 97 Doorn, 2001). Large variabilities have been observed even among cultivars, such as the 98 Asian buttercup (Kenza et al., 2000; Scariot et al., 2009), rose (Chamani et al., 2005), 99 and peony (Hoffman et al., 2010; Eason et al., 2002). Flowers that respond to low 100 concentrations of exogenous ethylene are probably those in which ethylene is naturally

101 involved in senescence (Reid and Wu, 1992). In this study, ethylene hastened the 102 symptoms of natural senescence in all species. However, a climacteric pattern was 103 observed only in poppy (ethylene up to 12  $\mu$ L L<sup>-1</sup> at day 5) while in peony and 104 sunflower ethylene production was not detected (data not shown).

Anemone cut flowers lose their decorative value in a very short period. Pedicels 105 elongate excessively and petals wilt, lose their color, become translucent, and abscise 106 107 (Meir, 2007). Previous studies have already demonstrated the ethylene sensitivity of the anemone (Armitage, 1993) and the efficacy of ethylene antagonists (silver nitrate and 108 109 silver thiosulphate, STS) in extending its vase life and preventing petal abscission (Sharifani et al., 2005). In this study, the application of  $\beta$ -CD-NS complex was found to 110 prolong cut flower aesthetic quality and longevity for up to 1.5 additional days, (Fig. 1), 111 112 slow stage 1 and 2 of flower opening (Tab. 1), and completely prevent abscission (Fig. 2) more effectively than gaseous 1-MCP. Petal rolling over and stem folding (Sharifani 113 114 et al., 2005), which usually occur in anemone cut flowers, were also limited.

115 Asian buttercup flower senescence is characterized by loss of turgor and petal colour change followed by petal wilting and drop when lightly touched. Previous tests of the 116 effects of silver thiosulphate (STS) and aminooxyacetic acid (AOA) on buttercup 117 postharvest performance have yielded unsatisfactory results in enhancing vase life 118 (Shahri et al., 2011). In this study, ethylene antagonists were effective in delaying 119 120 senescence symptoms with  $\beta$ -CD-NS complex; they outperformed gaseous 1-MCP as 121 demonstrated by delay of complete senescence (9.7 versus 8 days; Fig. 1) and original 122 flower colour maintenance (data not shown).

Previous studies of sunflower have shown exogenous ethylene applications result in short vase life, flower abscission (Redman et al., 2002), and in largely, wilted basal foliage immediately after anthesis (Pallez et al., 2002; Castro et al., 2011). Furthermore,

ethylene antagonist treatment has failed to limit or delay senescence (Redman et al., 2002). In this study, chlorophyll degradation was restrained by ethylene antagonist application. At day 5, the SPAD value was  $26.2 \pm 0.63$  units for control flowers and on average  $33.7 \pm 0.19$  units for flowers treated with ethylene antagonists. β-CD-NS complex was more effective than gaseous 1-MCP at prolonging vase life (9 *versus* 7 days, respectively; Fig. 1) and at slowing flower opening (stages 1 and 2; Tab. 1).

132 In rose, flower opening seemed to be affected by exogenous ethylene (Reid et al., 1989). Ethylene induced abscission of fully turgescent, non-senescent petals or the 133 entire corolla (Serek et al., 2006), which suggests that anti-ethylene compounds could 134 retard the behavior (Serek et al., 1995b) and extend vase life (Mor et al., 1989). 135 Precedent studies have noted decreased leaf and bud abscission by application of 1-136 MCP along with increased display life in both ethylene-contaminated and ethylene-free 137 138 air (Muller et al., 2000; Serek et al., 1994). This study showed β-CD-NS complex performed better than commercial 1-MCP as evidenced by extension of ornamental rose 139 140 quality (5.0 and 3.5 days, respectively; Fig. 1), and slowed flower opening (stages 2 and 141 3; Tab. 1).

Peony has been found by Jia et al. (2008) to have a high ethylene sensitivity; they correlated it to an increased flower diameter during senescence. On the other hand, Hoffman et al. (2010) have found 1-MCP ineffective at increasing storage longevity and quality of cut peony flowers. Our study results show that both 1-MCP applications significantly increased the longevity of peony to almost 2 days (7 days) beyond that of the control (5 days; Fig. 1), and limited abscission (Fig. 2). Also, β-CD-NS complex outperformed the commercial product in slowing flower opening (Fig. 2).

The vase life of poppy is usually 5 days in untreated flowers; it ends when petals shatter or crinkle, discolor or turn brown, and the stem collapses (Dole et al., 2009). This research indicated that gaseous 1-MCP was ineffective at delaying these symptoms of senescence while  $\beta$ -CD-NS complex, on the other hand, was shown to extend ornamental flower quality for up to 5.8 days and increased the vase life to as long as 8.7 days (Fig. 1). Moreover, this compound significantly reduced petal abscission (Fig. 2) and retarded flower opening (Tab. 1). Therefore,  $\beta$ -CD-NS complex use could increase the commercial market for poppy cut flowers, albeit one that will continue to be limited by high production costs (Dole and Greer, 2009).

In conclusion, β-CD-NS 1:8 was shown to enhance the efficacy of 1-MCP in six
ethylene-sensitive species, as was previously seen in carnation (Seglie et al., 2011a, b).
1-MCP is a high unstable and reactive gas that very quickly dimerizes even at room
temperature. This dimer has not antiethylenic activity. Most likely β-CD-NS stabilize to
a great extent the included 1-MCP thus preserving its properties.

163 The present study opens the possibilities of commercial use of the β-CD-NS complex164 in the floriculture industry.

165

## 166 Acknowledgements

The authors gratefully acknowledge the Flower Market of Sanremo (UCFLOR),
growers Lino Bloise, La Marca S. S., and Mediterranean Flowers S. S. A., for their
provision of cut flowers, and AgroFresh, Inc. (Rohm and Haas) for providing
SmartFresh<sup>™</sup>. This research was funded by the Liguria Region project number
94750059241.

- Armitage, A.M., Laushman, J.M., 1993. Specialty cut flowers: the production of
  annuals, perennials, bulbs, and woody plants for fresh and dried cut flowers. Timber
  press Inc. Portland, Oregon.
- 177 Castro, A.M.C., Sato, O., Rando, E.M., Santos, K.H.D., Silva, D.P.Da, Chimbo, JR.A.,
- Carvalho, F.K., Braga, C.DeL., 2011. Uso de fontes de nitrogênio visando retarder a
  senescência em girassol ornamental. Scientia Agraria Paranaenis 10, 83-90.
- 180 Chanami, E., Khalighi, A. Joyce, D.C., Irving, D.E., Zamani, Z.A., Mostofi, Y., Kafi,
- 181 M., 2005. Ethylene and anti-ethylene treatment effects on cut 'First Red' rose. J.
  182 Applied Hortic. 7, 3-7.
- 183 Dole, J.M., Fonteno, W.C., Blankenship, S. L., 2005. Comparison of silver thiosulphate
- 184 with 1-methylcyclopropene on 19 cut flower taxa. Acta Hortic. 682, 249-256.
- 185 Dole, J.M., Viloria, Z., Fanelli, F.L., Fonteno, W., 2009. Postharvest evaluation of cut
- dahlia, linaria, poppy, rudbeckia, trachelium, and zinnia. HortTechnology. 19, 593-600.
- Dole, J.M., Greer, L., 2009. Production protocol development for greenhouse cut
   *Linaria, Lupinus*, and *Papaver* flowers. Sci. Hortic. 122, 233-237.
- 190 Eason, J., Pinkney, T., Heyes, J., Brash, D., Bycroft, B., 2002. Effect of storage
- temperature and harvest bud maturity on bud opening and vase life of *Paeonia lactiflora* cultivars. New Zeal. J. Crop Hort. 30, 61-67.
- Guo, W., Chen, R., Dong, R., 2004. The postharvest characteristic and water balance of
  some cultivars of tree-peony cut flowers. Sci. Silvae Sin. 40, 89-93.
- 195 Hoffman, G.D., Mattinson, D.S., Fellman, J.K., 2010. 1-MCP shortens peony vase life.
- 196 Acta Hort. 857, 169-178.
- 197 Kenza, M., Umiel, N., Borochov, A., 2000. The involvement of ethylene in senescence
- 198 of ranunculus cut flowers. Postharvest Biol. Technol. 19. 287-290.

- Jia, P., Zhou, L., Guo, W., Wang, L., Dong, L., 2008. Postharvest behavior and
  endogenous ethylene pattern of *Paeonia suffrutticosa* cut flowers. Acta Hort. 768,
  445-450.
- 202 Martínez-Romero, D., Bailén, G., Serrano, M., Guillén, F., Valverde, J.M., Zapata, P.,
- Castillo, S., Valero, D., 2007. Tools to maintain postharvest fruit and vegetable
  quality through the inhibition of ethylene action: a review. Crit. Rev. Food Sci. 47,
  543-560.
- 206 Meir, S., Salim, S., Chernov, Z., Philosoph-Hadas, S., 2007. Quality improvement of
- 207 cut flowers and potted plants with postharvest treatments based on various cytokinins208 and auxins. Acta Hort. 755, 143-154.
- Mor, Y., Johnson, F., Faragher, J.D., 1989. Preserving the quality of cold-stored rose
  flowers with ethylene antagonists. Hortic. Sci. 24, 640-641.
- 211 Muller, R., Sisler, E.C., Serek, M., 2000. Stress induced ethylene production, ethylene
- binding, and the response to the ethylene action inhibitor 1-MCP in miniature roses.
- 213 Sci. Hortic. 83, 51-59.
- Pallez, L.C., Dole, J.M., Whipker, B.E., 2002. Production and postproduction studies
  with potted sunflowers. HortTechonology 12, 206-210.
- Redman, P.B., Dole, J.M., Maness, N.O., Anderson, J.A., 2002. Postharvest handling of
  nine specialty cut flower species. Sci. Hortic. 92, 293-303.
- Reid, M.S., Evans, R.Y., Dodge, L.L., Mor, Y., 1989. Ethylene and silver thiosulphate
- influences opening of cut rose flower. J. Am. Soc. Hortic. Sci. 114, 436-440.
- Reid, M.S., Wu, M.J., 1992. Ethylene and flower senescence. Plant Growth Regul. 11,
  37-43.

- Scariot, V., Larcher, F., Caser, M., Costa, E., Beruto, M., Devecchi, M., 2009. Flower
  longevity in ten cultivars of cut *Ranunculus asiaticus* L. as affected by ethylene and
  ethylene inhibitors. Europ. J. Hort. Sci. 74, 137-142.
- 225 Seglie, L., Martina, K., Devecchi, M., Roggero, C., Trotta F., Scariot, V., 2011a. The
- effects of 1-MCP in cyclodextrin-based nanosponges to improve the vase life of
   *Dianthus caryophyllus* cut flowers. Postharvest Biol. Technol. 59, 200-205.
- Seglie, L., Martina, K., Devecchi, M., Roggero, C., Trotta F., Scariot, V., 2011b. βcyclodextrin-based nanosponges as carriers for 1-MCP in extending the postharvest
  longevity of carnation cut flowers. An evaluation of different degrees of crosslinking. Plant Growth Regul. 65, 505-511.
- 232 Seglie, L., Spadaro, D., Trotta, F., Devecchi, M., Gullino, M.L., Scariot, V., 2012. Use
- of 1-methylcylopropene in cyclodextrin-based nanosponges to control grey mould
  caused by *Botrytis cinerea* on *Dianthus caryophyllus* cut flowers. Postharvest Biol.
  Technol. 64, 55-57.
- Serek, M., Sisler, E.C., Reid, M.S., 1994. Novel gaseous ethylene binding inhibitor
  prevents ethylene effects in potted flowering plants. J. Am. Hort. Sci. 119, 12301233.
- Serek, M., Sisler, E.C., Reid, M.S., 1995a. Effects of 1-MCP on the vase life and
  ethylene response of cut flowers. Plant Growth Regul. 16, 93-97.
- Serek, M., Sisler, E.C., Reid, M.S., 1995b. 1-methylcyclopropene, a novel gaseous
  inhibitor of ethylene action, improves the vase life of fruits, cut flowers and potted
  plants. Acta Hort. 394, 337-346.
- 244 Serek, M., Sisler, E.C., 2005. Impact of 1-MCP on postharvest quality of ornamentals.
- In: APEC Symp. Quality Management of Postharvest System Proceedings, 121-128.

246	Serek, M., Sisler, E.C., Frello, S., Sriskandarajah, S., 2006. Postharvest technologies for
247	extending the shelf life of ornamental crops. Int. J. Postharvest Technol. Inn. 1, 69-
248	75.

- Sharifani, M., Parizadeh, M., Mashayekhi, K., 2005. The effects of pre-storage on
  postharvest quality of cut anemone (*Anemone coronaria*) flowers. Acta Hort. 682,
  701-707.
- van Doorn, W.G., 2001. Categories of petal senescence and abscission: a re-evaluation.
  Ann. Bot. 87, 447-456.
- 254 Woltering, E.J., van Doorn, W.G., 1988. Role of ethylene in senescence of petals--

255 morphological and taxonomical relationship. J. Exp. Bot. 39, 1605-1616.

256 257

#### **Figure captions**

- **Fig. 1** Longevity of six species of cut flowers according to four treatments: placement in a vase suspension of  $\beta$ -CD-NS complex (0.25  $\mu$ L L<sup>-1</sup>), exposure to gaseous 1-MCP (0.25  $\mu$ L L<sup>-1</sup>) for 6 h, exposure to exogenous ethylene (1 ± 0.2  $\mu$ L L<sup>-1</sup>) for 24 h, and placement in tap water. The count of days to reach the initial stage of senescence (Level 1, line) and complete senescence (Level 2, dotted line) was measured.
- \*Mean separation within bars by the Ryan-Einot-Gabriel-Welsch's multiple stepdown F
  (REGW-F) test, P≤0.001.

266

**Fig. 2** Petal abscission in six flower species according to four treatments: placement in a vase suspension of  $\beta$ -CD-NS complex (0.25  $\mu$ L L<sup>-1</sup>), exposure to gaseous 1-MCP (0.25  $\mu$ L L<sup>-1</sup>, for 6 h, exposure to exogenous ethylene (1 ± 0.2  $\mu$ L L<sup>-1</sup>) for 24 h, and placement

- 270 in tap water. For each species, abscission (%) was calculated as the total number of
- abscised petals/the mean number of petals in three flowers \* 100.
- \*Mean separation within bars by the Ryan-Einot-Gabriel-Welsch's multiple stepdown F
- 273 (REGW-F) test, P≤0.001.
- 274