

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Molecular typization of *Arthroderma benhamiae*, a zoonotic agent of epidemic dermatophytosis in Central Europe

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1531130> since 2015-12-03T12:53:57Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Molecular typization of *Arthroderma benhamiae*, a zoonotic agent of epidemic dermatophytosis in Central Europe

Mycoses

Special Issue: 7th Trends in Medical Mycology, 9–12 October 2015, Lisbon, Portugal

[Volume 58, Issue Supplement S4](#), pages 51–226, October 2015

A. Cmokova,¹ M. Kolarik,² T. Vetrovsky,³ S. Dobiasova,⁴ R. Dobias,⁵ D. Stubbe,⁶ M. Skorepova,⁷ P. Lyskova,⁸ L. Hoyer,⁹ N. Mallatova,¹⁰ R. Kano,¹¹ P. Nenoff,¹² S. Uhrlab,¹² A. Peano,¹³ J. Koubkova,¹⁴ K. Mencl,¹⁵ H. Janouskovcova¹⁵ and V. Hubka¹

¹Charles University in Prague, Prague, Czech Republic; ²Institute of Microbiology, Czech Academy of Science, Prague, Czech Republic; ³Institute of Microbiology of the AS CR, Prague, Czech Republic; ⁴Institute of Public Health, Ostrava, Czech Republic; ⁵Institute of Health in Ostrava, Ostrava, Czech Republic; ⁶Scientific Institute of Public Health, Brussels, Belgium; ⁷Charles University in Prague, First Faculty of Medicine, Prague, Czech Republic; ⁸Institute of Health in Usti nad Labem, Prague, Czech Republic; ⁹University of Illinois at Urbana-Champaign, College of Veterinary Medicine, Prague, Czech Republic; ¹⁰Hospital Ceske Budejovice, Ceske Budejovice, Czech Republic; ¹¹Nihon University School of Veterinary Medicine, Fujisawa, Japan; ¹²Laboratory of medical microbiology, M€olbis, Germany; ¹³Universit_a degli Studi di Torino, Facoltà di Medicina Veterinaria, Turin, Italy; ¹⁴Clinical Veterinary Laboratory Labvet, Prague, Czech Republic and ¹⁵Pardubice Regional Hospital, Inc., Pardubice, Czech Republic

Objectives

The aim of this study was to find informative molecular markers to evaluate population structure of dermatophyte species *Arthroderma benhamiae* in the studied area. One of the main issues was to determine whether the current outbreak of dermatophytosis caused by *A. benhamiae* in Central Europe is caused by a new more virulent genotype which is with the high success rate transmitted to humans or whether other causes must be considered (e.g. high prevalence of the pathogen in animal husbandry).

Methods A total number of 262 *A. benhamiae* strains associated with cases of human and animal dermatophytosis from the Czech Republic (CZ), Belgium, UK, Germany, Italy, Japan and USA were used for analysis. Ten microsatellite markers were developed and used for typization of *A. benhamiae* strains together with sequence analysis of ITS region rDNA and glyceraldehyde-phosphate dehydrogenase gene. Bayesian inference analysis and distance methods were applied on the datasets to reveal the genetic variation of *A. benhamiae* population. PCR-based determination of mating idiomorphs was also performed in all strains. **Results** The sequence analysis based on two genes revealed presence of five genotypes. Most common genotype (n = 195, 74.4%) was characterized by yellow colony reverse on MEA agar with the exception of isolates from dogs in North America which showed different phenotype. Only MAT1-1 idiomorph was amplified in isolates of this genotype from Europe and low variability in microsatellites data was recorded. In contrast, the isolates of this genotype from North America were highly variable by microsatellite data and isolates of both mating type genes were revealed. The second most common genotype (n = 47, 17.9%) was characterized by dominant red (mating type MAT2) or brown (MAT1) colony reverse on MEA agar. The third genotype (n = 16), closely related to second genotype, included all strains from Japan and some from the CZ and Belgium and was characterized by both red and yellow colony reverse and MAT1-1 idiomorph. The remaining two genotypes were each represented by only two isolates from the CZ.

Conclusion *A. benhamiae* is a new emerging pathogen in the CZ and some other countries in Central and Western Europe region. According to recent studies this species is the most important zoophilic dermatophyte in the CZ causing almost 23% of tinea corporis (median age of patients – 10 years, females - 70%) and 29% of tinea capitis infections. The majority of infections are transmitted from guinea pigs and other rodents. It is probable that the most common genotype of *A. benhamiae* responsible for the outbreak of infections in the Central Europe (agent of 79.7% of human infections in the CZ) spread clonally. This hypothesis is also supported by uniform phenotype and mating type in European strains. The american strains of the same genotype (can be separated by microsatellite data) however showed different phenotype and presence of both mating type genes indicating sexual reproduction. Other genotypes contribute only marginally on the outbreak.