



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Footstep-triggered grain flows on the lee side of a desert sand dune (Erg Chebbi, Morocco)

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/1542195 since 2016-01-08T13:01:34Z
Published version:
DOI:10.1007/s00531-015-1246-3
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on: Questa è la versione dell'autore dell'opera: International J. of Earth Sciences, 104, 8, 2015, DOI: 10.1007/s00531-015-1246-3 ovvero Barale, Springer, 2015, pagg. 2213-2214

> *The definitive version is available at:* La versione definitiva è disponibile alla URL: http://link.springer.com/article/10.1007%2Fs00531-015-1246-3

1	Footstep-triggered grain flows on the lee side of a desert sand dune (Erg Chebbi, Morocco)
2	
3	Luca Barale
4	Dipartimento di Scienze della Terra, Via Valperga Caluso 35, 10125 Torino (Italy). luca.barale@hotmail.it
5	
6	
7	Acknowledgements
8	The author thanks the reviewers Nick Lancaster and Nigel P. Mountney, whose useful suggestions and
9	constructive criticisms really improved the manuscript. Carlo Bertok, Niccolò Dainelli, Alessandro Ellero,
10	Andrea Irace, Paolo Nafissi, Giuseppe Ottria, Fabrizio Piana, and Youssef Bakabou are kindly acknowledged
11	for sharing fieldwork in the Moroccan High Atlas during three field campaigns in 2013–2014.

12 (without abstract)

13 Grain flows — or grain avalanches — are important depositional processes on the lee side of aeolian sand 14 dunes. Sediment deposition on the lee side of aeolian dune bedforms is mainly represented by saltating sand 15 that is driven by the wind over the brink of the lee face and is deposited by grainfall as the flow separates at 16 the brink (e.g., Allen 1970; Hunter 1977, 1985; Anderson 1988; McDonald and Anderson 1995). Grain flows 17 occur when sediment deposition on the lee side causes it to reach the angle of repose of sand, so that further 18 sediment deposition produces the failure of the slope and the initiation of a grain flow (e.g., Allen 1970; 19 Hunter 1977, 1985; McDonald and Anderson 1995). The angle of repose for loose, non-cemented dry sand 20 grains that are well sorted and rounded, as is typical of aeolian sands, is about 34° (e.g., Carrigy 1970; 21 Fryberger and Schenk 1981; McDonald and Anderson 1995). In recent years, different studies have 22 investigated the frequency and the magnitude of grain flows in relation to the morphology and dimensions of 23 the bedforms on which they develop and to the speed and direction of the wind (e.g., Nickling et al. 2002; 24 Breton et al. 2008; Sutton et al. 2013; Romain and Mountney 2014). A better understanding of the processes 25 that control the development of grain flows and their resultant deposits has important implications in the 26 modelling of aeolian sandstone reservoirs, as grain-flow deposits typically have significantly higher 27 porosities and permeabilities than associated deposits (grainfall laminae and wind-ripple beds), due to the 28 typically loose packing of well sorted, well rounded and highly spherical grains (e.g., Prosser and Maskall 29 1993; Howell and Mountney 2001). In addition to the above cited sedimentary processes, lee-side failures 30 and consequent grain flows can be also triggered by animals (or persons) walking on the lee side or on the 31 crest of a dune bedform (e.g., McKee 1947; Andreotti 2004; Vriend et al. 2007). An example of dinosaur-32 triggered grain flows has been described by Loope (2006) in the aeolian deposits of the Lower Jurassic 33 Navajo Sandstone (Arizona–Utah, USA).

The photograph shows a series of tongue-shaped grain-flow lobes on the lee side of an aeolian sand dune at the northern edge of Erg Chebbi (Errachidia Province, eastern Morocco). The lobes are up to 1.5 m long and 0.4–0.5 m wide, and have a maximum thickness of 30–40 mm. Different generations of grain flows can be distinguished, with later, smaller grain flows locally developed from the partial reworking of earlier grainflow lobes. The regular spacing of the grain-flow lobes, and their indentation directly on the dune crestline, indicate that the grain flows were probably triggered by a person walking along the dune crest, whose 40 footsteps produced a series of localized slope failures at the top of the lee side. Wind ripples, striking almost 41 perpendicular to the crest of the dune, are present on the dune apron. A winding beetle trail, about 40 mm 42 wide, is visible in the left part of the image. The photograph was taken on November 12th, 2014 (coordinates: 43 31°12'38.7" N, 3°59'28.6" W).

44

45 References

- 46 Allen J (1970) The avalanching of granular solids on dune and similar slopes. J Geol 78:326–351
- 47 Anderson RS (1988) The pattern of grainfall deposition in the lee of aeolian dunes. Sedimentology 35:175–
- 48 188
- 49 Andreotti B (2004) The song of dunes as a wave-particle mode locking. Phys Rev Lett 93(23):238001
- 50 Breton C, Lancaster N, Nickling WJ (2008) Magnitude and frequency of grain flows on a desert sand dune.
- 51 Geomorphology 95:518–523
- 52 Carrigy MA (1970) Experiments on the angles of repose of granular materials. Sedimentology 14:147–158
- 53 Fryberger SG, Schenk C (1981) Wind sedimentation tunnel experiments on the origins of aeolian strata.
- 54 Sedimentology 28:805–821
- 55 Howell JA, Mountney NP (2001) Aeolian grain flow architecture: hard data for reservoir models and
- 56 implications for red bed sequence stratigraphy. Petrol Geosci 7:51–56
- 57 Hunter RE (1977) Basic types of stratification in small eolian dunes. Sedimentology 24:361–388
- 58 Hunter RE (1985) A kinematic model for the structure of lee-side deposits. Sedimentology 32:409–422
- 59 Loope DB (2006) Dry-season tracks in dinosaur-triggered grainflows. Palaios 21:132–142
- 60 McDonald RR, Anderson RS (1995) Experimental verification of aeolian saltation and lee side deposition
- 61 models. Sedimentology 42:39–56
- 62 McKee ED (1947) Experiments on the development of tracks in fine cross-bedded sand. J Sed Petrol 17:23-
- 63 28
- 64 Nickling WG, McKenna Neuman C, Lancaster N (2002) Grainfall processes in the lee of transverse dunes,
- 65 Silver Peak, Nevada. Sedimentology 49:191–209
- 66 Prosser DJ, Maskall R (1993) Permeability variation within aeolian sandstone: a case study using core cut
- 67 sub-parallel to slipface bedding, the Auk Field, Central North Sea, UK. Geol Soc Spec Publ 73:377–397

- 68 Romain HG, Mountney NP (2014) Reconstruction of three-dimensional eolian dune architecture from one-
- 69 dimensional core data through adoption of analog data from outcrop. AAPG Bull 98(1):1–22
- 70 Sutton SLF, McKenna Neuman C, Nickling WJ (2013) Avalanche grainflow on a simulated aeolian dune. J
- 71 Geophys Res-Earth 118:1767–1776
- 72 Vriend NM, Hunt ML, Clayton RW, Brennen CE, Brantley KS, Ruiz-Angulo A (2007) Solving the mystery
- 73 of booming sand dunes. Geophys Res Lett 34:L16306

