

The Bethe Ansatz and the Tzitzéica-Bullough-Dodd equation

Patrick Dorey¹, Simone Faldella², Stefano Negro³ and Roberto Tateo³

¹*Department of Mathematical Sciences, Durham University,
South Road, Durham DH1 3LE, United Kingdom*

²*Institut de Mathématiques de Bourgogne, Université de Bourgogne,
9 av. Alain Savary - B.P. 47 870, 21078 Dijon, France*

³*Dip. di Fisica and INFN, Università di Torino,
Via P. Giuria 1, 10125 Torino, Italy*

E-mails:

p.e.dorey@durham.ac.uk, simone.faldella@u-bourgogne.fr,
negro@to.infn.it, tateo@to.infn.it

Abstract

The theory of classically integrable nonlinear wave equations, and the Bethe Ansatz systems describing massive quantum field theories defined on an infinite cylinder, are related by an important mathematical correspondence that still lacks a satisfactory physical interpretation. In this paper we shall describe this link for the case of the classical and quantum versions of the (Tzitzéica-)Bullough-Dodd model.

PACS: 03.65.-Ge, 11.15.Tk, 11.25.HF, 11.55.DS.

Keywords: Spectral problems, affine Toda field theory, Bethe Ansatz.

1 Introduction

An interesting correspondence was discovered in [1] between the functional approach to the spectral theory of ordinary differential equations, mainly developed by Sibuya, Voros and collaborators [2, 3], and a series of works by Bazhanov, Lukyanov and Zamolodchikov [4, 5, 6], where earlier results of Baxter [7] were extended to the conformal field theories governing the continuum limits of certain integrable models (IM) on a 2d lattice.

It turns out that the initial observation of [1] was the first hint of a very general mathematical scheme – an ‘ODE/IM correspondence’ – involving wide classes of multi-parameter ordinary differential equations. Their generalized eigenvalue problems turn out to be constrained by the same Bethe Ansatz equations as arise in the conformal field theory limits of certain integrable vertex models, related to Lie algebras [8, 9, 10, 11, 12, 13, 14].

A precise relationship between elements of this new scheme and the \mathcal{PT} -symmetric quantum mechanical models of Bender, Boettcher and Meisinger [15, 16] was established in [17]. This has led to proofs of spectral reality in certain cases [18], that were further generalised in [19], and insights into the loss of reality in others [20]. Applications of the correspondence are also being found in various problems arising in condensed matter physics [21, 22, 23].

A generalization of this correspondence to off-critical, or massive, quantum field theories was subsequently obtained by Gaiotto, Moore and Neitzke [24] following a different chain of discoveries, relating to string and gauge theories. As a net outcome of a surprising series of mathematical connections, also bringing in the AdS/CFT correspondence, gluon scattering amplitudes in $\mathcal{N} = 4$ Super Yang-Mills can be now studied using powerful tools from the theory of integrable models [25].

A further important step for the understanding of the relationship between the AdS/CFT-related results and the original formulation of the ODE/IM correspondence was made by Lukyanov and Zamolodchikov in [26]. They showed how to link the classical sinh-Gordon equation to the quantum massive sine(h)-Gordon model, generalizing the earlier works on the ODE/IM correspondence. In doing this, they brought into play all the formalism developed in more than fifty years of soliton theory and the study of exactly-solvable nonlinear partial differential equations (of which sinh-Gordon is an exemplar), in particular the role played by the association between a nonlinear classical field theory and certain sets of linear differential equations.

In this paper we shall discuss another example of an off-critical generalization of the ODE/IM correspondence, which will be built starting from the definition of a particular model on the classical side. This model is often referred to as the Bullough-Dodd model [27], though the special properties of the corresponding partial differential equation were noticed as long ago as 1907, by Tzitzéica [28]. Like the sinh-Gordon model, it is an example of an affine Toda field theory, or 2d

Toda chain, as will be explained in §2. The quantum version of the Bullough-Dodd model, also known as the Izergin-Korepin model [29], plays an important role in framework of massive 2D quantum field theories corresponding to integrable perturbations of the minimal series of conformal field theories [30]. Together with its purely-imaginary coupling version it has been directly studied using the exact S-matrix approach [31]. It is also related to the scaling limit of important statistical-mechanical systems such as the q-state Potts models [32] and the dilute A_n models [33].

The analysis in this paper follows the steps taken in [11] and [26]. Some further details of the Bullough-Dodd model and the ODE/IM correspondence can be found in [34], while a more comprehensive presentation encompassing the extension to other affine Toda field theories will be the subject of a forthcoming publication.

2 The Bullough-Dodd model

The one dimensional Toda chain is defined as:

$$\partial_t^2 \eta^k = 2e^{(2\eta^{k+1} - 2\eta^k)} - 2e^{(2\eta^k - 2\eta^{k-1})}, \quad (2.1)$$

where k enumerates and labels the fields. This definition can be simply modified to yield the $(1 + 1)$ -dimensional variant of (2.1) [35]:

$$\partial_t^2 \eta^k - \partial_x^2 \eta^k = 2e^{(2\eta^{k+1} - 2\eta^k)} - 2e^{(2\eta^k - 2\eta^{k-1})}. \quad (2.2)$$

Varying the periodicity conditions and symmetries of the fields η^k , the system (2.2) leads to a whole family of non linear equations which are solvable by means of the Inverse Scattering Transform (see for example [36]). This involves an associated linear system, which has to reproduce the nonlinear chain through a compatibility condition. It turns out that the correct choice for (2.2) is [35]:

$$\begin{cases} X\Psi \equiv (\partial_t + V - i\lambda C_1 - i\lambda^{-1}C_2)\Psi & = 0 \\ T\Psi \equiv (\partial_x + W + i\lambda C_1 - i\lambda^{-1}C_2)\Psi & = 0 \end{cases} \quad (2.3)$$

where λ is the *spectral parameter*,

$$\begin{aligned} V_{ij} &= \partial_t \eta_i \delta_{ij}; \quad W_{ij} = \partial_x \eta_i \delta_{ij}; \quad C_{1ij} = C_{2ji} = C_j \delta_{i-1,j}, \\ C_i &= e^{(\eta^{i+1} - \eta^i)} \end{aligned} \quad (2.4)$$

and δ_{ij} is the Kronecker symbol which, in a case of a closed chain consisting of N elements, is defined as:

$$\delta_{ij} = \begin{cases} 1, & \text{when } i \equiv j \pmod{N}; \\ 0, & \text{in all other cases.} \end{cases} \quad (2.5)$$

It is also possible to give a Lagrangian formulation for the system (2.2):

$$\mathcal{L} = \sum_{k=1}^N \left(\frac{1}{2} \partial_\mu \eta^k \partial^\mu \eta^k - e^{(2\eta^k - 2\eta^{k-1})} + 1 \right) \quad (2.6)$$

with $\mu \in (t, x)$. Then it is straightforward to verify that:

$$\partial_\mu \frac{\partial \mathcal{L}}{\partial (\partial_\mu \eta^k)} - \frac{\partial \mathcal{L}}{\partial \eta^k} = 0 \implies \partial_t^2 \eta^k - \partial_x^2 \eta^k = 2e^{(2\eta^{k+1} - 2\eta^k)} - 2e^{(2\eta^k - 2\eta^{k-1})}.$$

Typically the further condition $\sum_k \eta^k = 0$ is imposed on the fields*, which results in the affine Toda field theory based on the $a_{N-1}^{(1)}$ affine Dynkin diagram. For $N = 2$, with $\eta^1 = -\eta^2 = \eta$, the equations then reduce to

$$\eta_{tt} - \eta_{xx} + 4 \sinh(2\eta) = 0 \quad \text{(the sinh-Gordon equation)}. \quad (2.7)$$

For $N = 3$, and with $\eta^3 = -\eta^1 - \eta^2$,

$$\begin{aligned} \eta_{tt}^1 - \eta_{xx}^1 + 2e^{(2\eta^1 - 2\eta^2)} - 2e^{(-4\eta^1 - 2\eta^2)} &= 0, \\ \eta_{tt}^2 - \eta_{xx}^2 + 2e^{(4\eta^1 + 2\eta^2)} - 2e^{(2\eta^1 - 2\eta^2)} &= 0, \end{aligned} \quad (2.8)$$

which are the equations of motion for the $a_2^{(1)}$ affine Toda field theory. The \mathbb{Z}_2 symmetry of the $a_2^{(1)}$ affine Dynkin diagram allows a further symmetry to be imposed, namely $\eta^1 = -\eta^2 = \eta$. Then (2.8) becomes

$$\eta_{tt} - \eta_{xx} + 2e^{4\eta} - 2e^{-2\eta} = 0 \quad \text{(the Bullough-Dodd equation)}. \quad (2.9)$$

Equation (2.9) can also be recovered directly from the compatibility of 3×3 matrix operators which are defined as [35]

$$T = \partial_x + \begin{pmatrix} \eta_t & -i\lambda^{-1}e^{-\eta} & i\lambda e^{2\eta} \\ i\lambda e^{-\eta} & 0 & -i\lambda^{-1}e^{-\eta} \\ -i\lambda^{-1}e^{2\eta} & i\lambda e^{-\eta} & -\eta_t \end{pmatrix} \quad (2.10)$$

$$X = \partial_t + \begin{pmatrix} \eta_x & i\lambda^{-1}e^{-\eta} & i\lambda e^{2\eta} \\ i\lambda e^{-\eta} & 0 & i\lambda^{-1}e^{-\eta} \\ i\lambda^{-1}e^{2\eta} & i\lambda e^{-\eta} & -\eta_x \end{pmatrix}. \quad (2.11)$$

If we were to introduce light-cone coordinates as

$$z = x + t \quad \bar{z} = x - t, \quad (2.12)$$

*This condition has the effect of excluding a zero mode in the ‘‘mass spectrum’’ of the theory:

$$m_n^2 = 4 \sin^2(\pi n/N), \quad n = 0, 1, \dots, N-1.$$

and redefine $\eta \rightarrow \eta/2$, (2.9) would become

$$\partial_z \partial_{\bar{z}} \eta(z, \bar{z}) + e^{-\eta(z, \bar{z})} - e^{2\eta(z, \bar{z})} = 0. \quad (2.13)$$

The nonlinear equation (2.13) is the starting point for the construction of objects which are in correspondence with the quantum world. However it is important to stress that, for this purpose, the coordinates z and \bar{z} should be considered as living in an ‘auxillary space’ which is *not* the same as the space on which the quantum field theory under discussion will be defined. In fact it will be better to adopt a Euclidean metric on this auxillary space so that z and \bar{z} are complex conjugates of each other, though this will often be relaxed and z and \bar{z} regarded as independent complex variables. To pave the way to the correspondence, it is also convenient to adopt a modified version of (2.13). This will be referred to as the modified Bullough-Dodd (mBD) equation and is

$$\partial_z \partial_{\bar{z}} \eta + e^{-\eta(z, \bar{z})} - p(z, s^{3M}) p(\bar{z}, s^{3M}) e^{2\eta(z, \bar{z})} = 0, \quad (\text{the mBD equation}) \quad (2.14)$$

where $M > 0$ and the function $p(z, s^{3M})$ is defined to be

$$p(z, s^{3M}) = z^{3M} - s^{3M}, \quad (2.15)$$

and s will turn out to take the role of a scale parameter. Equations (2.13) and (2.14) are related by a change of variables and a redefinition of the field, since, defining

$$w(z) = \int^z \sqrt{p(z')} dz' \quad \text{and} \quad \bar{w}(\bar{z}) = \int^{\bar{z}} \sqrt{p(\bar{z}')} d\bar{z}', \quad (2.16)$$

and shifting the field as $\eta \rightarrow \eta + \frac{1}{3} \ln(p(z)p(\bar{z}))$, the mBD equation reduces to the Bullough-Dodd equation. The mBD model has in general no explicit rotational symmetry. Instead it has a discrete symmetry dictated by the form of the ‘potential’ function $p(z)$:

$$z \rightarrow e^{\frac{2\pi i}{3M}} z, \quad \bar{z} \rightarrow e^{-\frac{2\pi i}{3M}} \bar{z}. \quad (2.17)$$

The solutions of the mBD equation (2.14) which will be relevant for the current analysis must respect this symmetry, be continuous at every finite nonzero z , \bar{z} , and grow more slowly than exponentially at $z, \bar{z} \rightarrow \infty$. As in [26] for the (modified) sinh-Gordon case, we will consider a one-parameter family of such solutions, characterized by a logarithmic behaviour near the origin. Before listing in detail the exact conditions which are required, polar coordinates can be introduced to show these properties more neatly:

$$z = \rho e^{i\phi}, \quad \bar{z} = \rho e^{-i\phi}. \quad (2.18)$$

In most the following z and \bar{z} will be treated as independent complex variables, so the solution η will be a function of the independent variables (ρ, ϕ) , according to (2.18).

It is now possible to list the properties of the sought-after solutions:

(i) periodicity:

$$\eta(\rho, \phi + \frac{2\pi}{3M}) = \eta(\rho, \phi); \quad (2.19)$$

or, better, the solutions $\eta(\rho, \phi)$ are single-valued functions on a cone with the apex angle $\frac{2\pi}{3M}$,

$$\mathcal{C}_{\frac{2\pi}{3M}} : \quad \phi \sim \phi + \frac{2\pi}{3M}, \quad 0 \leq \rho < \infty; \quad (2.20)$$

(ii) the solutions $\eta(\rho, \phi)$ are real-valued for real ρ and ϕ , and finite everywhere on the cone $\mathcal{C}_{\frac{2\pi}{3M}}$, except for the apex $\rho = 0$;

(iii) large- ρ asymptotic:

$$\eta(\rho, \phi) = -2M \ln \rho + o(1) \quad \text{as } \rho \rightarrow \infty; \quad (2.21)$$

(iv) small- ρ asymptotic:

$$\eta(\rho, \phi) = -2g \ln(\rho) + O(1) \quad \text{for } -1 < g < 1/2. \quad (2.22)$$

The factors $-2M$ and $-2g$ in (2.21) and (2.22) have been chosen, for future convenience, to be consistent with [11] and [26].

As in [26], one can start from equation (2.22) and develop an expansion for $z, \bar{z} \sim 0$ of the form

$$\begin{aligned} \eta = & -g \ln(z\bar{z}) + \eta_0 + \sum_{k=1}^{\infty} \gamma_k (z^{3Mk} + \bar{z}^{3Mk}) \\ & - \frac{e^{-\eta_0}}{(g+1)^2} (z\bar{z})^{g+1} - \frac{s^{6M} e^{2\eta_0}}{(-2g+1)^2} (z\bar{z})^{-2g+1} + \dots \end{aligned} \quad (2.23)$$

where η_0, γ_k are integration constants, all remaining terms omitted in the expansion being uniquely determined once these constants are given. These constants are not new parameters, but should be fixed by demanding the consistency of this expansion with the remaining conditions (i)-(iii) above.

The expansion (2.23) remains valid with z and \bar{z} regarded as independent complex variables and, for later analysis, it will be useful to record the form of (2.22) in the so-called *light-cone* limit $\bar{z} \rightarrow 0$ (with fixed z):

$$\eta \sim -g \ln(z\bar{z}) + \eta_0 + \gamma(z) \quad (2.24)$$

where $\gamma(z) = \sum_{k=1}^{\infty} \gamma_k z^{3Mk}$.

At this stage it is possible to define the linear problem associated to the mBD equation

$$\mathbf{D}\Psi = 0, \quad \bar{\mathbf{D}}\Psi = 0, \quad (2.25)$$

where \mathbf{D} and $\bar{\mathbf{D}}$ are components of an $sl(3)$ connection and can be found by rearranging, in light-cone coordinates, the matrix operators (2.10) with an appropriate redefinition of the spectral parameter λ and after the introduction of the potential $p(z)$:

$$\mathbf{D} = \partial_z + \begin{pmatrix} \frac{1}{2}\partial_z\eta & 0 & \lambda e^\eta p(z) \\ -\lambda e^{-\frac{1}{2}\eta} & 0 & 0 \\ 0 & -\lambda e^{-\frac{1}{2}\eta} & -\frac{1}{2}\partial_z\eta \end{pmatrix}, \quad (2.26)$$

$$\bar{\mathbf{D}} = \partial_{\bar{z}} + \begin{pmatrix} -\frac{1}{2}\partial_{\bar{z}}\eta & \lambda^{-1}e^{-\frac{1}{2}\eta} & 0 \\ 0 & 0 & \lambda^{-1}e^{-\frac{1}{2}\eta} \\ \lambda^{-1}p(\bar{z})e^\eta & 0 & \frac{1}{2}\partial_{\bar{z}}\eta \end{pmatrix}. \quad (2.27)$$

It is possible to deal directly with this system of six equations on Ψ to get information about the solutions and their asymptotics, but it turns out to be simpler to deal with a reduction of (2.26), (2.27) to two third-order differential equations. Moreover this step will be necessary for the continuation of the analysis and to show the connection between the original nonlinear problem and a previously-studied spectral problem. This reduction can be implemented defining a vector solution to equations (2.26), (2.27)[†]:

$$\Psi = \begin{pmatrix} \lambda^{-\frac{1}{2}}e^{\eta/2}\partial_z(e^\eta\partial_z(e^{-\eta}\psi)) \\ -\lambda^{\frac{1}{2}}e^\eta\partial_z(e^{-\eta}\psi) \\ \lambda^{\frac{3}{2}}e^{-\eta/2}\psi \end{pmatrix} = \begin{pmatrix} \lambda^{-\frac{3}{2}}e^{-\eta/2}\bar{\psi} \\ -\lambda^{-\frac{1}{2}}e^\eta\partial_{\bar{z}}(e^{-\eta}\bar{\psi}) \\ \lambda^{\frac{1}{2}}e^{\eta/2}\partial_{\bar{z}}(e^{-\eta}\bar{\psi}) \end{pmatrix}. \quad (2.28)$$

The equality between each row of the two parentheses in (2.28) must always hold, and will be useful to find large- ρ asymptotics of (2.25). Applying \mathbf{D} and $\bar{\mathbf{D}}$ to the first and the second vectors in (2.28) respectively, it is simple to find the system of two third-order linear differential equations which constitutes a compatibility condition on ψ and $\bar{\psi}$ so that Ψ really is a solution of (2.25). The two equations are:

$$\partial_z^3\psi - ((\partial_z\eta)^2 + 2\partial_z^2\eta)\partial_z\psi + (\lambda^3 p(z) - \partial_z\eta\partial_z^2\eta - \partial_z^3\eta)\psi = 0; \quad (2.29)$$

$$\partial_{\bar{z}}^3\bar{\psi} - ((\partial_{\bar{z}}\eta)^2 + 2\partial_{\bar{z}}^2\eta)\partial_{\bar{z}}\bar{\psi} + (\lambda^{-3}p(\bar{z}) - \partial_{\bar{z}}\eta\partial_{\bar{z}}^2\eta - \partial_{\bar{z}}^3\eta)\bar{\psi} = 0. \quad (2.30)$$

The next step is to determine the $\rho \rightarrow 0$ asymptotics of solutions to (2.25). To do this it is convenient to focus attention on eq. (2.29), though the result would not change if eq. (2.30) was taken as the starting point instead. This equation is entirely in z , except for the appearance of \bar{z} in $\eta(z, \bar{z})$ that, for now, can be considered as a simple parametric dependence. Finding an asymptotic solution $\psi(z)$ enables a solution $\bar{\Psi}$ of (2.25) for $\rho \rightarrow 0$ to be determined through (2.28).

[†]This is simply recovered solving each of the systems (2.25) respect to two out of three components of the vector Ψ .

Substituting the asymptotic form (2.22) of η into (2.29) and considering the $\rho \rightarrow 0$ limit (actually $z \rightarrow 0$) brings equation (2.29) into the form

$$\partial_z^3 \psi - \frac{1}{z^2} g(g+2) \partial_z \psi + \frac{1}{z^3} g(g+2) \psi = 0,$$

and seeking solutions of the form $\psi = z^\mu + \dots$ the problem is reduced to the solution of the following indicial equation:

$$(\mu - 1)[\mu(\mu - 2) - g(g + 2)] = 0.$$

This equation gives three different leading behaviours in the $z \rightarrow 0$ limit:

$$\psi_+ \sim z^{-g}, \quad \psi_0 \sim z, \quad \psi_- \sim z^{g+2}. \quad (2.31)$$

These expressions give the leading behaviours and can be adjusted with appropriate multiplicative constants; once inserted into (2.28) in the usual $\rho \rightarrow 0$ limit, they provide the three asymptotic solutions Ψ to (2.25). They are:

$$\Psi_+ \sim \begin{pmatrix} 0 \\ 0 \\ e^{(-i\phi-\theta)g} \end{pmatrix}, \quad \Psi_0 \sim \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \Psi_- \sim \begin{pmatrix} e^{(i\phi+\theta)g} \\ 0 \\ 0 \end{pmatrix} \quad (\rho \rightarrow 0) \quad (2.32)$$

with $\lambda = e^\theta$. In order to fix the values of the constants introduced above a particular symmetry condition has been considered. In fact, although $\eta(\rho, \phi)$ is a single-valued function on the cone (2.20), the connection components (2.26) and (2.27) are not. Instead, the linear problem (2.25) is invariant with respect to the transformation

$$\Omega : \quad \phi \rightarrow \phi + \frac{2\pi}{3M}, \quad \theta \rightarrow \theta - \frac{2\pi i}{3M}, \quad (2.33)$$

involving the shift of the spectral parameter θ (from now on θ and λ will be referred as spectral parameters interchangeably, keeping in mind that $\lambda = e^\theta$). Imposing this symmetry, it is straightforward to find (2.32).

On the other hand it is also possible to give the large- ρ asymptotics, and for this it is best to use a semiclassical approximation. As will be explained more formally below, assuming that θ is real, a generic solution of (2.29) grows exponentially at $\rho \rightarrow \infty$, but [2, 11] there are special ('subdominant') solutions that decay in the sector

$$-\frac{4\pi}{3M+3} < \phi < \frac{4\pi}{3M+3}. \quad (2.34)$$

In order to build such a solution to (2.25), the first step is the specification of a $\rho \rightarrow \infty$ asymptotic for (2.29), just as done before for the solution in the region near $\rho = 0$, again remembering to substitute the large- ρ solution of (2.14). In this particular limit equation (2.29) reduces to a sum of a third order derivative

and a ‘potential’ $Q(z)$ which will fix the leading term of the subdominant WKB solution:

$$Q(z) = e^{3\theta}(z^{3M} - s^{3M}). \quad (2.35)$$

It follows from (2.35) that the WKB-like solution has, for $M > 1/2$, the form:

$$\psi \sim c_1 z^{-M} \exp\left(-\frac{z^{M+1}}{M+1} e^\theta + f(\bar{z})\right), \quad (2.36)$$

where c_1 is an arbitrary constant and the term $f(\bar{z})$ in the exponential carries the dependence on \bar{z} of ψ . To retrieve a more exact form of (2.36), it is convenient to seek a solution of (2.30) of the form

$$\bar{\psi} \sim c_2 \bar{z}^{-M} \exp\left(-\frac{\bar{z}^{M+1}}{M+1} e^{-\theta} + g(z)\right) \quad (2.37)$$

and then by inserting both the results in (2.28) the compatibility between the two solutions will fix these integration-generated functions. This yields:

$$c_2 = c_1 e^\theta = c e^\theta, \quad f(\bar{z}) = -\frac{\bar{z}^{M+1}}{M+1} e^{-\theta}, \quad g(z) = -\frac{z^{M+1}}{M+1} e^\theta, \quad (2.38)$$

and the final semiclassical solution of (2.29) (in polar coordinates) is

$$\psi \stackrel{\text{WKB}}{\sim} c e^{-\theta} z^{-M} \exp\left(-\frac{2\rho^{M+1}}{M+1} \cosh(\theta + i(M+1)\phi)\right). \quad (2.39)$$

By inserting (2.39) into the first of (2.28) it is then straightforward to obtain

$$\Psi \stackrel{\text{WKB}}{\sim} e^{\frac{\theta}{2}} \begin{pmatrix} e^{i\phi M} \\ 1 \\ e^{-i\phi M} \end{pmatrix} \exp\left(-\frac{2\rho^{M+1}}{M+1} \cosh(\theta + i(M+1)\phi)\right), \quad \rho \rightarrow \infty. \quad (2.40)$$

Since the functions $\{\Psi_+, \Psi_0, \Psi_-\}$ form a basis in the space of solutions of linear problem (2.25), a linear relation of the following form must hold:

$$\Psi = Q^+(\theta, s)\Psi_+ + Q^0(\theta, s)\Psi_0 + Q^-(\theta, s)\Psi_-, \quad (2.41)$$

where the coefficients $Q^{\pm,0}$, which are independent of the variables ρ and ϕ , are functions of the spectral parameter θ and of s , as well as of the parameter g (this parameter has been temporarily omitted). These coefficients will coincide (up to an overall constant) with Q -functions of a 2d massive quantum field theory related to the Lie algebra A_2 , or $su(3)$.

3 The conformal limit

Before proceeding with the advertised correspondence, it is useful to work a little more on equation (2.29). The problem studied up to now has an important connection with a known spectral problem of a particular third-order differential equation. In order to build an appropriate analysis and to understand where the various steps come from, (2.29) will be reduced to this spectral problem and a brief review of previously-obtained results will be given.

Recall that equation (2.29) is of the form

$$\partial_z^3 \psi - ((\partial_z \eta)^2 + 2\partial_z^2 \eta) \partial_z \psi + (\lambda^3 p(z, s^{3M}) - \partial_z \eta \partial_z^2 \eta - \partial_z^3 \eta) \psi = 0,$$

and, as was said above, \bar{z} plays the role of a simple parameter, so it is possible to take the so-called light-cone limit $\bar{z} \rightarrow 0$ (or, if (2.30) had been chosen as starting point then the light-cone limit would have involved z instead of \bar{z}) in which η assumes the form (2.24). After that, the limit $z \sim s \rightarrow 0$, $\theta \rightarrow +\infty$ can be taken, with the combinations

$$x = e^{\frac{\theta}{M+1}} z, \quad E = s^{3M} e^{\frac{3\theta M}{M+1}} \quad (3.42)$$

kept finite while

$$\tilde{x} = e^{-\frac{\theta}{M+1}} \bar{z} \rightarrow 0, \quad \tilde{E} = s^{3M} e^{-\frac{3\theta M}{M+1}} \rightarrow 0. \quad (3.43)$$

This limit is a particular scaling limit because z , s are sent to zero and θ to infinity, but they are rearranged in combinations such that the new ‘variables’ x and E do not diverge or collapse. This is an important turning point of the theory, since it transforms the correspondence between an ODE and a massive QFT into a correspondence with a conformal field theory [11] (which is massless). For this reason the limit (3.43) can be called the *conformal limit*, with s (or, more precisely, s^{M+1}) taking the role of the mass scale.

Inserting (3.42) in (2.29) and taking the limits just discussed, equation (2.29) reduces to

$$\left[\partial_x^3 - g(g+2) \left(\frac{1}{x^2} \partial_x - \frac{1}{x^3} \right) + p(x, E) \right] y(x, E, g) = 0, \quad (3.44)$$

where y has been used instead of ψ to avoid confusion among the various solutions. Motivated by the results of [11] and [2], a generalization of a theorem due to Sibuya for second-order equations can now be proposed:

Conjecture 3.1 *Equation (3.44) has a solution $y(x, E, g)$ with the following properties:*

- (i) *y is an entire function of (x, E) though, due to the branch point in the potential at $x = 0$, x must in general be considered to live on a suitable cover of the punctured complex plane;*

(ii) $y, y' = \partial_x y$ and $y'' = \partial_x^2 y$ admit, for $M > 1/2$, the asymptotic representations

$$y \sim x^{-M} e^{-\frac{1}{M+1}x^{M+1}}, \quad y' \sim -e^{-\frac{1}{M+1}x^{M+1}}, \quad y'' \sim x^{-M} e^{-\frac{1}{M+1}x^{M+1}}, \quad (3.45)$$

as $x \rightarrow \infty$ in the sector

$$|\arg(x)| < \frac{4\pi}{3M+3}; \quad (3.46)$$

(iii) y is uniquely fixed by the properties i) and ii).

The behaviour of the solution (3.45) follows from the more general formula

$$y(x, E) \sim Q(x, E)^{-1/3} \exp\left(-\int_{x_0}^x Q(x, t)^{1/3} dt\right), \quad (3.47)$$

where y solves a generic equation $\partial_x^3 y + Q(x, E)y = 0$. This is the analogue of a WKB approximation for a solution of a Schrödinger equation. It is possible to define rotated solutions, by analogy with [13], in order to construct bases of solutions for (3.42). For general values of k , define

$$y_k(x, E, g) = \omega^k y(\omega^{-k}x, \omega^{-3Mk}E, g), \quad (3.48)$$

with

$$\omega = \exp\left(\frac{2\pi i}{3M+3}\right). \quad (3.49)$$

Substituting in, y_k solves

$$\partial_x^3 y_k - g(g+2)\left(\frac{1}{x^2}\partial_x - \frac{1}{x^3}\right)y_k + e^{-2k\pi i}p(x, E)y_k = 0, \quad (3.50)$$

and so for $k \in \mathbb{Z}$, (3.48) provides a potentially-new solution to (3.42). However for now it is convenient to leave k arbitrary since fractional values will also be needed. We also define the *Stokes sectors*

$$\mathcal{S}_k : \quad \left| \arg(x) - \frac{2\pi k}{3M+3} \right| < \frac{\pi}{3M+3}. \quad (3.51)$$

Some discussion of the dominance and subdominance of solutions is also required, since, for a third-order problem, this is a little more complicated than the second-order case. For this we largely follow [13]. First, the behaviour specified by conjecture 3.1 lies in the sectors $\mathcal{S}_{-\frac{3}{2}} \cup \mathcal{S}_{-\frac{1}{2}} \cup \mathcal{S}_{\frac{1}{2}} \cup \mathcal{S}_{\frac{3}{2}}$. Furthermore, in general, there are three types of asymptotic solution for large $|x|$. Aside the well-behaved leading term $x^{-M} \exp(-x^{M+1}/(M+1))$, there are also solutions that behave as $x^{-M} \exp(e^{\pm\pi i/3}x^{M+1}/(M+1))$. (This can be traced to the three third roots of -1 , which are $-1, e^{\pi i/3}$ and $e^{-\pi i/3}$). According to the sector is being considered, either

one or two of these solutions tend to zero for large $|x|$. The *subdominant* solution in any given sector will be defined to be the one that tends to zero fastest in that sector (it might also be called maximally subdominant). Finally, y_k , up to overall scalar factor, is characterized as the solution of (3.42) which is subdominant in the sector \mathcal{S}_k . The asymptotic (3.45) along with the definition (3.48) imply

$$\begin{aligned} y_k &\sim \omega^{k(M+1)} x^{-M} e^{-\frac{x^{M+1}}{M+1} \omega^{-(M+1)k}}, & y'_k &\sim -e^{-\frac{x^{M+1}}{M+1} \omega^{-(M+1)k}}, \\ y''_k &\sim \omega^{-k(M+1)} x^M e^{-\frac{x^{M+1}}{M+1} \omega^{-(M+1)k}}, \end{aligned} \quad (3.52)$$

for $|x| \rightarrow \infty$ with

$$x \in \mathcal{S}_{k-\frac{3}{2}} \cup \mathcal{S}_{k-\frac{1}{2}} \cup \mathcal{S}_{k+\frac{1}{2}} \cup \mathcal{S}_{k+\frac{3}{2}}. \quad (3.53)$$

In the region formed by the two sectors $\mathcal{S}_{k+\frac{1}{2}} \cup \mathcal{S}_{k+\frac{3}{2}}$, the three asymptotics y_k , y_{k+1} and y_{k+2} live together and comparing their forms, given by (3.52), they turn out to be independent of each other. A further proof of this fact can be built defining the *Wronskian* of the three solutions:

$$W_{k_1, k_2, k_3} = W[y_{k_1}, y_{k_2}, y_{k_3}] \quad (3.54)$$

where the 3×3 Wronskian is defined as

$$W[f, g, h] = \det \begin{pmatrix} f & f' & f'' \\ g & g' & g'' \\ h & h' & h'' \end{pmatrix}. \quad (3.55)$$

For f , g and h solving (3.42), $W[f, g, h]$ is independent of x , and f , g and h are linearly independent if and only if $W[f, g, h] \neq 0$. In order to demonstrate the independence of y_k , y_{k+1} and y_{k+2} , it is helpful first to compute $(k_1, k_2, k_3) = (-1, 0, 1)$

$$W_{-1, 0, 1} = -3i\sqrt{3} \quad (3.56)$$

and then, using the general result

$$W_{k_1+b, k_2+b, k_3+b}(E) = W_{k_1, k_2, k_3}(\omega^{-3Mb} E), \quad (3.57)$$

we see that $W_{k, k+1, k+2}$ is always nonzero, thus confirming the independence of $\{y_k, y_{k+1}, y_{k+2}\}$.

The final step for this section is to show how to build new solutions to (3.42) using the y_k 's. Supposing that k_1 and k_2 differ by an integer so that $e^{-2k_1\pi i} = e^{-2k_2\pi i} = e^{-2k\pi i}$, it can be checked that the function

$$z_{k_1, k_2}(x, E, g) = y_{k_1} y'_{k_2} - y_{k_2} y'_{k_1}, \quad (3.58)$$

which is actually a 2×2 Wronskian, provides a solution of

$$\partial_x^3 z_{k_1, k_2} - g(g+2) \left(\frac{1}{x^2} \partial_x - \frac{1}{x^3} \right) z_{k_1, k_2} - e^{-2k\pi i} (x^{3M} - E) z_{k_1, k_2}. \quad (3.59)$$

Equation (3.59) is the adjoint of (3.50) and if k is shifted by a half-integer, then z_{k_1, k_2} becomes a solution of the original equation (3.50)

$$\partial_x^3 z_{k_1 + \frac{1}{2}, k_2 + \frac{1}{2}} - g(g+2) \left(\frac{1}{x^2} \partial_x - \frac{1}{x^3} \right) z_{k_1 + \frac{1}{2}, k_2 + \frac{1}{2}} + e^{-2k\pi i} (x^{3M} - E) z_{k_1 + \frac{1}{2}, k_2 + \frac{1}{2}}. \quad (3.60)$$

For $|k_1 - k_2| < 3$, the regions (3.53) for $k_1 = k_2 = k$ overlap, and it is possible to get an asymptotic for z_{k_1, k_2} from (3.52). In particular, for $k = 1, 2, 3$ we have

$$z_{-k/2, k/2}(x, E, g) \sim 2i \sin(\pi k/3) x^{-M} e^{-2 \cos(\pi k/3) \frac{1}{M+1} x^{M+1}}, \quad x \rightarrow \infty. \quad (3.61)$$

Considering (3.59) and (3.61), for $k = 1$ it follows, by uniqueness of solutions, that

$$z_{-\frac{1}{2}, \frac{1}{2}}(x, E, g) = i\sqrt{3}y(x, E, g). \quad (3.62)$$

This result is limited to the ($k = 1$)-case, as for other values of the parameter is not possible to retrieve enough information to uniquely pin the function down. On the other hand (3.62) is useful to determine the *Bethe Ansatz equations*, but this will be shown in the next section, starting from the original problem (2.29). Now that the analyticity properties of solutions of (3.42) have been studied, it is possible to return to (2.29) and apply to that, with the appropriate caution, all the results that have been found in this section.

4 Bethe Ansatz equations

In this section, Bethe Ansatz equations will be built starting from certain functional relations involving the coefficients Q^\pm already introduced at the end of §2. In order to get there it is necessary to apply the same analysis described in §3 to reproduce similar results for the original equation (2.29), which is repeated below for convenience:

$$\partial_x^3 \psi - ((\partial_x \eta)^2 + 2\partial_x^2 \eta) \partial_x \psi + (p(x, E) - \partial_x \eta \partial_x^2 \eta - \partial_x^3 \eta) \psi = 0, \quad (4.63)$$

$$\partial_{\tilde{x}}^3 \bar{\psi} - ((\partial_{\tilde{x}} \eta)^2 + 2\partial_{\tilde{x}}^2 \eta) \partial_{\tilde{x}} \bar{\psi} + (p(\tilde{x}, \tilde{E}) - \partial_{\tilde{x}} \eta \partial_{\tilde{x}}^2 \eta - \partial_{\tilde{x}}^3 \eta) \bar{\psi} = 0, \quad (4.64)$$

where

$$x = ze^{\theta/(M+1)}, \quad \tilde{x} = \bar{z}e^{-\theta/(M+1)}, \quad E = s^{3M} e^{3\theta M/(M+1)}, \quad \tilde{E} = s^{3M} e^{-3\theta M/(M+1)}, \quad (4.65)$$

and

$$\psi \equiv \psi(x, \tilde{x}, E, \tilde{E}, g), \quad \bar{\psi} \equiv \bar{\psi}(x, \tilde{x}, E, \tilde{E}, g). \quad (4.66)$$

First of all, the rotated solutions, analogues to (3.48), have to be defined. In order to do this, the periodicity of $\eta(\rho, \phi)$ has to be exploited since it is the only

information, besides the asymptotic behaviours, which is known. It turns out that the right form is

$$\psi_k(x, \tilde{x}, E, \tilde{E}, g) = \omega^k \psi(\omega^{-k}x, \omega^k \tilde{x}, \omega^{-3Mk}E, \omega^{3Mk} \tilde{E}, g), \quad (4.67)$$

where

$$\omega = \exp\left(\frac{2\pi i}{3M+3}\right). \quad (4.68)$$

The function (4.67) solves

$$\partial_x^3 \psi_k - ((\partial_x \eta)^2 + 2\partial_x^2 \eta) \partial_x \psi_k + (e^{-2k\pi i} p(x, E) - \partial_x \eta \partial_x^2 \eta - \partial_x^3 \eta) \psi_k = 0, \quad (4.69)$$

where, analogously to (3.50), a $e^{-2k\pi i}$ -term appears. In general arbitrary integer or half-integer values for k will be considered, and the definition of Stokes sectors remains unchanged (see (3.51)). Following the discussion of §3, it is straightforward to find the $|x| \rightarrow \infty$ behaviour of the rotated solutions (4.67), (keeping in mind that \tilde{x} goes to $\omega^k \tilde{x}$) and setting

$$c = e^{\theta/(M+1)} \quad (4.70)$$

$$\begin{aligned} \psi_k &\sim \omega^{k(M+1)} x^{-M} \exp\left[-\frac{x^{M+1}}{M+1} \omega^{-k(M+1)} - \frac{\tilde{x}^{M+1}}{M+1} \omega^{k(M+1)}\right], \\ \psi'_k &\sim -\exp\left[-\frac{x^{M+1}}{M+1} \omega^{-k(M+1)} - \frac{\tilde{x}^{M+1}}{M+1} \omega^{k(M+1)}\right], \\ \psi''_k &\sim \omega^{-k(M+1)} x^M \exp\left[-\frac{x^{M+1}}{M+1} \omega^{-k(M+1)} - \frac{\tilde{x}^{M+1}}{M+1} \omega^{k(M+1)}\right]. \end{aligned} \quad (4.71)$$

(the derivatives are on x only.)

Now, the three generic solutions $\{\psi_k, \psi_{k+1}, \psi_{k+2}\}$ form a basis for the solutions of eq.(2.29). Again, the proof of this property comes from the evaluation of the Wronskian for $(k_1, k_2, k_3) = (-1, 0, 1)$

$$W_{-1,0,1} = W[\psi_{-1}, \psi_0, \psi_1] = -3i\sqrt{3}, \quad (4.72)$$

which shows that these three solutions are independent.

The next step is to define the analogues of the functions (3.58) that, here, will be denoted as u_{k_1, k_2} so as to avoid confusion with the variable z . They read as

$$u_{k_1, k_2}(x, \tilde{x}, E, \tilde{E}, g) = [\psi_{k_1} \psi'_{k_2} - \psi_{k_2} \psi'_{k_1}](x, \tilde{x}, E, \tilde{E}, g), \quad (4.73)$$

where, again, the \bar{z} -dependence has been omitted. The solutions (4.73) solve

$$\partial_x^3 u_{k_1, k_2} - ((\partial_x \eta)^2 + 2\partial_x^2 \eta) \partial_x u_{k_1, k_2} - \left(e^{-2\pi i k} p(x, E) + \partial_x \eta \partial_x^2 \eta + \partial_x^3 \eta\right) u_{k_1, k_2} = 0, \quad (4.74)$$

and, also in this case, it is possible to evaluate (4.73) for $k_1 = -k_2 = -\frac{k}{2}$

$$u_{-k/2,k/2} \sim 2i \sin\left(\frac{\pi}{3}k\right)x^{-M} \exp\left[-2 \cos\left(\frac{\pi}{3}k\right)\left(\frac{x^{M+1}}{M+1} + \frac{\tilde{x}^{M+1}}{M+1}\right)\right], \quad (4.75)$$

which, when compared with (4.71) for $k = 1$, gives the identification

$$u_{-1/2,1/2} = i\sqrt{3}\psi. \quad (4.76)$$

To get the Bethe Ansatz equations (BAE) eq.(4.76) is used, exploiting the relation

$$u_{-1/2,1/2} = \psi_{-\frac{1}{2}}\psi'_{\frac{1}{2}} - \psi_{\frac{1}{2}}\psi'_{-\frac{1}{2}} = i\sqrt{3}\psi_0. \quad (4.77)$$

At this point the solutions ψ_k can be expressed in the basis of the $|z| \rightarrow 0$ solutions (2.31) with the appropriate normalization to capture the correct behaviour in the conformal limit

$$\psi_k = Q_k^+(E, \tilde{E})\chi_k^+ + Q_k^0(E, \tilde{E})\chi_k^0 + Q_k^-(E, \tilde{E})\chi_k^-, \quad (4.78)$$

where

$$Q_k^{\pm,0}(E, \tilde{E}) = Q^{\pm,0}(\omega^{-k3M}E, \omega^{k3M}\tilde{E}), \quad (4.79)$$

and

$$\begin{cases} \chi_k^+ \sim \omega^{k(g+1)}x^{-g} \\ \chi_k^0 \sim x \\ \chi_k^- \sim \omega^{-k(g+1)}x^{g+2}. \end{cases} \quad (4.80)$$

The functions $Q^{\pm,0}(E, \tilde{E})$, introduced in (4.78), are the massive off-critical analogue of the CFT-related functions $Q^{\pm,0}(E)$ [13], [8]. Inserting the expression (4.78) into (4.77) and considering, at first, the terms proportional to x^{-g} , a functional relation containing Q^+ and Q^0 is obtained

$$\begin{aligned} i\sqrt{3}Q_0^+ &= Q_{-1/2}^+Q_{1/2}^0\omega^{-(g+1)/2} - gQ_{1/2}^+Q_{-1/2}^0\omega^{(g+1)/2} + \\ &- Q_{1/2}^+Q_{-1/2}^0\omega^{(g+1)/2} + gQ_{-1/2}^+Q_{1/2}^0\omega^{-(g+1)/2} \\ &= (g+1)\left(Q_{-1/2}^+Q_{1/2}^0\omega^{-(g+1)/2} - Q_{1/2}^+Q_{-1/2}^0\omega^{(g+1)/2}\right) \end{aligned} \quad (4.81)$$

with the appropriate ω rotation factors. Equation (4.81) can be written in terms of the variable $\theta = (M+1)/(3M) \ln E$ as:

$$i\sqrt{3}Q^+(\theta) = (g+1)\left(Q^+(\theta + i\frac{\pi}{3})Q^0(\theta - i\frac{\pi}{3})\omega^{-\frac{g+1}{2}} - \omega^{\frac{g+1}{2}}Q^+(\theta - i\frac{\pi}{3})Q^0(\theta + i\frac{\pi}{3})\right) \quad (4.82)$$

(with s and g kept constant.)

Considering (4.82) evaluated at $\theta = \theta_n + i\pi/3$ and $\theta = \theta_n - i\pi/3$ with $Q^+(\theta_n) = 0$ to get

$$i\sqrt{3}Q^+(\theta_n + i\pi/3) = (g+1)\left(Q^+(\theta_n + i2\pi/3)Q^0(\theta_n)\omega^{-\frac{g+1}{2}}\right) \quad (4.83)$$

and

$$i\sqrt{3}Q^+(\theta_n - i\pi/3) = -(g+1) \left(\omega^{\frac{g+1}{2}} Q^+(\theta_n - i\frac{2\pi}{3}) Q^0(\theta_n) \right) \quad (4.84)$$

respectively, and taking the ratio, the sought-after Bethe Ansatz equations are obtained [11]

$$\frac{Q^+(\theta_n + i2\pi/3) Q^+(\theta_n - i\pi/3)}{Q^+(\theta_n - i2\pi/3) Q^+(\theta_n + i\pi/3)} = -\omega^{(g+1)}. \quad (4.85)$$

The same method can be used to find BAE for the Q^- functions. In conclusion: by defining θ^\pm the zeroes relative to Q^\pm , the Bethe Ansatz equations for the two spectral functions are:

$$\frac{Q^\pm(\theta_n^\pm + i2\pi/3) Q^\pm(\theta_n^\pm - i\pi/3)}{Q^\pm(\theta_n^\pm - i2\pi/3) Q^\pm(\theta_n^\pm + i\pi/3)} = -\omega^{\pm(g+1)}. \quad (4.86)$$

The Bethe Ansatz equations (4.86) can be transformed into the nonlinear integral equation given in section 6 of [11] for the ground state energy of the Izergin-Korepin model or, equivalently, of the scaling q -state Potts and tricritical Potts models on a cylinder geometry. This family of systems plays a key role in the study of integrable quantum field theories in 1+1 dimensions and statistical mechanical models in 2 dimensions, and is related to the $\phi_{1,2}$, $\phi_{2,1}$ and $\phi_{1,5}$ integrable deformations of minimal conformal field theories. The Ising model in external magnetic field, with its E_8 -related mass spectrum [37], is perhaps the most physically-interesting system in the family. A schematic dictionary of the correspondence between the classical objects introduced here and the ‘quantum field theory world’ is given in Table 1.

classical	modified Bullough-Dodd		QFT on a cylinder
	θ	\Leftrightarrow	particle rapidity
	s^{M+1}	\Leftrightarrow	Rm (R : circumference, m : mass gap)
	g	\Leftrightarrow	twist parameter
	M	\Leftrightarrow	‘ q ’ in the q -state Potts model


Table 1: The dictionary

5 Conclusions

The main objective of this article was to discuss a particular generalization of the so-called ODE/IM correspondence to the massive case. The steps taken by Lukyanov and Zamolodchikov in [26] clarified how massive generalisations of the ODE/IM correspondence should be built. The results of [26] concern the relation between the classical and the quantum versions of the sinh-Gordon model. Adapting their discussion it has been possible, here, to relate the classical (Titzeica-)

Bullough-Dodd field equation to the Izergin-Korepin massive quantum field theory. The discussion parallels precisely the conformal case treated in [11] and links a certain linear problem to a Bethe Ansatz system associated with the a_2 algebra. The Bethe Ansatz equations (4.86) and the demonstration of a link with the conformal field theory limit case of [11], described in section 3, are the main results of the paper.

After the first discovery of this type of correspondence, it was striking how certain functional relations, typically emerging in the QFT domain, encoded spectral data. In the work of Lukyanov and Zamolodchikov, this rich structure was enlarged to contain also the relation between certain nonlinear partial differential equations and linear spectral problems, thereby giving further useful insights into the general structure of the theory. Here, further support of the validity of this scheme has been given, enlarging the number of working cases of the correspondence. The Bullough-Dodd equation has been chosen since, with the sinh-Gordon equation, it is simplest representative of the affine Toda field theories. At this stage it is fairly clear how a more general correspondence scheme could be developed starting from Toda field theories based on more general Lie algebras. The general scheme is as follows, where the arrows have been annotated with labels of sections in this paper and a reference, to indicate where the corresponding step is further discussed for the particular case of the Bullough-Dodd equation:



Another direction for future work is the generalization of the correspondence between the Bethe Ansatz and classical integrable systems to non relativistically-invariant models such as the KdV equation and its hierarchy, and the generalization from integrable field theories to integrable lattice models.

In conclusion, the connection between these two, originally disconnected, domains of mathematics and theoretical physics gives a hint of a bigger scheme in the wide framework of classical and quantum integrability.

Acknowledgements –

This project was partially supported by an INFN grant PI11, an STFC rolling grant ST/G000433/1 and by the Italian MIUR-PRIN contract 2009KHZKRX-007 “Symmetries of the Universe and of the Fundamental Interactions”.

References

- [1] P. Dorey and R. Tateo, ‘Anharmonic oscillators, the thermodynamic Bethe ansatz, and nonlinear integral equations’, *J. Phys. A* **32**, L419 (1999) [arXiv:hep-th/9812211].
- [2] Y. Sibuya, ‘Global theory of a second-order linear ordinary differential equation with polynomial coefficient’, Amsterdam: North-Holland, 1975.
- [3] A. Voros, ‘Semi-classical correspondence and exact results: the case of the spectra of homogeneous Schrödinger operators’, *J. Physique Lett.* **43** (1982) L1.
- [4] V.V. Bazhanov, S.L. Lukyanov and A.B. Zamolodchikov, ‘Integrable structure of conformal field theory, quantum KdV theory and thermodynamic Bethe ansatz’, *Commun. Math. Phys.* **177** (1996) 381 [hep-th/9412229].
- [5] V.V. Bazhanov, S.L. Lukyanov and A.B. Zamolodchikov, ‘Integrable Structure of Conformal Field Theory II. Q-operator and DDV equation’, *Commun. Math. Phys.* **190** (1997) 247 [arXiv:hep-th/9604044].
- [6] V.V. Bazhanov, S.L. Lukyanov and A.B. Zamolodchikov, ‘Integrable structure of conformal field theory. III: The Yang-Baxter relation’, *Commun. Math. Phys.* **200** (1999) 297 [arXiv:hep-th/9805008].
- [7] R.J. Baxter, ‘Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain; 1. Some fundamental eigenvectors’, *Ann. Phys.* **76** (1973) 1–24; ‘2. Equivalence to a generalized ice-type model’, *Ann. Phys.* **76** (1973) 25–47; ‘3. Eigenvectors of the transfer matrix and Hamiltonian’, *Ann. Phys.* **76** (1973) 48–71.
- [8] V.V. Bazhanov, S.L. Lukyanov and A.B. Zamolodchikov, ‘Spectral determinants for Schroedinger equation and Q-operators of conformal field theory’, *J. Statist. Phys.* **102** (2001) 567 [arXiv:hep-th/9812247].
- [9] J. Suzuki, ‘Anharmonic oscillators, spectral determinant and short exact sequence of $U_q(\widehat{sl}(2))$ ’, *J. Phys. A* **32** (1999) L183 [arXiv:hep-th/9902053].
- [10] J. Suzuki, ‘Functional relations in Stokes multipliers and solvable models related to $U_q(A_n^{(1)})$ ’, *J. Phys. A* **33** (2000) 3507 [arXiv:hep-th/9910215].
- [11] P. Dorey and R. Tateo, ‘Differential equations and integrable models: The SU(3) case’, *Nucl. Phys. B* **571** (2000) 583 [Erratum-ibid. B **603** (2001) 582] [arXiv:hep-th/9910102].
- [12] P. Dorey, C. Dunning, D. Masoero, J. Suzuki and R. Tateo, ‘Pseudo-differential equations, and the Bethe ansatz for the classical Lie algebras’, *Nucl. Phys. B* **772** (2007) 249 [arXiv:hep-th/0612298].

- [13] P. Dorey, C. Dunning and R. Tateo, ‘The ODE/IM Correspondence’, *J. Phys. A* **40** (2007) R205 [arXiv:hep-th/0703066].
- [14] P. Dorey, C. Dunning and R. Tateo, ‘From PT-symmetric quantum mechanics to conformal field theory’, *Pramana* **73**, 217 (2009) [arXiv:0906.1130 [hep-th]].
- [15] C.M. Bender and S. Boettcher, ‘Real spectra in non-hermitian Hamiltonians having \mathcal{PT} symmetry’, *Phys. Rev. Lett.* **80** (1998) 5243 [arXiv:physics/9712001].
- [16] C.M. Bender, S. Boettcher and P.N. Meisinger, ‘ \mathcal{PT} symmetric quantum mechanics’, *J. Math. Phys.* **40** (1999) 2201 [arXiv:quant-ph/9809072].
- [17] P. Dorey and R. Tateo, ‘On the relation between Stokes multipliers and the T-Q systems of conformal field theory’, *Nucl. Phys. B* **563** (1999) 573 [Erratum-ibid. B **603** (2001) 581] [arXiv:hep-th/9906219].
- [18] P. Dorey, C. Dunning and R. Tateo, ‘Spectral equivalences, Bethe Ansatz equations, and reality properties in PT-symmetric quantum mechanics’, *J. Phys. A* **34** (2001) 5679 [arXiv:hep-th/0103051].
- [19] K.C. Shin, ‘On the reality of the eigenvalues for a class of PT-symmetric oscillators’, *Commun. Math. Phys.* **229** (2002) 543 [arXiv:math-ph/0201013].
- [20] P. Dorey, C. Dunning, A. Lishman and R. Tateo, ‘PT symmetry breaking and exceptional points for a class of inhomogeneous complex potentials’, *J. Phys. A* **42** (2009) 465302 [arXiv:0907.3673 [hep-th]].
- [21] V.V. Bazhanov, S.L. Lukyanov and A.M. Tsvelik, ‘Analytical results for the Coqblin-Schrieffer model with generalized magnetic fields’, *Phys. Rev. B* **68** (2003) 094427 [arXiv:cond-mat/0305237].
- [22] V. Gritsev, E. Altman, E. Demler and A. Polkovnikov, ‘Full quantum distribution of contrast in interference experiments between interacting one dimensional Bose liquids’, *Nature Physics* **2** (2006) 705 [arXiv:cond-mat/0602475].
- [23] A. Lamacraft and P. Fendley, ‘Order parameter statistics in the critical quantum Ising chain’, *Phys. Rev. Lett.* **100** (2008) 165706 [arXiv:0802.1246 [cond-mat]].
- [24] D. Gaiotto, G.W. Moore and A. Neitzke, ‘Wall-crossing, Hitchin Systems, and the WKB Approximation’ [arXiv:0907.3987 [hep-th]].
- [25] L.F. Alday, D. Gaiotto and J. Maldacena, ‘Thermodynamic Bubble Ansatz’, *JHEP* **1109** (2011) 032 [arXiv:0911.4708 [hep-th]].
- [26] S.L. Lukyanov and A. B. Zamolodchikov, ‘Quantum Sine(h)-Gordon Model and Classical Integrable Equations’, *JHEP* **1007** (2010) 008 [arXiv:1003.5333 [math-ph]].

- [27] R.K. Dodd and R.K. Bullough, ‘Polynomial Conserved Densities for the Sine-Gordon Equations’, Proc. Roy. Soc. Lond. **A352** (1977) 481.
- [28] G. Tzitzeica, ‘Sur une Nouvelle Classe de Surfaces’, C. R. Acad. Sci. Paris, **144** (1907) 1257.
- [29] A.G. Izergin and V.E. Korepin, The inverse scattering method approach to the quantum Shabat-Mikhailov model, Comm. Math. Phys. **79** (1981) 303331
- [30] A.B. Zamolodchikov, Integrable Field Theory from Conformal Field Theory, Proceedings of Taniguchi Symposium, Kyoto (1988)
- [31] F.A. Smirnov, ‘Exact S matrices for $\phi(1,2)$ perturbed minimal models of conformal field theory’, Int. J. Mod. Phys. A **6**, 1407 (1991).
- [32] R.B. Potts, ‘Some generalized order-disorder transformations’, Proc. Cambridge Phil. Soc. **48** (1952) 106.
- [33] S.O. Warnaar, B. Nienhuis and K.A. Seaton, ‘New construction of solvable lattice models including an Ising model in a field’, Phys. Rev. Lett. **69**, 710 (1992).
- [34] S. Faldella, ‘Correspondence between Classical and Quantum Integrability’, (Master Thesis in “Physics of Fundamental Interactions” July 2011, University of Turin, Italy).
- [35] A.V. Mikhaïlov, ‘Integrability of a two-dimensional generalization of the Toda chain’, JETP Letters, **30(7)** (1979) 414.
- [36] M. Ablowitz and P. Clarkson, ‘Solitons, Nonlinear Evolution Equations and Inverse Scattering’, Cambridge University Press, 1991.
- [37] A.B. Zamolodchikov, ‘Integrals of Motion and S Matrix of the (Scaled) $T=\bar{T}(c)$ Ising Model with Magnetic Field’, Int. J. Mod. Phys. A **4** (1989) 4235.