

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Italian aquaculture and the diffusion of alien species: costs and benefits

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1554549> since 2017-07-27T14:09:25Z

Published version:

DOI:10.1111/are.12997

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 Italian aquaculture and the diffusion of alien species: costs and benefits

2

3 Benedetto Sicuro^{1*}, Martina Tarantola¹, Emanuela Valle¹

4 ¹Department of Veterinary Sciences, University of Torino (ITALY)

5 L.go Paolo Braccini 2

6 10095 Grugliasco (Torino)

7 ITALY

8

9 *Corresponding author email: benedetto.sicuro@unito.it; telephone number

10 +390116709260; fax: +390116709240

11

12 Running title: Italian aq. al. species costs and benef.

13 keywords: alien fish, alien aquatic invertebrates, wels catfish, ornamental aquaculture,

14 aquaculture impact, Italy

15

16

17

18

19

20

21

22

23

24

25

26

27

28 **Abstract**

29 The aim of this review is to clarify the relation between Italian aquaculture and the
30 introduction on alien species in Italy. In Italy the most common aquatic alien species are
31 rainbow trout and Manila clam and they represent main economic product of Italian
32 aquaculture. Wels catfish and red clawed crayfish have been voluntarily or involuntarily
33 introduced for aquaculture scope and they are now the most invasive alien aquatic
34 species. Other alien naturalized species, as some salmonid species, are important
35 economic resources in North Italy for inland professional fishery and are considered
36 worthwhile of conservation. Thus, aquaculture is directly or indirectly responsible of the
37 introduction of several alien species in Italy, but its effect on alien species diffusion is
38 peculiar, as few successfully farmed species have become rapidly common. Until now
39 conventional aquaculture has played main role in the diffusion of alien species in Italy,
40 while ornamental aquaculture will be the main source of alien introductions in the
41 future.

42

43

44

45

46

47

48 **Introduction**

49 Biological invasions are indissolubly related with human society (Simberloff , Martin,
50 Genovesi, Maris, Wardle, Aronson, Courchamp, Galil, Garcia-Berthou, Pascal, Pysek,
51 Sousa, Tabacchi & Vila 2013; Bellard, Thuiller, Leroy, Genovesi, Bakkenesk &
52 Courchamp 2013; Essl, Dullinger, Rabitsch, Hulme, Hülber, Jarošíke, Kleinbauer,
53 Krausmann, Kühn, Nentwig, Vilà, Genovesi, Gherardi, Desprez-Loustau, Roques &
54 Pyšek 2011). In particular, in the aquatic ecosystems these invasions are caused by alien
55 species that are species introduced to areas beyond their natural range of distribution by
56 humans, directly or indirectly. Aquaculture plays a key role in alien species diffusion at
57 international level (Casal 2006; De Silva, Nguyen, Turchini, Amarasinghe & Abery
58 2009; Diana 2009; Turchini & De Silva 2008), FAO database on introductions of
59 aquatic species (DIAS) shows that aquaculture is worldwide the most often cited reason
60 for alien fish species introduction (FAO, 2009), (Gherardi, Bertolino, Bodon, Casellato,
61 Cianfanelli, Ferraguti, Lori, Mura, Nocita, Riccardi, Rossetti, Rota, Scalera, Zerunian &
62 Tricarico 2008). Aquaculture can also increase the diffusion parasites from reared fish
63 to wild ones (Krkosek, Gottesfeld, Proctor, Rolston, Carr-Harris, & Lewis 2007). From
64 the quantitative point of view, aquaculture and restocking activities are the main causes
65 of introduction of alien species in Europe, as showed by IMPASSE (Impacts of Alien
66 Species in Aquaculture) project funded by EU Sixth Framework Programme in 2008.
67 Some alien species were initially reared and successively escaped in the wild, as
68 goldfish or catfish or red claw crayfish, others unintentionally transferred with target
69 farmed species as wels catfish, others deliberately released in the natural environment,
70 as Manila clams. In Italy, the greater part of freshwater alien species have been
71 introduced primarily for angling and secondarily for aquaculture scopes, while marine
72 alien species have arrived by international maritime trade or penetration through Suez

73 Canal (lessepsian species) (Andaloro, Falautano, Perzia, Maricchiolo & Castriota 2012).
74 Consequently, the geographical distribution of aquatic alien species in Italy markedly
75 differs in marine and freshwater ecosystems. The great number of marine species were
76 introduced in the 1980s and 1990s in the lagoon of Venice (NE Italy) and in the gulf of
77 Taranto (SE Italy) that are considered “hot spot” of xenodiversity. These introductions
78 were mainly caused by the intercontinental naval traffic due to tourism, commerce and
79 the presence of an Italian Navy base in Taranto (Felline, Caricato, Cutignano, Gorbi,
80 Lionetto, Mollo, Regoli & Terlizzi 2012; Occhipinti-Ambrogi, Marchini, Cantone,
81 Castelli, Chimenz, Cormaci, Frogliola, Furnari, Gambi, Giaccone, Giangrande, Gravili,
82 Mastrototaro, Mazziotti, Orsi-Relini & Piraino 2011). Moreover, it should be
83 considered that Italy has about 7,000 km of coastline and a central position in
84 Mediterranean, these facts naturally expose to marine alien species invasion. Between
85 1945 and 2009, 165 alien marine species have been recorded in Italy, mostly originating
86 from tropical regions of the world (Occhipinti-Ambrogi *et al.* 2011) while 112
87 established alien aquatic species have been recorded in inland waters (Gherardi *et al.*
88 2008). The Italian Ministry of the Environment has published an atlas of Mediterranean
89 alien species (http://www.sidimar.tutelamare.it/distribuzione_alieni.jsp).
90 Meroplanktonic microscopic larvae naturally favor the spread of aquatic alien species,
91 both in marine than in inland water ecosystems. Marine species are often transferred
92 along great distances, principally by means ballast waters and keel fouling. The
93 voluntary release of fish for recreational fishing or aquaculture, in particular in
94 freshwater ecosystems, is particularly diffused in Italy where controls on legal and
95 illegal stocking have been largely ineffective in the past (Gherardi *et al.* 2008).
96 Moreover, ornamental aquaculture is an emerging sector that has caused several alien
97 species introduction, particularly in western countries (Rhyne, Tlustý, Schofield,

98 Kaufman, Morris & Bruckner 2012). In Italy, the shortage of clear rules, the increase of
99 internet trade (Mazza *et al.* 2015) and the continuously increasing number of
100 ornamental species favors illegal trade and international black market (Mazza,
101 Tricarico, Genovesi & Gherardi 2013). In general terms, it is very well known that
102 animal farming is one of the principal direct or indirect cause of diffusion of alien
103 species, but it should be also remembered the economic relevance of animal farming
104 and aquaculture (Diana 2009; Perdikaris & Paschos, 2010). Considering the close
105 relation between the introduction of aquatic alien animal species and aquaculture
106 activities, this review is principally focused on the clarification of the role of Italian
107 aquaculture in the diffusion of alien species in Italy.

108

109 **Freshwater fish**

110 In general, the most common alien Italian species of freshwater fish is the rainbow trout
111 (*Onchorynchus mykiss*) (Table 1). It was introduced in the beginning of 20th century and
112 currently is the first species for Italian aquaculture; in 2013 the rainbow trout production
113 in Italy accounted for 136.5 Mln € (Italian Association of fish farmers, [http://www.api-
114 online.it/index.cfm/en/home](http://www.api-online.it/index.cfm/en/home)). Rainbow trout doesn't make natural reproduction in Italy,
115 consequently its presence is exclusively dependent from artificial restocking, with
116 exception of a population in Trentino region (NE Italy). Its impact on native salmonids
117 species is related to competition with native species and potential diffusion of diseases.
118 Salmonids are generally considered high value species in Italy and several introductions
119 have been attempted in the last decades. Some species of salmonids have been
120 introduced in the late 60's in some alpine lakes and rivers in North Italy, such as arctic
121 charr (*Salvelinus alpinus*), native to Artic region and Brook trout (*Salvelinus fontinalis*)
122 native to North America (Magnea, Sciascia, Paparella, Tiberti & Provenziale 2013).

123 European whitefish, *Coregonus lavaretus* and *C. macrophthalmus*, are zooplanktophagus
124 salmonid imported in Italy from North Europe at the beginning of 20th century and
125 currently acclimatized in some great lakes in the Italian alpine region (Regione
126 Lombardia, 2014) (Figure 2). All the catfish currently present in Italy are alien, some
127 species are naturalized and farmed. Black bullhead catfish (*Ameiurus melas*) and
128 channel catfish (*Ictalurus punctatus*) native to North America and brown bullhead
129 catfish (*A. nebulosus*), native to North America, colonized Italian freshwater
130 ecosystems. Black bullhead catfish and channel catfish and are important species for
131 aquaculture in Central Italy. Wels catfish (*Silurus glanis*) was reported as accidentally
132 escaped from angling ponds in 1956 (Gandolfi & Giannini, 1979). After an initial phase
133 of expansion, in the 1980s it became established in the Po basin in North Italy
134 (Gandolfi, Zerunian, Torricelli & Marconato 1991) and in the Tiber Arno rivers, in
135 Central Italy (Figure 1). Several cyprinids have been recently introduced in Italy as the
136 common bream (*Abramis brama*) and *Pseudorasbora parva* (Volta, Jeppesen, Leoni,
137 Campi, Sala, Garibaldi, Lauridsen & Winfield 2013). The common roach (*Rutilus*
138 *rutius*), recently introduced by Danube basin, has stably colonized Maggiore and
139 Lugano lakes, that are among the bigger Italian lakes (Regione Lombardia, 2012). The
140 most invasive cyprinid is the asp (*Aspius aspius*) (Zerunian, Goltara, Schipani & Boz
141 2009), introduced in Italy from central Europe, the only example of carnivorous
142 cyprinid that stably colonized some areas of Po river basin and currently considered a
143 species highly appreciated for recreational fishing. Few species of Asian alien ciprinids
144 are occasionally farmed in Italy: grass carp (*Ctenopharingodon idellus*), bighead carp
145 (*Hypophthalmichthys nobilis*) and silver carp (*Hypophthalmichthys molitrix*).
146 Largemouth bass (*Micropterus salmoides*) has been introduced in Italy at the beginning

147 of 20th century, it is an active predator that inhabits lentic waters (Marinelli, Scalici &
148 Gibertini 2007; Zerunian *et al.* 2009).

149 **Marine fish**

150 Due to the mentioned geographical reasons, the situation of alien marine fish species is
151 radically different from freshwater and there are several alien fish species with low or
152 unknown impact. Until December 2011, 48 fish alien species have been recorded along
153 Italian coasts, of which only one species is probably originated by aquaculture
154 (Occhipinti-Ambrogi *et al.* 2011). Between the alien marine species, some Carangidae
155 species recently arrived along Italian coasts, as *Seriola carpenteri*, *S. fasciata* and
156 particularly *S. rivoliiana* (Andaloro *et al.* 2012), are potentially interesting for farming,
157 similarly to *S. dumerilii*, fish currently farmed in Japan. In other Mediterranean
158 countries, the presence of some alien species is already considered an opportunity for
159 aquaculture, for example in Cyprus, where a lessepsian fish, rabbit fish (*Siganus*
160 *rivulatus*) is actively farmed (Stephanou & Georgiou, 2000). Differently from
161 freshwaters, marine finfish aquaculture in Italy is based on Mediterranean species:
162 European sea bass (*Dicentrarchus labrax*) and gilthead sea bass (*Sparus aurata*).

163

164 **Freshwater invertebrates**

165 One of the most invasive alien species recently introduced in Europe and in Italy is the
166 red swamp crayfish (*Procambarus clarkii*) (Aquiloni *et al.* 2010), that is listed among
167 the 100 worst invasive alien species (DAISIE 2008). It that has been introduced for
168 farming in the 1980s from Atlantic coasts of North America (Figure 1). Between
169 bivalves, a particularly invasive species is zebra mussel (*Dreissena polymorpha*) that
170 was introduced in Garda lake at the end of '60 and successively had a rapid diffusion in

171 other Italian lakes (Binelli *et al.*, 1997; Lancioni and Gaino, 2006). Asian basket clam,
172 (*Corbicula fluminea*), native to Southeast Asia has become a serious treat for freshwater
173 bivalves of North America and successively introduced in Europe (Araujo, Moreno &
174 Ramos 1993). Similarly to red swamp crayfish, *C. fluminea* is also listed among the 100
175 worst invasive alien species. It was initially found in the Po River in the late 1990's and
176 rapidly invaded many waterways in Northern Italy and arrived in the largest Italian
177 lakes, Garda lake and Maggiore lake (Kamburska, Lauceri, Beltrami, Boggero,
178 Cardeccia, Guarneri, Manca & Riccardi 2013b). Another freshwater alien bivalve
179 introduced in Italy is Chinese pond mussel (*Sinanodonta woodiana*) which is an
180 unionids endemic to China (Colomba, Liberto, Reitano, Grasso, Di Franco & Sparacio
181 2013; Lancioni & Gaino 2006; Kamburska, Lauceri & Riccardi 2013a). Between
182 freshwater gastropods, the most invasive species is probably the mud snail
183 *Potamopyrgus antipodarum*, native to New Zealand, that has colonized all regions of
184 Italy, except Sardinia (Mazza, Agostini, Aquiloni, Cianfanelli, Tricarico & Gherardi
185 2011), reaching in some cases densities of 800,000 individuals/m².

186 **Marine invertebrates**

187 Marine alien invertebrate species are the largest and diversified group of alien species in
188 Italy. These include crustaceans, ascidians, ctenophore, bryozoans, cnidarians,
189 decapods, mollusks and polychaetes for a total number of 401 species (Andaloro *et al.*
190 2012). The lagoon of Venice is the Italian hotspot for marine invertebrate xenodiversity:
191 39 species, including 9 mollusks, and 9 crustaceans. Crocetta (2012) stated that marine
192 mollusks is the group with the highest number of marine alien species in Italy: 35
193 species including 18 Gastropoda, 16 Bivalvia and 1 Cephalopoda (up to December
194 2010). Few species of bivalves have been intentionally introduced for aquaculture: the

195 oysters *Crassostrea gigas*, *C. angularis*, *Saccostrea glomerata* and the Manila clam
196 (*R. philippinarum*).

197 **Bivalves**

198 The most common marine alien Italian bivalve is the Manila clam (*R. philippinarum*)
199 that was deliberately introduced for aquaculture in 1983 in the Venice lagoon and
200 quickly substituted the Italian clam species (*Venus gallina*) (Chiesa, Nonnis Marzano,
201 Minervini, De Lucrezia, Baccarani, Bordignon, Poli, Ravagnan & Argese 2011;
202 Pranovi, Franceschini, Casale, Zucchetta, Torricelli & Giovanardi 2006). Today Manila
203 clam is the first species for Italian aquaculture in value and in 2011 accounted for 222
204 Mln €. After its introduction, Manila clams rapidly became an important economic
205 resource and Italy is first producer country in Europe and second in the world, after
206 China (Perdikaris & Pascos 2010; Chiesa *et al.* 2011). Its typical farming method is a
207 capture - based aquaculture and the Manila clam aquaculture is based on harvesting
208 from naturalized wild population, principally along the coasts of North East Italy,
209 Veneto and Emilia Romagna regions. Other marine bivalve species introduced for
210 aquaculture are the Pacific oysters (*Crassostrea gigas*) and (*C. angulata*), but their
211 diffusion has been not successful as Manila clam and their distribution in the wild is
212 limited to some areas in northern and central Italy (Macali, Conde, Smeriglio, Mariottini
213 P Crocetta 2013). The Asian date mussel (*Arcuatula senhousia*), previously known as
214 *Musculista senhousia* or *Musculista senhousia*, is a alien marine mytilid that was
215 introduced in the early 1990s in the northeastern Adriatic Sea and currently is spread in
216 the Po River deltaic area, northeastern Italy (Mistri 2004; Munari 2008). This mussel
217 sometimes reach densities up 10,000 individuals/sqm in the northern Adriatic, where it
218 is threatening Adriatic shellfish farms (Crocetta 2012). The soft-shell clams (*Mya*
219 *arenaria*), a bivalve native of New England coasts, where it is commercially important

220 for fisheries and aquaculture, is considered among the 100 Mediterranean worst
221 invasive and outcompetes native bivalves (Crocetta & Turolla 2011). Fortunately it is
222 suitable for human consumption and its harvesting can represent a valid method for
223 contrasting its diffusion. The rayed pearl oyster (*Pinctada radiata*), commonly known
224 as pearl oyster, has recently colonized the coasts of a small island in South Italy, Linosa
225 island (Lodola, Nicolini, Savini, Deidun & Occhipinti-Ambrogi 2013). In Greece, *P.*
226 *imbricata* was intentionally introduced during the last century for aquaculture purposes
227 and has established in the wild.

228 **Other marine invertebrates**

229 In Italy there are several alien marine species of invertebrates and, between these, the
230 blue crab (*Callinectes sapidus*) (Thessalou –Legaki, Aydogan, Bekas, Bilge, Boyaci,
231 Brunelli, Circosta, Crocetta, Durucan, Erdem, Ergolavou, Filiz, Fois, Gouva, Kapiris,
232 Katsanevakis, Kljajić, Konstantinidis, Konstantinou, Koutsogiannopoulos, Lamon,
233 Mačić, Mazzette, Meloni, Mureddu, Paschos, Perdikaris, Piras, Poursanidis, Ramos-
234 Esplá, Rosso, Sordino, Sperone, Sterioti, Taşkin, Toscano, Tripepi, Tsiakkiros &
235 Zenetos 2012) recently observed in South Italy (Mancinelli, Carrozzo, Costantini,
236 Rossi, Marini & Pinna 2013) is spreading. The blue crab comes from the Western
237 Atlantic coast of North America and it was reported for the first time in Mediterranean
238 in 1931. Its first presence was dated back to 1971 and successively in Venice lagoon
239 and in Apulia in January 2001 (Florio, Breber, Scirocco, Specchiulli, Cilenti & Lumare
240 2008). This species has a great economic value along southern U.S. Atlantic coasts
241 where farming initiatives are starting. Another portunid, the chinese mitten crab
242 (*Eriocheir sinensis*) has been found also in the Venice lagoon (Mizzan 2005).
243 Polychaete anellids are introduced unintentionally by ballast waters or for bait trade and
244 represent a numerous group of marine alien invertebrates, between them *Branchiomma*

245 *bairdi* that has been recently found along Tyrrhenian coasts, reaching very high densities
246 (c.a. 400 individuals/m²), particularly in polluted areas (Arias, Giangrande, Gambi &
247 Anadon 2013). Another example of invasive sabellid polychaete is *Branchiomma*
248 *luctuosum*, that outcompetes the Mediterranean native tubeworm *Sabella spallanzanii*
249 (Occhipinti *et al.* 2011). One species of alien ctenophore jellyfish, recently found along
250 Italian coasts, is *Mnemiopsis leidyi* that is an effective plankton feeder particularly active
251 on fish eggs and larvae, native to Atlantic coast of US (Boero 2013).

252

253 **Amphibians and aquatic reptiles**

254 Alien amphibians and aquatic reptiles represents a limited phenomenon in Italy and
255 their presence is almost completely caused by ornamental aquaculture. At least eight
256 species of alien amphibians and 13 species of aquatic alien reptiles have been
257 introduced in Italy, but only five species introduced during the last century are
258 considered invasive (Ficetola, Thuiller & Padoa-Schioppa 2009). Main alien reptiles
259 diffused in Italy are red-eared slider, (*Trachemys scripta elegans* Wield, 1839) and
260 yellow-eared slider (*T. s. scripta*), as well as hybrids between them (Agosta & Parolini
261 1999). Among amphibians, American bullfrog (*Lithobates catesbeianus*) was
262 introduced in Italy during the 1930s for food. Moreover green frogs (genus *Pelophylax*),
263 were introduced for farming purposes (Ficetola *et al.* 2009) and their real diffusion is
264 difficult to assess, being able to hybridize with native green frog species.

265

266 **Impact of aquatic alien species**

267 It is widely recognized that the diffusion of alien species result in a loss of biodiversity
268 and sometimes local extinctions (Diana 2009; Ribeiro & Leunda 2012). In Italy, the
269 most common alien species among fish and mollusks, *i.e* rainbow trout and Manila
270 clam, generate a great economic income being the first and second species of Italian
271 aquaculture for economic value (Italian Association of fish farmers, [http://www.api-](http://www.api-online.it/index.cfm/en/home)
272 [online.it/index.cfm/en/home](http://www.api-online.it/index.cfm/en/home)) (Table 1). Due to the success of rainbow farming, Italian
273 inland aquaculture is dominated by alien species, as happens in neighbor countries
274 (Perdikaris & Pascos 2010; Turchini & De Silva 2008). For instance, in 2011 the 92.3
275 % of quantity (that corresponded to 84.2% of value) of freshwater fish farmed
276 production was based on alien species. Naturalized alien salmonids, mainly spread in
277 northern Italian inland waters (whitefish, arctic char and brook trout) (Figure 2), are
278 important species for recreational and professional fisheries (Table 1) and did not
279 apparently affect lake productivity (Zerunian *et al.* 2009). European whitefish is one of
280 the dominant fish species in the bigger Italian lakes (Garda and Maggiore lakes) and it
281 is considered a typical product of local gastronomy (Regione Lombardia 2012). Being
282 unable to make natural reproduction, rainbow trout is occasionally released in controlled
283 conditions for recreational fishing, without significant effect on autochthonous
284 salmonids, with the exception of one population in Trentino region that encompass
285 natural reproduction. Other species as wels catfish (*S. glanis*) and red swamp crayfish
286 have an tremendous impact on local aquatic fauna and in the freshwater ecosystems.
287 Wels catfish is an active predator and its presence in Italy caused several problems in
288 the Po River basin to native species (*i.e.* *Alburnus arborella* and *Scardinius*
289 *erythrophthalmus*), causing in some cases local extinctions (Castaldelli, Pluchinotta,
290 Milardi, Lanzoni, Giari, Rossi & Fano 2013; Zerunian *et al.* 2009). Wels catfish
291 outcompetes native predators, the northern pike (*Esox lucius*) and European perch

292 (*Perca fluviatilis*), that have become extremely rare after its introduction, thus affecting
293 recreational and professional fishing activities. Wels catfish impact is also evident on
294 those fish species that lay eggs on bottom vegetation, as tench (*Tinca tinca*) (Milillio,
295 2012). The diffusion of wels catfish induced also decline in other alien species as the
296 largemouth bass (*Micropterus salmoides*, Lacépède), black bullhead catfish, (*Ameiurus*
297 *melas* Rafinesque, 1820) and goldfish, (*Carassius auratus* L.) that were appreciated for
298 farming or recreational fishing. Red swamp crayfish has a high ecological plasticity and
299 it reduces biomass and species richness of macroinvertebrates, inducing drastic habitat
300 changes (Liu, Guo, Ke, Wang & Li 2011; Scalici, Chiesa, Scuderi, Celauro & Gibertini
301 2010). This species can outcompete the autochthonous crayfish, white-clawed crayfish
302 (*Austropotamobius pallipes*), and introduced in Italy the crayfish pest, a fungal disease
303 carried by *Aphanomices astaci*, lethal for European species of crayfish (Aquiloni *et al.*
304 2010). Red swamp crayfish make burrows in the pond and causes a noticeable physical
305 impact on freshwater ecosystems (Scalici *et al.* 2010; Garzoli, Paganelli, Rodolfi,
306 Savini, Moretto, Occhipinti-Ambrogi & Picco 2014; Scalera and Zaghi, 2004; Barbaresi
307 *et al.* 2004). Some marine alien or lessepsian fish species belonging to Tetraodontids
308 such as *Sphoeroides pachygaster* and *Lagocephalus sceleratus* are known for their
309 toxicity due to the neurotoxin tetrodotoxin that may produce paralysis of the diaphragm
310 and death (Andaloro *et al.* 2012) and they can also introduce pathogens for other fish
311 species. Uncontrolled harvests of Pacific oysters (*Crassostrea gigas*) eventually
312 contaminated by toxic algae can lead to diseases in humans. Between freshwater
313 bivalves, zebra mussel can accumulate biotoxins and sometimes favors the bloom of
314 *Microcystis aeruginosa*, a toxic microalga (Mazza *et al.* 2013). Similarly to the red
315 clawed crayfish, alien amphibians can spread fungal diseases as chytridiomycosis and
316 *Batrachochytrium dendrobatidis*, that can cause autochthonous amphibian decline

317 (Ficetola & Scali 2010). Concerning aquatic reptiles, sliders outcompetes for food and
318 basking places the autochthonous European pond turtle (*Emys orbicularis*) (Ficetola *et*
319 *al.* 2009) and, being active predators, sliders have heavy impact on fish and amphibians
320 species, in particular in those species that lay eggs in shallow waters (Ferri & Soccini
321 2003).

322 **Impact of exotic species and aquaculture: an ecological perspective**

323

324 Dominance and diversity are two key concepts of basic ecology that define the relations
325 between species in natural ecosystems and in particular they represent the ecosystems
326 answer to environmental perturbations. These concepts can be adopted in this context in
327 order to better explain the relation between aquaculture and diffusion of exotic species.
328 The role of aquaculture has been dual, in fact few successfully farmed species have
329 become rapidly common, these species can be considered “xenodominant”, while the
330 species involuntarily introduced are much more numerous, never reach considerable
331 amounts and increase xenodiversity. The diffusion of species such as tilapia or carps has
332 been enormously increased for farming purposes (Diana 2009). Conventional
333 aquaculture (and in an historic perspective, any traditional zootechnic activity) acts as a
334 kind of key of success for few alien species and in several cases the introductions of
335 freshwater fish have resulted in great societal benefits (Diana 2009; Gozlan 2008).
336 Ornamental aquaculture and escaped species increase xenodiversity. Therefore,
337 xenodominance is the first ecological effect of successful alien species and
338 xenodominant species have beneficial economic effect on local communities, as the
339 rainbow trout or Manila clam.

340

341 **Management of alien species: status and perspective**

342 Until this moment the main solution proposed for alien invasive species has been the
343 control of international trade or containment measures against diffusion of alien. Most
344 common proposed measures regard on one hand rules and international directives that
345 discipline the import of ornamental species and farmed animals, on the other hand
346 suggestions on alternative methods against unattended transfers, such as the substitution
347 of ballast waters in open ocean areas and the removal of fouling organisms, for marine
348 species. In Italy main conservation efforts and/or management project have been
349 focused on freshwater ecosystems that are much more fragile (Gherardi *et al.* 2008).
350 Upon our knowledge, there are not management initiatives on marine fish species.
351 Freshwater ecosystems lacking native piscivorous fishes, or being highly altered by
352 humans, appear to be the most vulnerable to biological invasions (Casal 2006;
353 Castaldelli *et al.* 2013). This vulnerability could also increase in the future, as
354 consequence of climatic conditions change and it is likely that will favor the spread of
355 introduced species not only in Europe, but also in northeastern North America, and
356 Oceania (Bellard *et al.* 2013). In these last two decades, several UE projects in North
357 Italy, have been focused on freshwater alien species management, in particular targeted
358 against wels catfish diffusion. Between 2010 and 2012, an intensive fishing activity in
359 collaboration with local associations of non-professional fisherman produced a
360 significant harvest of wels catfish biomass.
361 (http://www.progettosiluro.altervista.org/opuscolo_siluro.pdf). Other containment
362 measures have been adopted for wels catfish as selective fishing and exportation of live
363 fish toward eastern European markets (Castaldelli *et al.* 2013), exportation in areas
364 where wels catfish angling is a rooted touristic activity, (as Slovenia) and promoting
365 direct consumption. Containment strategies have been proposed for red swamp crayfish

366 control (Garzoli *et al.* 2014) and two LIFE projects have been recently funded for this
367 purpose: RARITY (http://www.life-rarity.eu/pages/rarity_en.htm) and SOS TUSCAN
368 WETLANDS (<http://www.life-sostuscanwetlands.eu/index.php/it/?lang=en>). The
369 management of alien salmonids is completely different from above mentioned species,
370 some of these naturalized species are important for local freshwater fishery and
371 consequently worth of conservation and protection. In Trentino region (NE Italy) the
372 arctic charr is even protected by a UE disciplinary (Official Journal of the European
373 Union, September 24th, 2013) and it is considered typical product of regional
374 gastronomy. Re-introduction projects have been carried out in areas where these alien
375 salmonids decreased, due to over exploitation and environmental pollution.

376 Economic opportunities can arise from alien species presence thus partially
377 transforming the nuisance into a opportunity such as aquaculture, recreational fishing
378 and fishing tourism. In Italy fishing tourism is a modern and well known activity
379 (Alberini, Zanatta & Rosato 2007; Picchi, Scalera & Zaghi 2006; Scalera & Zaghi
380 2004) and in 2012, in the Lombardy region, a significant increase of fishing tourism
381 practiced by German fisherman has been noticed, as a consequence of wels catfish
382 increase. Fisherman often collaborated in conservation projects as happened during
383 LIFE projects in 2003 and 2004 in Emilia region, but, more interestingly, they are
384 available to pay extra admission fees in order to do selective fishing in protected areas.
385 It should be remembered that recreational fishing in Italy regards nowadays almost 2
386 Mln persons only in freshwaters.

387 Novel products and activities can be developed from alien species, as happens in
388 France, Croatia and Poland, where wels catfish is farmed (Turchini *et al.*, 2008;
389 Ulikowski, Szczepkowski & Szczepkowska 2003), or Cyprus where rabittfish is farmed
390 in marine cages (Stephanou & Georgiou, 2000). Selective fishing targeted on low-value

391 species as pumpkinseed bass (*Lepomis gibbosus*), can be eventually oriented for the
392 production of forage/bait fish, following the experience of bluegill sunfish (*Lepomis*
393 *macrochirus*) farming in southern US states. The recent diffusion of alien blue crab (*C.*
394 *sapidus*) along Italian coasts could be turned from ecological trouble into opportunity,
395 by crab farming under controlled conditions, considering that along eastern coasts of
396 U.S. it is one of most expensive seafood. Freshwater mollusks, when not directly
397 suitable for human consumption, are potentially utilizable for farmed animal feeds
398 (Sicuro, Abete, Forneris, Mioletti, Panini & Amedeo 2010). Zebra mussel farming plant
399 has been proposed as a potential system for a bioremediation project in the Baltic area
400 (Schernewski, Stybel & Neumann 2012). Chinese pond mussel (*S. woodiana*) in Central
401 Italy (Tuscany region) was used in 2003 for freshwater pearl production
402 (http://prometeo.adm.unipi.it/temp/4_Pisa.pdf). As showed until this point, potential
403 utilization of exotic species is an attractive option, but it must be handled with care as
404 potential exploitation could promote the further diffusion of exotic.

405 As regards as reptiles species, the ban to import *T. s. scripta* was imposed by EU since
406 1997 via the Protection of Species of Wild Fauna and Flora by Regulating Trade. Only
407 one of the two subspecies of exotic sliders (*T. s. scripta* and *T. s. elegans*) is curiously
408 considered dangerous, while other species can be legally imported and traded. This fact
409 has not biological reason as the two subspecies have similar ecological features and
410 hybrids are fecund and spread in the wild.

411 **Conclusions**

412 In Italy several alien aquatic species are currently present and some of them have
413 documented relevant ecological and economic impact, both negative than positive.

414 From 2009, the adoption of the EU regulation 708/2007, concerning the use of alien and
415 locally absent species in aquaculture, has disciplined this sector. However, it is not
416 sufficient to prevent the diffusion of aquatic alien species as Italian aquaculture
417 encompasses a multiplicity of aspects that include conventional finfish aquaculture,
418 bivalve aquaculture, ornamental aquaculture and the relation with introduction of alien
419 species is consequently a multiple facets issue. Basically, there are two main groups of
420 exotic aquatic species originated by aquaculture: the invasive ones with wels catfish and
421 Louisiana red swamp crayfish and another heterogeneous group of exotic species that
422 have beneficial effects. The most common Italian aquatic alien species are rainbow trout
423 and Manila clams and they represent main economic products of Italian aquaculture.
424 Moreover other species of naturalized salmonids are important economic resources for
425 inland professional fishery in North Italy. As regard as ornamental aquaculture,
426 including aquatic reptiles and amphibians, the consequences on alien species diffusion
427 are currently more limited, but it should not be underestimated in perspective. The
428 introduction of new exotic species in Italy is almost completely forbidden by a recent
429 Italian law (n.1143/2014, applied from January 2015), consequently ornamental
430 aquaculture will be the main future source of exotic aquatic species diffusion. In this
431 context, the internet trade (currently uncontrolled) will represent the real challenge of
432 future. In this moment Italian ornamental aquaculture is almost exclusively based on
433 alien species, in freshwater and marine species, moreover there are few ornamental
434 aquaculture farm in Italy and it can be estimated that more than 90% of this sector is
435 dominated by importation from foreign countries. Advisable solution could be the
436 encouraging of locally – reared ornamental species farming, that in this moment is not
437 yet economically convenient in Italy.

438

439 **Acknowledgments**

440 The authors are grateful to dr. Pierluca Costa and prof. Pierpaolo Mussa for their critical
441 revision of manuscript.

442

443 **References**

444 Agosta F. & Parolini L. (1999) Autoecologia e rapporti sinecologici di popolazioni
445 introdotte in Lombardia di *Trachemys scripta elegans*. Dati preliminari. Atti 2° Congr.
446 Naz.le SHI, Praia del Mare, 6-10 ottobre 1998. *Rivista di Idrobiologia* **38**: 421-430

447 Alberini A., Zanatta V. & Rosato P. (2007) Combining actual and contingent behavior
448 to estimate the value of sports fishing in the Lagoon of Venice. *Ecological Economics* **6**
449 **1**: 530 – 541

450 Andaloro F., Falautano M., Perzia P., Maricchiolo C. & Castriota L. (2012)
451 Identification and distribution of non-indigenous species in the Mediterranean Sea: the
452 Italian challenge. *Aliens: The Invasive Species Bulletin* **32**: 13-19.

453 Aquiloni L., Tricarico E. & Gherardi F. (2010) Crayfish in Italy: distribution, threats
454 and management. *International Aquatic Research* **2**: 1-14

455 Araujo R., Moreno D. & Ramos M.A. (1993) The Asiatic clam *Corbicula fluminea*
456 (Müller, 1774) (Bivalvia: Corbiculidae) in Europe. *American Malacological Bulletin*
457 **10**: 39–49

458 Arias A., Giangrande A., Gambi M. & Anadon N. (2013) Biology and new records of
459 the invasive species *Branchiomma bairdi* (Annelida: Sabellidae) in the Mediterranean
460 Sea. *Mediterranean Marine Science* **14**,(1): 162-171

- 461 Barbaresi S., Tricarico E. & Gherardi F. (2004) Factors inducing the intense burrowing
462 activity by the red swamp crayfish, *Procambarus clarki*, an invasive species.
463 *Naturwissenschaften* **91**: 342-345
- 464 Bellard C., Thuiller W., Leroy B., Genovesi P., Bakkenesk M. & Courchamp F. (2013)
465 Will climate change promote future invasions? *Global Change Biology* **19**: 3740–3748
- 466 Boero F. (2013) Review of jellyfish blooms in the Mediterranean and Black Sea.
467 Studies and Reviews. General Fisheries Commission for the Mediterranean. No. 92.
468 Rome, 53 p.
- 469 Casal C.M.V. (2006) Global documentation of fish introductions: the growing crisis
470 and recommendations for action *Biological Invasions* **8**: 3–11
- 471 Castaldelli G., Pluchinotta A., Milardi M., Lanzoni M., Giari L., Rossi R. & Fano E.A.
472 (2013) Introduction of exotic fish species and decline of native species in the lower Po
473 basin, north-eastern Italy. *Aquatic Conservation: Marine and Freshwater Ecosystems*
474 **23**: 405–417
- 475 Chiesa S., Nonnis Marzano F., Minervini G., De Lucrezia D., Baccarani G., Bordignon
476 G., Poli I., Ravagnan G. & Argese E. (2011) The invasive Manila clam *Ruditapes*
477 *philippinarum* (Adams and Reeve, 1850) in Northern Adriatic Sea: Population genetics
478 assessed by an integrated molecular approach. *Fisheries Research* **110**: 259–267
- 479 Colomba M.S., Liberto F., Reitano A., Grasso R., Di Franco D. & Sparacio I. (2013) On
480 the presence of *Dreissena polymorpha* Pallas, 1771 and *Sinanodonta woodiana*
481 *woodiana* (Lea, 1834) in Sicily (Bivalvia). *Biodiversity* **4** (4): 571-580

- 482 Crocetta F. (2012) Marine alien Mollusca in Italy: a critical review and state of the
483 knowledge. *Journal of the Marine Biological Association of the United Kingdom* **92**,
484 (6): 1357-1365.
- 485 Crocetta F. & Turolla E. (2011) *Mya arenaria* Linné, 1758 (Mollusca: Bivalvia) in the
486 Mediterranean Sea: its distribution revisited. *Journal of Biological Research-*
487 *Thessaloniki*. **16**: 188 – 193
- 488 De Silva S.S., Nguyen T.T.T., Turchini G.M., Amarasinghe U.S. & Abery N.W. (2009)
489 Alien species in aquaculture and biodiversity: a paradox in food production. *Ambio* **38**
490 (1): 24–28.
- 491 Diana J.S. (2009) Aquaculture Production and Biodiversity Conservation. *BioScience*
492 **59** (1): 27–38.
- 493 Essl F., Dullinger S., Rabitsch W., Hulme P.E., Hülber K., Jarošíke V., Kleinbauer I.,
494 Krausmann F., Kühn I., Nentwig W., Vilà M., Genovesi P., Gherardi F., Desprez-
495 Loustau M., Roques A. & Pyšek P. (2011) Socioeconomic legacy yields an invasion
496 debt. *Proceedings of National Academy of Sciences* **108** (1): 203–207.
- 497 FAO(2009) (Food and Agriculture Organization of the United Nations) DIAS. FAO
498 Database on Introductions of Aquatic Species. Available from URL:
499 <http://www.fao.org/fishery/dias/en>.
- 500 Felling S., Caricato R., Cutignano A., Gorbi S., Lionetto M.G., Mollo E., Regoli F. &
501 Terlizzi A. (2012) Subtle Effects of Biological Invasions: Cellular and Physiological
502 Responses of Fish Eating the Exotic Pest *Caulerpa racemosa*. *PLoS ONE* **7**(6): 38763 –
503 38769

- 504 Ferri V. & Soccini C. (2003) Riproduzione di *Trachemys scripta elegans* in condizioni
505 semi-naturali in Lombardia (Italia settentrionale). *Natura Bresciana* **33**: 89-92.
- 506 Ficetola G.F., Thuiller W. & Padoa-Schioppa E. (2009) From introduction to the
507 establishment of alien species: bioclimatic differences between presence and re
508 production localities in the elidersi turtle. *Diversity and Distributions*. **15**: 108-116.
- 509 Ficetola G.F. & Scali S. (2010) Invasive amphibians and reptiles in Italy. In: *Atti VIII*
510 *Congresso Nazionale Societas Herpetologica Italica*. (ed. by L. Di Tizio, A. Di Cerbo,
511 N. Di Francesco, A. Cameli), pp. 335-340. Ianieri Edizioni, Pescara (*in Italian*).
- 512 Florio M., Breber P., Scirocco, T., Specchiulli A., Cilenti L. & Lumare L. (2008) Exotic
513 species in Lesina and Varano Lakes: Gargano National Park (Italy). *Transitional Waters*
514 *Bulletin*. **2**:69–79
- 515 Gandolfi G., Zerunian S., Torricelli P. & Marconato A. (1991) Pesci delle Acque
516 Interne Italiane. Istituto Poligrafico e Zecca dello. Stato: Roma pp 189 – *In Italian*
- 517 Gandolfi G. & Giannini M. (1979) La presenza di *Silurus glanis* nel fiume Po
518 (*Osteichthyes Siluridae*). *Natura* **70**: 3–6.
- 519 Garzoli L., Paganelli D., Rodolfi M., Savini D, Moretto M., Occhipinti-Ambrogi A. &
520 Picco A.M. (2014) First evidence of microfungal “extra oomph” in the invasive red
521 swamp crayfish *Procambarus clarki*. *Aquatic Invasions* **9** (1): 47–58
- 522 Gherardi F., Bertolino S., Bodon M., Casellato S., Cianfanelli S., Ferraguti M., Lori E.,
523 Mura G., Nocita A., Riccardi N., Rossetti G., Rota E., Scalera R., Zerunian S. &
524 Tricarico E. (2008) Animal xenodiversity in Italian inland waters: distribution, modes of
525 arrival, and pathways. *Biological Invasions* **10**:435–454

- 526 Gozlan R. E. (2008) Introduction of non-native freshwater fish: is it all bad? *Fish and*
527 *Fisheries* **9** (1):106–115
- 528 Kamburska L., Lauceri R. & Riccardi N. (2013a) Establishment of a new alien species
529 in Lake Maggiore (Northern Italy): *Anodonta* (*Sinanodonta*) *woodiana* (Lea, 1834)
530 (*Bivalvia*: *Unionidae*). *Aquatic Invasions* **8** (1): 111–116.
- 531 Kamburska L., Lauceri R., Beltrami M., Boggero A., Cardeccia A., Guarneri I., Manca
532 M. & Riccardi N. (2013b) Establishment of *Corbicula fluminea* (O.F. Müller, 1774) in
533 Lake Maggiore: a spatial approach to trace the invasion dynamics. *Bioinvasions*
534 *Records* **2** (2): 105–117.
- 535 Krkosek M., Gottesfeld A., Proctor B., Rolston D., Carr-Harris C. & Lewis M.A. (2007)
536 Effects of host migration, diversity and aquaculture on sea lice threats to Pacific salmon
537 populations. *Proceeding of the Royal Society B* **274**: 3141–3149.
- 538 Lancioni T. & Gaino E. (2006) The invasive zebra mussel *Dreissena polymorpha* in
539 Lake Trasimeno (Central Italy): Distribution and reproduction. *Italian Journal of*
540 *Zoology* **73**(4): 335–346.
- 541 Liu X., Guo Z., Ke Z., Wang S. & Li Y. (2011) Increasing Potential Risk of a Global
542 Aquatic Invader in Europe in Contrast to Other Continents under Future Climate
543 Change. *PLoS ONE* **6**(3): e18429
- 544 Lodola A., Nicolini L., Savini D., Deidun A. & Occhipinti-Ambrogi A. (2013) Range
545 expansion and biometric features of *Pinctada imbricata radiata* (*Bivalvia*: *Pteriidae*)
546 around Linosa Island, Central Mediterranean Sea (Italy). *Italian Journal of Zoology* **80**
547 (2): 303-312

- 548 Macali A., Conde A., Smeriglio C., Mariottini P. & Crocetta F. (2013) The evolution of
549 the molluscan biota of Sabaudia Lake: a matter of human history. *Scientia Marina*
550 **77**(4): 649-662
- 551 Magnea U., Sciascia R., Paparella F., R. Tiberti & Provenzale A. (2013) A model for
552 high-altitude alpine lake ecosystems and the effect of introduced fish. *Ecological*
553 *Modelling* **251**: 211 – 220
- 554 Mancinelli G., Carrozzo L., Costantini M.L. , Rossi L., Marini G. & Pinna M. (2013)
555 Occurrence of the Atlantic blue crab *Callinectes sapidus* Rathbun, 1896 in two
556 Mediterranean coastal habitats: Temporary visitor or permanent resident? *Estuarine*
557 *Coastal and Shelf Science* **135**, 46-56
- 558 Marinelli A., Scalici M., & Gibertini G. (2007) Diet and reproduction of largemouth
559 bass in a recently introduced population, lake Bracciano (central Italy). *Bulletin*
560 *Francais Pêche Pisciculture* **385**: 53-68
- 561 Mazza G., Agostini N., Aquiloni L., Cianfanelli S, Tricarico E. & Gherardi F. (2011)
562 Ecological characterisation of streams invaded by the New Zealand mud snail
563 *Potamopyrgus antipodarum* (Gray 1843): the case study of a National Park in Italy.
564 *Ethology Ecology & Evolution*, **23** (2): 151-164
- 565 Mazza G., Tricarico E., Genovesi P. & Gherardi F. (2013) Biological invaders are
566 threats to human health: an overview. *Ethology Ecology & Evolution*. **26** (2-3): 112 -
567 119
- 568 Mazza G., Aquiloni L., Inghilesi A.F., Giuliani C., Lazzaro L., Ferretti G., Lastrucci L.,
569 Foggi B. & Tricarico E. (2015) Aliens just a click away: the online aquarium trade in
570 Italy. *Management of Biological Invasions* **6**:253-261.

- 571 Milillio S. (2012) Diffusione del ‘*Silurus ganis*’ nell’area del Po, gestione economica di
572 una specie alloctona e turismo sostenibile Ms thesis 62 pp
- 573 Mistri M. (2004) Prey preference of *Carcinus aestuarii*: possible implications with the
574 control of an invasive mytilid and Manila clam culture in a northern Adriatic lagoon.
575 *Aquaculture* **230** (1): pp. 261-272
- 576 Mizzan L. (2005) *Rhithropanopeus harrisi* (Gould, 1841) (Crustacea, Decapoda,
577 Panopeidae) and *Eriocheir sinensis* H. Milne Edwards, 1854 (Crustacea, Decapoda,
578 Grapsidae): two new exotic crabs in the Venetian Lagoon. *Bollettino del Museo Civico*
579 *di Storia Naturale di Venezia* **56**: 89-95
- 580 Munari C. (2008) Effects of the exotic invader *Musculista senhousia* on benthic
581 communities of two Mediterranean lagoons. *Hydrobiologia* **611**:29–43
- 582 Occhipinti-Ambrogi A., Marchini A., Cantone G., Castelli A., Chimenz C., Cormaci
583 M., Froggia C., Furnari G., Gambi M.C., Giaccone G., Giangrande A., Gravili C.,
584 Mastrototaro F., Mazziotti C., Orsi-Relini L. & Piraino S. (2011) Alien species along
585 the Italian coasts: an overview. *Biological Invasions* **13**, (2): 531-532
- 586 Perdikaris C. & Paschos I. (2010) Organic aquaculture in Greece: a brief review.
587 *Reviews in Aquaculture* **2**: 102–105
- 588 Picchi S., Scalera R. & Zaghi D. (2006) Il bilancio di LIFE Natura in Italia - Indicazioni
589 e prospettive per il futuro”. Ministero dell’Ambiente e della Tutela del Territorio -
590 Direzione Generale Protezione della Natura. Roma. 224.pp.
- 591 Pranovi F., Franceschini G., Casale M., Zucchetta M., Torricelli P. & Giovanardi O.
592 (2006) An ecological imbalance induced by a non-native species: the Manila clam in the
593 Venice Lagoon. *Biological Invasions* **8**: 595–609

- 594 Regione Lombardia (2014) Censimento della fauna ittica nei laghi alpini nel territorio
595 della regione Lombardia pp. 25. Available from URL:
596 [http://www.lavoro.regione.lombardia.it/shared/ccurl/456/284/Censimento%20fauna%20](http://www.lavoro.regione.lombardia.it/shared/ccurl/456/284/Censimento%20fauna%20ittica%20-%20avanzamento%20febbraio%202014.pdf)
597 [ittica%20-%20avanzamento%20febbraio%202014.pdf](http://www.lavoro.regione.lombardia.it/shared/ccurl/456/284/Censimento%20fauna%20ittica%20-%20avanzamento%20febbraio%202014.pdf) [in Italian]
- 598 Regione Lombardia. (2012) Programma della pesca e dell'acquacoltura della regione
599 Lombardia 2012-2014. N. 294, 240 pp. Available from URL:
600 www.arca.regione.lombardia.it/.../burl%2048%20SO%2029.11.2012.pdf [in Italian]
- 601 Rhyne A.L., Tlusty M.F., Schofield P.J., Kaufman L., Morris J.A. Jr. & Bruckner A.W.
602 (2012) Revealing the Appetite of the Marine Aquarium Fish Trade: The Volume and
603 Biodiversity of Fish Imported into the United States. *PLoS ONE* **7**(5): 1-9
- 604 Ribeiro F. & Leunda P.M. (2012) Non-native fish impacts on Mediterranean freshwater
605 ecosystems: current knowledge and research needs. *Fisheries Management and Ecology*
606 **19**: 142–156
- 607 Scalera R. & Zaghi D. (2004) LIFE Focus / Alien species and nature conservation in the
608 EU. The role of the LIFE program. Luxembourg: Office for Official Publications of the
609 European Communities 56 pp.
- 610 Scalici M., Chiesa S., Scuderi S., Celauro D. & Gibertini G. (2010) Population structure
611 and dynamics of *Procambarus clarkii* (Girard, 1852) in a Mediterranean brackish
612 wetland (Central Italy). *Biological Invasions* **12**: 1415–1425
- 613 Schernewski G., Stybel N. & Neumann T. (2012) Zebra Mussel Farming in the
614 Szczecin (Oder) Lagoon: Water-Quality Objectives and Cost-Effectiveness. *Ecology*
615 *and Society* **17**(2): 4- 12

- 616 Sicuro B., Abete C., Forneris G., Mioletti S., Panini E. & Amedeo S. (2010) Potential
617 utilisation of farmed freshwater mussels (*Anodonta anatina* and *Unio mancus*) in North
618 West Italy. *Cuban Journal of Agricultural Science*. **44** (4): 409 – 417
- 619 Simberloff D., Martin J, Genovesi P., Maris V., Wardle D.A., Aronson J., Courchamp
620 F., Galil B., Garcia-Berthou E., Pascal M., Pysek P., Sousa R., Tabacchi E. & Vila M.
621 (2013) Impacts of biological invasions: what's what and the way forward. *Trends in*
622 *Ecology & Evolution*. **28**, (1): 58 – 66
- 623 Stephanou D. & Georgiou G. (2000) Recent experiences on the culture of rabbitfish
624 *Siganus rivulatus* in Cyprus. Recent advances in Mediterranean aquaculture finfish
625 species diversification. Zaragoza :*CIHEAM Cahiers Options Méditerranéennes* **47**:
626 295-301. Available at <http://om.ciheam.org/om/pdf/c47/00600630.pdf>.
- 627 Thessalou –Legaki M., Aydogan O., Bekas P., Bilge G., Boyaci Y.Ö., Brunelli E.,
628 Circosta V., Crocetta F., Durucan F., Erdem M., Ergolavou F., Filiz H., Fois F., Gouva
629 E., Kapiris K., Katsanevakis S., Kljajić Z., Konstantinidis E., Konstantinou G.,
630 Koutsogiannopoulos D., Lamon S., Mačić V., Mazzette R., Meloni D., Mureddu A.,
631 Paschos I., Perdikaris C., Piras F., Poursanidis D., Ramos-Esplá A.A., Rosso A.,
632 Sordino P., Sperone E., Sterioti A., Taşkin E., Toscano F., Tripepi S., Tsiakkios L. &
633 Zenetos A. (2012) New Mediterranean Biodiversity Records (December 2012).
634 *Mediterr. Mar. Sci.* **13**: 312-327
- 635 Turchini G.M. & De Silva S.S. (2008) Bio-economical and ethical impacts of alien
636 finfish culture in European inland waters. *Aquaculture International* **16** (3): 243–272.
- 637 Ulikowski D., Szczepkowski M. & Szczepkowska B. (2003) Preliminary studies of
638 intensive wels catfish (*Silurus glanis* L.) and sturgeon (*Acipenser* sp.) pond cultivation.
639 *Archives of Polish Fisheries* **11** (2): 295-300.

640 Volta P., Jeppesen E., Leoni B., Campi B., Sala P., Garibaldi L., Lauridsen T.L. &
641 Winfield I.J. (2013) Recent invasion by a non-native cyprinid (common bream *Abramis*
642 *brama*) is followed by major changes in the ecological quality of a shallow lake in
643 southern Europe. *Biological Invasions* **15** (9): 2065-2079

644 Zerunian S., Goltara A., Schipani I. & Boz B. (2009) Adeguamento dell'Indice di Stato
645 delle Comunità Ittiche alla Direttiva Quadro sulle Acque. *Biologia Ambientale* **23** (2):
646 15-30.

647

648

649

650

651

652 Table 1. List of species of Italian aquaculture in 2011 (Italian Association of fish
 653 farmers, <http://www.api-online.it/index.cfm/en/home>)

Species	tons	Mln €	Origin
Rainbow trout (<i>Onchorynchus mykiss</i>)	41000	149.7	A
European seabass (<i>Dicentrarchus labrax</i>)	8700	64	I
Gilthead seabream (<i>Sparus aurata</i>)	9700	72	I
Gray mullets (mugilids)	3500	9.8	I
Sturgeons (Acipenserids)	1660	14.8	I + A
Eel (<i>Anguilla anguilla</i>)	1100	11.4	I
Common carp (<i>Cyprinus carpio</i>)	750	2.9	I
Catfish (<i>Ictalurus sp.</i>)	550	3.3	A
Arctic charr (<i>Salvelinus alpinus</i>)	400	1.6	A
Meagre (<i>Argyrosomus regius</i>)	300	2.1	I
Other fish species	5150	32.4	I + A
Blue mussels (<i>Mytilus galloprovincialis</i>)	100000	78	I
Manila clams (<i>Ruditapes philippinarum</i>)	33000	144	A
TOTAL	205810	586	

654

655 A, exotic species; I, autochthonous species; I + A autochthonous + exotic species

656


657

658

659

660 Figure 1. Wels catfish and red swamp crayfish distribution

661



662

663

664

665

666

667

668

669

670 Figure 2. Naturalized exotic salmonid distribution



- Brook trout (*Salvelinus fontinalis*)
- European whitefish (*Coregonus lavaretus*)
and (*C. macrophthalmus*)
- # Arctic charr (*Salvelinus alpinus*)

671

672

673

674

675