



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

*Placenta, Volume 36, Issue 9, Page A27, 2015, © 2015 Published by Elsevier Inc.,
<http://dx.doi.org/10.1016/j.placenta.2015.07.265>*

Placenta derived mesenchymal stem cells (PDMSCs) modulate HIF-1a, VEGF and JunB genes in ovarian cancer cells.

Domenica Giuffrida, Cristian Zenerino, Rossella Barrile, Anna Maria Nuzzo, Tullia Todros, Alessandro Rolfo

Dept. of Surgical Sciences, University of Turin, Turin, Italy

Ovarian cancer (OC) has the highest mortality rate of all gynecological cancers. We previously demonstrated that PDMSCs produce soluble factors able to inhibit OC cell cycle and growth through a pathway that still remains to be determined. Hypoxia Inducible Factor-1 alpha (HIF-1a), main player in cellular response to hypoxia, promotes the expression of VEGF, responsible for neo-angiogenesis. HIF-1a over-expression has been described in most human malignancies and it is linked to bad cancer prognosis. Activating Protein-1 (AP-1) molecules are pivotal regulators of OC cell cycle progression and metastatization. Herein, we investigated HIF-1a, VEGF and JunB expression in OC ES-2 and SKOV-3 cells treated by PDMSCs conditioned medium (CM) in order to verify our hypothesis that PDMSCs could act through the modulation of HIF-1a and anti-proliferative AP-1 molecules.

Methods: PDMSCs were isolated from physiological human term placentae (n=12). CM will be produced by incubating PDMSCs for 24-48 hours in DMEM medium. CM will be harvested and filtered to remove cellular debris. ES-2 and SKOV-3 cells were treated by 24/48h CMs for 24h or 48h and mRNA was isolated. Expression of HIF-1a, VEGF and JunB was assessed by Real Time PCR.

Results: HIF-1a mRNA levels were significantly decreased in both ES-2 and SKOV-3 cells treated by 24/48h PDMSCs-CMs for 24/48h. HIF-1a decrease was accompanied by VEGF down-regulation at 48h in ES-2 and at 24h in SKOV-3 cells. Interestingly, CM promoted anti-proliferative JunB mRNA accumulation in ES-2 at 24h/48h, while it induced JunB down-regulation in SKOV-3 at both time points.

Conclusions: We demonstrated, for the first time to our knowledge, that healthy PDMSCs-CM treatment exerts its anti-tumoral activity through HIF-1 α pathway down-regulation and anti-proliferative JunB accumulation in malignant ES-2 OC cells. JunB down-regulation in moderately differentiated/less aggressive SKOV-3 PDMSCs-treated cells requires further investigation.

The definitive version is available at:

[http://www.placentajournal.org/article/S0143-4004\(15\)01234-5/fulltext](http://www.placentajournal.org/article/S0143-4004(15)01234-5/fulltext)