



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Phase III study (MONET1) of motesanib plus carboplatin/paclitaxel in patients with advanced nonsquamous nonsmall-cell lung cancer (NSCLC): Asian subgroup analysis

This is the author's manuscript

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/150328 since 2016-06-10T11:10:28Z

Published version:

DOI:10.1093/annonc/mdt552

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

Questa è la versione dell'autore dell'opera:

Ann Oncol. 2014 Feb;25(2):529-36. doi: 10.1093/annonc/mdt552. Epub 2014 Jan 13.

Phase III study (MONET1) of motesanib plus carboplatin/paclitaxel in patients with advanced nonsquamous nonsmall-cell lung cancer (NSCLC): Asian subgroup analysis.

Kubota K, Ichinose Y, Scagliotti G, Spigel D, Kim JH, Shinkai T, Takeda K, Kim SW, Hsia TC, Li RK, Tiangco BJ, Yau S, Lim WT, Yao B, Hei YJ, Park K.

La versione definitiva è disponibile alla URL:

http://annonc.oxfordjournals.org/content/25/2/529.long

Phase III study (MONET1) of motesanib plus carboplatin/paclitaxel in patients with advanced nonsquamous nonsmall-cell lung cancer (NSCLC): Asian subgroup analysis

Kubota K, Ichinose Y, Scagliotti G, Spigel D, Kim JH, Shinkai T, Takeda K, Kim SW, Hsia TC, Li RK, Tiangco BJ, Yau S, Lim WT, Yao B, Hei YJ, Park K.

Abstract

Background This preplanned subset analysis of the phase III MONET1 study aimed to determine whether motesanib combined with carboplatin/paclitaxel (C/P) would result in improved overall survival (OS) versus chemotherapy alone, in a subset of Asian patients with nonsquamous nonsmall-cell lung cancer (NSCLC).

Patients and methods Patients with nonsquamous NSCLC (stage IIIB/IV or recurrent) and no prior systemic therapy for advanced disease were randomized to IV carboplatin (AUC, 6 mg/ml min) and paclitaxel (200 mg/m²) for up to six 3-week cycles, plus either oral motesanib 125 mg q.d. or placebo. Primary end point was OS; secondary end points included progression-free survival (PFS), objective response rate (ORR), and safety.

Results Two hundred twenty-seven Asian patients from MONET1 were included in this descriptive analysis. Median OS was 20.9 months in the motesanib plus C/P arm and 14.5 months in the placebo plus C/P arm (P = 0.0223); median PFS was 7.0 and 5.3 months, respectively, (P = 0.0004); and ORR was 62% and 27%, respectively, (P < 0.0001). Grade ≥ 3 adverse events were more common in the motesanib plus C/P arm versus placebo plus C/P (79% versus 61%).

Conclusion In this preplanned subset analysis of Asian patients with nonsquamous NSCLC, motesanib plus C/P significantly improved OS, PFS, and ORR versus placebo plus C/P.

Clinical trial number NCT00460317.

Key words

Asia, carboplatin, motesanib, NSCLC, paclitaxel

Introduction

The incidence of lung cancer is high (and increasing) in Asia [1], with an estimated agestandardized incidence of 22.3 per 100 000 [2]. Nonsmall-cell lung cancer (NSCLC) accounts for 80%-85% of all cases of lung cancer [3].

Cigarette smoking is estimated to cause 85%–90% of lung cancers in the USA [4]. In large parts of Asia, the associated hazards and peak of smoking-related deaths may not be fully realized for at least two decades [5].

Currently, platinum-based doublet combination chemotherapy is the first-line treatment of choice for NSCLC; however, the prognosis for patients with advanced NSCLC is poor with a median overall survival (OS) of 8–11 months [3]. Bevacizumab, a targeted antiangiogenic agent, in combination with carboplatin/paclitaxel (C/P) significantly improved OS versus chemotherapy alone in patients with recurrent or advanced NSCLC [6]. Multitargeted antiangiogenic agents may provide advantages over agents with single targets [7].

Motesanib is a highly selective, oral small-molecule multikinase inhibitor of vascular endothelial growth factor receptor (VEGFR)-1, -2, -3, platelet-derived growth factor receptor (PDGFR), and Kit [8]. Motesanib has demonstrated antitumor activity as monotherapy in advanced solid tumors [9, 10]. In a phase II study in advanced nonsquamous NSCLC, motesanib plus C/P was estimated to be as efficacious as bevacizumab plus C/P [11].

However, in the phase III MONET1 study, OS was not significantly improved with motesanib plus C/P compared with C/P alone in patients with advanced nonsquamous NSCLC, or in a subset of patients with adenocarcinoma [12]. In a retrospective population-based study of NSCLC in the USA, Asian ethnicity was found to be an independent favorable prognostic factor for OS in NSCLC regardless of smoking status [13].

The objectives of the present study were to assess OS, progression-free survival (PFS), and adverse events (AEs) in an exploratory subgroup analysis of Asian patients with advanced nonsquamous NSCLC who received motesanib plus C/P in the MONET1 study, compared with placebo plus C/P.

materials and methods

study design

MONET1 was an international, double-blind, placebo-controlled, randomized phase III study to evaluate the efficacy and tolerability of motesanib when administered with C/P. Full details of the methodology have been previously reported [12]. In brief, patients received carboplatin (AUC, 6 mg/ml min) and paclitaxel (200 mg/m²) on day 1 of each 3-week cycle up to six cycles and were randomized 1:1 to also receive oral motesanib 125 mg q.d. (arm A) or placebo (arm B). Treatment continued until disease progression, unacceptable toxicity, or withdrawal of consent.

The primary end point was OS; secondary end points included PFS, overall response rate (ORR), and the incidence of AEs.

patients

Patients aged ≥ 18 years with histologically confirmed unresectable stage IIIB NSCLC with pericardial or pleural effusion or stage IV/recurrent nonsquamous NSCLC were eligible for inclusion. Other inclusion criteria included measurable or nonmeasurable disease by modified Response Evaluation Criteria in Solid Tumors (RECIST) [14]; Eastern Cooperative Oncology Group (ECOG) performance status 0 or 1; and life expectancy ≥ 3 months. Key exclusion criteria included central nervous system metastases. Study procedures were approved by an independent ethics committee/institutional review board at each study site, and all patients provided written informed consent.

assessments

Tumor assessments were made via computed tomography (CT) or magnetic resonance imaging scans (assessed per modified RECIST v1.0). AEs were graded by National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE), version 3.0.

statistical analysis

For the present exploratory subanalysis, the Asian subset included patients enrolled in East Asia; Japan, Korea, Philippines, Hong Kong, Taiwan, or Singapore. Non-Asian patients were those enrolled in the European Union, US, Canada, Australia, South America, Turkey, Ukraine, India, Israel, or Russia.

P-values are descriptive only due to the exploratory nature of this subanalysis. The full analysis set (all randomized patients) was used for all efficacy analyses. For OS and PFS, arms A and B were compared using a two-sided stratified log-rank test; stratified Cox proportional hazards model was used to define HR. For ORR, the treatment arms were compared using the stratified Cochran–Mantel–Haenszel test. The safety analysis set (all randomized patients who received at least one dose of motesanib) was used for all analyses of safety.

results

This analysis included 227 Asian patients with nonsquamous NSCLC who were randomized to receive motesanib (n = 110; arm A) or placebo (n = 117; arm B) in addition to C/P (Table <u>1</u>). Approximately 50% of patients (n = 106) were Japanese (arm A: n = 55; arm B: n = 51); and almost 90% (n = 198) had adenocarcinoma. Baseline demographics and disease characteristics of the Asian patients were compared with the total population treated in the MONET1 study (N = 1090). A higher percentage of Asian patients were never smokers compared with the overall population.

Table 1.

Baseline demographics and disease characteristics

Efficacy and safety analyses of the Asian patients were compared with non-Asian patients (n = 863). At the time of analysis, 139 patients had died and 88 continued on study. Asian and non-Asian patients received a median of 164 and 106 days of motesanib, respectively, versus 125 and 126 days of placebo. In Asian patients, median daily dose of motesanib was 121.6 mg compared with 125.0 mg in the non-Asian population. Median follow-up was 63 weeks.

survival outcomes

In Asian patients, treatment with motesanib plus C/P was associated with improved OS (median 20.9 versus 14.5 months [P = 0.0223]; Figure <u>1</u>A) and PFS (7.0 versus 5.3 months [P = 0.0004]; Figure <u>1</u>C) compared with placebo plus C/P. No difference between arms was observed in non-Asian patients for OS (Figure <u>1</u>B) or PFS (Figure <u>1</u>D).



Figure 1.

Overall (A and B) and progression-free (C and D) survival.

response

All 227 Asian patients were assessable for response by modified RECIST (Table <u>2</u>). In Asian patients, the ORR was 62% for motesanib plus C/P and 27% for placebo plus C/P (P < 0.0001). The difference in ORR between treatment arms was more evident in Asian than non-Asian patients.

Median duration of response among responders in the Asian population was 5.8 months for motesanib plus C/P and 4.4 months for placebo plus C/P.

Table 2.

Overall response by RECIST

adverse events

The safety analysis included 222 Asian patients compared with 850 non-Asian patients (Table <u>3</u>). Motesanib/placebo-related AEs were more common in the Asian subgroup, compared with the non-Asian subgroup. Motesanib treatment was discontinued due to AEs in 26% of Asian and 32% of non-Asian patients. The frequency of motesanib/placebo-related serious AEs appeared similar between the Asian and non-Asian patients; however, more treatment-emergent grade 5 AEs were reported in the non-Asian patients compared with the Asian patients.

Table 3.

Safety overview in Asian and non-Asian patients

The most common treatment-emergent AEs in the Asian subgroup are shown in Table <u>4</u>. Neutropenia (44% versus 28%), hypertension (51% versus 9%), thrombocytopenia (21% versus 15%), diarrhea (63% versus 33%), and vomiting (38% versus 30%) occurred more frequently in the motesanib plus C/P group compared with the placebo plus C/P group. In addition, 9% of patients in the motesanib group had gallbladder-related AEs (cholecystitis, cholecystitis acute, gallbladder enlargement, gallbladder edema) compared with 2% in the placebo group. Hypothyroidism was observed in 6% of patients receiving motesanib plus C/P versus <1% in those receiving placebo plus C/P. There was a low incidence of bleeding/thrombosis events, and no difference was observed between the two treatment arms.

Table 4.

Treatment-emergent AEs (any grade) in Asian patients ($\geq 10\%$ in either arm)

discussion

This subset analysis of MONET1 found that in Asian patients with advanced nonsquamous NSCLC, motesanib plus C/P treatment was associated with increased OS (P < 0.05), PFS (P < 0.001), and ORR (P < 0.0001) compared with C/P alone, whereas in the non-Asian subgroup there were no significant differences in OS and PFS, and a substantially smaller increase in ORR (P = 0.0033).

The reason for the differential effect of motesanib in the Asian subpopulation is unclear; however, a number of factors may be involved. There is evidence of epidemiologic, etiologic, and pharmacogenomic differences between Asian and Caucasian lung cancer patients [15], which may affect the impact of motesanib added to C/P. A higher rate of epidermal growth factor receptor (EGFR) mutations was found in Asian compared with Caucasian or non-Asian populations [15], and EGFR mutations have been associated with better outcomes following treatment with EGFR tyrosine kinase inhibitors, such as gefitinib [16]. Additionally, there is known interaction between the EGFR and VEGFR signaling pathways in NSCLC [17] with EGFR regulating VEGF production. Of the VEGF pathway inhibitors, currently bevacizumab plus C/P is the only combination to have demonstrated an OS benefit [6]. A subgroup analysis of bevacizumab plus chemotherapy in Asian patients suggested improved outcomes with addition of bevacizumab [18].

A differential rate of EGFR mutation in Asian versus non-Asian patients may have had an effect on the impact of motesanib in MONET1; however, EGFR mutational status was not determined and, therefore, no analyses were carried out that took this factor into account.

Similar results were observed in a recent phase III study of sunitinib plus erlotinib versus placebo plus erlotinib in patients with refractory NSCLC [19]. In the overall study population, OS and PFS were not improved in patients who received combination therapy compared with those who received erlotinib alone whereas an exploratory subgroup analysis in patients with Asian ethnicity showed significantly improved OS and lower PFS [19]. EGFR mutational status was not established in the majority of patients, and thus analyses were not presented to determine whether this factor was associated with the differential efficacy seen in Asian patients.

Another factor possibly affecting the findings from our subset analysis is the differential effectiveness of C/P seen in patients with advanced NSCLC in Japan and the USA, which may be associated with pharmacogenomic differences between these populations affecting paclitaxel disposition and DNA repair [20]. PFS and OS were longer, but rates of neutropenia were significantly higher with C/P in Japanese versus American patients, possibly associated with specific genetic differences between populations. Notably, the addition of motesanib to C/P has been associated with a moderate increase in paclitaxel exposure [21]. Enhancement of this effect due to specific pharmacogenomic factors in Asian patients may have contributed to differential activity and toxicity related to paclitaxel exposure in the present analysis.

Asian [13] and specifically Japanese [22] and Korean [23] ethnicity are independent favorable prognostic factors for OS compared with white/Caucasian ethnicity in patients with NSCLC. Associated with the pharmacogenetic and pharmacodynamic differences discussed above, there are differences in treatment guidelines for NSCLC between Asian and Western countries [15]. Based on market share, the most commonly used first-line regimen for advanced NSCLC in Japan is C/P, followed by gefitinib [15]; in Korea, the authors' experience suggests that gemcitabine/cisplatin is more widely used than C/P [24, 25]. Thus, the findings of this analysis, showing an association with improved outcomes with the addition of motesanib to C/P, may be of particular importance. The findings from other studies and analyses in Asian patients provide support for the activity and tolerability of angiogenesis inhibitors in combination with chemotherapy for first-line NSCLC [26–28].

Duration of motesanib treatment was longer (and continued for longer than placebo) in Asian than in non-Asian patients in MONET1. Motesanib plus C/P may be better tolerated by Asian versus non-Asian patients, supported by the higher rate of treatment discontinuations due to AEs in the latter population. Alternatively, responding patients may have stayed on therapy for longer as a consequence of the observed responses. The incidence of any-grade and grade \geq 3 motesanib-related AEs appeared higher among Asian patients in the motesanib plus C/P arm, associated with this longer duration of therapy. In contrast, the rate of grade 5 AEs in the motesanib arm appeared higher in non-Asian patients, although this finding was also seen in the placebo arm.

A number of AEs were observed more frequently in the motesanib versus placebo arm, including hypertension, neutropenia, diarrhea, nausea, rash, and gallbladder-related AEs, consistent with previous studies of motesanib, and other angiogenesis inhibitors for NSCLC [11, 21, 27, 28]. There did not appear to be any excess of drug-related mortality in Asian patients versus non-Asian patients. Bleeding complications were low compared with those observed in a previous study of motesanib in NSCLC [11] and were consistent with studies of other angiogenesis inhibitors [26, 27]. In addition, bleeding was not increased in the motesanib plus C/P arm compared with placebo plus C/P.

In summary, in this exploratory subset analysis of Asian patients with nonsquamous NSCLC, treatment with motesanib plus C/P was associated with improved OS, PFS, and ORR compared

with placebo plus C/P. A phase III study in Asian patients is ongoing to confirm these findings (JPRN-JapicCTI-121887).

funding

This study was funded by Amgen, Inc.

disclosure

KP is a consultant/has an advisory role for Amgen, Inc. KK (Taiho, Lilly, AstraZeneca, Daiichi-Sankyo) and GS (Lilly, AstraZeneca, Roche Diagnostics) have received honoraria. YI has received research funding from Amgen, Inc. and Takeda Bio Development Center Ltd. BY and YJH are employees and shareholders of Amgen, Inc. All remaining authors have declared no conflicts of interest.

acknowledgements

The authors thank the patients who participated in this study and their families. They also acknowledge Catherine Crookes of FireKite for writing assistance in the development of this manuscript, which was funded by Millennium: The Takeda Oncology Company.

references

- 1. Wu YL, Park K, Soo RA, et al INSPIRE: a phase III study of the BLP25 liposome vaccine (L-BLP25) in Asian patients with unresectable stage III non-small cell lung cancer. BMC Cancer 2011;11:430.
- 2. GLOBOCAN 2008. 2008. Cancer Incidence, Mortality and Prevalence Worldwide in 2008 <u>http://globocan.iarc.fr/</u> (25 October 2013, date last accessed).
- 3. Dempke WC, Suto T, Reck M Targeted therapies for non-small cell lung cancer. Lung Cancer 2010;67:257-74.
- 4. Freedman ND, Leitzmann MF,Hollenbeck AR, et al *Cigarette smoking and subsequent risk of lung cancer in men and women: analysis of a prospective cohort study. Lancet Oncol 2008;9:649-56.*
- 5. Huxley R, Jamrozik K, Lam TH, et al Impact of smoking and smoking cessation on lung cancer mortality in the Asia-Pacific region. Am J Epidemiol 2007;165:1280-6.
- 6. Sandler A, Gray R, Perry MC, et al Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. N Engl J Med 2006;355:2542-50.
- 7. Scagliotti GV, Govindan R Targeting angiogenesis with multitargeted tyrosine kinase inhibitors in the treatment of non-small cell lung cancer. Oncologist 2010;15:436-46.
- 8. Polverino A, Coxon A, Starnes C, et al

AMG 706, an oral, multikinase inhibitor that selectively targets vascular endothelial growth factor, platelet-derived growth factor, and kit receptors, potently inhibits angiogenesis and induces regression in tumor xenografts. Cancer Res 2006;66:8715-21.

- 9. Rosen LS, Kurzrock R, Mulay M, et al Safety, pharmacokinetics, and efficacy of AMG 706, an oral multikinase inhibitor, in patients with advanced solid tumors. J Clin Oncol 2007;25:2369-76.
- 10. Sherman SI, Wirth LJ, Droz JP, et al *Motesanib diphosphate in progressive differentiated thyroid cancer. N Engl J Med 2008;359:31-42.*
- 11. Blumenschein GR Jr., Kabbinavar F, Menon H, et al A phase II, multicenter, open-label randomized study of motesanib or bevacizumab in combination with paclitaxel and carboplatin for advanced nonsquamous non-small-cell lung cancer. Ann Oncol 2011;22:2057-67.
- 12. Scagliotti GV, Vynnychenko I, Park K, et al International, randomized, placebo-controlled, double-blind phase III study of motesanib plus carboplatin/paclitaxel in patients with advanced nonsquamous non-small-cell lung cancer: MONET1. J Clin Oncol 2012;30:2829-36.
- 13. Ou SH, Ziogas A, Zell JA Asian ethnicity is a favorable prognostic factor for overall survival in non-small cell lung cancer (NSCLC) and is independent of smoking status. J Thorac Oncol 2009;4:1083-93.
 - 14. Therasse P, Arbuck SG, Eisenhauer EA, et al New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. J Natl Cancer Inst 2000;92:205-16.
 - 15. Saijo N, Fukuoka M, Thongprasert S, et al Lung cancer working group report. Jpn J Clin Oncol 2010;40(Suppl 1):i7-12.
 - 16. Mitsudomi T, Kosaka T, Endoh H, et al Mutations of the epidermal growth factor receptor gene predict prolonged survival after gefitinib treatment in patients with non-small-cell lung cancer with postoperative recurrence. J Clin Oncol 2005;23:2513-20.
 - 17. Ciardiello F, Troiani T, Bianco R, et al Interaction between the epidermal growth factor receptor (EGFR) and the vascular endothelial growth factor (VEGF) pathways: a rational approach for multi-target anticancer therapy. Ann Oncol 2006;17(Suppl 7):vii109-14.
 - 18. Mok TS, Hsia TC, Tsai CM, et al *Efficacy of bevacizumab with cisplatin and gemcitabine in Asian patients with advanced or recurrent non-squamous non-small cell lung cancer who have not received prior chemotherapy: a substudy of the Avastin in Lung trial. Asia Pac J Clin Oncol 2011;7(Suppl 2):4-12.*
 - 19. Scagliotti GV, Krzakowski M, Szczesna A, et al Sunitinib plus erlotinib versus placebo plus erlotinib in patients with previously treated advanced non-small-cell lung cancer: a phase III trial. J Clin Oncol 2012;30:2070-8.
 - 20. Gandara DR, Kawaguchi T, Crowley J, et al Japanese-US common-arm analysis of paclitaxel plus carboplatin in advanced non-small-cell lung cancer: a model for assessing population-related pharmacogenomics. J Clin Oncol 2009;27:3540-6.

- 21. Blumenschein GR Jr., Reckamp K, Stephenson GJ, et al Phase 1b study of motesanib, an oral angiogenesis inhibitor, in combination with carboplatin/paclitaxel and/or panitumumab for the treatment of advanced non-small cell lung cancer. Clin Cancer Res 2010;16:279-90.
- 22. Kawaguchi T, Matsumura A, Fukai S, et al

Japanese ethnicity compared with Caucasian ethnicity and never-smoking status are independent favorable prognostic factors for overall survival in non-small cell lung cancer: a collaborative epidemiologic study of the National Hospital Organization Study Group for Lung Cancer (NHSGLC) in Japan and a Southern California Regional Cancer Registry databases. J Thorac Oncol 2010;5:1001-10.

23. Ahn MJ, Lee J, Park YH, et al

Korean ethnicity as compared with white ethnicity is an independent favorable prognostic factor for overall survival in non-small cell lung cancer before and after the oral epidermal growth factor receptor tyrosine kinase inhibitor era. J Thorac Oncol 2010;5:1185-96.

24. Kim JH, Kim SY, Jung KH, et al

Randomized phase II study of gemcitabine plus cisplatin versus etoposide plus cisplatin for the treatment of locally advanced or metastatic non-small cell lung cancer: Korean Cancer Study Group experience. Lung Cancer 2006;52:75-81.

25. Park JO, Kim SW, Ahn JS, et al

Phase III trial of two versus four additional cycles in patients who are nonprogressive after two cycles of platinum-based chemotherapy in non small-cell lung cancer. J Clin Oncol 2007;25:5233-9.

- 26. Okamoto I, Yoshioka H, Takeda K, et al Phase I clinical study of the angiogenesis inhibitor TSU-68 combined with carboplatin and paclitaxel in chemotherapy-naive patients with advanced non-small cell lung cancer. J Thorac Oncol 2012;7:427-33.
- 27. Tsai CM, Au JS, Chang GC, et al Safety and efficacy of first-line bevacizumab with chemotherapy in Asian patients with advanced nonsquamous NSCLC: results from the phase IV MO19390 (SAiL) study. J Thorac Oncol 2011;6:1092-.

28. Zhao X, Mei K, Cai X, et al

A randomized phase II study of recombinant human endostatin plus gemcitabine/cisplatin compared with gemcitabine/cisplatin alone as first-line therapy in advanced non-small-cell lung cancer. Invest New Drugs 2012;30:1144-9.