

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Liquid biopsies to evaluate early therapeutic response in colorectal cancer

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1566487> since 2016-06-14T16:06:20Z

Published version:

DOI:10.1093/annonc/mdv228

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Ann Oncol. 2015 Aug;26(8):1525-7. doi: 10.1093/annonc/mdv228

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://annonc.oxfordjournals.org/content/26/8/1525.long>

Liquid biopsies to evaluate early therapeutic response in colorectal cancer

Clara Montagut^{1,2}, Giulia Siravegna^{3,4,5} and Alberto Bardelli^{3,4}

¹ Medical Oncology Department, Hospital del Mar, Barcelona, Spain ²Cancer Research Program, FIMIM, Hospital del Mar, Barcelona, Spain ³ University of Torino, Department of Oncology, SP 142, KM 3.95, 10060 Candiolo, Torino, Italy; ⁴ Candiolo Cancer Institute – FPO, IRCCS, Candiolo, Torino, Italy; ⁵ FIRC Institute of Molecular Oncology (IFOM), Milano, Italy.

Tumor burden and response to therapy are presently assessed by the Response Evaluation Criteria In Solid Tumors (RECIST), a standardized method for determining therapeutic response to anticancer therapy using imaging-driven measurements of lesion size. This is often accompanied by blood-based test of protein biomarkers, such as CEA in colorectal cancer and CA125 in ovarian cancers [1, 2].

The introduction of RECIST in clinical trials and daily clinical practice represented an historical step forward when it was implemented almost two decades ago [3]. More recently, advances in anti-cancer therapies and biological understanding of cancer have highlighted limitations of RECIST to assess drug response and sparked the pursuit of new methodologies to measure tumor burden. New standards are already being used in certain settings such as PET-TC to assess response in lymphoma, modified RECIST criteria in gastrointestinal stromal tumors, immunological criteria for immunotherapy or morphological response for antiangiogenic drugs [4-6]. In essence, what clinicians and patients seek is a method to assess whether drug administration is effective on tumor cells, how long it takes to do so and when it stops working. In addition, it would be preferable to continuously monitor therapeutic response, rather than wait for imaging assessments that are performed several weeks (sometimes months) after initiation of therapy.

In the current issue, Tie et al explore the potential of circulating tumor DNA in evaluating tumour burden and monitoring/predicting response to chemotherapy at an early stage of treatment [7]. Their study is based on previous evidences that tumor-derived DNA is shed into the bloodstream by cancer cell [8]. Although the presence of circulating tumor DNA in the blood of patients has been known for decades, only recent advances in genomics allowed the detection and quantification of cancer-related molecular alterations with high specificity and sensitivity [9, 10].

Using colorectal cancer as a model system, the study started by identifying tumor-specific somatic variants (mutations) in tissue specimens obtained at diagnosis, and exploiting these to monitor tumor burden non-invasively in the blood. The authors sequenced a panel of 15 genes frequently mutated in mCRC, and at least one mutation was identified in the tumor tissue of 98% of the cases. The study shows a sensitivity of basal ctDNA detection as high as 92%, and a close correlation between ctDNA quantification and initial tumor burden assessed by CT-scan. Next, the authors analysed outlier patients in which ctDNA did not correlate with tumor burden as assessed by RECIST criteria. They found a false positive and a false negative case, in which CT-scan failed to truly express the real load of the tumor. This evidence, albeit limited to a few cases, suggests that ctDNA may prove superior to RECIST criteria when it comes to assess tumor burden. The study also confirms earlier observations that ctDNA is superior to CEA as a blood marker of tumor load. This is likely due to fact that somatic genomic variants in oncogenes and tumor suppressor genes are tumor-specific and to the fact that ctDNA has a shorter half-life than CEA.

Up until this point the findings are not novel, as previous studies have highlighted that somatic mutations can be used to monitor tumour burden in colorectal and other cancer types [8, 11]. Rapid increases in ctDNA levels are also known to correlate with disease progression and declines in ctDNA levels with successful pharmacological or surgical treatment [11-14].

The innovative section and the central finding of the current report is the evidence that the modulation of cancer mutations levels in circulating free DNA appears to anticipate response to therapy assessed by the classical RECIST criteria (Figure 1). Tie et al report a correlation between changes in cfDNA mutation levels after one cycle of chemotherapy and response by RECIST criteria 8-12 weeks after the onset of therapy, and more importantly a trend toward a correlation between changes in ctDNA and progression free survival (PFS). Interestingly, although very early changes (d+3) in ctDNA did not correlate with response, a spike in ctDNA was observed in 3 patients with a striking response, suggesting that early changes in mutational loads in plasma may reflect DNA release into the bloodstream and may be a used to identify the best responders.

A test that anticipates response to a given drug early during the course of treatment without having to wait for image changes in the CT-scan weeks after the onset of therapy, potentially allows for an early switch in treatment, avoiding unnecessary side effects, enhancing efficacy and minimizing costs. Moreover, recent evidence shows

that early tumor shrinkage (ETS) in the treatment of metastatic colorectal cancer correlates with longer survival.

What are the implications of this work? The encouraging results in this paper support the use of ctDNA as an early indicator of tumor response in metastatic colorectal cancer patients. However, given the small size of the cohort, these data should be viewed as hypothesis-generating and must be replicated in the context of large clinical trials to show a clinically and statistically significant impact on survival of patients. From a methodological standpoint, the approach is relatively easy to translate in daily clinical practice. The approach requires availability of tumor-tissue (surgery or biopsy) to identify patient specific mutations. While this is rarely a limitation for colorectal cancer patients, in other tumor types such as lung tumors access to lesions is more difficult and may sometimes be a limitation.

The idea of assessing response to treatment in cancer patients using mutation levels is reminiscent of the use of viral load to monitor therapy response in HIV patients. Will ctDNA substitute RECIST in a near future? A number of reasons suggest that imaging will always be necessary to evaluate the site of metastatic disease, the resectability of tumoral lesions or the clinical compromise of adjacent anatomic structures. Therefore ctDNA and RECIST assessment will likely complement each other, and together with other methods (i.e., functional imaging) will define new integrated standards for evaluation of therapy in oncology. As highlighted by this study, the opportunities offered by circulating free tumour DNA are only starting to be explored but is already clear that liquid biopsies are here to stay and will become a central asset in the management of cancer patients.

Funding: Associazione Italiana per la Ricerca sul Cancro (AIRC) IG grant no. 12812 (A.B.); AIRC 2010 Special Program Molecular Clinical Oncology 5 per mille, project no. 9970 (A.B.); FPRC 5 per mille 2010 and 2011 Ministero della Salute (A.B.); Ministero dell'Istruzione, dell'Università e della Ricerca, progetto PRIN (A.B). RD12/0036/0051, PI12/00989, PI12/00680, PT13/0010/0005 and 2014 SGR 740 grants and by the Xarxa de Banc de Tumors de Catalunya (C.M.)

Disclosure: The authors have declared no conflicts of interest.'

REFERENCES

1. Engelen MJ, de Bruijn HW, Hollema H et al. Serum CA 125, carcinoembryonic antigen, and CA 19-9 as tumor markers in borderline ovarian tumors. *Gynecol Oncol* 2000; 78: 16-20.
2. McKeown E, Nelson DW, Johnson EK et al. Current approaches and challenges for monitoring treatment response in colon and rectal cancer. *J Cancer* 2014; 5: 31-43.
3. Eisenhauer EA, Therasse P, Bogaerts J et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009; 45: 228-247.
4. Cheson BD, Fisher RI, Barrington SF et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol* 2014; 32: 3059-3068.
5. Wolchok JD, Hoos A, O'Day S et al. Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. *Clin Cancer Res* 2009; 15: 7412-7420.
6. Chun YS, Vauthey JN, Boonsirikamchai P et al. Association of computed tomography morphologic criteria with pathologic response and survival in patients treated with bevacizumab for colorectal liver metastases. *JAMA* 2009; 302: 2338-2344.
7. Tie J, Kinde I, Wang Y et al. Circulating Tumor DNA as an Early Marker of Therapeutic Response in Patients with Metastatic Colorectal Cancer. *Ann Oncol* 2015.
8. Bettgowda C, Sausen M, Leary RJ et al. Detection of circulating tumor DNA in early- and late-stage human malignancies. *Sci Transl Med* 2014; 6: 224ra224.
9. Diaz LA, Bardelli A. Liquid biopsies: genotyping circulating tumor DNA. *J Clin Oncol* 2014; 32: 579-586.
10. Siravegna G, Bardelli A. Genotyping cell-free tumor DNA in the blood to detect residual disease and drug resistance. *Genome Biol* 2014; 15: 449.
11. Diehl F, Schmidt K, Choti MA et al. Circulating mutant DNA to assess tumor dynamics. *Nat Med* 2008; 14: 985-990.
12. Diaz LA, Williams RT, Wu J et al. The molecular evolution of acquired resistance to targeted EGFR blockade in colorectal cancers. *Nature* 2012; 486: 537-540.
13. Misale S, Yaeger R, Hobor S et al. Emergence of KRAS mutations and acquired resistance to anti-EGFR therapy in colorectal cancer. *Nature* 2012; 486: 532-536.
14. Beaver JA, Jelovac D, Balukrishna S et al. Detection of Cancer DNA in Plasma of Early Stage Breast Cancer Patients. *Clin Cancer Res* 2014.

